



Constructing exact sequences on non-conforming discrete spaces

Martin Campos Pinto

► To cite this version:

Martin Campos Pinto. Constructing exact sequences on non-conforming discrete spaces. Comptes rendus de l'Académie des sciences. Série I, Mathématique, 2016, 354. hal-01167425

HAL Id: hal-01167425

<https://hal.science/hal-01167425>

Submitted on 24 Jun 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Constructing exact sequences on non-conforming discrete spaces

Martin Campos Pinto ^{a,b},

^a*CNRS, UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France*

^b*Sorbonne Universités, UPMC Univ Paris 06, UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France*

Received 17 March 2015

Presented by Olivier Pironneau

Abstract

In this note we propose a general procedure to construct exact sequences on non-conforming function spaces and we show how this construction can be used to derive a proper discrete Gauss law for structure-preserving discontinuous Galerkin (DG) approximations to the time-dependent 2d Maxwell equations. *To cite this article: M. Campos Pinto, C. R. Acad. Sci. Paris, Ser. I *** (2015).*

Résumé

Construction de suites exactes sur des espaces discrets non-conformes. Dans cette note nous proposons un procédé général pour construire des suites exactes sur des espaces non-conformes, et nous montrons comment ce procédé peut servir à écrire une loi de Gauss discrète convenable dans le cadre d'approximations Galerkin discontinues (DG) des équations de Maxwell temporelles en 2d. *Pour citer cet article : M. Campos Pinto, C. R. Acad. Sci. Paris, Ser. I *** (2015).*

Version française abrégée

Etant donné une suite exacte discrète $V_h^0 \xrightarrow{d_h^0} V_h^1 \xrightarrow{d_h^1} V_h^2$ qui sera composée en pratique d'espaces conformes, il est possible de construire une nouvelle suite exacte sur un espace arbitraire $\tilde{V}_h^1$ par la procédure suivante.

- En s'appuyant sur un projecteur $\mathcal{P}_h^1$ vers l'espace conforme V_h^1 , on pose $\tilde{d}_h^1 := d_h^1 \mathcal{P}_h^1$. Le noyau de ce nouvel opérateur est essentiellement composé de l'image de d_h^0 et du noyau de $\mathcal{P}_h^1$.
- Pour définir un bon opérateur $\tilde{d}_h^0$ on étend l'opérateur d_h^0 à un espace produit du type $V_h^0 \times \tilde{V}_h^1$ de façon à atteindre tous les éléments de ce nouveau noyau.

Email address: campos@ann.jussieu.fr (Martin Campos Pinto).

L'application au cadre des schémas DG pour les équations de Maxwell 2d se fait alors au moyen de l'observation suivante : lorsque l'opérateur $\mathcal{P}_h^1$ est obtenu par moyennisation locale des degrés de liberté d'arêtes dans une extension aux fonctions régulières par morceaux de l'interpolant canonique sur les éléments finis de Nédélec en 2d, la construction esquissée plus haut permet de retrouver l'opérateur rotationnel discret standard du schéma DG à flux centré.

La suite exacte ainsi construite fait alors intervenir un nouvel opérateur discret de type gradient dans un cadre Galerkin discontinu, qui par dualité définit un nouvel opérateur de divergence discrète. Conformément au programme de travail décrit dans [7] en vue d'obtenir des schémas généraux préservant la structure pour les équations de Maxwell temporelles, cet opérateur divergence a les propriétés requises pour définir une loi de Gauss discrète devant être vérifiée par un schéma DG préservant la charge.

1. Introduction

Exact sequences of function spaces with differential operators such as gradient, curl or divergence play a major role in the construction and analysis of mixed finite element approximations of physical problems. This is especially true in computational electromagnetics since the works of Bossavit [5,6] who has pointed out the importance of preserving at the discrete level the natural structure of Maxwell's equations in terms of de Rham diagrams of differential forms [14], and in the recent decades the study of structure-preserving schemes has become a field of its own with numerous developments, see e.g. [2,13,10,3,16,11,1,4,9].

In the context of Hilbert spaces, de Rham diagrams take the form of sequences like

$$H^1(\Omega) \xrightarrow{\text{grad}} H(\text{curl}; \Omega) \xrightarrow{\text{curl}} H(\text{div}; \Omega) \xrightarrow{\text{div}} L^2(\Omega) \quad (1)$$

which, under some smoothness and topological assumptions on the domain Ω , are *exact* in the sense that the range of each operator coincides with the kernel of the next one in the sequence. Many fundamental properties of finite-element discretizations, such as the spectral correctness of Maxwell's eigenvalue problem are then related to the question whether the underlying function spaces preserve this property.

To our knowledge, most of the available work in the field is concerned with *conforming* methods where the finite-dimensional function spaces are embedded in the above sequence (1). In this note we propose a general procedure to construct exact sequences on non-conforming function spaces, and we show how this construction can be used to derive a proper discrete Gauss law in the context of structure-preserving discontinuous Galerkin (DG) approximations of the time-dependent Maxwell equations in 2d, following the approach outlined in [7].

2. Abstract construction of exact sequences on arbitrary spaces

In this section we provide a general procedure to build an exact sequence on an arbitrary discrete space, starting from a known exact sequence. To this end we consider an abstract sequence of the form

$$V_h^0 \xrightarrow{d_h^0} V_h^1 \xrightarrow{d_h^1} V_h^2 \quad (2)$$

that is assumed exact in the sense that

$$\ker d_h^1 = d_h^0 V_h^0. \quad (3)$$

The typical situation that we have in mind is that of conforming spaces V_h^k (i.e., embedded in a sequence of Hilbert spaces such as (1)) and operators d_h^k defined as the restriction of the corresponding differential

operators. Now, if $\tilde{V}_h^1$ is a finite-dimensional space that is a subset of neither V_h^1 nor the Hilbert space associated to the differential operator d_h^1 in the continuous sequence (1), we can build a new operator $\tilde{d}_h^1$ approximating d_h^1 on $\tilde{V}_h^1$ by considering a projection on V_h^1 ,

$$\mathcal{P}_h^1 : \hat{V}_h^1 \rightarrow V_h^1 \quad \text{where} \quad \hat{V}_h^1 := V_h^1 + \tilde{V}_h^1 \quad (4)$$

(here the sum is necessary since $\mathcal{P}_h^1$ must be defined on V_h^1), and by setting

$$\tilde{d}_h^1 := d_h^1 \mathcal{P}_h^1|_{\tilde{V}_h^1} : \tilde{V}_h^1 \rightarrow V_h^2. \quad (5)$$

In order to build an exact sequence involving $\tilde{V}_h^1$ we need to characterize the kernel of the latter operator.

Lemma 2.1 *It holds*

$$\ker \tilde{d}_h^1 = \left(d_h^0 V_h^0 \oplus (I - \mathcal{P}_h^1) \tilde{V}_h^1 \right) \cap \tilde{V}_h^1.$$

Proof: The inclusion $\supset$ is easily verified by applying $\tilde{d}_h^1$, indeed this operator coincides with d_h^1 on $d_h^0 V_h^0$ since the latter is in V_h^1 . To verify next the inclusion $\subset$ we take $u \in \ker \tilde{d}_h^1$. Then $\mathcal{P}_h^1 u$ is in V_h^1 and also in $\ker d_h^1$, hence in $d_h^0 V_h^0$ according to the exact sequence property (3). In particular, we have

$$u = \mathcal{P}_h^1 u + (I - \mathcal{P}_h^1)u \in d_h^0 V_h^0 \oplus (I - \mathcal{P}_h^1) \tilde{V}_h^1$$

where we have used that $u \in \tilde{V}_h^1$, and the inclusion follows. Here the direct sum is verified by applying $\mathcal{P}_h^1$ to some $u \in d_h^0 V_h^0 \cap (I - \mathcal{P}_h^1) \tilde{V}_h^1$: clearly $\mathcal{P}_h^1 u = 0$, and since $d_h^0 V_h^0 \subset V_h^1$ we also have $u = \mathcal{P}_h^1 u$. $\square$

Now that the kernel of $\tilde{d}_h^1$ is characterized, we may define an approximation of d_h^0 on the product space $V_h^0 \times \tilde{V}_h^1$. Specifically, we set

$$\tilde{d}_h^0 : \hat{V}_h^0 \rightarrow \hat{V}_h^1, \quad (\varphi, u) \mapsto d_h^0 \varphi + (I - \mathcal{P}_h^1)u, \quad \text{where} \quad \hat{V}_h^0 := V_h^0 \times \tilde{V}_h^1. \quad (6)$$

The proper operator on $\tilde{V}_h^1$ is then obtained by a simple restriction

$$\tilde{d}_h^0 := \tilde{d}_h^0|_{\tilde{V}_h^0} : \tilde{V}_h^0 \rightarrow \tilde{V}_h^1 \quad \text{where} \quad \tilde{V}_h^0 := (\tilde{d}_h^0)^{-1}(\tilde{V}_h^1) = \{(\varphi, u) \in \hat{V}_h^0 : \tilde{d}_h^0(\varphi, u) \in \tilde{V}_h^1\}, \quad (7)$$

indeed this yields the following result.

Theorem 2.2 *The sequence*

$$\tilde{V}_h^0 \xrightarrow{\tilde{d}_h^0} \tilde{V}_h^1 \xrightarrow{\tilde{d}_h^1} V_h^2 \quad (8)$$

is exact, in the sense that $\tilde{d}_h^0 \tilde{V}_h^0 = \ker \tilde{d}_h^1$.

Proof: The claimed relation is straightforward to verify, using the definition of $\tilde{d}_h^0$ and Lemma 2.1. $\square$

Remark 1 *If we let $\hat{d}_h^1 := d_h^1 \mathcal{P}_h^1$ then the sequence*

$$\hat{V}_h^0 \xrightarrow{\hat{d}_h^0} \hat{V}_h^1 \xrightarrow{\hat{d}_h^1} V_h^2 \quad (9)$$

is also exact, and it remains so if we set $\hat{V}_h^0 := V_h^0 \times \hat{V}_h^1$ in (6), since $(I - \mathcal{P}_h^1) \hat{V}_h^1 = (I - \mathcal{P}_h^1) \tilde{V}_h^1$.

Remark 2 *If $V_h^1 \subset \tilde{V}_h^1$, the above construction simplifies, as the sequences (8) and (9) coincide.*

3. Application to the centered DG discretization of the 2d Maxwell system

The abstract construction described above can be applied to the design of structure-preserving Maxwell schemes based on fully discontinuous finite-element spaces, following the program outlined in [7, Sec. 4]. In

this short note we illustrate this construction in the case of a discontinuous Galerkin (DG) discretization of the normalized 2d Maxwell system

$$\begin{cases} \partial_t B + \operatorname{curl} \mathbf{E} = 0 \\ \partial_t \mathbf{E} - \operatorname{curl} B = -\mathbf{J} \end{cases} \quad (10)$$

on some bounded domain Ω of $\mathbb{R}^2$. Here we assume that Ω is simply-connected, Lipschitz and partitioned by a geometrically conforming triangulation $\mathcal{T}_h$ with set of edges denoted by $\mathcal{E}_h$.

3.1. A reference exact sequence with conforming spaces

To begin with we consider the conforming sequence

$$V_h^0 := \mathcal{L}_p(\Omega, \mathcal{T}_h) \xrightarrow{\mathbf{grad}} V_h^1 := \mathcal{N}_{p-1}(\Omega, \mathcal{T}_h) \xrightarrow{\operatorname{curl}} V_h^2 := \mathbb{P}_{p-1}(\mathcal{T}_h) \quad (11)$$

where

$$\mathcal{L}_p(\Omega, \mathcal{T}_h) := \mathbb{P}_p(\mathcal{T}_h) \cap H^1(\Omega) \quad (12)$$

denotes the continuous ‘‘Lagrange’’ elements of total degree $\leq p$,

$$\mathcal{N}_{p-1}(\Omega, \mathcal{T}_h) := \mathcal{N}_{p-1}(\mathcal{T}_h) \cap H(\operatorname{curl}; \Omega) \quad \text{with} \quad \mathcal{N}_{p-1}(T) := \mathbb{P}_{p-1}(T)^2 \oplus \begin{pmatrix} -y \\ x \end{pmatrix} \mathbb{P}_{p-1}^{\operatorname{hom}}(T) \quad (13)$$

is the first-kind Nédélec space of order p (introduced in [15] for the 3d setting, see also [4, Sec. 2.3]), and

$$\mathbb{P}_{p-1}(\mathcal{T}_h) := \{v \in L^2(\Omega) : v|_T \in \mathbb{P}_{p-1}(T), T \in \mathcal{T}_h\} \quad (14)$$

are the discontinuous elements of degree $\leq p-1$. As is well known, (11) is indeed exact in the sense that

$$\operatorname{Im} \mathbf{grad}_h = \ker \operatorname{curl}_h \quad (15)$$

holds with $\mathbf{grad}_h := \mathbf{grad}|_{V_h^0}$ and $\operatorname{curl}_h := \operatorname{curl}|_{V_h^1}$, which correspond to d_h^0 and d_h^1 in our abstract construction. Relation (15) is usually derived from the continuous exact sequence

$$H^1(\Omega) \xrightarrow{\mathbf{grad}} H(\operatorname{curl}; \Omega) \xrightarrow{\operatorname{curl}} L^2(\Omega) \quad (16)$$

by using commuting diagrams of projection operators (see e.g. [4]) but a direct argument can be given here: If $\mathbf{u} = \mathbf{v} + \begin{pmatrix} -y \\ x \end{pmatrix} v \in \mathcal{N}_{p-1}$ with $\mathbf{v} \in \mathbb{P}_{p-1}^2$ and $v \in \mathbb{P}_{p-1}^{\operatorname{hom}}$, then $\operatorname{curl} \mathbf{u} = \operatorname{curl} \mathbf{v} + (p+1)v$. Now assume $\operatorname{curl} \mathbf{u} = 0$. Since $\operatorname{curl} \mathbf{v}$ clearly belongs to $\mathbb{P}_{p-2}$, this yields $v = 0$ and hence $\mathbf{u} \in \mathbb{P}_{p-1}^2$. Using this and the exact sequence (16) we find $\mathbf{u} = \mathbf{grad} \varphi$ for some φ which must belong to $\mathbb{P}_p$, hence (15).

3.2. A new exact sequence with discontinuous Galerkin spaces

We next consider replacing the Nédélec space V_h^1 by a fully discontinuous piecewise polynomial space

$$\tilde{V}_h^1 := \mathbb{P}_{p-1}(\mathcal{T}_h)^2 \not\subset H(\operatorname{curl}; \Omega).$$

To construct an exact sequence based on this discontinuous space, following Section 2 we need a projection operator $\mathcal{P}_h^1$ on V_h^1 . Here a natural choice is to extend the standard finite element interpolation based on the 2d Nédélec degrees of freedom which correspond to (see, e.g., [4, Sec. 2.3])

$$\begin{cases} \mathcal{M}_T^1 = \{\mathbf{u} \mapsto \int_T \mathbf{u} \cdot \mathbf{v} : \mathbf{v} \in \mathcal{P}_{p-2}(T)^2\}, & T \in \mathcal{T}_h \\ \mathcal{M}_e^1 = \{\mathbf{u} \mapsto \int_e (\mathbf{n}_e \times \mathbf{u})v : v \in \mathcal{P}_{p-1}(e)\}, & e \in \mathcal{E}_h \end{cases} \quad (17)$$

by using local averages on the edges. Specifically, we define

$$\mathcal{P}_h^1 : \hat{V}_h^1 \rightarrow V_h^1 = \mathcal{N}_{p-1}(\Omega, \mathcal{T}_h) \quad \text{where} \quad \hat{V}_h^1 := \tilde{V}_h^1 + V_h^1 \quad (18)$$

by the relations

$$\begin{cases} \mathcal{M}_T^1(\mathcal{P}_h^1 \mathbf{u}) = \mathcal{M}_T^1(\mathbf{u}), & T \in \mathcal{T}_h \\ \mathcal{M}_e^1(\mathcal{P}_h^1 \mathbf{u}) = \mathcal{M}_e^1(\{\mathbf{u}\}_e), & e \in \mathcal{E}_h. \end{cases} \quad (19)$$

Here we have used the standard notation for the averages on interior and boundary edges, namely

$$\{\mathbf{u}\}_e := \frac{1}{2}(\mathbf{u}|_{T_e^-} + \mathbf{u}|_{T_e^+})|_e \quad \text{and} \quad \{\mathbf{u}\}_e := (\mathbf{u}|_{T_e})|_e \quad \text{respectively.} \quad (20)$$

This projection operator has several interesting properties. First it is local, in the sense that the degrees of freedom of $\mathcal{P}_h^1 \mathbf{u}$ associated to some mesh element (i.e., edge or triangle) only depend on the values of $\mathbf{u}$ on the neighboring mesh elements. In particular, $\mathcal{P}_h^1$ can be implemented with a sparse matrix. Second, the non-conforming curl operator $\tilde{d}_h^1$ corresponding to (5), that we shall now denote

$$\text{curl}_h^{\text{dg}} := \text{curl} \mathcal{P}_h^1|_{\tilde{V}_h^1} \quad (21)$$

coincides with the standard DG curl with centered fluxes [12]. Indeed a straightforward computation using Green formulas and the form of the degrees of freedom (17) yields the following result.

Lemma 3.1 *The non-conforming curl operator defined on $\tilde{V}_h^1 = \mathbb{P}_{p-1}(\mathcal{T}_h)^2$ by (18)-(21) satisfies*

$$\langle \text{curl} \mathcal{P}_h^1 \mathbf{u}, v \rangle = \sum_{T \in \mathcal{T}_h} \langle \mathbf{u}, \text{curl} v \rangle_T - \sum_{e \in \mathcal{E}_h} \langle \{\mathbf{u}\}_e, [v] \rangle_e \quad \text{for all} \quad v \in \mathbb{P}_{p-1}(\mathcal{T}_h) \quad (22)$$

with a standard notation for the averages (20) and tangential jumps on interior and boundary edges, i.e.

$$[v]_e := (\mathbf{n}_e^- \times v|_{T_e^-} + \mathbf{n}_e^+ \times v|_{T_e^+})|_e \quad \text{and} \quad [v]_e := (\mathbf{n}_e \times v|_{T_e})|_e \quad \text{respectively.} \quad (23)$$

An exact sequence is then provided by the abstract construction from Section 2. Specifically, setting

$$\mathbf{grad}_h^{\text{dg}} : \tilde{V}_h^0 \rightarrow \tilde{V}_h^1, \quad (\varphi, \mathbf{u}) \mapsto \mathbf{grad} \varphi + (I - \mathcal{P}_h^1) \mathbf{u}, \quad (24)$$

with

$$\tilde{V}_h^0 := \{(\varphi, \mathbf{u}) \in V_h^0 \times \tilde{V}_h^1 : \mathbf{grad} \varphi + (I - \mathcal{P}_h^1) \mathbf{u} \in \tilde{V}_h^1\} \quad (25)$$

we find that the sequence

$$\tilde{V}_h^0 \xrightarrow{\mathbf{grad}_h^{\text{dg}}} \tilde{V}_h^1 \xrightarrow{\text{curl}_h^{\text{dg}}} V_h^2 \quad (26)$$

is exact in the sense that

$$\text{Im} \mathbf{grad}_h^{\text{dg}} = \ker \text{curl}_h^{\text{dg}}. \quad (27)$$

3.3. A proper discrete Gauss law for the DG discretization of the 2d Maxwell system

Following [7, Sec. 4] the above construction can be used to derive a proper discrete Gauss law of the form

$$\text{div}_h^{\text{dg}} \tilde{\mathbf{E}}_h = \tilde{\rho}_h \quad (28)$$

(involving some approximation $\tilde{\rho}_h$ to the exact charge density ρ) that should be preserved by a structure-preserving DG scheme for the time-dependent problem (10). To do so we define as a discrete divergence operator the adjoint of the above gradient operator (24) with a minus sign, namely

$$\text{div}_h^{\text{dg}} := -(\mathbf{grad}_h^{\text{dg}})^* : \tilde{V}_h^1 \rightarrow \tilde{V}_h^0. \quad (29)$$

In particular, if ρ_h is an approximation to ρ in the continuous finite-element space $V_h^0 = \mathcal{L}_p(\Omega, \mathcal{T}_h)$, then setting $\tilde{\rho}_h := (\rho_h, 0) \in \tilde{V}_h^0$ allows to rewrite the discrete Gauss law (28) with test functions in $\tilde{V}_h^0$, as

$$\langle \tilde{\mathbf{E}}_h, -\mathbf{grad} \varphi \rangle + \langle \tilde{\mathbf{E}}_h, (\mathcal{P}_h^1 - I)\mathbf{u} \rangle = \langle \rho_h, \varphi \rangle \quad \text{for all } (\varphi, \mathbf{u}) \in \tilde{V}_h^0. \quad (30)$$

Finally we point out that such a discrete Gauss law is well suited to the design of charge-conserving DG-particle schemes, and we refer to the forthcoming article [8] for more details.

Acknowledgements

The author acknowledges stimulating discussions with Eric Sonnendrücker and Mathieu Lutz.

References

- [1] D.N. Arnold, R.S. Falk, and R. Winther. Finite element exterior calculus: from Hodge theory to numerical stability. *Bull. Amer. Math. Soc.(NS)*, 47(2):281–354, 2010.
- [2] D. Boffi. A note on the deRham complex and a discrete compactness property. *Applied Mathematics Letters*, 14(1):33–38, January 2001.
- [3] D. Boffi. Compatible Discretizations for Eigenvalue Problems. In *Compatible Spatial Discretizations*, pages 121–142. Springer New York, New York, NY, 2006.
- [4] D. Boffi, F. Brezzi, and M. Fortin. *Mixed finite element methods and applications*, volume 44 of *Springer Series in Computational Mathematics*. Springer, 2013.
- [5] A. Bossavit. Whitney forms: a class of finite elements for three-dimensional computations in electromagnetism. In *Physical Science, Measurement and Instrumentation, Management and Education - Reviews, IEE Proceedings A*, pages 493–500, 1988.
- [6] A. Bossavit. Un nouveau point de vue sur les éléments mixtes. *Matapli (Bull. Soc. Math. Appl. Industr.)*, 20:23–35, 1989. (French).
- [7] M. Campos Pinto, M. Mounier, and E. Sonnendrücker. Handling the divergence constraints in Maxwell and Vlasov-Maxwell simulations. submitted, 2015.
- [8] M. Campos Pinto and E. Sonnendrücker. Compatible Maxwell solvers with particles in 2d. I: conforming and non-conforming schemes with a strong Faraday law. In preparation.
- [9] M. Campos Pinto and E. Sonnendrücker. Gauss-compatible Galerkin schemes for time-dependent Maxwell equations. <http://hal.upmc.fr/hal-00969326>, 2014.
- [10] M. Costabel and M. Dauge. Computation of resonance frequencies for Maxwell equations in non-smooth domains. In *Topics in computational wave propagation*, pages 125–161. Springer, Berlin, 2003.
- [11] L. Demkowicz. Polynomial Exact Sequences and Projection-Based Interpolation with Application to Maxwell Equations. In *Mixed finite elements, compatibility conditions, and applications*. Springer-Verlag, Berlin; Fondazione C.I.M.E., Florence, 2008.
- [12] L. Fezoui, S. Lanteri, S. Lohrengel, and S. Piperno. Convergence and stability of a discontinuous Galerkin time-domain method for the 3D heterogeneous Maxwell equations on unstructured meshes. *ESAIM: Mathematical Modelling and Numerical Analysis*, 39(6):1149–1176, November 2005.
- [13] R. Hiptmair. Finite elements in computational electromagnetism. *Acta Numerica*, 11:237–339, 2002.
- [14] P.R. Kotiuga. *Hodge Decompositions and Computational Electromagnetics*. PhD thesis, Mc Gill University Montréal, Canada, October 1984.
- [15] J.-C. Nédélec. Mixed finite elements in $\mathbf{R}^3$. *Numerische Mathematik*, 35(3):315–341, 1980.
- [16] D.A. White, J.M. Koning, and R.N. Rieben. Development and application of compatible discretizations of Maxwell’s equations. In *Compatible Spatial Discretizations*, pages 209–234. Springer, New York, 2006.