

Supplementary Material for “Adaptive kernel estimation of the baseline function in the Cox model, with high-dimensional covariates”

Agathe Guilloux

Laboratoire de Statistique Théorique et Appliquée,
Université Pierre et Marie Curie - Paris 6
e-mail : `agathe.guilloux@upmc.fr`

Sarah Lemler

Laboratoire de Mathématiques et Modélisation d'Évry UMR CNRS 8071- USC INRA,
Université d'Évry Val d'Essonne, France
e-mail : `sarah.lemler@genopole.cnrs.fr`

Marie-Luce Taupin

Laboratoire de Mathématiques et Modélisation d'Évry UMR CNRS 8071- USC INRA,
Université d'Évry Val d'Essonne, France
e-mail : `marie-luce.taupin@genopole.cnrs.fr`
Unité MaIAGE, INRA Jouy-En-Josas, France

1 Technical lemmas

1.1 Proof of Lemma 6.2

From Definition (20), we have $\alpha_h(t) = K_h * \alpha_0(t)$ for all $t \in [0, \tau]$ so that

$$\mathbb{E}[\|\bar{\alpha}_h - \alpha_0\|_2^2] = \mathbb{E}[\|\bar{\alpha}_h - \alpha_h\|_2^2] + \|\alpha_h - \alpha_0\|_2^2.$$

The first term of the right part of this equality can be rewritten as

$$\mathbb{E}[\|\bar{\alpha}_h - \alpha_h\|_2^2] = \int_0^\tau \text{Var}[\bar{\alpha}_h(t)] dt.$$

It remains to bound $\text{Var}[\bar{\alpha}_h(t)]$:

$$\begin{aligned}\text{Var}[\bar{\alpha}_h(t)] &= \frac{1}{n} \text{Var} \left[\int_0^\tau K_h(t-u) \frac{1}{S(u, \boldsymbol{\beta}_0)} dN_1(u) \right] \\ &\leq \frac{1}{n} \mathbb{E} \left[\left(\int_0^\tau K_h(t-u) \frac{1}{S(u, \boldsymbol{\beta}_0)} dN_1(u) \right)^2 \right].\end{aligned}$$

We apply the Doob-Meyer decomposition $N_1 = M_1 + \Lambda_1$:

$$\begin{aligned}\text{Var}[\bar{\alpha}_h(t)] &\leq \frac{2}{n} \mathbb{E} \left[\left(\int_0^\tau \frac{K_h(t-u)}{S(u, \boldsymbol{\beta}_0)} dM_1(u) \right)^2 \right] \\ &\quad + \frac{2}{n} \mathbb{E} \left[\left(\int_0^\tau \frac{K_h(t-u)}{S(u, \boldsymbol{\beta}_0)} \alpha_0(u) e^{\boldsymbol{\beta}_0^T \mathbf{Z}_1} Y_1(u) du \right)^2 \right]\end{aligned}$$

Then, we have

$$\mathbb{E} \left[\left(\int_0^\tau \frac{K_h(t-u)}{S(u, \boldsymbol{\beta}_0)} dM_1(u) \right)^2 \right] \leq \mathbb{E} \left[\int_0^\tau \frac{K_h^2(t-u)}{S^2(u, \boldsymbol{\beta}_0)} \alpha_0(u) e^{\boldsymbol{\beta}_0^T \mathbf{Z}_1} Y_1(u) du \right],$$

and we finally get that

$$\mathbb{E} \left[\left(\int_0^\tau \frac{K_h(t-u)}{S(u, \boldsymbol{\beta}_0)} dM_1(u) \right)^2 \right] \leq \frac{\mathbb{E}[e^{\boldsymbol{\beta}_0^T \mathbf{Z}_1}] \|\alpha_0\|_{\infty, \tau} \tau \|K\|_{\mathbb{L}^2(\mathbb{R})}^2}{c_S^2 h}.$$

Thus, the integrated variance term of the pseudo-estimator is bounded by

$$\int_0^\tau \text{Var}[\bar{\alpha}_h(t)] dt \leq \frac{2 \|\alpha_0\|_{\infty, \tau} \tau}{c_S^2} (\mathbb{E}[e^{\boldsymbol{\beta}_0^T \mathbf{Z}_1}] + \|\alpha_0\|_{\infty, \tau} \mathbb{E}[e^{2\boldsymbol{\beta}_0^T \mathbf{Z}_1}]_\tau) \frac{\|K\|_{\mathbb{L}^2(\mathbb{R})}^2}{nh},$$

which gives a bound of order $1/nh$.

Gathering the bias term $\|\alpha_h - \alpha_0\|_2^2$ and the bound on variance term gives Inequality (22) in Lemma 6.2. □

1.2 Proof of Lemma 6.3

The proof of Lemma 6.3 relies on an additional lemmas. First, write

$$\hat{\alpha}_h^{\hat{\boldsymbol{\beta}}}(t) - \bar{\alpha}_h(t) = \frac{1}{nh} \sum_{i=1}^n \int_0^\tau \frac{S(u, \boldsymbol{\beta}_0) - S_n(u, \hat{\boldsymbol{\beta}})}{S(u, \boldsymbol{\beta}_0) S_n(u, \hat{\boldsymbol{\beta}})} K\left(\frac{t-u}{h}\right) \mathbf{1}_{\{\bar{Y}(u) > 0\}} dN_i(u).$$

We study the difference process $\hat{\alpha}_h^{\hat{\boldsymbol{\beta}}} - \bar{\alpha}_h$ on Ω_k , defined by (32) and on its complement. From Lemma 6.6, the process $\hat{\alpha}_h^{\hat{\boldsymbol{\beta}}} - \bar{\alpha}_h$ is controled on Ω_k^c . The following lemma allows to bound the difference process on Ω_k .

Lemma 1.1. *Under Assumptions 3.4.(ii)-(iii) , 3.5.(i)-(iii), 3.1 and 3.2, for any $k \in \mathbb{N}$, we have*

$$\mathbb{E}[|\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h|_2^2 \mathbf{1}(\Omega_k)] \leq c_4(s) \frac{\log(n^k p)}{n},$$

where c_4 is a constant depending on $B, |\beta_0|_1, R, \|\alpha_0\|_{\infty, \tau}, c_S, \|K\|_{\mathbb{L}^2(\mathbb{R})}, \tau$ and s the sparsity index of β_0 .

Gathering Lemmas 6.6 and 1.1, we finally get that, for a fixed k

$$\mathbb{E}[|\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h|_2^2] \leq c(s) \frac{\log(n^k p)}{n},$$

with $c(s)$ a constant depending on $B, |\beta_0|_1, R, \|\alpha_0\|_{\infty, \tau}, c_S, \|K\|_{\mathbb{L}^2(\mathbb{R})}, \tau$ and s the sparsity index of β_0 . and Lemma 6.3 is then proved. Let now prove the Lemmas 6.6 and 1.1.

1.3 Proof of Lemma 6.6 :

We have to bound $\mathbb{E}[|\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h|_2^2 \mathbf{1}(\Omega_k^c)]$, which is equal to

$$\mathbb{E}[|\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h|_2^2 \mathbf{1}(\Omega_k^c)] = \int_0^\tau \mathbb{E}[(\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h)^2(t) \mathbf{1}(\Omega_k^c)] dt.$$

First, let us focus on $\mathbb{E}[(\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h)^2(t) \mathbf{1}(\Omega_k^c)]$ defined by

$$\mathbb{E} \left[\left(\frac{1}{n} \sum_{i=1}^n \int_0^\tau K_h(t-u) \mathbf{1}_{\{\bar{Y}(u) > 0\}} \frac{S(u, \beta_0) - S_n(u, \hat{\beta})}{S(u, \beta_0) S_n(u, \hat{\beta})} dN_i(t) \right)^2 \mathbf{1}(\Omega_k^c) \right].$$

From Assumptions 3.1, $\hat{\beta}$ belongs to a ball $\mathcal{B}(0, R)$ and $|\beta_0|_1 < \infty$, so we have the following bound

$$S_n(u, \hat{\beta}) - S(u, \beta_0) \leq \frac{1}{n} \sum_{i=1}^n e^{\beta_0^T Z_i} e^{(\hat{\beta} - \beta_0)^T Z_i} \leq e^{2B|\beta_0|_1} e^{BR}. \quad (1)$$

For sake of simplicity, let us denote $C(B, R, |\beta_0|_1)$ the bound in (1). From $\mathbf{1}_{\{\bar{Y}(u) > 0\}}$ in the definition of $\hat{\alpha}_h^{\hat{\beta}}$, there exists $i_0 \in \{1, \dots, n\}$ such that $Y_{i_0} \neq 0$, so that from Assumption 3.4.(i)

$$S_n(u, \hat{\beta}) \geq \frac{1}{n} e^{-B|\beta_0|_1} e^{-B|\hat{\beta} - \beta_0|_1} \geq \frac{1}{n} e^{-2B|\beta_0|_1} e^{-BR}. \quad (2)$$

Combining (1) and (2), for $\tilde{C}(B, R, |\beta_0|_1) = e^{8B|\beta_0|_1} e^{4BR}$, we obtain the following bound

$$\mathbb{E}[(\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h)^2(t) \mathbf{1}(\Omega_k^c)] \leq \tilde{C}(B, R, |\beta_0|_1) \frac{n^2}{c_S^2} \mathbb{E} \left[\left(\int_0^\tau K_h(t-u) dN_1(u) \right)^2 \mathbf{1}(\Omega_k^c) \right]. \quad (3)$$

From Assumption 3.5.(i) and the Cauchy-Schwarz Inequality, we get

$$\begin{aligned}\mathbb{E}\left[\left(\int_0^\tau K_h(t-u)dN_1(u)\right)^2 \mathbf{1}(\Omega_k^c)\right] &\leq \frac{\|K\|_\infty^2}{h^2} \mathbb{E}[N_1(\tau)^2 \mathbf{1}(\Omega_k^c)] \\ &\leq \frac{\|K\|_\infty^2}{h^2} \mathbb{E}[N_1(\tau)^4]^{1/2} \mathbb{P}(\Omega_k^c)^{1/2}.\end{aligned}\quad (4)$$

From the Doob-Meyer decomposition and the B urkholder Inequality 2.4, we deduce that $\mathbb{E}[N_1(\tau)] < \infty$.

Now we focus on $\int_0^\tau \mathbb{E}[(\hat{\alpha}_h^\beta - \bar{\alpha}_h)^2(t) \mathbf{1}(\Omega_k^c)] dt$. From the two bounds (3) and (4) obtained above, we have

$$\int_0^\tau \mathbb{E}[(\hat{\alpha}_h^\beta - \bar{\alpha}_h)^2(t) \mathbf{1}(\Omega_k^c)] dt \leq \tilde{C}(B, R, |\beta_0|_1) \frac{n^2}{c_S^2} \left(\frac{\|K\|_\infty^2}{h^2} \mathbb{E}^{1/2}[N_1(\tau)^4] \sqrt{\mathbb{P}(\Omega_k^c)} \right).$$

Let introduce the following lemma that gives a bound for $\mathbb{P}(\Omega_k^c)$.

Lemma 1.2. *Under Assumptions 3.1 and 3.2, for all $k \in \mathbb{N}$, there exists $n_0 \in \mathbb{N}$, such that for $n > n_0$ we have*

$$\mathbb{P}[\Omega_k^c] \leq c_2 n^{-k}, \quad (5)$$

where c_2 is a constant depending on B , $|\beta_0|_1$ and s .

From Lemma 1.2 and the fact that $h^{-1} \leq n$ from Assumption 3.5.(ii), we get that

$$\int_0^\tau \mathbb{E}[(\hat{\alpha}_h^\beta - \bar{\alpha}_h)^2(t) \mathbf{1}(\Omega_k^c)] dt \leq C(B, |\beta_0|_1, R, \|\alpha_0\|_{\infty, \tau}, c_S, \tau, \|K\|_\infty) n^{4-k/2},$$

where $C(B, |\beta_0|_1, R, \|\alpha_0\|_{\infty, \tau}, c_S, \tau, \|K\|_\infty)$ is a constant depending on elements in brackets. Finally, we obtain

$$\mathbb{E}[\|\hat{\alpha}_h^\beta - \bar{\alpha}_h\|_2^2 \mathbf{1}(\Omega_k^c)] \leq C(B, |\beta_0|_1, R, \|\alpha_0\|_{\infty, \tau}, c_S, \tau, \|K\|_\infty) n^{4-k/2},$$

which ends the proof of Lemma 6.6. \square

1.4 Proof of Lemma 1.1 :

On Ω_k , we have

$$\mathbb{E}[(\hat{\alpha}_h^\beta - \bar{\alpha}_h)^2(t) \mathbf{1}(\Omega_k)] \leq \frac{16B^2 e^{4B|\beta_0|_1} e^{2BR} C^2(s) \log(pn^k)}{c_S^2 n} \mathbb{E}\left[\left(\frac{1}{n} \sum_{i=1}^n \int_0^\tau \frac{|K_h(t-u)|}{S(u, \beta_0)} dN_i(u)\right)^2\right].$$

Then, from the Doob-Meyer decomposition, we deduce that

$$\begin{aligned}\mathbb{E}\left[\left(\frac{1}{n} \sum_{i=1}^n \int_0^\tau \frac{|K_h(t-u)|}{S(u, \beta_0)} dN_i(u)\right)^2\right] &\leq \frac{2}{n} \mathbb{E}\left[\left(\int_0^\tau \frac{K_h(t-u)}{S(u, \beta_0)} dM_1(u)\right)^2\right] \\ &\quad + \frac{2}{n} \mathbb{E}\left[\left(\int_0^\tau \frac{K_h(t-u)}{S(u, \beta_0)} \alpha_0(u) e^{\beta_0^T Z_1} Y_1(u) du\right)^2\right] \\ &\leq \frac{2\|\alpha_0\|_{\infty, \tau}}{c_S^2} (\mathbb{E}[e^{\beta_0^T Z_1}] + \|\alpha_0\|_{\infty, \tau} \mathbb{E}[e^{2\beta_0^T Z_1}]) \frac{\|K\|_{L^2(\mathbb{R})}^2}{c_S^2}.\end{aligned}$$

We finally obtain

$$\mathbb{E}[\|\hat{\alpha}_h^{\hat{\beta}} - \bar{\alpha}_h\|_{2,h}^2 \mathbf{1}(\Omega_k)] \leq C(c_S, \|\alpha_0\|_{\infty, \tau}, |\beta_0|_1, B, R, \|K\|_{\mathbb{L}^2(\mathbb{R})}, s) \frac{\log(pn^k)}{n}. \quad (6)$$

□

1.5 Proof of Lemma 1.2

In order to calculate $\mathbb{P}(\Omega_k^c)$, let us begin by study the set $\Omega_{H,k}$ defined by (30). Let us introduce the two following sets

$$\begin{aligned} \Omega_1 &:= \left\{ \omega : \forall u \in [0, \tau], |S_n(u, \hat{\beta}) - S_n(u, \beta_0)| \leq B e^{BR} e^{2B|\beta_0|_1} C(s) \sqrt{\frac{\log(pn^k)}{n}} \right\}, \\ \Omega_2 &:= \left\{ \omega : \forall u \in [0, \tau], |S_n(u, \beta_0) - S(u, \beta_0)| \leq B e^{BR} e^{2B|\beta_0|_1} C(s) \sqrt{\frac{\log(pn^k)}{n}} \right\}. \end{aligned}$$

We have $\Omega_{H,k} \supset \Omega_1 \cap \Omega_2$. We begin to calculate $\mathbb{P}(\Omega_1^c)$. By definition, we have

$$\begin{aligned} |S_n(u, \hat{\beta}) - S_n(u, \beta_0)| &= \left| \frac{1}{n} \sum_{i=1}^n (e^{\hat{\beta}^T \mathbf{Z}_i} - e^{\beta_0^T \mathbf{Z}_i}) Y_i(u) \right| \\ &\leq e^{B|\beta_0|_1} |e^{B|\hat{\beta} - \beta_0|_1} - 1| \end{aligned}$$

Under Assumptions 3.1 and 3.2, from Proposition 3.3, there exists a constant $c > 0$ such that, with probability larger than $1 - cn^{-k}$,

$$|\hat{\beta} - \beta_0|_1 \leq C(s) \sqrt{\frac{\log(pn^k)}{n}}.$$

So, with probability larger than $1 - cn^{-k}$, using that $|e^x - e^y| \leq |x - y|e^{x \vee y}$ for all x, y , we have

$$\begin{aligned} |S_n(u, \hat{\beta}) - S_n(u, \beta_0)| &\leq e^{B|\beta_0|_1} B |\hat{\beta} - \beta_0|_1 e^{B|\hat{\beta} - \beta_0|_1} \\ &\leq B e^{BR} e^{2B|\beta_0|_1} C(s) \sqrt{\frac{\log(pn^k)}{n}}. \end{aligned}$$

We deduce that

$$\mathbb{P}(\Omega_1^c) \leq cn^{-k}. \quad (7)$$

To calculate $\mathbb{P}(\Omega_2^c)$, we remark that

$$n(S_n(u, \beta_0) - S(u, \beta_0)) = \sum_{i=1}^n \left(e^{\beta_0^T \mathbf{Z}_i} Y_i(u) - \mathbb{E}[e^{\beta_0^T \mathbf{Z}_i} Y_i(u)] \right).$$

As $0 \leq e^{\beta_0^T Z_1} Y_1(u) \leq e^{B|\beta_0|_1}$, we apply a Hoeffding inequality:

$$\mathbb{P}\left(|S_n(u, \beta_0) - S(u, \beta_0)| \geq \frac{y}{n}\right) \leq 2 \exp\left(-\frac{2y^2}{ne^{2B|\beta_0|_1}}\right),$$

and with $y = Be^{BR}e^{2B|\beta_0|_1}C(s)\sqrt{n \log(pn^k)}/2$, we finally get

$$\begin{aligned} \mathbb{P}\left(|S_n(u, \beta_0) - S(u, \beta_0)| \geq C(s)Be^{BR}e^{2B|\beta_0|_1}\sqrt{\frac{\log(pn^k)}{n}}\right) \\ \leq 2 \exp\left(-\frac{2B^2e^{2BR}e^{4B|\beta_0|_1}C^2(s)\log(pn^k)}{e^{2B|\beta_0|_1}}\right) \\ \leq \frac{2}{pn^k}. \end{aligned}$$

We conclude that there exists a constant $c_7 > 0$ such that

$$\mathbb{P}(\Omega_2^c) \leq c_7 n^{-k}. \quad (8)$$

Gathering (7) and (8), we obtain

$$\mathbb{P}(\Omega_{H,k}^c) \leq \mathbb{P}(\Omega_1^c) + \mathbb{P}(\Omega_2^c) \leq \tilde{c} n^{-k}, \quad (9)$$

where $\tilde{c} > 0$ is a constant. It remains to calculate $\mathbb{P}(\Omega_{S_n}^c)$, with $\Omega_{S_n}^c$ defined by (31), to obtain $\mathbb{P}(\Omega_k^c)$. We decompose

$$S_n(u, \hat{\beta}) - S(u, \beta_0) = S_n(u, \hat{\beta}) - S_n(u, \beta_0) + S_n(u, \beta_0) - S(u, \beta_0).$$

On $\Omega_1 \cap \Omega_2$,

$$S_n(u, \hat{\beta}) - S_n(u, \beta_0) \geq -2Be^{BR}e^{2B|\beta_0|_1}C(s)\sqrt{\frac{\log(pn^k)}{n}} \in (-\infty, 0)$$

So for n large enough, we have that $S_n(u, \hat{\beta}) - S_n(u, \beta_0) \geq -c_S/2$. For n large enough, $\Omega_1 \cap \Omega_2 \subset \Omega_{S_n}$, and

$$\mathbb{P}(\Omega_{S_n}^c) \leq \mathbb{P}(\Omega_1^c) + \mathbb{P}(\Omega_2^c) \leq \tilde{c} n^{-k}. \quad (10)$$

Gathering (9) and (10), we finally obtain for n large enough that $\mathbb{P}(\Omega_k^c) \leq c_2 n^{-k}$, where c_2 is a constant depending on B , $|\beta_0|_1$ and s . \square

2 Classical results

In this appendix, some classical technical lemmas and a theorem needed for the proofs of the two main theorems of the chapter, are listed. We do not give the proofs of these well-known results but we give the references where to find their proofs.

2.1 A Cauchy-Schwarz Inequality

The following lemma gives a useful inequality concerning integrals with respect to the counting process N .

Lemma 2.1 (Cauchy-Schwarz). *For all function g bounded on $[0, \tau]$,*

$$N(\tau) \int_{\tau_1}^{\tau_2} g^2(s) dN(s) \geq \left(\int_{\tau_1}^{\tau_2} g(s) dN(s) \right)^2, \quad 0 \leq \tau_1 \leq \tau_2 \leq \tau$$

We refer to [Bouaziz et al. \(2013\)](#) for the proof of this lemma.

2.2 Young Inequality

The following lemma provides an inequality that bounds a norm of the convolution product of two functions by a product of norms of each function.

Lemma 2.2 (Young Inequality). *Let $p, q \in [1, +\infty)$ such that $1/p + 1/q \geq 1$. If $s \in \mathbb{L}^p(\mathbb{R})$ and $t \in \mathbb{L}^q(\mathbb{R})$, then s and t are convolvable. Moreover, if $1/r = 1/p + 1/q - 1$, then $f * g \in \mathbb{L}^r(\mathbb{R})$ and*

$$\|s * t\|_r \leq \|s\|_p \|t\|_q$$

This convolution inequality is proved in [Hirsch and Lacombe \(1999\)](#) (Theorem 3.4 p.149).

2.3 Talagrand Inequality

The following Talagrand Inequality is a concentration inequality that allows to control the supremum of an empirical process.

Theorem 2.3 (Talagrand Inequality). *Let ξ_1, \dots, ξ_n be independent random values, and let*

$$\nu_{n,\xi}(f) = \frac{1}{n} \sum_{i=1}^n \{f(\xi_i) - \mathbb{E}[f(\xi_i)]\}.$$

Then, for a countable class of functions \mathcal{F} uniformly bounded and $\varepsilon > 0$, we have

$$\mathbb{E} \left[\left\{ \sup_{f \in \mathcal{F}} \nu_{n,\xi}^2(f) - 2(1 + 2\varepsilon^2)H^2 \right\}_+ \right] \leq \frac{4}{d} \left(\frac{W}{n} e^{-d\varepsilon^2 \frac{nH^2}{W}} + \frac{98M^2}{dn^2\varphi^2(\varepsilon)} e^{-\frac{2d\varphi(\varepsilon)\varepsilon}{\tau\sqrt{2}} \frac{nH}{M}} \right),$$

with $\varphi(\varepsilon) = \sqrt{1 + \varepsilon^2} - 1$, $d = 1/6$ and

$$\sup_{f \in \mathcal{F}} \|f\|_\infty \leq M, \quad \mathbb{E} \left[\sup_{f \in \mathcal{F}} |\nu_{n,\xi}(f)| \right] \leq H, \quad \sup_{f \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^n \text{Var}[f(\xi)] \leq W.$$

This theorem is a useful corollary from the classical Talagrand established by [Talagrand \(1996\)](#). The proof of Theorem 2.3 can be found in [Comte et al. \(2008\)](#) (Lemma 6.1). The proof of the theorem follows from a concentration Inequality in [Klein and Rio \(2005\)](#) and arguments that can be found in [Birgé and Massart \(1998\)](#).

2.4 A classical inequality: the B urkholder Inequality

The last technical result is a B urkholder Inequality that gives a norm relation between a martingale and its optional process. We refer to [Liptser and Shirayev \(1989\)](#) p.75, for the proof of this result.

Theorem 2.4 (B urkholder Inequality). *If $M = (M_t, \mathcal{F}_t)_{t \geq 0}$ is a martingale, then there are universal constants γ_b and κ_b (independent of M) such that for every $t \geq 0$*

$$\gamma_b \|\sqrt{[M]_t}\|_2 \leq \|M_t\|_2 \leq \kappa_b \|\sqrt{[M]_t}\|_2,$$

where $[M]_t$ is the quadratic variation of M_t .

This theorem is used to prove Lemma 6.5 and in the oracle inequalities of Theorem 4.1, the constants depend on κ_b .

References

- L. Birg e and P. Massart. Minimum contrast estimators on sieves: exponential bounds and rates of convergence. *Bernoulli*, 4(3):329–375, 1998.
- O. Bouaziz, F. Comte, and A. Guillaou. Nonparametric estimation of the intensity function of a recurrent event process. *Statistica Sinica*, 23(2):635–665, 2013.
- F. Comte, J. Dedecker, and M.L. Taupin. Adaptive density deconvolution with dependent inputs. *Mathematical Methods of Statistics*, 17(2):87–112, 2008.
- F. Hirsch and G. Lacombe. *Elements of functional analysis*, volume 192. Springer, 1999.
- T. Klein and E. Rio. Concentration around the mean for maxima of empirical processes. *The Annals of Probability*, 33(3):1060–1077, 2005.
- R. Sh. Liptser and A. N. Shirayev. *Theory of martingales*, volume 49 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1989. ISBN 0-7923-0395-4. URL <http://dx.doi.org/10.1007/978-94-009-2438-3>. Translated from the Russian by K. Dzhaparidze [Kacha Dzhaparidze].
- M. Talagrand. New concentration inequalities in product spaces. *Inventiones mathematicae*, 126(3):505–563, 1996.