



HAL
open science

Exponential functionals of spectrally one-sided lévy processes conditioned to stay positive

Grégoire Véchambre

► **To cite this version:**

Grégoire Véchambre. Exponential functionals of spectrally one-sided lévy processes conditioned to stay positive. 2017. hal-01175612v4

HAL Id: hal-01175612

<https://hal.science/hal-01175612v4>

Preprint submitted on 7 Feb 2017 (v4), last revised 12 Feb 2018 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXPONENTIAL FUNCTIONALS OF SPECTRALLY ONE-SIDED LÉVY PROCESSES CONDITIONED TO STAY POSITIVE

GRÉGOIRE VÉCHAMBRE

ABSTRACT. We study the properties of the exponential functional $\int_0^{+\infty} e^{-X^\uparrow(t)} dt$ where X^\uparrow is a spectrally one-sided Lévy process conditioned to stay positive. In particular, we study finiteness, self-decomposability, existence of finite exponential moments, asymptotic tail at 0 and smoothness of the density.

1. INTRODUCTION

We consider a spectrally negative Lévy process V that is not the opposite of a subordinator. We denote its Laplace exponent by Ψ_V :

$$\forall t, \lambda \geq 0, \mathbb{E} \left[e^{\lambda V(t)} \right] = e^{t\Psi_V(\lambda)}.$$

In the case where V drifts to $-\infty$, it is well known that its Laplace exponent admits a non trivial zero that we denote here by κ , $\kappa := \inf\{\lambda > 0, \Psi_V(\lambda) = 0\}$. If V does not drift to $-\infty$, then 0 is the only zero of Ψ_V on $[0, +\infty[$ so we put $\kappa := 0$ in this case. We denote by (Q, γ, ν) the generating triplet of V . According to the Lévy-Kintchine formula, Ψ_V can be expressed as

$$\Psi_V(\lambda) = \frac{Q}{2}\lambda^2 - \gamma\lambda + \int_{-\infty}^0 (e^{\lambda x} - 1 - \lambda x \mathbf{1}_{|x|<1}) \nu(dx). \quad (1.1)$$

We also consider Z , a spectrally positive Lévy process drifting to $+\infty$ with unbounded variation.

We are interested in the exponential functionals of V and Z conditioned to stay positive,

$$I(V^\uparrow) := \int_0^{+\infty} e^{-V^\uparrow(t)} dt \quad \text{and} \quad I(Z^\uparrow) := \int_0^{+\infty} e^{-Z^\uparrow(t)} dt.$$

For both we study finiteness, exponential moments, and the asymptotic tail at 0. For $I(V^\uparrow)$, we use its property of self-decomposability to get precise estimates on the asymptotic tail at 0 and properties of the density. The exponential functional $I(V^\uparrow)$ has already appeared in the study of self-similar Markov processes with no positive jumps by Bertoin, Caballero [5] and Pardo [13]. In both those articles is proved the self-decomposability of the functional $I(V^\uparrow)$. Moreover, in [5], the authors also compute the moments of small positive order of this functional. In [13], the asymptotic almost sure behavior at 0 and $+\infty$ of a positive self-similar Markov process is linked to the left tail of $I(V^\uparrow)$. As a consequence, it seems that the study of the left tail of $I(V^\uparrow)$ in the present paper can be put in relation with the results of [13] to understand the almost sure behavior of positive self-similar Markov processes with no positive jumps.

Date: 07/02/2017 à 12:28:14.

2010 Mathematics Subject Classification. 60G51.

Key words and phrases. Lévy processes conditioned to stay positive, exponential functionals, self-decomposable distributions.

Our first motivation here is to extend to spectrally one-sided Lévy processes conditioned to stay positive the general study of exponential functionals of Lévy processes. Those functionals have been widely studied because of their importance in probability theory. For example they are fundamental in the study of diffusions in random environments and appear in many applications such as the study of self-similar Markov processes and mathematical finance, see Bertoin, Yor [4] and Pardo, Rivero [16] for surveys on those functionals and their applications. For a general Lévy process, equivalent conditions for the finiteness of the exponential functional are given in [4], the asymptotic tail at $+\infty$ of the functional is studied in [23], [11], [24], [25] (see also [4], [16]), the absolute continuity is proved in [9], [17], and properties of the density (such as regularity) are studied in [12], [17], [21]. Recently, factorization identities for exponential functionals of Lévy processes have been proved in [14], [20], [21] (see also [16]). They are helpful for the study of some properties of exponential functionals such as unimodality, smoothness of the density, complete monotonicity of the density, and they allow to derive explicite representations of the density in some special cases. Moreover, these factorization identities give a central place to the study of exponential functionals of subordinators and of spectrally negative Lévy processes, since the latter two are the two typical factors in which a general exponential functional is decomposed. A fruitful approach in the study of exponential functionals of Lévy processes is to consider their Mellin transform that satisfies a functional equation, see [12], [11], [10]. The solution of this equation in term of a generalized Weierstrass product is given in [21]. The equation on the Mellin transform allows to compute the entire moments of exponential functionals in some cases, see [4]. The relation between self-similar Markov processes and exponential functionals of Lévy processes can also be used to study the latter, see [18], [19]. In this paper we highlight and exploit some links between exponential functionals of Lévy processes and exponential functionals of Lévy processes conditioned to stay positive. Also, as a by-product of our approach, we retrieve some known results about the exponential functionals of spectrally negative Lévy processes.

Our second motivation is the possibility to apply our results to the study of diffusions in spectrally negative Lévy environments. Such processes, introduced by Brox [7] when the environment is given by a brownian motion, have been specifically studied for the spectrally negative Lévy case by Singh [28]. In [2], Andreatti, Devulder and the author prove that the supremum of the local time, \mathcal{L}_X^* , of a diffusion in a drifted brownian environment converges in distribution. They express the limit law in term of a bidimensional stable subordinator that depends on an exponential functional of the environment conditioned to stay positive. In order to generalize their results to a diffusion in a spectrally negative Lévy environment, knowledge on the exponential functionals involved is needed. These are precisely exponential functionals of the environment (which is spectrally negative) and its dual (which is spectrally positive) conditioned to stay positive.

Finally, we have hints that the almost sure asymptotic behavior of \mathcal{L}_X^* , for a diffusion in the spectrally negative Lévy environment V , is crucially linked to the right and left tails of the distribution of $I(V^\uparrow)$. This is why we study these tails here and give for the left tail a precise asymptotic estimate when it is possible, in particular when Ψ_V is regularly varying. For the right tail, we are mainly interested in the existence of some finite exponential moments. The application of the present work to diffusions in random environments is a work in preparation by the author [30], [29].

For a process A and a borelian set S we denote

$$\tau(A, S) := \inf \{t \geq 0, A(t) \in S\}, \quad \mathcal{R}(A, S) := \sup \{t \geq 0, A(t) \in S\},$$

where by convention $\inf \emptyset = \infty$. We shall only write $\tau(A, x)$ (respectively $\mathcal{R}(A, x)$) instead of $\tau(A, \{x\})$ (respectively $\mathcal{R}(A, \{x\})$) and $\tau(A, x+)$ instead of $\tau(A, [x, +\infty[)$. For example, since V^\uparrow has no positive jumps, we see that it reaches each positive level continuously : $\forall x >$

0, $\tau(V^\uparrow, x+) = \tau(V^\uparrow, x)$ and since moreover V^\uparrow converges to $+\infty$ we have $\forall x > 0$, $\mathcal{R}(V^\uparrow, [0, x]) = \mathcal{R}(V^\uparrow, x)$.

If A is Markovian and $x \in \mathbb{R}$ we denote A_x for the process A starting from x . For A_0 we shall only write A . For any (possibly random) time $T > 0$, we write A^T for the process A shifted and centered at time T : $\forall s \geq 0$, $A^T(s) := A(T + s) - A(T)$.

Let W be the scale function of V , defined as in Section VII.2 of Bertoin [3]. It satisfies

$$\forall 0 < x < y, \mathbb{P}(\tau(V_x, y) < \tau(V_x,] - \infty, 0]) = W(x)/W(x + y).$$

According to Theorem VII.8 in [3], this function is continuous, increasing, and for any $\lambda > \kappa$,

$$\int_0^{+\infty} e^{-\lambda x} W(x) dx = \frac{1}{\Psi_V(\lambda)} < +\infty.$$

We now explain how the scale function W allows to define V^\uparrow , that is, V conditioned to stay positive (see [3], Section VII.3 for proofs and more details). The relation

$$\forall t \geq 0, x, y > 0, p_t^\uparrow(x, dy) := W(y) \mathbb{P}\left(V_x(t) \in dy, \inf_{[0, t]} V_x > 0\right) / W(x),$$

defines a Markovian semigroup. For any $x > 0$, we denote by V_x^\uparrow the Markovian process starting from x , taking values in $]0, +\infty[$ and whose transition semigroup is p_t^\uparrow . V_x^\uparrow is commonly called *V conditioned to stay positive starting from x*. This is justified by the fact that for any $y > x > 0$ the process $(V_x^\uparrow(t), 0 \leq t \leq \tau(V_x^\uparrow, y))$ is equal in distribution to the process $(V_x(t), 0 \leq t \leq \tau(V_x, y))$ conditionally on $\{\tau(V_x, y) < \tau(V_x,] - \infty, 0])\}$. In the case where V drifts to $+\infty$ this relation remains true after replacing the hitting time of y by $+\infty$. In the cases where V oscillates or drifts to $-\infty$, it is not possible to condition in the usual sens V_x to remain positive for ever, because this occurs with a probability equal to 0. However, even in these cases, V_x^\uparrow has infinite life-time. It is moreover possible to show that in all cases V_x^\uparrow converges almost surely to $+\infty$.

It is possible to show that there exists a process V_0^\uparrow , that we denote by V^\uparrow , starting from 0 and which is the limit in distribution of the processes V_x^\uparrow as x goes to 0. V^\uparrow is called *V conditioned to stay positive*, it is a Feller process whose restriction to $]0, +\infty[$ of the transition laws is given by p_t^\uparrow . Moreover, note that for any positive x , we have from the Markov property and the absence of positive jumps that the process V^\uparrow , shifted at $\tau(V^\uparrow, x)$, is equal in law to V_x^\uparrow .

In the case where V drifts to $-\infty$, we recall the definition of V conditioned to drift to $+\infty$ (see [3], Section VII.1 for proofs and more details). Note that, since κ is a zero of Ψ_V , the process $e^{\kappa V(t)}$ is a martingale with expectation 1. This allows to define the convolution semigroup of probability measures $(p_t^\sharp)_{t \geq 0}$ via the relation

$$p_t^\sharp(dy) := e^{\kappa y} \mathbb{P}(V(t) \in dy).$$

The convolution semigroup $(p_t^\sharp)_{t \geq 0}$ is associated to a spectrally negative Lévy process that we denote by V^\sharp and which is commonly called *V conditioned to drift to $+\infty$* . This is justified by the fact that

- V^\sharp drifts to $+\infty$,
- For any $y > 0$ the process $(V^\sharp(t), 0 \leq t \leq \tau(V^\sharp, y))$ is equal in distribution to the process $(V(t), 0 \leq t \leq \tau(V, y))$ conditionally on $\{\tau(V, y) < +\infty\}$.

It is not difficult to see that the Laplace exponent Ψ_{V^\sharp} of V^\sharp is obtained by translation of Ψ_V : $\Psi_{V^\sharp} = \Psi_V(\kappa + \cdot)$. As a consequence $\Psi'_{V^\sharp}(0) > 0$ which, according to Corollary VII.2 in [3],

proves that V^\sharp drifts to $+\infty$ as stated above. It is also proven that $V^\uparrow = (V^\sharp)^\uparrow$ and this is why, for convenience, we often work with V^\sharp instead of V .

In order to do our proofs in a systematic way, we define V^\sharp to be V conditioned to drift to $+\infty$ in the case where V drifts to $-\infty$ and *only* V in the other cases (when V oscillates or drifts to $+\infty$). As a consequence, V^\sharp always denotes a spectrally negative Lévy process that does not drifts to $-\infty$ (it oscillates if V does and it drifts to $+\infty$ if V drifts to $+\infty$ or $-\infty$) and that satisfies $\Psi_{V^\sharp} = \Psi_V(\kappa + \cdot)$. In any case we have that for all $0 < x < y$, $(V_x^\uparrow(t), 0 \leq t \leq \tau(V_x^\uparrow, y))$ is equal in law to $(V_x^\sharp(t), 0 \leq t \leq \tau(V_x^\sharp, y))$ conditionally on $\{\tau(V_x^\sharp, y) < \tau(V_x^\sharp,]-\infty, 0])\}$. Note that the same identity is true with V instead of V^\sharp , but the advantage of dealing with V^\sharp is that $\tau(V_x^\sharp, y)$ is always finite (while $\tau(V_x, y)$ can possibly be infinite) which simplifies the argumentation.

Let us now recall some facts about Z^\uparrow , that is, Z conditioned to stay positive. By assuming unbounded variation for Z we excluded the case where Z is a subordinator. In this case, Z would stay positive and $I(Z^\uparrow)$ would be only $I(Z)$ which has already been widely studied. It is already known to be finite and have some finite exponential moments (see for example Theorem 2 in [4]), so Theorem 1.13 below is already known in the subordinator case. Since, in our case, $-Z$ is spectrally negative and not the opposite of a subordinator (we denote by κ_Z the non-trivial zero of Ψ_{-Z}), it is regular for $]0, +\infty[$ according to Theorem VII.1 in [3], so Z is regular for $] -\infty, 0[$. Moreover, Z drifts to $+\infty$. We can thus define the Markov family $(Z_x^\uparrow, x > 0)$ as in Doney [8], Chapter 8. It can be seen from there that for any $x > 0$ the process Z_x^\uparrow is Markovian and has infinite life-time (here we use the hypothesis that Z drifts to $+\infty$). Then, note from Theorem 25 in [8] that Z_0^\uparrow , that we denote by Z^\uparrow , is well defined if and only if Z is regular for $]0, +\infty[$. According to Corollary VII.5 in [3] applied to the dual of Z , the latter is equivalent to the fact that Z has unbounded variation. This is why we have made this assumption.

1.1. Results. In the special case of the exponential functional of a drifted brownian motion conditioned to stay positive, all the properties that are studied here are already known. We discuss this case in the next subsection.

Our first result is the finiteness of $I(V^\uparrow)$ and the fact that it admits exponential moments.

Theorem 1.1. *The random variable $I(V^\uparrow)$ is almost surely finite and*

$$\forall \lambda < \Psi_V(\kappa + 1), \mathbb{E} \left[e^{\lambda I(V^\uparrow)} \right] < +\infty. \quad (1.2)$$

Moreover, there are two positive constants $K_1 > K_2 > 0$ such that for all x large enough

$$\exp(-K_1 x) \leq \mathbb{P} \left(I(V^\uparrow) \geq x \right) \leq \exp(-K_2 x). \quad (1.3)$$

Note that from (1.2), K_2 in (1.3) can be chosen as close as we want to $\Psi_V(\kappa + 1)$.

If V drifts to $+\infty$, is non-arithmetic, is such that the Laplace transform of $V(1)$ extends on the negative half-line, and if there is $\alpha > 0$ such that $\Psi_V(-\alpha) = 0$ and $\Psi_V'(-\alpha) \in]-\infty, 0[$ then, according to the results of [23] and [11] (see also [4], [16]), we have that the right tail of $I(V)$ is of order $x^{-\alpha}$. If the Lévy measure of V is subexponential or convolution equivalent, then the right tail of $I(V)$ is given by the results of [11] and [25] (see also [16]). In any case $I(V)$ has no exponential moments (see for example Proposition 5.11(1) in [22]) so the existence of finite exponential moments for $I(V^\uparrow)$ given by Theorem 1.1 contrasts with the right tail known for $I(V)$. This difference between the two tails comes from the fact that V^\uparrow stays positive so the right tail of $I(V^\uparrow)$ is determined by the right tail of the local time of V^\uparrow at positive levels, which is exponential by the Markov property. To the contrary, the absolute minimum of V is negative

and has a strong influence on the right tail of $I(V)$ (see for example [25] for comparisons between the right tail of the exponential functional and the right tail of the absolute minimum).

Then, a fundamental point of our study is Proposition 3.2 which says that for any positive y , $I(V^\uparrow)$ satisfies the random affine equation

$$I(V^\uparrow) \stackrel{\mathcal{L}}{=} A^y + e^{-y}I(V^\uparrow), \quad (1.4)$$

where A^y is independent of the second term and will be specified later. From (1.4) we can notice that $I(V^\uparrow)$ is a positive self-decomposable random variable and is therefore infinitely divisible, absolutely continuous and unimodal. It is well known that the exponential functional $I(V)$ of a spectrally negative Lévy process V is also self-decomposable (as long as it is finite). It can be seen by splitting the trajectory at $\tau(V, y)$, the first passage time at y (see also Remark 1 in Rivero [25] for an explicit representation of $I(V)$ as the stationary distribution of an Ornstein-Uhlenbeck type process, and Section 5 in Patie, Savov [22] where $I(V)$ is characterized via its Lévy triplet). As we said, the right tail of $I(V)$ (when the latter is finite) never has exponential moments while, by Theorem 1.1, the right tail of $I(V^\uparrow)$ always has. It follows that the class of self-decomposable random variables that can be represented by $I(V^\uparrow)$, for some spectrally negative Lévy process V , is disjoint from the class of self-decomposable random variables that can be represented by $I(V)$, for some spectrally negative Lévy process V . However, the following results (especially Proposition 1.10) show that the asymptotic tails at 0 of these two classes of self-decomposable random variables are comparable. Another consequence of (1.4) is that for any positive y , $I(V^\uparrow)$ can be written as the random series

$$I(V^\uparrow) \stackrel{\mathcal{L}}{=} \sum_{k \geq 0} e^{-ky} A_k^y,$$

where the random variables A_k^y are *iid* and have the same law as A^y . The novelty of our approach is mainly to study the exponential functional $I(V^\uparrow)$ via the random coefficient A^y of which we decompose the law into two convolution factors that are accessible via their Laplace transforms. This decomposition is thus a useful tool for the study of $I(V^\uparrow)$ and is the base of the proofs of many results we present below.

Our next results link the asymptotic behavior of Ψ_V with the left tail of $I(V^\uparrow)$.

Theorem 1.2. *Assume that there is $\alpha > 1$ and a positive constant C such that for all λ large enough we have $\Psi_V(\lambda) \leq C\lambda^\alpha$. Then for all $\delta \in]0, 1[$ and x small enough we have*

$$\mathbb{P}\left(I(V^\uparrow) \leq x\right) \leq \exp\left(-\delta(\alpha - 1)/(Cx)^{1/(\alpha-1)}\right). \quad (1.5)$$

Assume that there is $\alpha > 1$ and a positive constant c such that for all λ large enough we have $\Psi_V(\lambda) \geq c\lambda^\alpha$. Then for all $\delta > 1$ and x small enough we have

$$\mathbb{P}\left(I(V^\uparrow) \leq x\right) \geq \exp\left(-\delta\alpha^\alpha/(\alpha-1)/(cx)^{1/(\alpha-1)}\right). \quad (1.6)$$

Let us now recall how is usually quantified the asymptotic behavior of Ψ_V . We define, as in [3] page 94,

$$\begin{aligned} \sigma &:= \sup \left\{ \alpha \geq 0, \lim_{\lambda \rightarrow +\infty} \lambda^{-\alpha} \Psi_V(\lambda) = \infty \right\}, \\ \beta &:= \inf \left\{ \alpha \geq 0, \lim_{\lambda \rightarrow +\infty} \lambda^{-\alpha} \Psi_V(\lambda) = 0 \right\}. \end{aligned}$$

Recall that $\Psi_{V^\#}(\cdot) = \Psi_V(\kappa + \cdot)$, so σ and β are identical whether they are defined from $\Psi_{V^\#}$ or Ψ_V .

If Ψ_V has α -regular variation for $\alpha \in [1, 2]$, for example if V is a (drifted or not) α -stable Lévy process with no positive jumps, we have $\sigma = \beta = \alpha$. Recall that Q is the brownian component of V , it is well known that $\Psi_V(\lambda)/\lambda^2$ converges to $Q/2$ when λ goes to infinity. As a consequence : when $Q > 0$, Ψ_V has 2-regular variation, and when $Q = 0$, $1 \leq \sigma \leq \beta \leq 2$, where $1 \leq \sigma$ comes from the convexity of Ψ_V .

Remark 1.3. *When V has bounded variation, we know (see for example [3] Section I.1) that the brownian component of V is null, the Lévy measure ν of V satisfies $\int_{-1}^0 |x|\nu(dx) < +\infty$ and $\gamma^* := -\gamma - \int_{-1}^0 x\nu(dx)$, the factor of λ in the expression of $\Psi_V(\lambda)$, is positive (otherwise V would be the opposite of a subordinator). It is thus easy to see that in this case $\Psi_V(\lambda)/\lambda$ converges to γ^* when λ goes to infinity, so $\sigma = \beta = 1$. In the next theorem, we sometimes assume that $\sigma > 1$, the reader should be aware that it excludes the case where V has bounded variation. However, this case is quite easy and shall be treated in the remarks.*

We are now ready to state a general result on the asymptotic tails at 0 of $I(V^\uparrow)$:

Theorem 1.4. *We have*

$$\forall \beta' > \beta, \lim_{x \rightarrow 0} x^{1/(\beta'-1)} \log \left(\mathbb{P} \left(I(V^\uparrow) \leq x \right) \right) = -\infty, \quad (1.7)$$

$$\text{if } \sigma > 1, \forall \sigma' \in]1, \sigma[, \lim_{x \rightarrow 0} x^{1/(\sigma'-1)} \log \left(\mathbb{P} \left(I(V^\uparrow) \leq x \right) \right) = 0. \quad (1.8)$$

Theorem 1.4 gives for $\mathbb{P}(I(V^\uparrow) \leq x)$ a lower bound involving σ and an upper bound involving β . In the case where Ψ_V is regularly varying we can naturally expect a more precise result. Let us introduce the notation $\varphi_V(x) := \inf\{\lambda \geq 0, \Psi_V(\lambda + \kappa)/\lambda > x\}$, where we recall that $\kappa = 0$ in the case where V drifts to $+\infty$ or oscillates.

Theorem 1.5. *Assume that Ψ_V has α -regular variation at $+\infty$ for some $\alpha \in]1, 2]$. Then,*

$$\log \left(\mathbb{P} \left(I(V^\uparrow) \leq x \right) \right) \underset{x \rightarrow 0}{\sim} -(\alpha - 1) \varphi_V \left(\frac{1}{x} \right).$$

This result has the following corollary :

Corollary 1.6. *We assume that there is a positive constant C and $\alpha \in]1, 2]$ such that $\Psi_V(\lambda) \sim_{\lambda \rightarrow +\infty} C\lambda^\alpha$, then*

$$\log \left(\mathbb{P} \left(I(V^\uparrow) \leq x \right) \right) \underset{x \rightarrow 0}{\sim} -\frac{\alpha - 1}{(Cx)^{\frac{1}{\alpha-1}}}.$$

The above result is true in particular when, for some $\alpha \in]1, 2]$, V is an α -stable spectrally negative Lévy process (with adjonction or not of a drift). In particular, it agrees exactly with the tail (1.14) given in the next subsection for the particular case of a drifted brownian motion. Also, we note that Corollary 1.6 has a particular interest for applications to diffusions in random environments. Indeed, it allows to determine the exact value of the lim sup of the renormalized supremum of the local time for a diffusion in a Lévy environment V that satisfies the hypothesis of the Corollary, see [29].

Remark 1.7. *Since $\Psi_V(\lambda)/\lambda^2$ has always a finite limit at $+\infty$ we get, from Theorem 1.2, that there is always a positive constant K (depending on V) such that for x small enough*

$$\mathbb{P} \left(I(V^\uparrow) \leq x \right) \leq e^{-K/x}.$$

Remark 1.8. Note that Theorem 1.4 holds when $\beta = 1$ and $1/(\beta' - 1)$ can then equal any number in $]0, +\infty[$. When V has bounded variation, we even have a stronger result : the support of the distribution of $I(V^\uparrow)$ is $[1/\gamma^*, +\infty[$, so $\mathbb{P}(I(V^\uparrow) \leq x)$ is null for x small enough. This fact is the counterpart of Proposition 5.10(2) from [22].

Remark 1.9. Recall that $I(V^\uparrow)$ is unimodal. If 0 was a mode, then we would have $\mathbb{P}(I(V^\uparrow) \leq x) \geq cx$ for some positive constant c and x small enough, which is incompatible with, for example, Remark 1.7. As a consequence the density of $I(V^\uparrow)$ is non-decreasing on a neighborhood of 0. This allows to remark that Theorems 1.2, 1.4, 1.5, Corollary 1.6 and Remarks 1.7, 1.8 are true for the density of $I(V^\uparrow)$ in place of the repartition function $\mathbb{P}(I(V^\uparrow) \leq \cdot)$.

When V drifts to $+\infty$ we compare the left tails of $I(V^\uparrow)$ and $I(V)$ in the next proposition.

Proposition 1.10. If V drifts to $+\infty$, there is a positive constant c such that for all positive ϵ and x small enough,

$$\mathbb{P}(I(V) \leq x) \leq \mathbb{P}(I(V^\uparrow) \leq x) \leq \mathbb{P}(I(V) \leq (1 + \epsilon)x) / c\epsilon x.$$

This implies that all the results we prove for the left tail of $I(V^\uparrow)$ (Theorems 1.2, 1.4, 1.5, Corollary 1.6 and Remarks 1.7, 1.8) are true for $I(V)$ in place of $I(V^\uparrow)$. In particular, the combination of Theorem 1.5 with Proposition 1.10 allows to retrieve Proposition 1 of Pardo [15]. Reciprocally, we can combine Proposition 1.10 with the very precise result recently obtained by Patie and Savov [22], Theorem 5.24, to prove another result on the left tail of $I(V^\uparrow)$ when V does not oscillate and has unbounded variation.

Proposition 1.11. Assume that V has unbounded variation and does not oscillate. There are positive constants K_1 and K_2 such that for any positive $\delta > 1$ and x small enough,

$$\begin{aligned} K_1(\delta - 1) \sqrt{\varphi'_V\left(\frac{\delta}{x}\right)} \exp\left(-\int_{\Psi'_V(\kappa)}^{\delta/x} \frac{\varphi_V(r)}{r} dr\right) &\leq \mathbb{P}(I(V^\uparrow) \leq x) \\ &\leq \frac{K_2}{(\delta - 1)x} \sqrt{\varphi'_V\left(\frac{1}{\delta x}\right)} \exp\left(-\int_{\Psi'_V(\kappa)}^{1/\delta x} \frac{\varphi_V(r)}{r} dr\right). \end{aligned}$$

It seems that our previous results on the left tail of $I(V^\uparrow)$ could be proved from Proposition 1.11 when V has unbounded variation and does not oscillate. Moreover, it seems to give a more precise estimate compared to Theorem 1.4 in the case where we do not have assumptions on the asymptotic behavior of Ψ_V . However, since Proposition 1.11 follows from the results that are known for the left tail of $I(V)$, and not from a direct study of $I(V^\uparrow)$, it dramatically excludes the case where V oscillates. This is why we emphasize on the method developed for the previous theorems that treats indistinctly the three cases of the three possible behaviors for V .

Considering $I(V^\uparrow)$ from the point of view of self-decomposable random variables, we can combine our results on the left tail of $I(V^\uparrow)$ with the fine results of Sato, Yamazato [27] to get the following Theorem. It proves in particular the smoothness of the density of $I(V^\uparrow)$ when V has unbounded variation.

Theorem 1.12. Assume that V has unbounded variation, then

- $I(V^\uparrow)$ is a self-decomposable random variable of type I_6 , in the terminology of [27],
- The density of $I(V^\uparrow)$ belongs to the Schwartz space. All its derivatives converge to 0 at $+\infty$ and 0,
- The repartition function $x \mapsto \mathbb{P}(I(V^\uparrow) \leq x)$ is log-concave on $]0, +\infty[$.

We note that, in the class I of self-decomposable random variables (still in the terminology of [27]), the subclasses I_6 and I_7 are the one for which the less is known about the left tails.

In [22], the exponential functional $I(V)$ is also studied (when V is assumed to drift to $+\infty$) as a self-decomposable random variable using the fine tools developed by [27]. More precisely, they give a representation of the Lévy triplet of $I(V)$ which allows to determine its type as a self-decomposable random variable and the exact regularity of its density. In Lemma 5.14(2) of [22], they prove in particular that $I(V)$ is of type I_6 if and only if V has non-zero Gaussian component or infinite Lévy measure. This is also the necessary and sufficient condition under which the density of $I(V)$ is of class C^∞ , according to Proposition 5.10(3) of [22]. In our case, we believe that the same condition is also equivalent with the fact that $I(V^\uparrow)$ is of type I_6 and with the infinite differentiability its density. However, the proof requires some further developments in the study of A^y and could be made in a subsequent study. Also, note that even though we expect to have the same regularity for $I(V^\uparrow)$ as for $I(V)$ (when the latter is finite), the density of $I(V)$ never belongs to the Schwartz space for it is too heavy-tailed.

In the spectrally positive case we first prove the finiteness and the existence of some finite exponential moments. We can state the result as follows :

Theorem 1.13. *The random variable $I(Z^\uparrow)$ is almost surely finite and admits some finite exponential moments.*

Remark 1.14. *Many results are already known for the exponential functional $I(Z)$. In Patie [19] (see also Corollary 2.1 in [14] and Corollary 1.3 in [20]) a decomposition of its density into a power series is given, from which follows smoothness and precise asymptotics at $+\infty$ for the density of $I(Z)$ and its derivatives. In particular, the right tail of $I(Z)$ is of order $x^{-\alpha}$ where α is the non-trivial zero of the Laplace exponent of $-Z$ (this fact is also a consequence of the results of [23], [11], see also [4], [16]). The existence of finite exponential moments for $I(Z^\uparrow)$ given by Theorem 1.13 thus contrasts with the right tail known for $I(Z)$. Here again, this difference between the two tails comes from the fact that Z^\uparrow stays positive whereas the absolute minimum of Z is negative (it follows an exponential distribution with parameter α) and has a strong influence on the right tail of $I(Z)$.*

For the left tail of $I(Z^\uparrow)$, we are only interested in the case where Z possesses positive jumps. Indeed, if the Lévy measure of Z is the zero measure then it is known, from the Lévy-Khintchine formula, that Z is a drifted brownian motion. The exact asymptotic tail at 0 of $I(Z^\uparrow)$ is then given by Corollary 1.6.

We now provide estimates for the asymptotic tail at 0 of $I(Z^\uparrow)$. They show that the tail of $I(Z^\uparrow)$ is heavier than the one given for $I(V^\uparrow)$ and this comes from the positive jumps. We first compare with the non-conditioned case.

Proposition 1.15. *$I(Z)$ is stochastically greater than $I(Z^\uparrow)$. As a consequence,*

$$\forall x > 0, \mathbb{P}(I(Z) \leq x) \leq \mathbb{P}(I(Z^\uparrow) \leq x).$$

Proposition 1.15 has to be put in relation with results about left tails of exponential functionals of Lévy processes. We now provide such results (that are all applicable to Z) for Lévy processes that possess positive jumps. Let Y be a Lévy process drifting to $+\infty$ and let π denote its Lévy measure. We also define $\bar{\pi}(x) := \pi([x, +\infty[)$ for all $x > 0$. In the following theorem we link the left tail of the exponential functional $I(Y)$ with the Lévy measure π .

Theorem 1.16. *There is a positive constant C such that for all x small enough,*

$$\mathbb{P}(I(Y) \leq x) \geq Cx\bar{\pi}(\log(1/x)). \tag{1.9}$$

Another bound is required, for example in the case where the support of π is bounded from above or in the case where $\bar{\pi}$ decreases too rapidly. Assume that $\pi(\cdot \cap]0, +\infty[)$ is non-zero, denote by S the supremum of its support (note that $S = +\infty$ if this support is not bounded from above). We fix any $c > 1/2S$ (where by convention, $1/2\infty = 0$). Then, for all x small enough we have

$$\mathbb{P}(I(Y) \leq x) \geq e^{-c(\log(x))^2}. \quad (1.10)$$

In particular, if $Y = \alpha N$ where N is a standard Poisson process and $\alpha > 0$, then we have

$$-\log(\mathbb{P}(I(\alpha N) \leq x)) \underset{x \rightarrow 0}{\sim} \frac{1}{2\alpha} (\log(x))^2. \quad (1.11)$$

Note that the tail in (1.11) does not depend on the parameter of the standard Poisson process.

It is interesting to put in relation Remark 1.7 and the combination Theorem 1.16-Proposition 1.15. We see that in the presence of positive jumps, the asymptotic tail at 0 of $I(Z^\uparrow)$ (or $I(Z)$) is always at least of order $e^{-c(\log(x))^2}$ for some constant c . To the contrary, in the absence of positive jumps, the asymptotic tail at 0 of the exponential functional is always at most of order $e^{-c/x}$ for some constant c . The existence of jumps thus plays an important role for the asymptotic tail at 0 of an exponential functional.

Also, note that (1.10) gives exactly the minimal possible order of $\mathbb{P}(I(Y) \leq x)$, for Y lying among Lévy processes drifting to $+\infty$ with positive jumps. Indeed, (1.10) is always satisfied for such Y and (1.11) shows that this bound is optimal for the exponential functional of a multiple of a standard Poisson process.

When $\bar{\pi}$ does not decrease too rapidly then (1.9) is preferable to (1.10). Actually, we can precise (1.9) in the spectrally positive case under a regularity hypothesis on π . Let us introduce the following assumption on π that is made in [17] to study the asymptotic tails of exponential functionals of subordinators :

$$(A) : \quad \exists \alpha > 0 \text{ s.t. } \forall y \in \mathbb{R}, \quad \frac{\bar{\pi}(x+y)}{\bar{\pi}(x)} \underset{x \rightarrow +\infty}{\longrightarrow} e^{-\alpha y}.$$

Our last result precises (1.9) when Y is spectrally positive and π satisfies (A). It complements the study [19] of densities of exponential functionals of spectrally positive Lévy processes and we prove it as a consequence of the factorization identity in [14] together with a result of [17].

Proposition 1.17. *Assume that Y is spectrally positive and (A) is satisfied for some $\alpha > 0$. We denote by κ_Y the non-trivial zero of the Laplace exponent of $-Y$, by H the ascending ladder height process associated with Y , and by m_Y the density of $I(Y)$. Then,*

$$m_Y(x) \underset{x \rightarrow 0}{\sim} \frac{\mathbb{E}[I(H)^{-\alpha}] \Gamma(\alpha + \kappa_Y)}{\Gamma(\kappa_Y)} \bar{\pi}(\log(1/x)).$$

The study of the spectrally positive case does not go as far as the study of the spectrally negative case. The reason for this is twofold. First, we do not have, in the spectrally positive case, a decomposition of the law of $I(Z^\uparrow)$ as in (1.4), which deprives us of an important tool for the study. Secondly we do not need, in the applications, the results on the exponential functional to be as precise, in the spectrally positive case, as in the spectrally negative case. Indeed, in the study of a diffusion in a spectrally negative Lévy environment V drifting to $-\infty$, a random variable \mathcal{R} appears. Its law is the convolution of the laws of $I(V^\uparrow)$ and $I(\hat{V}^\uparrow)$, where $\hat{V} := -V$ is the dual process of V and is thus spectrally positive. The combination of the above theorems shows that for some things the behavior of $I(V^\uparrow)$ is dominant in the study of \mathcal{R} when V has jumps. In particular, the asymptotic tail at 0 of \mathcal{R} is the same as the one of $I(V^\uparrow)$.

The rest of the paper is organized as follows. In Section 2 we prove some preliminary results on V^\uparrow . In Section 3 we prove Theorem 1.1 and establish Proposition 3.2, the decomposition of the law of $I(V^\uparrow)$. In Section 4 we prove Theorems 1.2, 1.4 and 1.5 by studying the asymptotic behavior of the Laplace transform of $I(V^\uparrow)$, and we establish Propositions 1.10 and 1.11. In Section 5 we prove Theorem 1.12 looking at $I(V^\uparrow)$ as a self-decomposable random variable. Section 6 is devoted to the spectrally positive case, it contains the proofs of Theorem 1.13 and Proposition 1.15. In Section 7 we study the asymptotic tails at 0 of exponential functionals in the presence of positive jumps, this is where we prove Theorem 1.16 and Proposition 1.17.

1.2. The example of drifted brownian motion conditioned to stay positive. The most simple case is the intersection of the spectrally positive and the spectrally negative case, that is, when V is a drifted brownian motion. All the results mentioned here are already known in this case. We define the κ -drifted brownian motion by $W_\kappa(t) := W(t) - \frac{\kappa}{2}t$, where W is a standard brownian motion. It is known that the two processes W_κ^\uparrow and $W_{-\kappa}^\uparrow$ are equal in law. This follows, for example, from the expression of the generator of W_κ^\uparrow , or from the fact that for positive κ , the Laplace exponent of $W_\kappa^\#$ is equal to the Laplace exponent of $W_{-\kappa}$, so the processes conditioned to stay positive have the same law. We thus only consider positive κ .

It is known (see (4.6) in Androletti, Devulder [1], see also Lemma 6.6 in [2]) that $I(W_k^\uparrow)$ is almost surely finite and has Laplace transform

$$\mathbb{E}[e^{-\lambda I(W_k^\uparrow)}] = \frac{1}{2^\kappa \Gamma(1+\kappa)} \frac{(2\sqrt{2\lambda})^\kappa}{I_\kappa(2\sqrt{2\lambda})},$$

where I_κ is a modified Bessel function. This expression can also be written

$$\mathbb{E}[e^{-\lambda I(W_k^\uparrow)}] = \frac{1}{\Gamma(1+\kappa)} \frac{1}{\sum_{j=0}^{+\infty} \frac{(2\lambda)^j}{j! \Gamma(1+j+\kappa)}}. \quad (1.12)$$

It is easy to see that it can be analytically extended in a neighborhood of 0, so the random variable $I(W_k^\uparrow)$ admits some finite exponential moments.

An easy calculation on the asymptotic of (1.12) when λ goes to infinity yields

$$-\log \left(\mathbb{E}[e^{-\lambda I(W_k^\uparrow)}] \right) \underset{\lambda \rightarrow +\infty}{\sim} 2\sqrt{2\lambda}. \quad (1.13)$$

Combining (1.13) and De Bruijn's Theorem (see Theorem 4.12.9 in [6]) we get

$$-\log \left(\mathbb{P} \left(I(W_k^\uparrow) \leq x \right) \right) \underset{x \rightarrow 0}{\sim} \frac{2}{x}. \quad (1.14)$$

This estimate can be seen as a particular case of Corollary 1.6 (when it is applied in the case of a drifted brownian motion).

As the expression (1.12) extends to the imaginary line, we get the expression of the characteristic function of $I(W_k^\uparrow)$ which can be proved, using estimates on modified Bessel functions, to belong to the Schwartz space. Therefore, the density of $I(W_k^\uparrow)$, which is the Fourier transform of its characteristic function, belongs to the Schwartz space, which agrees with Theorem 1.12.

2. PRELIMINARY RESULTS ON V^\uparrow

2.1. Exponential functionals and excursion theory. We fix $y > 0$. In this subsection, we use excursions to prove that the integrals of exponential $V^\uparrow(\tau(V^\uparrow, y) + \cdot)$ or $(V^\#(t), 0 \leq t \leq \tau(V^\#, y))$ stopped at their last passage time at y and 0 respectively are equal in law to some subordinators stopped at independent exponential random variables.

It is easy to see that the regularity of $\{y\}$ for the markovian processes $V_y^\#$ and V_y^\uparrow is equivalent to the regularity of $\{0\}$ for V (or $V^\#$) which in turn, according to Corollary VII.5 in [3], is equivalent to the fact that V has unbounded variation. The property of $\{y\}$ being instantaneous for $V_y^\#$ and V_y^\uparrow is equivalent to the same property of $\{0\}$ for V , but this is a well known property of spectrally negative Lévy processes that are not opposite of subordinators. $\{y\}$ is thus always instantaneous for $V_y^\#$ and V_y^\uparrow and the only alternative is whether it is regular or not, which corresponds to the fact that V has or not unbounded variation.

We apply excursion theory away from y (see Section IV of [3]). Let us denote by L_y^\uparrow (respectively $L_y^\#$) a local time at y of the process V_y^\uparrow (respectively $V_y^\#$) and η_y^\uparrow (respectively $\eta_y^\#$) the associated excursion measure. We denote $\eta_0^\#$ for $\eta_0^\#$. The inverse of the local time $L_y^{\uparrow,-1}$ (respectively $L_y^{\#,-1}$) is a subordinator and V_y^\uparrow (respectively $V_y^\#$) can be represented as a Poisson point process on the set of excursions, with intensity measure η_y^\uparrow (respectively $\eta_y^\#$). Note that this is also true in the irregular case (when V has bounded variation) if the local time L_y^\uparrow (respectively $L_y^\#$) is defined artificially as in [3], Section IV.5. In this case, the excursion measure is proportional to the law of the first excursion and in particular the total mass of the excursion measure is finite.

Given $\xi : [0, \zeta] \rightarrow \mathbb{R}$ an excursion away from y , we define $\zeta(\xi)$ to be its life-time, $H_y(\xi) := \sup_{[0, \zeta(\xi)]} \xi - y$ its height and $G(\xi) := \int_0^{\zeta(\xi)} e^{-\xi(t)} dt$.

For any $h > 0$, we consider IP_h , FP_h and N_h three subsets that make a partition of the excursions of $V^\#$ away from y . These three subsets are respectively : the set of excursions higher than h that stay positive before reaching $y + h$, the set of excursions of height smaller than h that stay positive, the set of excursions that reach $]-\infty, 0]$ before ever reaching $y + h$:

$$IP_h := \{\xi, H_y(\xi) \geq h, \forall t \in [0, \tau(\xi, y + h)], \xi(t) > 0\},$$

$$FP_h := \{\xi, H_y(\xi) < h, \forall t \in [0, \zeta(\xi)], \xi(t) > 0\}, \quad N_h := \{\xi, \tau(\xi,]-\infty, 0]) < \tau(\xi, y + h)\}.$$

IP_∞ , FP_∞ and N_∞ are defined as the monotone limits of the sets IP_h , FP_h and N_h : IP_∞ is the set of infinite excursions that stay positive, FP_∞ is the set of finite excursions that stay positive and N_∞ is the set of excursions that reach $]-\infty, 0]$. $\eta_y^\#(IP_\infty)$ and $\eta_y^\#(N_\infty)$ are always finite whereas $\eta_y^\#(FP_\infty)$ is infinite in the regular case (when V has unbounded variation). Also, note that $\eta_y^\#(IP_\infty) = 0$ if V oscillates.

Lemma 2.1. *Let y be positive and let S be a pure jump subordinator with Lévy measure $G\eta_y^\#(\cdot \cap FP_\infty)$, the image measure of $\eta_y^\#(\cdot \cap FP_\infty)$ by G . Let T be an exponential random variable with parameter $\eta_y^\#(IP_\infty) + \eta_y^\#(N_\infty)$, independent of S . We have*

$$\int_{\tau(V^\uparrow, y)}^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt \stackrel{\mathcal{L}}{=} S_T, \quad (2.15)$$

where $\tau(\cdot, \cdot)$ and $\mathcal{R}(\cdot, \cdot)$ are defined in the introduction.

Proof. $V^\uparrow(\tau(V^\uparrow, y) + \cdot)$ has the same law as V_y^\uparrow , from the Markov property applied to V^\uparrow at time $\tau(V^\uparrow, y)$ and the absence of positive jumps. As a consequence, $\int_{\tau(V^\uparrow, y)}^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt$ is equal in law to $\int_0^{\mathcal{R}(V_y^\uparrow, y)} e^{-V_y^\uparrow(t)} dt$ and we are left to prove the result for the latter.

Then, let us fix $h > 0$. As it is mentioned in the introduction, $(V_y^\uparrow(t), 0 \leq t \leq \tau(V_y^\uparrow, y + h))$ is equal in law to $(V_y^\#(t), 0 \leq t \leq \tau(V_y^\#, y + h))$ conditionally on $\{\tau(V_y^\#, y + h) < \tau(V_y^\#,]-\infty, 0])\}$.

$(V_y^\sharp(t), 0 \leq t \leq \tau(V_y^\sharp, y+h))$ can be built from a Poisson point process on the set of excursions with intensity measure η_y^\sharp . $(V_y^\uparrow(t), 0 \leq t \leq \tau(V_y^\uparrow, y+h))$ can be built from this same process, conditioned not to have jumps in N_h before its first jump in IP_h . In other words, we build $(V_y^\uparrow(t), 0 \leq t \leq \tau(V_y^\uparrow, y+h))$ from the process of jumps in FP_h stopped at the exponential time (that has parameter $\eta_y^\sharp(IP_h) + \eta_y^\sharp(N_h)$) at which occurs the first jump in $IP_h \cup N$ and conditionally to the fact that this jump belongs to IP_h . Then, the process of jumps in FP_h and in $IP_h \cup N$ are independent and by a property of Poisson point processes, the fact that the first jump in $IP_h \cup N$ belongs to IP_h is independent of the time when this jump occurs. As a consequence, $(V_y^\uparrow(t), 0 \leq t \leq \tau(V_y^\uparrow, y+h))$ is built from a Poisson point process with intensity measure $\eta_y^\sharp(\cdot \cap FP_h)$, until an independent exponential time, T_h , of parameter $\eta_y^\sharp(IP_h) + \eta_y^\sharp(N_h)$ where we pick, independently, a jump following the law $\eta_y^\sharp(\cdot \cap IP_h)/\eta_y^\sharp(IP_h)$ (and we only keep the part of this excursion that is before its hitting time of $y+h$).

Let $\mathcal{R}^h(V_y^\uparrow, y)$ be the last passage time of V_y^\uparrow at y before $\tau(V_y^\uparrow, y+h)$:

$$\mathcal{R}^h(V_y^\uparrow, y) := \sup\{t \in [0, \tau(V_y^\uparrow, y+h)], V_y^\uparrow(t) = y\}.$$

From above, if $(p_s)_{s \geq 0}$ is a Poisson point process in FP_∞ with intensity measure $\eta_y^\sharp(\cdot \cap FP_\infty)$ and if T_h is an independent exponential random variable with parameter $\eta_y^\sharp(IP_h) + \eta_y^\sharp(N_h)$, then $(V_y^\uparrow(t), 0 \leq t \leq \mathcal{R}^h(V_y^\uparrow, y))$ is built by putting aside the excursions of the process $(p_s \mathbb{1}_{p_s \in FP_h}, 0 \leq s \leq T_h)$. Since V_y^\uparrow converges almost surely to $+\infty$, $\mathcal{R}^h(V_y^\uparrow, y)$ converges almost surely to $\mathcal{R}(V_y^\uparrow, y)$, the last passage time at y , when h goes to infinity. On the other hand, $\eta_y^\sharp(IP_h) + \eta_y^\sharp(N_h)$ converges to $\eta_y^\sharp(IP_\infty) + \eta_y^\sharp(N_\infty)$ when h goes to infinity. As a consequence T_h converges to an exponential random variable T with parameter $\eta_y^\sharp(IP_\infty) + \eta_y^\sharp(N_\infty) > 0$. Also, FP_h increases to FP_∞ when h goes to infinity. Then, identifying the limits when h goes to infinity of both $(V_y^\uparrow(t), 0 \leq t \leq \mathcal{R}^h(V_y^\uparrow, y))$ and $(p_s \mathbb{1}_{p_s \in FP_h}, 0 \leq s \leq T_h)$, we get that $(V_y^\uparrow(t), 0 \leq t \leq \mathcal{R}(V_y^\uparrow, y))$ is built by putting aside the excursions of the process $(p_s, 0 \leq s \leq T)$, where T is an exponential random variable with parameter $\eta_y^\sharp(IP_\infty) + \eta_y^\sharp(N_\infty)$ and is independent from $(p_s, s \geq 0)$.

Now, remark that $\int_0^{\mathcal{R}(V_y^\uparrow, y)} e^{-V_y^\uparrow(t)} dt$ is the sum of the images by G of the excursions of $(V_y^\uparrow(t), 0 \leq t \leq \mathcal{R}(V_y^\uparrow, y))$ away from y . We thus have

$$\int_0^{\mathcal{R}(V_y^\uparrow, y)} e^{-V_y^\uparrow(t)} dt \stackrel{\mathcal{L}}{=} \sum_{0 < s < T} G(p_s). \quad (2.16)$$

By properties of Poisson point processes, the process in the right hand side, $\sum_{0 < s < T} G(p_s)$, is the sum of the jumps of a Poisson point process on \mathbb{R}_+ , with intensity measure $G\eta_y^\sharp(\cdot \cap FP_\infty)$. Thus, from the Lévy-Ito decomposition, it has the same law as the subordinator S , which yields the result. \square

Remark 2.2. *In the case where V has bounded variation, the total mass of η_y^\sharp is finite so S is only a compound Poisson process. In particular S_T is then null with a positive probability.*

$y > 0$ is still fixed (and arbitrary), let $\mathcal{R}^y(V^\sharp, 0)$ be the last passage time of V^\sharp at 0 before $\tau(V^\sharp, y)$:

$$\mathcal{R}^y(V^\sharp, 0) := \sup\{t \in [0, \tau(V^\sharp, y)], V^\sharp(t) = 0\}.$$

In order to study the trajectory of V^\sharp before $\mathcal{R}^y(V^\sharp, 0)$, we now consider excursions away from 0. Let I_y and F_y denote respectively the subset of excursions higher than y and lower than y :

$$I_y := \{\xi, H_0(\xi) \geq y\}, \quad F_y := \{\xi, H_0(\xi) < y\}.$$

A simpler proof as for Lemma 2.1 gives the following lemma.

Lemma 2.3. *Let S be a pure jump subordinator with Lévy measure $G\eta^\sharp(\cdot \cap F_y)$, the image measure of $\eta^\sharp(\cdot \cap F_y)$ by G . Let T be an exponential random variable with parameter $\eta^\sharp(I_y)$ which is independent of S . We have*

$$\int_0^{\mathcal{R}^y(V^\sharp, 0)} e^{-V^\sharp(t)} dt \stackrel{\mathcal{L}}{=} S_T. \quad (2.17)$$

In the case where V does not oscillate, V^\sharp drifts to $+\infty$ and we need to study the trajectory before $\mathcal{R}(V^\sharp, 0)$, the last passage time of V^\sharp at 0. We still consider excursions away from 0. Let I and F denote respectively the subsets of infinite and finite excursions :

$$I := \{\xi, \zeta(\xi) = +\infty\}, \quad F := \{\xi, \zeta(\xi) < +\infty\}.$$

A similar proof as for Lemma 2.1 gives the following lemma.

Lemma 2.4. *We assume that V does not oscillate. Let S be a pure jump subordinator with Lévy measure $G\eta^\sharp(\cdot \cap F)$, the image measure of $\eta^\sharp(\cdot \cap F)$ by G . Let T be an exponential random variable with parameter $\eta^\sharp(I)$ which is independent of S . We have*

$$\int_0^{\mathcal{R}(V^\sharp, 0)} e^{-V^\sharp(t)} dt \stackrel{\mathcal{L}}{=} S_T. \quad (2.18)$$

2.2. V^\uparrow and V^\sharp shifted at a last passage time. To obtain the decomposition (1.4) of the law of $I(V^\uparrow)$, we split V^\uparrow at its last passage time at a point y and obtain two independent trajectories that we can identify.

Lemma 2.5. (Corollary VII.19 of [3])

For any positive y , the two trajectories

$$\left(V^\uparrow(t), 0 \leq t \leq \mathcal{R}(V^\uparrow, y) \right) \text{ and } \left(V^\uparrow(t + \mathcal{R}(V^\uparrow, y)) - y, t \geq 0 \right)$$

are independent and the second is equal in law to V^\uparrow .

Lemma 2.6. • The two trajectories $(V^\sharp(t), 0 \leq t \leq \mathcal{R}^y(V^\sharp, 0))$

$$\text{and } \left(V^\sharp(t + \mathcal{R}^y(V^\sharp, 0)), 0 \leq t \leq \tau(V^\sharp, y) - \mathcal{R}^y(V^\sharp, 0) \right)$$

are independent and the second is equal in law to $(V^\uparrow(t), 0 \leq t \leq \tau(V^\uparrow, y))$. As a consequence we have $\tau(V^\uparrow, y) \stackrel{\mathcal{L}}{=} \tau(V^\sharp, y) - \mathcal{R}^y(V^\sharp, 0) \leq \tau(V^\sharp, y)$.

• We assume that V does not oscillate. The two trajectories

$$\left(V^\sharp(t), 0 \leq t \leq \mathcal{R}(V^\sharp, 0) \right) \text{ and } \left(V^\sharp(t + \mathcal{R}(V^\sharp, 0)), t \geq 0 \right)$$

are independent and the second is equal in law to V^\uparrow .

Proof. We fix $y > 0$ and $a \in]0, y[$. Let us denote by $(e(s), s \geq 0)$ the process of excursions of V^\sharp away from 0. Recall the notations I_y and F_y . $T_y := \inf\{s \geq 0, e(s) \in I_y\}$ is the time when occurs the first excursion higher than y and ξ_y denotes this excursion.

Decomposing V^\sharp as its excursions away from 0, we see that $\mathcal{R}^y(V^\sharp, 0)$ is the instant when begins the first excursion higher than y , so

$$\left(V^\sharp(t + \mathcal{R}^y(V^\sharp, 0)), 0 \leq t \leq \tau(V^\sharp, y) - \mathcal{R}^y(V^\sharp, 0) \right) = (\xi_y(t), 0 \leq t \leq \tau(\xi_y, y)). \quad (2.19)$$

$(V^\sharp(t), 0 \leq t \leq \mathcal{R}^y(V^\sharp, 0))$ is thus a function of $(e(s)\mathbb{1}_{e(s) \in F_y}, 0 \leq s \leq T_y)$ while $(V^\sharp(t + \mathcal{R}^y(V^\sharp, 0)), 0 \leq t \leq \tau(V^\sharp, y) - \mathcal{R}^y(V^\sharp, 0))$ is a function of ξ_y . By properties of Poisson point processes, T_y is an exponential random variable independent of ξ_y and the process of finite excursions $(e(s)\mathbb{1}_{e(s) \in F_y}, s \geq 0)$ is also independent of ξ_y . Therefore the objects $(e(s)\mathbb{1}_{e(s) \in F_y}, 0 \leq s \leq T_y)$ and ξ_y are independent. From this independence we deduce that

$$\left(V^\sharp(t), 0 \leq t \leq \mathcal{R}^y(V^\sharp, 0) \right) \perp\!\!\!\perp \left(V^\sharp(t + \mathcal{R}^y(V^\sharp, 0)), 0 \leq t \leq \tau(V^\sharp, y) - \mathcal{R}^y(V^\sharp, 0) \right),$$

which is the required independence. It only remains to prove that the right hand side in (2.19) has the same law as $(V^\uparrow(t), 0 \leq t \leq \tau(V^\uparrow, y))$.

Using the Markov property at time $\tau(\xi_a, a)$, for an excursion $\xi_a \in I_a$, we have that $\xi_a(\cdot + \tau(\xi_a, a))$ equals in law V_a^\sharp killed when it ever reaches 0.

Since $I_y \subset I_a$ we can apply this to an excursion $\xi_y \in I_y$ and get that $(\xi_y(t + \tau(\xi_y, a)), 0 \leq t \leq \tau(\xi_y, y) - \tau(\xi_y, a))$ is equal in law to $(V_a^\sharp(t), 0 \leq t \leq \tau(V_a^\sharp, y))$ conditioned to reach y before 0. Since V_a^\sharp has no positive jumps, reaching y before 0 is the same as reaching y before $]-\infty, 0]$.

As we mentioned in the introduction, $(V_a^\sharp(t), 0 \leq t \leq \tau(V_a^\sharp, y))$ conditioned to reach y before $]-\infty, 0]$ is equal in law to $(V_a^\uparrow(t), 0 \leq t \leq \tau(V_a^\uparrow, y))$. Putting all this together we get

$$(\xi_y(t + \tau(\xi_y, a)), 0 \leq t \leq \tau(\xi_y, y) - \tau(\xi_y, a)) \stackrel{\mathcal{L}}{=} \left(V_a^\uparrow(t), 0 \leq t \leq \tau(V_a^\uparrow, y) \right).$$

Since $\tau(\xi_y, a)$ converges almost surely to 0 when a goes to 0 and V_a^\uparrow converges in distribution to V^\uparrow according to Proposition VII.14 in [3], we can let a go to 0 in both members and get

$$(\xi_y(t), 0 \leq t \leq \tau(\xi_y, y)) \stackrel{\mathcal{L}}{=} \left(V^\uparrow(t), 0 \leq t \leq \tau(V^\uparrow, y) \right). \quad (2.20)$$

As a consequence the right hand side in (2.19) has the same law as $(V^\uparrow(t), 0 \leq t \leq \tau(V^\uparrow, y))$, which concludes the proof of the first point of the lemma.

We now assume that V does not oscillate (so that V^\sharp drifts to $+\infty$) and we prove the second point. For the independence, the arguments of the proof of the first point can be repeated, just replacing y by $+\infty$ (we consider ξ_∞ , the infinite excursion away from 0, instead of the first excursion higher than y). To prove that $(V^\sharp(t + \mathcal{R}(V^\sharp, 0)), t \geq 0)$ is equal in law to V^\uparrow , it suffices to prove that ξ_∞ is equal in law to V^\uparrow . Let y be finite, we know from the proof of the first point that (2.20) is true for any excursion in I_y . Since $\xi_\infty \in I_y$ we have

$$(\xi_\infty(t), 0 \leq t \leq \tau(\xi_\infty, y)) \stackrel{\mathcal{L}}{=} \left(V^\uparrow(t), 0 \leq t \leq \tau(V^\uparrow, y) \right).$$

Since y is arbitrary and $\tau(V^\uparrow, y)$ converges almost surely to $+\infty$ when y goes to $+\infty$, we get

$$(\xi_\infty(t), t \geq 0) \stackrel{\mathcal{L}}{=} \left(V^\uparrow(t), t \geq 0 \right),$$

which gives the result. □

3. FINITENESS, EXPONENTIAL MOMENTS, AND SELF-DECOMPOSABILITY

3.1. Finiteness and exponential moments : Proof of Theorem 1.1. We are grateful to an anonymous referee for the following proof that is considerably simpler than the one given by the author in the previous versions of this paper.

Proof. of Theorem 1.1 (the finiteness of $I(V^\uparrow)$ and (1.2))

The idea of the proof is to provide finite upper bounds for the moments of $I(V^\uparrow)$. The first step is to prove that $\mathbb{E}[I(V^\uparrow)] < +\infty$. Using Fubini's Theorem and Corollary VII.16 of [3] we have

$$\begin{aligned}\mathbb{E}[I(V^\uparrow)] &= \int_0^{+\infty} \mathbb{E}[e^{-V^\uparrow(t)}] dt = \int_0^{+\infty} \int_0^{+\infty} e^{-y} \mathbb{P}(V^\uparrow(t) \in dy) dt \\ &= \int_0^{+\infty} \int_0^{+\infty} e^{-y} \frac{yW(y)}{t} \mathbb{P}(V(t) \in dy) dt.\end{aligned}$$

Now, using Corollary VII.3 of [3] we get

$$\mathbb{E}[I(V^\uparrow)] = \int_0^{+\infty} \int_0^{+\infty} e^{-y} W(y) \mathbb{P}(\tau(V, y) \in dt) dy = \int_0^{+\infty} e^{-y} W(y) \mathbb{P}(\tau(V, y) < +\infty) dy.$$

Since $\mathbb{P}(\tau(V, y) < +\infty) = \mathbb{P}(\sup_{[0, +\infty[} V \geq y) = e^{-\kappa y}$ we obtain

$$\mathbb{E}[I(V^\uparrow)] = \int_0^{+\infty} e^{-(1+\kappa)y} W(y) dy = \frac{1}{\Psi_V(\kappa + 1)} < +\infty.$$

The finiteness in the above expression comes from the fact that $\int_0^{+\infty} e^{-\lambda y} W(y) dy < +\infty$ for $\lambda > \kappa$. As a consequence, the exponential functional $I(V^\uparrow)$ is almost surely finite and has finite expectation.

We now turn to the proof of the finiteness of the Laplace transform. We proceed by bounding the moments of the exponential functional. For any $x \geq 0$ let us define $h(x) := \mathbb{E}[I(V_x^\uparrow)]$. For any $x > 0$ we have

$$I(V^\uparrow) = \int_0^{+\infty} e^{-V^\uparrow(t)} dt \geq \int_{\tau(V, x)}^{+\infty} e^{-V^\uparrow(t)} dt \stackrel{\mathcal{L}}{=} \int_0^{+\infty} e^{-V_x^\uparrow(t)} dt = I(V_x^\uparrow).$$

For the equality in law in the above expression we have used the Markov property for V^\uparrow at time $\tau(V, x)$. As a consequence we have

$$\forall x > 0, h(x) = \mathbb{E}[I(V_x^\uparrow)] \leq \mathbb{E}[I(V^\uparrow)] = 1/\Psi_V(\kappa + 1) < +\infty. \quad (3.21)$$

Now, note that for any $k \geq 1$,

$$\begin{aligned}\mathbb{E} \left[\left(I(V^\uparrow) \right)^k \right] &= \mathbb{E} \left[\int_0^{+\infty} \dots \int_0^{+\infty} e^{-V^\uparrow(t_1)} \times \dots \times e^{-V^\uparrow(t_k)} dt_1 \dots dt_k \right] \\ &= k! \mathbb{E} \left[\int_{0 \leq t_1 < \dots < t_k} e^{-V^\uparrow(t_1)} \times \dots \times e^{-V^\uparrow(t_k)} dt_1 \dots dt_k \right],\end{aligned}$$

so that

$$\forall k \geq 1, \mathbb{E} \left[\left(I(V^\uparrow) \right)^k \right] / k! = \mathbb{E} \left[\int_{0 \leq t_1 < \dots < t_k} e^{-V^\uparrow(t_1)} \times \dots \times e^{-V^\uparrow(t_k)} dt_1 \dots dt_k \right]. \quad (3.22)$$

Let us prove by induction that for any $k \geq 1$,

$$\mathbb{E} \left[\left(I(V^\uparrow) \right)^k \right] \leq k! \left(\mathbb{E}[I(V^\uparrow)] \right)^k < +\infty. \quad (3.23)$$

(3.23) is clearly true for $k = 1$. Let us assume that it is true for some arbitrary rank k . According to (3.22), $\mathbb{E}[(I(V^\uparrow))^{k+1}]/(k+1)!$ equals

$$\begin{aligned} & \mathbb{E} \left[\int_{0 \leq t_1 < \dots < t_{k+1}} e^{-V^\uparrow(t_1)} \times \dots \times e^{-V^\uparrow(t_k)} \times e^{-V^\uparrow(t_{k+1})} dt_1 \dots dt_k dt_{k+1} \right] \\ &= \mathbb{E} \left[\int_{0 \leq t_1 < \dots < t_k} e^{-V^\uparrow(t_1)} \times \dots \times e^{-V^\uparrow(t_k)} \left(\int_{t_k}^{+\infty} e^{-V^\uparrow(s)} ds \right) dt_1 \dots dt_k \right] \\ &= \mathbb{E} \left[\int_{0 \leq t_1 < \dots < t_k} e^{-V^\uparrow(t_1)} \times \dots \times e^{-V^\uparrow(t_k)} \mathbb{E} \left[\int_{t_k}^{+\infty} e^{-V^\uparrow(s)} ds \mid \sigma(V^\uparrow(u), 0 \leq u \leq t_k) \right] dt_1 \dots dt_k \right]. \end{aligned}$$

From the Markov property at time t_k , the conditional expectation in the above expression equals $h(V^\uparrow(t_k))$ which, according to (3.21), is almost surely less than $\mathbb{E}[I(V^\uparrow)]$. We thus get

$$\begin{aligned} \mathbb{E}[(I(V^\uparrow))^{k+1}]/(k+1)! &\leq \mathbb{E}[I(V^\uparrow)] \times \mathbb{E} \left[\int_0^{0 \leq t_1 < \dots < t_k} e^{-V^\uparrow(t_1)} \times \dots \times e^{-V^\uparrow(t_k)} dt_1 \dots dt_k \right] \\ &= \mathbb{E}[I(V^\uparrow)] \times \mathbb{E} \left[(I(V^\uparrow))^k / k! \right] \leq \left(\mathbb{E}[I(V^\uparrow)] \right)^{k+1}, \end{aligned}$$

where we have used (3.22) and the induction hypothesis. Thus the induction is proved. As a consequence, for all $\lambda \in]0, 1/\mathbb{E}[I(V^\uparrow)][=]0, \Psi_V(\kappa + 1)[$ we have

$$\mathbb{E} \left[e^{\lambda I(V^\uparrow)} \right] = \sum_{k \geq 0} \lambda^k \mathbb{E} \left[(I(V^\uparrow))^k / k! \right] \leq \sum_{k \geq 0} \left(\lambda \mathbb{E}[I(V^\uparrow)] \right)^k < +\infty.$$

The finiteness of the Laplace transform is obvious for $\lambda \leq 0$, so (1.2) is proved. \square

The argument of the above proof shows that, if $I = \int_0^{+\infty} f(A(t)) dt$ where f is a positive function and A is a positive càd-làg Markovian process with no positive jumps, then I admits some finite exponential moments if and only if I has finite expectation.

Remark 3.1. *If V does not oscillate, the finiteness of $I(V^\uparrow)$ can be derived as a consequence of Lemma 2.6. Indeed, from the second statement of Lemma 2.6 we have*

$$I(V^\uparrow) \stackrel{\mathcal{L}}{=} \int_0^{+\infty} e^{-V^\sharp(t + \mathcal{R}(V^\sharp, 0))} dt = \int_{\mathcal{R}(V^\sharp, 0)}^{+\infty} e^{-V^\sharp(t)} dt \leq \int_0^{+\infty} e^{-V^\sharp(t)} dt = I(V^\sharp).$$

Then, V^\sharp drifts to $+\infty$ so Theorem 1 in [4] ensures that $I(V^\sharp)$ is almost surely finite which yields the result. Note that, if one would wish to study $I(V^\uparrow)$ using the above inequality, one would lose all the relevant information about the right tail of the functional. Indeed, the right tail of $I(V^\uparrow)$ is at most exponential while the right tail of $I(V^\sharp)$ is at least inverse polynomial, as we said in the introduction. However, the above inequality leads to a relevant comparison of the left tails of $I(V^\uparrow)$ and $I(V^\sharp)$ in Proposition 1.10.

3.2. Decomposition of the law of $I(V^\uparrow)$. In this subsection, we prove that the law of $I(V^\uparrow)$ is solution of the random affine equation (1.4) and we give a decomposition of its non-trivial coefficient A^y . This is a key point of our analysis of the law of $I(V^\uparrow)$. The self-decomposability of $I(V^\uparrow)$, which is a direct consequence of the next proposition, has already been proved in [5] and [13] using the same argument. However we give a proof for the sake of completeness, and because we go further, by decomposing the variable A^y appearing in the relation of self-decomposability.

Proposition 3.2. *For any $y > 0$, the law of $I(V^\uparrow)$ satisfies the random affine equation*

$$I(V^\uparrow) \stackrel{\mathcal{L}}{=} \int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt + S_T + e^{-y} I(\tilde{V}^\uparrow). \quad (3.24)$$

The three terms of the right hand side are independent, S_T is as in Lemma 2.1, and \tilde{V}^\uparrow is an independent copy of V^\uparrow . We define

$$A^y := \int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt + S_T$$

to lighten notations. As a consequence, $I(V^\uparrow)$ has the same law as the sum of the random series

$$I(V^\uparrow) \stackrel{\mathcal{L}}{=} \sum_{k \geq 0} e^{-ky} A_k^y, \quad (3.25)$$

where the random variables A_k^y are iid and have the same law as A^y .

Let us remark that, according to Theorem VII.18 in [3], A^y is equal in law to $e^{-y} \int_0^{\tau(V, y)} e^{V(t)} dt$ conditionally on $\{\tau(V, y) < +\infty\}$.

Remark 3.3. The almost sure convergence of the random series in (3.25) is a consequence of the almost sure finiteness, given by Theorem 1.1, of the positive random variable $I(V^\uparrow)$. Also, it is a well known fact on random power series with iid coefficients that their radius of convergence is almost surely equal to a constant belonging to $\{0, 1\}$. Since this constant, in the case of the power series in (3.25), is greater than e^{-y} , we deduce that it equals 1.

Proof. of Proposition 3.2

We fix $y > 0$. As V^\uparrow has no positive jumps and goes to infinity we have $\tau(V^\uparrow, y) \leq \mathcal{R}(V^\uparrow, y) < +\infty$ and $V^\uparrow(\tau(V^\uparrow, y)) = V^\uparrow(\mathcal{R}(V^\uparrow, y)) = y$. We write

$$\begin{aligned} I(V^\uparrow) &= \int_0^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt + \int_{\mathcal{R}(V^\uparrow, y)}^{+\infty} e^{-V^\uparrow(t)} dt \\ &= \int_0^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt + e^{-y} \int_0^{+\infty} e^{-(V^\uparrow(t+\mathcal{R}(V^\uparrow, y))-y)} dt \\ &\stackrel{\mathcal{L}}{=} \int_0^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt + e^{-y} I(\tilde{V}^\uparrow). \end{aligned}$$

We have used Lemma 2.5 for the last equality in which \tilde{V}^\uparrow is an independent copy of V^\uparrow . We now decompose

$$\int_0^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt = \int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt + \int_{\tau(V^\uparrow, y)}^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt. \quad (3.26)$$

Since $V^\uparrow(\tau(V^\uparrow, y)) = y$, combining with the Markov property at time $\tau(V^\uparrow, y)$, the two terms in the right hand side of (3.26) are independent :

$$\int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt \perp\!\!\!\perp \int_{\tau(V^\uparrow, y)}^{\mathcal{R}(V^\uparrow, y)} e^{-V^\uparrow(t)} dt.$$

Now, thanks to Lemma 2.1, the second term has the same law as S_T with S_T as in the lemma. This achieves the proof. \square

As a direct consequence of (3.25) we can now prove the remaining part of Theorem 1.1.

Proof. of Theorem 1.1 (1.3)

Let S and T be as in Proposition 3.2 and $\epsilon \in]0, \mathbb{E}[S_1][$ (we can see that $\mathbb{E}[S_1] < +\infty$ but this is not important here). Then we have

$$\mathbb{P}(S_T \geq x) \geq \mathbb{P}(S_{x/\epsilon} \geq x) \times \mathbb{P}(T \geq x/\epsilon).$$

The first factor in the right hand side converges to 1 thanks to the law of large numbers for Lévy processes (see for example Theorem 36.5 in Sato [26]). The second is equal to $e^{-px/\epsilon}$, where p is the parameter of the exponential random variable T . Therefore, we have an exponential lower bound for the right tail of S_T : $\mathbb{P}(S_T \geq t) \geq e^{-Cx}$ for some positive constant C , when x is large enough. We then notice from (3.25) that $I(V^\uparrow)$ is stochastically greater than S_T so we get an exponential lower bound for the right tail of $I(V^\uparrow)$, which proves the lower bound in (1.3). The upper bound in (1.3) follows from (1.2) and Markov's inequality which finishes the proof of Theorem 1.1. \square

Remark 3.4. *It is possible to prove the first part of Theorem 1.1 (the fact that $I(V^\uparrow)$ is finite and admits some finite exponential moments) by invoking (3.25) and proving that each of the two terms composing A^y admit some finite exponential moments.*

4. ASYMPTOTIC TAIL AT 0

First, let us prove that Theorem 1.2 easily implies Theorem 1.4 and prove Remark 1.8.

Proof. of Theorem 1.4

Assume that Theorem 1.2 is proved.

We first prove (1.7). Let us fix $\beta' > \beta$ and $\epsilon > 0$. From the definition of β we have that $\Psi_V(\lambda) \leq \epsilon \lambda^{\beta'}$ for all λ large enough. Using the first point of Theorem 1.2 with $C = \epsilon, \alpha = \beta'$ and $\delta = 1/2$ we deduce that

$$\limsup_{x \rightarrow 0} x^{1/(\beta'-1)} \log \left(\mathbb{P} \left(I(V^\uparrow) \leq x \right) \right) \leq -(\beta' - 1)/2\epsilon^{1/(\beta'-1)}.$$

Since ϵ can be chosen as small as we want we obtain (1.7).

We now assume that $\sigma > 1$ and prove (1.8). Let us fix $\sigma' \in]1, \sigma[$ and $M > 0$. From the definition of σ we have that $\Psi_V(\lambda) \geq M\lambda^{\sigma'}$ for all λ large enough. Using the second point of Theorem 1.2 with $c = M, \alpha = \sigma'$ and $\delta = 2$ we deduce that

$$0 \geq \liminf_{x \rightarrow 0} x^{1/(\sigma'-1)} \log \left(\mathbb{P} \left(I(V^\uparrow) \leq x \right) \right) \geq -2\sigma'^{\sigma'/(\sigma'-1)}/M^{1/(\sigma'-1)}.$$

Since M can be chosen as large as we want we obtain (1.8). \square

Proof. of Remark 1.8

Let us assume that V has bounded variation. As it can be seen from Remark 1.3, it is the difference of a positive drift γ^*t and a pure jump subordinator S_t : $\forall t > 0, V(t) = \gamma^*t - S_t \leq \gamma^*t$. Let us fix $y > 0$, we have almost surely

$$\int_0^{\tau(V,y)} e^{-V(t)} dt \geq \int_0^{\tau(V,y)} e^{-\gamma^*t} dt = \frac{1}{\gamma^*} \left(1 - e^{-\tau(V,y)} \right). \quad (4.27)$$

Since V has bounded variation, we have $\mathbb{P}(V(t) > 0, 0 \leq t \leq \tau(V,y)) > 0$ (see for example (47.1) in [26]) and we can see that $(V^\uparrow(t), 0 \leq t \leq \tau(V^\uparrow,y))$ is equal in law to $(V, 0 \leq t \leq \tau(V,y))$

conditioned in the usual sense to remain positive. Combining with (4.27), we see that almost surely

$$I(V^\uparrow) \geq \int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt \geq \frac{1}{\gamma^*} \left(1 - e^{-\tau(V^\uparrow, y)}\right). \quad (4.28)$$

Then, since $\tau(V^\uparrow, y)$ converges almost surely to $+\infty$ when y goes to $+\infty$, we deduce that $I(V^\uparrow)$ is more than the positive constant $1/\gamma^*$ almost surely.

Reciprocally, the support of S_t contains 0 because it is pure jump. Therefore, for any $\epsilon > 0$ and $y > 0$, with a positive probability, the left hand side in (4.27) can be ϵ -close to the right hand side while V stays positive on $[0, \tau(V^\uparrow, y)]$. Since $(V^\uparrow(t), 0 \leq t \leq \tau(V^\uparrow, y))$ is equal in law to $(V, 0 \leq t \leq \tau(V, y))$ conditioned in the usual sense to remain positive, we deduce that the second term in (4.28) can be ϵ -close to the third term with a positive probability. Then, note that almost surely $\tau(V^\uparrow, y) \geq y/\gamma^*$ (because it holds for $\tau(V, y)$) and that, from the Markov property, the event $\{\int_{\tau(V^\uparrow, y)}^{+\infty} e^{-V^\uparrow(t)} dt < \epsilon\}$ is independent from the event considered in (4.28) and has a positive probability provided y is large enough. As a consequence $I(V^\uparrow)$ can be 3ϵ -close to $1/\gamma^*$ with a positive probability so the support of the distribution of $I(V^\uparrow)$ contains $1/\gamma^*$. Then, from the properties of non-degenerate self-decomposable distributions, we deduce that the support of $I(V^\uparrow)$ is $[1/\gamma^*, +\infty[$.

□

In the next Subsection we exploit Proposition 3.2 to prepare the proofs of Theorems 1.2 and 1.5.

4.1. Laplace transform of $I(V^\uparrow)$. In order to prove asymptotic estimates on $\mathbb{P}(I(V^\uparrow) \leq x)$, we first study the Laplace transform of $I(V^\uparrow)$ via the series decomposition (3.25). It is thus natural that we need first to study the Laplace transform of A^y .

First, let us define a notation. V^\sharp is a spectrally negative Lévy process, so, according to Theorem VII.1 in [3], the process $\tau(V^\sharp, \cdot)$ is a subordinator which Laplace exponent Φ_{V^\sharp} is defined for $\lambda \geq 0$ by

$$\Phi_{V^\sharp}(\lambda) := -\log \left(\mathbb{E} \left[e^{-\lambda \tau(V^\sharp, 1)} \right] \right).$$

Moreover, we have $\Phi_{V^\sharp} = \Psi_{V^\sharp}^{-1}$.

Proposition 4.1. *We fix $y > 0$. Let A^y be as in Proposition 3.2, then, for all $\epsilon > 0$ and λ large enough we have*

$$(1 - \epsilon)y\Phi_{V^\sharp}(e^{-y}\lambda) \leq -\log \left(\mathbb{E} \left[e^{-\lambda A^y} \right] \right) \leq (1 + \epsilon)y\Phi_{V^\sharp}(\lambda). \quad (4.29)$$

Proof. According to the definition of A^y in Proposition 3.2, A^y can be decomposed as the sum of two independent random variables, one having the same law as $\int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt$ and another having the same law as S_T , defined as in Lemma 2.1. Let Φ_S be the Laplace exponent of the subordinator S :

$$\forall \lambda \geq 0, \Phi_S(\lambda) := -\log \left(\mathbb{E} \left[e^{-\lambda S_1} \right] \right). \quad (4.30)$$

We can see that the Laplace transform of the random variable S_T is given by

$$\forall \lambda \geq 0, \mathbb{E} \left[e^{-\lambda S_T} \right] = \frac{\eta_y^\sharp(IP_\infty) + \eta_y^\sharp(N_\infty)}{\eta_y^\sharp(IP_\infty) + \eta_y^\sharp(N_\infty) + \Phi_S(\lambda)}.$$

We thus have

$$\begin{aligned} -\log\left(\mathbb{E}\left[e^{-\lambda A^y}\right]\right) &= -\log\left(\mathbb{E}\left[\exp\left(-\lambda\int_0^{\tau(V^\uparrow,y)}e^{-V^\uparrow(t)}dt\right)\right]\right) - \log\left(\mathbb{E}\left[e^{-\lambda S_T}\right]\right) \\ &= -\log\left(\mathbb{E}\left[\exp\left(-\lambda\int_0^{\tau(V^\uparrow,y)}e^{-V^\uparrow(t)}dt\right)\right]\right) - \log\left(\frac{p}{p+\Phi_S(\lambda)}\right), \end{aligned} \quad (4.31)$$

where we denoted p for the constant $\eta_y^\sharp(IP_\infty) + \eta_y^\sharp(N_\infty)$. Using the fact that V^\uparrow is non-negative and the first point of Lemma 2.6 we have

$$\int_0^{\tau(V^\uparrow,y)}e^{-V^\uparrow(t)}dt \leq \tau(V^\uparrow,y) \stackrel{sto}{\leq} \tau(V^\sharp,y), \quad (4.32)$$

where $\stackrel{sto}{\leq}$ denotes a stochastic inequality. As a consequence of (4.32) and of the definition of Φ_{V^\sharp} we have

$$-\log\left(\mathbb{E}\left[\exp\left(-\lambda\int_0^{\tau(V^\uparrow,y)}e^{-V^\uparrow(t)}dt\right)\right]\right) \leq -\log\left(\mathbb{E}\left[e^{-\lambda\tau(V^\sharp,y)}\right]\right) = y\Phi_{V^\sharp}(\lambda).$$

Combining this inequality with (4.31) we obtain

$$-\log\left(\mathbb{E}\left[e^{-\lambda A^y}\right]\right) \leq y\Phi_{V^\sharp}(\lambda) - \log\left(\frac{p}{p+\Phi_S(\lambda)}\right). \quad (4.33)$$

Using the first point of Lemma 2.6 and Lemma 2.3 we have

$$-\log\left(\mathbb{E}\left[\exp\left(-\lambda\int_0^{\tau(V^\uparrow,y)}e^{-V^\uparrow(t)}dt\right)\right]\right) = -\log\left(\mathbb{E}\left[e^{-\lambda\int_0^{\tau(V^\sharp,y)}e^{-V^\sharp(t)}dt\right] / \mathbb{E}\left[e^{-\lambda\tilde{S}_T}\right]\right),$$

where \tilde{S}_T is as S_T from Lemma 2.3. Here again, if $\Phi_{\tilde{S}}$ denotes the Laplace exponent of the subordinator \tilde{S} as in (4.30) we have

$$\forall \lambda \geq 0, \mathbb{E}\left[e^{-\lambda\tilde{S}_T}\right] = \frac{\eta^\sharp(I_y)}{\eta^\sharp(I_y) + \Phi_{\tilde{S}}(\lambda)}.$$

Moreover we have

$$\int_0^{\tau(V^\sharp,y)}e^{-V^\sharp(t)}dt \geq e^{-y}\tau(V^\sharp,y).$$

Putting together the above three expressions and the definition of Φ_{V^\sharp} we obtain

$$-\log\left(\mathbb{E}\left[\exp\left(-\lambda\int_0^{\tau(V^\uparrow,y)}e^{-V^\uparrow(t)}dt\right)\right]\right) \geq y\Phi_{V^\sharp}(e^{-y}\lambda) + \log\left(\frac{\eta^\sharp(I_y)}{\eta^\sharp(I_y) + \Phi_{\tilde{S}}(\lambda)}\right).$$

Combining the above inequality with (4.31) and the fact that the term $-\log(p/(p+\Phi_S(\lambda)))$ is non-negative we get

$$-\log\left(\mathbb{E}\left[e^{-\lambda A^y}\right]\right) \geq y\Phi_{V^\sharp}(e^{-y}\lambda) + \log\left(\frac{\eta^\sharp(I_y)}{\eta^\sharp(I_y) + \Phi_{\tilde{S}}(\lambda)}\right). \quad (4.34)$$

According to the Lévy-Khintchine formula applied to the pure jump subordinator S , the Laplace exponent Φ_S can be written

$$\Phi_S(\lambda) = \int_0^{+\infty}(1 - e^{-\lambda x})\nu_S(dx).$$

By dominated convergence $\Phi_S(\lambda)/\lambda$ converges to 0 when λ goes to $+\infty$. As a consequence we have $\Phi_S(\lambda) \leq \lambda$ for large λ . Similarly we have $\Phi_{\tilde{S}}(\lambda) \leq \lambda$ for large λ . On the other hand, since $\Psi_{V^\sharp}(\lambda)/\lambda^2$ is bounded when λ goes to infinity, there is a positive constant c such that

$\Phi_{V^\#}(\lambda) \geq c\lambda^{1/2}$ for large λ . Combining all this with (4.33) and (4.34) we get (4.29) for any fixed $\epsilon > 0$ and λ large enough. \square

Proposition 4.2. *Assume that there is $\alpha \geq 1$ and a positive constant C such that for all λ large enough we have $\Psi_V(\lambda) \leq C\lambda^\alpha$, then we have*

$$\liminf_{\lambda \rightarrow +\infty} -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) / \lambda^{1/\alpha} \geq \alpha / C^{1/\alpha}. \quad (4.35)$$

Assume that there is $\alpha \geq 1$ and a positive constant c such that for all λ large enough we have $\Psi_V(\lambda) \geq c\lambda^\alpha$, then we have

$$\limsup_{\lambda \rightarrow +\infty} -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) / \lambda^{1/\alpha} \leq \alpha / c^{1/\alpha}. \quad (4.36)$$

Proof. Let us fix $y > 0$ for which we apply the decomposition (3.25). Let us denote

$$\mathcal{M}(\lambda) := -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) = \sum_{k \geq 0} -\log \left(\mathbb{E} \left[e^{-\lambda e^{-ky} A^y} \right] \right),$$

where the second equality comes from (3.25) and from the fact that the sequence $(A_k^y)_{k \geq 0}$ is *iid*.

The asymptotic behavior of $\mathcal{M}(\lambda)$ is thus related to the asymptotic behavior of $-\log(\mathbb{E}[e^{-\lambda A^y}])$. Unfortunately, Proposition 4.1 can not be applied simultaneously to all the terms of the sum defining $\mathcal{M}(\lambda)$. We separate this sum into three parts : a sum over a finite number of small indices for which we can apply (4.29) to each term, a sum over an infinite number of large indices that can be neglected, and a sum over the remaining indices (in finite number) that can be neglected too.

We now prove the second point of the proposition. We assume that there is $\alpha \geq 1$ and a positive constant c such that for all λ large enough we have $\Psi_V(\lambda) \geq c\lambda^\alpha$, and we prove (4.36). Let us fix $\delta > 1$. Since $\Psi_{V^\#}(\cdot) = \Psi_V(\kappa + \cdot)$ and $\Phi_{V^\#} = \Psi_{V^\#}^{-1}$, we have for all λ large enough that $\Phi_{V^\#}(\lambda) \leq \delta\lambda^{1/\alpha}/c^{1/\alpha}$.

According to Proposition 4.1 there exists $\lambda_\delta > 1$ such that (4.29) is satisfied with $\epsilon = \delta - 1$ for all $\lambda \geq \lambda_\delta$. By increasing λ_δ if necessary, we can also assume that $\Phi_{V^\#}(\lambda) \leq \delta\lambda^{1/\alpha}/c^{1/\alpha}$ for all $\lambda \geq \lambda_\delta$. Putting all this together we get

$$\lambda \geq \lambda_\delta \Rightarrow -\log \left(\mathbb{E} \left[e^{-\lambda A^y} \right] \right) \leq \delta^2 y \lambda^{1/\alpha} / c^{1/\alpha}. \quad (4.37)$$

Also, let us choose $M \in]0, 1[$ small enough so that

$$\forall \lambda \in [0, M], \quad 0 \leq -\log \left(\mathbb{E} \left[e^{-\lambda A^y} \right] \right) \leq 2\lambda \mathbb{E} [A^y]. \quad (4.38)$$

For any $\lambda > \lambda_\delta$ we define $n_1(\lambda) := \lfloor \log(\lambda/\lambda_\delta)/y \rfloor - 1$ and $n_2(\lambda) := \lfloor \log(\lambda/M)/y \rfloor$. From the definition of $\mathcal{M}(\lambda)$ we can write

$$\forall \lambda > \lambda_\delta, \quad \mathcal{M}(\lambda) = T_1(\lambda) + T_2(\lambda) + T_3(\lambda), \quad (4.39)$$

with

$$\begin{aligned} T_1(\lambda) &:= \sum_{k=0}^{n_1(\lambda)} -\log \left(\mathbb{E} \left[e^{-\lambda e^{-ky} A^y} \right] \right), \quad T_2(\lambda) := \sum_{k=n_1(\lambda)+1}^{n_2(\lambda)} -\log \left(\mathbb{E} \left[e^{-\lambda e^{-ky} A^y} \right] \right), \\ T_3(\lambda) &:= \sum_{k=n_2(\lambda)+1}^{+\infty} -\log \left(\mathbb{E} \left[e^{-\lambda e^{-ky} A^y} \right] \right). \end{aligned} \quad (4.40)$$

From the definition of $n_1(\lambda)$, (4.37) can be applied to each term of the sum defining $T_1(\lambda)$, we thus have

$$T_1(\lambda) \leq \delta^2 y (\lambda/c)^{1/\alpha} \sum_{k=0}^{n_1(\lambda)} e^{-ky/\alpha} = \delta^2 y (\lambda/c)^{1/\alpha} \frac{1 - e^{-y(n_1(\lambda)+1)/\alpha}}{1 - e^{-y/\alpha}}.$$

We get that for λ large enough

$$T_1(\lambda) \leq \delta^2 y (\lambda/c)^{1/\alpha} \frac{1}{1 - e^{-y/\alpha}}. \quad (4.41)$$

Using the monotony of the Laplace transform and the definitions of $n_1(\lambda)$ and $n_2(\lambda)$ we get

$$\begin{aligned} 0 \leq T_2(\lambda) &\leq -(n_2(\lambda) - n_1(\lambda)) \log \left(\mathbb{E} \left[e^{-\lambda e^{-(n_1(\lambda)+1)y} A^y} \right] \right) \\ &\leq -\frac{2y + \log(\lambda_\delta/M)}{y} \log \left(\mathbb{E} \left[e^{-\lambda_\delta e^{-1} A^y} \right] \right) < +\infty. \end{aligned} \quad (4.42)$$

From the definitions of $n_2(\lambda)$ and M we have

$$\begin{aligned} 0 \leq T_3(\lambda) &\leq 2\lambda \mathbb{E}[A^y] \sum_{k=n_2(\lambda)+1}^{+\infty} e^{-ky} = 2\lambda e^{-y(n_2(\lambda)+1)} \mathbb{E}[A^y] / (1 - e^{-y}) \\ &\leq 2M \mathbb{E}[A^y] / (1 - e^{-y}) < +\infty. \end{aligned} \quad (4.43)$$

Putting (4.41), (4.42), and (4.43) into (4.39) we obtain that for λ large enough

$$\limsup_{\lambda \rightarrow +\infty} \mathcal{M}(\lambda) / \lambda^{1/\alpha} \leq \frac{\delta^2 y}{c^{1/\alpha} (1 - e^{-y/\alpha})}.$$

Since, from the definition $\mathcal{M}(\lambda) = -\log(\mathbb{E}[e^{-\lambda I(V^\dagger)}])$ which does not depend on δ nor on y , we can let δ go to 1 and then y go to 0 in the above expression. We obtain

$$\limsup_{\lambda \rightarrow +\infty} -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) / \lambda^{1/\alpha} \leq \alpha / c^{1/\alpha}.$$

For the first point of the proposition, we proceed exactly as we just did for the second point, using the lower bound of (4.29) instead of the upper bound. In particular $\sum_{k=0}^{n_1(\lambda)} e^{-ky/\alpha}$ from the above argument is replaced by $\sum_{k=1}^{n_1(\lambda)+1} e^{-ky/\alpha}$. We finally get, for $y > 0$, $\delta \in]0, 1[$ and λ large enough,

$$T_1(\lambda) \geq \delta^2 y (\lambda/C)^{1/\alpha} e^{-y} \frac{1 - e^{-(n_1(\lambda)+1)y/\alpha}}{1 - e^{-y/\alpha}}.$$

Then, since $n_1(\lambda)$ converges toward $+\infty$ as λ goes to infinity, we get $\liminf_{\lambda \rightarrow +\infty} \mathcal{M}(\lambda) / \lambda^{1/\alpha} \geq \delta^2 y (\lambda/C)^{1/\alpha} e^{-y} / (1 - e^{-y/\alpha})$. Letting δ go to 1 and then y go to 0 we obtain

$$\liminf_{\lambda \rightarrow +\infty} -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) / \lambda^{1/\alpha} \geq \alpha / C^\alpha,$$

which terminates the proof. □

We also study what happens when Ψ_V has regular variation in the next proposition :

Proposition 4.3. *Assume that Ψ_V has α -regular variation at $+\infty$ for some $\alpha \in]1, 2]$. Then,*

$$-\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) \underset{\lambda \rightarrow +\infty}{\sim} \alpha \Phi_{V^\#}(\lambda). \quad (4.44)$$

Proof. The idea is to modify a little the proof of Proposition 4.2. Let us fix $y > 0$ and $\epsilon \in]0, 1/\alpha[$. We keep the notation

$$\mathcal{M}(\lambda) := -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) = \sum_{k \geq 0} -\log \left(\mathbb{E} \left[e^{-\lambda e^{-ky} A^y} \right] \right).$$

Since Ψ_V has α -regular variation, $\Phi_{V^\#}$ has $1/\alpha$ -regular variation so, thanks to Karamata representation theorem we can write

$$\Phi_{V^\#}(\lambda) = \lambda^{\frac{1}{\alpha}} \exp \left(c(\lambda) + \int_1^\lambda \frac{r(t)}{t} dt \right). \quad (4.45)$$

In this representation c is a bounded measurable function that converges to a constant c at $+\infty$, and r is a bounded measurable function that converges to 0 at $+\infty$. Let us chose $\lambda_\epsilon > 1$ large enough such that for all $\lambda \geq \lambda_\epsilon$: (4.29) is satisfied, $c - \epsilon \leq c(\lambda) \leq c + \epsilon$, and $|r(\lambda)| \leq \epsilon$. We also fix M as in (4.38) and for any $\lambda > \lambda_\epsilon$ we define $n_1(\lambda) := \lfloor \log(\lambda/\lambda_\delta)/y \rfloor - 1$ and $n_2(\lambda) := \lfloor \log(\lambda/M)/y \rfloor$. For $\lambda \geq \lambda_\epsilon$, we then still have $\mathcal{M}(\lambda) = T_1(\lambda) + T_2(\lambda) + T_3(\lambda)$ with $T_1(\lambda), T_2(\lambda)$ and $T_3(\lambda)$ as in (4.40). Using (4.29) in each term of the sum defining $T_1(\lambda)$, and then the representation (4.45) we get

$$\begin{aligned} \frac{T_1(\lambda)}{\Phi_{V^\#}(\lambda)} &\leq (1 + \epsilon)y \sum_{k=0}^{n_1(\lambda)} \frac{\Phi_{V^\#}(e^{-ky}\lambda)}{\Phi_{V^\#}(\lambda)} \\ &= (1 + \epsilon)y \sum_{k=0}^{n_1(\lambda)} \exp \left(-\frac{ky}{\alpha} + c(e^{-ky}\lambda) - c(\lambda) - \int_{e^{-ky}\lambda}^\lambda \frac{r(t)}{t} dt \right). \end{aligned}$$

For $k \leq n_1(\lambda)$, we have $e^{-ky}\lambda \geq \lambda_\epsilon$ so, according to the choice of λ_ϵ , we have $-2\epsilon \leq c(e^{-ky}\lambda) - c(\lambda) \leq 2\epsilon$ and $\epsilon \log(\lambda/e^{-ky}\lambda) \geq \int_{e^{-ky}\lambda}^\lambda \frac{r(t)}{t} dt \geq -\epsilon \log(\lambda/e^{-ky}\lambda) = -\epsilon ky$. As a consequence,

$$\begin{aligned} \frac{T_1(\lambda)}{\Phi_{V^\#}(\lambda)} &\leq (1 + \epsilon)y e^{2\epsilon} \sum_{k=0}^{n_1(\lambda)} \exp \left(-k \left(\frac{y}{\alpha} - \epsilon y \right) \right) \\ &= (1 + \epsilon)y e^{2\epsilon} \frac{1 - e^{-(n_1(\lambda)+1)(y/\alpha - \epsilon y)}}{1 - e^{-y/\alpha + \epsilon y}} \leq \frac{(1 + \epsilon)y e^{2\epsilon}}{1 - e^{-y/\alpha + \epsilon y}}. \end{aligned}$$

Then, $T_2(\lambda)$ and $T_3(\lambda)$ are bounded as in the proof of Proposition 4.2 so we get

$$\limsup_{\lambda \rightarrow +\infty} \mathcal{M}(\lambda)/\Phi_{V^\#}(\lambda) \leq \frac{(1 + \epsilon)y e^{2\epsilon}}{1 - e^{-y/\alpha + \epsilon y}}.$$

Here again, the left hand side does not depend on ϵ nor on y so we can let ϵ go to 0 and then y go to 0 in the above expression. We obtain

$$\limsup_{\lambda \rightarrow +\infty} -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) / \Phi_{V^\#}(\lambda) \leq \alpha.$$

To prove a lower bound, we proceed exactly as we just did, using the lower bound of (4.29). In particular $\sum_{k=0}^{n_1(\lambda)} \Phi_{V^\#}(e^{-ky}\lambda)/\Phi_{V^\#}(\lambda)$ from the above argument is replace by

$\sum_{k=1}^{n_1(\lambda)+1} \Phi_{V^\#}(e^{-ky}\lambda)/\Phi_{V^\#}(\lambda)$. We finally get

$$\frac{T_1(\lambda)}{\Phi_{V^\#}(\lambda)} \geq (1 - \epsilon)y e^{-2\epsilon - y} \frac{1 - e^{-(n_1(\lambda)+1)(y/\alpha + \epsilon y)}}{1 - e^{-y/\alpha - \epsilon y}}.$$

Then, since $n_1(\lambda)$ converges toward $+\infty$ as λ goes to infinity, we get $\liminf_{\lambda \rightarrow +\infty} \mathcal{M}(\lambda)/\Phi_{V^\#}(\lambda) \geq (1 - \epsilon)y e^{-2\epsilon - y}/(1 - e^{-y/\alpha - \epsilon y})$. Letting ϵ go to 0 and then y go to 0 we obtain

$$\liminf_{\lambda \rightarrow +\infty} -\log \left(\mathbb{E} \left[e^{-\lambda I(V^\dagger)} \right] \right) / \Phi_{V^\#}(\lambda) \geq \alpha,$$

which terminates the proof. \square

4.2. Tail at 0 of $I(V^\uparrow)$: proof of Theorems 1.2 and 1.5. We now prove Theorems 1.2 and 1.5 by using Propositions 4.2 and 4.3 together with the deep link that exists between the left tail of a random variable and the asymptotic behavior of its Laplace transform.

Proof. of Theorem 1.2

We assume that there is $\alpha > 1$ and a positive constant C such that for all λ large enough we have $\Psi_V(\lambda) \leq C\lambda^\alpha$. We fix $\delta \in]0, 1[$. According to the first point of Proposition 4.2, there exists $\lambda_\delta > 0$ such that for all $\lambda > \lambda_\delta$ we have

$$\mathbb{E} \left[e^{-\lambda I(V^\uparrow)} \right] \leq \exp \left(-\delta^{1-1/\alpha} \alpha \lambda^{1/\alpha} / C^{1/\alpha} \right). \quad (4.46)$$

Let us fix $x \in]0, \delta^{(\alpha-1)/\alpha} C^{-1/\alpha} \lambda_\delta^{(1-\alpha)/\alpha}[$. Using Markov's inequality and (4.46) we get that for any $\lambda > \lambda_\delta$,

$$\mathbb{P} \left(I(V^\uparrow) \leq x \right) \leq e^{\lambda x} \mathbb{E} \left[e^{-\lambda I(V^\uparrow)} \right] \leq \exp \left(\lambda x - \delta^{1-1/\alpha} \alpha \lambda^{1/\alpha} / C^{1/\alpha} \right).$$

Since $x \in]0, \delta^{(\alpha-1)/\alpha} C^{-1/\alpha} \lambda_\delta^{(1-\alpha)/\alpha}[$ we have $\delta C^{-1/(\alpha-1)} x^{-\alpha/(\alpha-1)} \geq \lambda_\delta$ so, in the above inequality, we can replace λ by $\delta C^{-1/(\alpha-1)} x^{-\alpha/(\alpha-1)}$. We obtain

$$\mathbb{P} \left(I(V^\uparrow) \leq x \right) \leq \exp \left(-\delta(\alpha-1)/(Cx)^{1/(\alpha-1)} \right),$$

which is (1.5).

We now prove the second point. Assume that there is $\alpha > 1$ and a positive constant c such that for all λ large enough we have $\Psi_V(\lambda) \geq c\lambda^\alpha$. We fix $\delta > 1$ and $r \in]0, 1[$. According to the second point of Proposition 4.2, there exists $\lambda_\delta > 0$ such that for all $\lambda > \lambda_\delta$ we have

$$\mathbb{E} \left[e^{-\lambda I(V^\uparrow)} \right] \geq \exp \left(-\delta^{r(\alpha-1)/\alpha} \alpha \lambda^{1/\alpha} / c^{1/\alpha} \right). \quad (4.47)$$

Let us fix $x \in]0, \delta^{(\alpha-1)/\alpha} \alpha c^{-1/\alpha} \lambda_\delta^{(1-\alpha)/\alpha}[$. Using (4.47) we get that for any $\lambda > \lambda_\delta$,

$$\begin{aligned} \exp \left(-\delta^{r(\alpha-1)/\alpha} \alpha \lambda^{1/\alpha} / c^{1/\alpha} \right) &\leq \mathbb{E} \left[e^{-\lambda I(V^\uparrow)} \right] = \mathbb{E} \left[e^{-\lambda I(V^\uparrow)} \mathbf{1}_{\{I(V^\uparrow) \leq x\}} \right] + \mathbb{E} \left[e^{-\lambda I(V^\uparrow)} \mathbf{1}_{\{I(V^\uparrow) > x\}} \right] \\ &\leq \mathbb{P} \left(I(V^\uparrow) \leq x \right) + \exp(-\lambda x), \end{aligned}$$

so we get

$$\mathbb{P} \left(I(V^\uparrow) \leq x \right) \geq \exp \left(-\delta^{r(\alpha-1)/\alpha} \alpha \lambda^{1/\alpha} / c^{1/\alpha} \right) - \exp(-\lambda x).$$

Since $x \in]0, \delta^{(\alpha-1)/\alpha} \alpha c^{-1/\alpha} \lambda_\delta^{(1-\alpha)/\alpha}[$ we have $\delta \alpha^{\alpha/(\alpha-1)} c^{-1/(\alpha-1)} x^{-\alpha/(\alpha-1)} \geq \lambda_\delta$ so, in the above inequality, we can replace λ by $\delta \alpha^{\alpha/(\alpha-1)} c^{-1/(\alpha-1)} x^{-\alpha/(\alpha-1)}$. We obtain

$$\mathbb{P} \left(I(V^\uparrow) \leq x \right) \geq \exp \left(-\delta^{(1+r(\alpha-1))/\alpha} \alpha^{\alpha/(\alpha-1)} / (cx)^{1/(\alpha-1)} \right) - \exp \left(-\delta \alpha^{\alpha/(\alpha-1)} / (cx)^{1/(\alpha-1)} \right).$$

Since $\delta > \delta^{(1+r(\alpha-1))/\alpha}$, the second term converges to 0 faster than the first one when x goes to 0. We thus get, for x small enough,

$$\mathbb{P} \left(I(V^\uparrow) \leq x \right) \geq \frac{1}{2} \exp \left(-\delta^{(1+r(\alpha-1))/\alpha} \alpha^{\alpha/(\alpha-1)} / (cx)^{1/(\alpha-1)} \right) \geq \exp \left(-\delta \alpha^{\alpha/(\alpha-1)} / (cx)^{1/(\alpha-1)} \right).$$

The last inequality is true for x small enough and comes from the fact that $\delta > \delta^{(1+r(\alpha-1))/\alpha}$. The above is precisely (1.6). \square

Proof. of Theorem 1.5

We assume that Ψ_V has α -regular variation at $+\infty$ for some $\alpha \in]1, 2]$. From Proposition 4.3 we deduce that

$$-\log \left(\mathbb{E} \left[e^{-\lambda I(V^\uparrow)} \right] \right) \underset{\lambda \rightarrow +\infty}{\sim} \alpha \Phi_{V^\#}(\lambda).$$

Then, since $\Phi_{V^\#}$ is the inverse function of $\Psi_V(\kappa + \cdot)$, it has $1/\alpha$ -regular variation at $+\infty$ and the application of De Bruijn's Theorem (see Theorem 4.12.9 in [6]) yields the result. \square

4.3. Connection between $I(V^\uparrow)$ and $I(V^\#)$: Proof of Propositions 1.10 and 1.11. In this subsection, we assume that V does not oscillate (so that $I(V^\#) < +\infty$) and we relate $I(V^\uparrow)$ and $I(V^\#)$. This allows to prove Proposition 1.10 and to combine it with Theorem 5.24 of [22] to prove Proposition 1.11. Note that in particular, when V drifts to $+\infty$, this is only a comparison between $I(V^\uparrow)$ and $I(V)$.

Proposition 4.4. *If V does not oscillates we have*

$$I(V^\#) \stackrel{\mathcal{L}}{=} S_T + I(V^\uparrow), \quad (4.48)$$

where the two terms of the right hand side are independent, and S_T is as in Lemma 2.4.

Thanks to Proposition 4.4, we can explain heuristically why the functionals $I(V^\uparrow)$ and $I(V^\#)$ have similar left tails but different right tails :

Remark 4.5. *In (4.48), the term S_T contains the contributions of the excursions of $V^\#$ that meet the half-line of negative real numbers, it is thus heavy tailed. On the other hand, the right tail of $I(V^\uparrow)$ is given by Theorem 1.1 and is at most exponential. This is why only the contribution of the term S_T is relevant regarding the right tail of $I(V^\#)$ and the two functionals have different right tails. For the left tails, the one of S_T is at least the left tail of an exponential random variable (see Lemma 4.6 below) while the left tail of $I(V^\uparrow)$ is lighter according to our preceding results. This is why the left tail of $I(V^\#)$ is mainly determined by the term $I(V^\uparrow)$ in (4.48).*

Proof. of Proposition 4.4

We write

$$\begin{aligned} I(V^\#) &= \int_0^{\mathcal{R}(V^\#, 0)} e^{-V^\#(t)} dt + \int_{\mathcal{R}(V^\#, 0)}^{+\infty} e^{-V^\#(t)} dt \\ &= \int_0^{\mathcal{R}(V^\#, 0)} e^{-V^\#(t)} dt + \int_0^{+\infty} e^{-V^\#(t + \mathcal{R}(V^\#, 0))} dt \\ &\stackrel{\mathcal{L}}{=} \int_0^{\mathcal{R}(V^\#, 0)} e^{-V^\#(t)} dt + I(V^\uparrow). \end{aligned} \quad (4.49)$$

We have used Lemma 2.6 for the last equality in which the two terms $\int_0^{\mathcal{R}(V^\#, 0)} e^{-V^\#(t)} dt$ and $I(V^\uparrow)$ are independent.

Now, thanks to Lemma 2.4, the term $\int_0^{\mathcal{R}(V^\#, 0)} e^{-V^\#(t)} dt$ has the same law as S_T with S_T as in the lemma. This achieves the proof. \square

Similarly to the proof of Proposition 3.2, we decomposed $I(V^\#)$ as the sum of two independent random variables, one having the same law as a subordinator stopped at an independent

exponential time and the other having the same law as $I(V^\uparrow)$. We now need a lemma about the asymptotic tail at 0 of a subordinator stopped at an independent exponential time.

Lemma 4.6. *Let S be a subordinator and T an independent exponential random variable, there exists a positive constant c such that for all x small enough*

$$\mathbb{P}(S_T < x) \geq cx.$$

Proof. We prove in fact a stronger result : the function $x \mapsto \mathbb{P}(S_T < x)$ is sub-additive, that is,

$$\forall x, y \geq 0, \mathbb{P}(S_T < x + y) \leq \mathbb{P}(S_T < x) + \mathbb{P}(S_T < y). \quad (4.50)$$

The lemma follows easily once (4.50) is proved. Recall from the introduction the notation $\tau(S, h+)$ for $\tau(S, [h, +\infty[)$. Let $x, y > 0$ (the case where $x = 0$ or $y = 0$ is obvious), we have

$$\begin{aligned} \mathbb{P}(S_T < x + y) &= \mathbb{P}(S_T < x) + \mathbb{P}(T \geq \tau(S, x+), T < \tau(S, (x + y)+)) \\ &\leq \mathbb{P}(S_T < x) + \mathbb{P}(T \geq \tau(S, x+), T < \tau(S, (S_{\tau(S, x+)} + y)+)). \end{aligned}$$

For the inequality we have used that $S_{\tau(S, x+)} \geq x$ almost surely. From the characteristic property of the exponential distribution and the Markov property applied to S at time $\tau(S, x+)$, the latter is equal to

$$= \mathbb{P}(S_T < x) + \mathbb{P}(T \geq \tau(S, x+)) \times \mathbb{P}(T < \tau(S, y+)).$$

Since $\mathbb{P}(T \geq \tau(S, x+)) \leq 1$ and $\mathbb{P}(T < \tau(S, y+)) = \mathbb{P}(S_T < y)$ we obtain (4.50). \square

We can now prove Proposition 1.10.

Proof. of Proposition 1.10.

We are in the case where V drifts to $+\infty$ so $V^\# = V$. Applying Proposition 4.4 we get

$$\mathbb{P}(I(V) \leq x) \leq \mathbb{P}(I(V^\uparrow) \leq x). \quad (4.51)$$

For $\epsilon > 0$, using the equality in law of Proposition 4.4 and Lemma 4.6 we obtain

$$\mathbb{P}(I(V) \leq (1 + \epsilon)x) \geq \mathbb{P}(I(V^\uparrow) \leq x) \times \mathbb{P}(S_T \leq \epsilon x) \geq c\epsilon x \mathbb{P}(I(V^\uparrow) \leq x),$$

for an appropriate constant $c > 0$ and x small enough. Combining with (4.51) we get the result. \square

Proof. of Proposition 1.11

If V does not oscillate and has unbounded variation, $V^\#$ drifts to $+\infty$, has Laplace exponent $\Psi_{V^\#} = \Psi_V(\kappa + \cdot)$, and $(V^\#)^\uparrow = V^\uparrow$. Note that, since $I(V^\#)$ is self-decomposable, it is unimodale. Then, the mode is not 0 (because of the continuity of the density or because of the argument of Remark 1.9 which is valid since Remark 1.7 is true for $I(V^\#)$). As a consequence, the density of $I(V^\#)$ is non-decreasing on a neighborhood of 0. Let us denote by $m^\#$ this density so that for any $x \geq 0$, $\mathbb{P}(I(V^\#) \leq x) = \int_0^x m^\#(z) dz$. Let us fix $\delta > 1$. The combination of Proposition 1.10 (applied to $V^\#$ and $\epsilon = \delta - 1$) with the non-decrease of $m^\#$ gives for all x small enough

$$\mathbb{P}(I(V^\uparrow) \leq x) \geq \mathbb{P}(I(V^\#) \leq x) \geq \int_{x/\delta}^x m^\#(z) dz \geq \left(1 - \frac{1}{\delta}\right) x m^\#\left(\frac{x}{\delta}\right),$$

and

$$\mathbb{P}(I(V^\uparrow) \leq x) \leq \frac{1}{c(\delta - 1)x} \mathbb{P}(I(V^\#) \leq \delta x) = \frac{1}{c(\delta - 1)x} \int_0^{\delta x} m^\#(z) dz \leq \frac{\delta}{c(\delta - 1)} m^\#(\delta x).$$

The constant c above is as in Proposition 1.10. Applying Theorem 5.24 of [22] to $m^\sharp(x/\delta)$ and $m^\sharp(\delta x)$ and combining with the above estimates we get the result. \square

Remark 4.7. *In Theorem 5.24 of [22] is given an equivalent for the left tail of $I(V)$ (not an inequality). However, it would not be possible to state this estimate for the left tail of $I(V^\uparrow)$, since the convolution factor S_T in Proposition 4.4 changes the equivalent of the left tail. This is why Propositions 4.4 and 1.10 only allow to transfer inequalities. Then, if one wants to deduce from Theorem 5.24 of [22] an equivalent for the log of the left tail of $I(V^\uparrow)$, one has to be aware that in Proposition 4.4, the upper bound for $\mathbb{P}(I(V^\uparrow) \leq x)$ is given in term of $\mathbb{P}(I(V) \leq (1+\epsilon)x)$ and not $\mathbb{P}(I(V) \leq x)$. Due to this, the asymptotic of the log of the left tail of $I(V)$ can be transferred to $I(V^\uparrow)$ only in the case of regular variation, but this case is already treated by Theorem 1.5 in full generality.*

5. SMOOTHNESS OF THE DENSITY, PROOF OF THEOREM 1.12

$I(V^\uparrow)$ is positive and infinitely divisible so it has no brownian component, its Lévy measure does not charge the negative half-line and has a structure of bounded variation. As a consequence, in the terminology of [27], $I(V^\uparrow)$ is of type I and $\int_0^1 k(u)du < +\infty$.

We assume that V has unbounded variation. Let γ_0 be as in the terminology of [27]. We first prove that $\gamma_0 = 0$ which is equivalent to proving that the support of $I(V^\uparrow)$ contains 0. Let us apply Proposition 3.2 for some $y > 0$. In view of (3.25) and of the definition of A^y , we only need to prove that both the support of $\int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt$ and the support of S_T contain 0.

First, note that, since V (and therefore V^\sharp) has unbounded variation, Corollary VII.5 in [3] yields that $\Psi_{V^\sharp}(\lambda)/\lambda \xrightarrow{\lambda \rightarrow +\infty} +\infty$ so $\Phi_{V^\sharp}(\lambda)/\lambda \xrightarrow{\lambda \rightarrow +\infty} 0$. Φ_{V^\sharp} is the Laplace exponent of the subordinator $\tau(V^\sharp, \cdot)$ which has, therefore, no drift component. As a consequence 0 belongs to the support of $\tau(V^\sharp, y)$ and, because of (4.32), to the support of $\int_0^{\tau(V^\uparrow, y)} e^{-V^\uparrow(t)} dt$.

Then, Lemma 4.6 proves that the distribution of S_T charges every neighborhood of 0 so, by definition, the support of S_T contains 0. We thus have $\gamma_0 = 0$.

If $I(V^\uparrow)$ was of type I_1, I_2, I_3, I_4 or I_5 , we could apply Theorem 1.6 of [27] with $n = 0$ and replace γ_0 by 0 in there. The estimate obtained would be in conflict with Remarks 1.9 and 1.7 which guarantee the existence of a constant K such that the density of $I(V^\uparrow)$ is bounded by $e^{-K/x}$ for x small enough. We thus deduce that $I(V^\uparrow)$ is not of type I_1, I_2, I_3, I_4 nor I_5 . Since it is neither of type I_7 (because $\int_0^1 k(u)du < +\infty$), we get that $I(V^\uparrow)$ is of type I_6 .

The fact that the density of $I(V^\uparrow)$ belongs to the Schwartz space is then a consequence of the type I_6 and of the following Lemma 5.1. The derivatives of all orders of this density converge to 0 when x goes to 0 since this density is of class \mathcal{C}^∞ and null on $] -\infty, 0[$.

The log-concavity of $x \mapsto \mathbb{P}(I(V^\uparrow) \leq x)$ on $]0, +\infty[$ is a consequence of the type I_6 and of Lemma 5.2 in [27].

It only remains to state and justify Lemma 5.1. Our argument to justify it extends a little the proof of Theorem 28.4 of [26].

Lemma 5.1. *Let X be a self-decomposable distribution of type I_6 or I_7 , in the terminology of [27]. If X admits finite moments of any positive order, then its density belongs to the Schwartz space.*

Proof. Since X is self-decomposable of type I6, its characteristic function can be represented as

$$\mathbb{E} \left[e^{i\xi X} \right] = \exp \left(\int_0^{+\infty} (e^{i\xi z} - 1) \frac{k(z)}{z} dz \right) =: \varphi_X(\xi).$$

Without loss of generality we have assumed that the drift component of X is null. In the above expression, k is a right-continuous non-increasing function such that $k(0+) = +\infty$. Let us fix an arbitrary $m > 0$ and ϵ_m such that $k(z) \geq m$ for all $z \in [0, \epsilon_m]$. It is plain from the Lévy-Kintchine formula that we can define an infinitely divisible random variable Y_m with characteristic function

$$\mathbb{E} \left[e^{i\xi Y_m} \right] = \exp \left(\int_0^{\epsilon_m} (e^{i\xi z} - 1) \frac{m}{z} dz \right) =: A_m(\xi),$$

and an infinitely divisible random variable Z_m with characteristic function

$$\mathbb{E} \left[e^{i\xi Z_m} \right] = \exp \left(\int_0^{+\infty} (e^{i\xi z} - 1) \frac{k(z) - m\mathbb{1}_{[0, \epsilon_m]}(z)}{z} dz \right) =: B_m(\xi).$$

If we assume Y_m and Z_m to be independent we have clearly $\varphi_X(\xi) = A_m(\xi) \times B_m(\xi)$, so $X \stackrel{\mathcal{L}}{=} Y_m + Z_m$. We first study the asymptotic behavior of the successive derivatives of A_m and B_m to provide a bound for $\varphi_X(\xi)$.

X admits finite moments of any positive order, Y_m and Z_m are positive and satisfy $Y_m + Z_m \stackrel{\mathcal{L}}{=} X$, we thus deduce that Y_m and Z_m admit finite moments of any positive order. As a consequence A_m and B_m are of class C^∞ and all their derivatives are bounded. In particular

$$\forall n \geq 0, \left| B_m^{(n)}(\xi) \right| = \mathcal{O}_{|\xi| \rightarrow +\infty} (1). \quad (5.52)$$

Then, note that

$$A'_m(\xi) = im \int_0^{\epsilon_m} e^{i\xi z} dz \times A_m(\xi) = \frac{m}{\xi} (e^{i\xi \epsilon_m} - 1) \times A_m(\xi).$$

Let us denote $C_m(\xi) := m(e^{i\xi \epsilon_m} - 1)/\xi$. It is easy to see that the derivatives of any order of C_m converge to 0 when $|\xi|$ goes to $+\infty$. Let us justify by strong induction that for any $n \geq 1$ we have

$$A_m^{(n)}(\xi)/A_m(\xi) \xrightarrow{|\xi| \rightarrow +\infty} 0. \quad (5.53)$$

When $n = 1$, this is equivalent to the convergence to 0 of $C_m(\xi)$, which is already known. Let us now fix $l \geq 1$ and assume that (5.53) is true for all $n \leq l$. Applying the Leibniz formula to derive l times the product $A_m^{(l)}(\xi) = C_m^{(l)}(\xi) \times A_m(\xi)$, we get

$$A_m^{(l+1)}(\xi)/A_m(\xi) = \sum_{j=0}^l C_l^j C_m^{(j)}(\xi) A_m^{(l-j)}(\xi)/A_m(\xi).$$

The latter converges to 0 when $|\xi|$ goes to $+\infty$ by the induction hypothesis and the fact that the derivatives of any order of C_m converge to 0 at infinity. Thus the induction is proved. We can see that $|A_m(\xi)| = \mathcal{O}_{|\xi| \rightarrow +\infty} (|\xi|^{-m})$ (see for example the proof of Theorem 28.4 in [26]). Combining with (5.53) this yields

$$\forall n \geq 0, \left| A_m^{(n)}(\xi) \right| = \mathcal{O}_{|\xi| \rightarrow +\infty} (|\xi|^{-m}). \quad (5.54)$$

From (5.52) and (5.54) applied with $n = 0$ we know that $|\varphi_X(\xi)| = \mathcal{O}_{|\xi| \rightarrow +\infty} (|\xi|^{-m})$. For any $n \geq 1$ we can apply the Leibniz formula to derive n times the product $\varphi_X(\xi) = A_m(\xi) \times B_m(\xi)$

and we get

$$\varphi_X^{(n)}(\xi) = \sum_{j=0}^n C_n^j A_m^{(j)}(\xi) B_m^{(n-j)}(\xi).$$

According to (5.52) and (5.54) we deduce that $\forall n \geq 0$, $|\varphi_X^{(n)}(\xi)| = \mathcal{O}_{|\xi| \rightarrow +\infty}(|\xi|^{-m})$. Then, since m is arbitrary, this implies that φ_X belongs to the Schwartz space. Since this space is stable by Fourier transform, X admits a density in the Schwartz space. \square

6. THE SPECTRALLY POSITIVE CASE

We now study the exponential functional of Z^\uparrow , where Z is a spectrally positive Lévy process drifting to $+\infty$ with unbounded variation.

Recall that, for any $x \geq 0$, Z_x^\uparrow must be seen as Z conditioned to stay positive and starting from x . Note that, since Z converges almost surely to infinity, for $x > 0$, Z_x^\uparrow is only Z_x conditioned in the usual sense to remain positive.

The idea is that adding a small term of negative drift to Z^\uparrow does not change its convergence to $+\infty$. Thus, Z^\uparrow is ultimately greater than a deterministic linear function for which the exponential functional is defined and deterministically bounded. The key point is then to control the time taken by Z^\uparrow to become greater than the linear function once and for good. We start with the following lemma.

Lemma 6.1. *For any $y > 0$, there exists $\epsilon > 0$ and positive constants c_1 and c_2 such that*

$$\forall s > 0, \mathbb{P}\left(\mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]) > s\right) \leq c_1 e^{-c_2 s}.$$

Proof. We fix $y > 0$. A spectrally negative Lévy process X drifts to $-\infty$ if and only if $\mathbb{E}[X(1)] < 0$ (see for example Corollary VII.2 in [3]). Z is a spectrally positive Lévy process drifting to $+\infty$ so taking the dual we get $\mathbb{E}[Z(1)] > 0$. Now $\mathbb{E}[(Z - \epsilon \cdot)(1)] = \mathbb{E}[Z(1)] - \epsilon$ which is positive for ϵ chosen small enough. This implies that $Z - \epsilon \cdot$ is also a spectrally positive Lévy process that drifts to $+\infty$ and which is not a subordinator. We have

$$\begin{aligned} \mathbb{P}\left(\mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]) > s\right) &= \mathbb{P}\left(\inf_{t \in [s, +\infty[} Z_y^\uparrow(t) - (y + \epsilon t) \leq 0\right) \\ &= \mathcal{C} \mathbb{P}\left(\inf_{t \in [s, +\infty[} Z_y(t) - (y + \epsilon t) \leq 0, \inf_{[0, +\infty[} Z_y > 0\right). \end{aligned}$$

We have set $\mathcal{C} := 1/\mathbb{P}(\inf_{[0, +\infty[} Z_y > 0) = 1/(1 - e^{-\kappa zy})$. The last equality comes from the fact that Z_y^\uparrow is only Z_y conditioned to stay positive in the usual sense. Now, noting that $Z_y \stackrel{\mathcal{L}}{=} y + Z$, we bound the above quantity by

$$\begin{aligned} &\mathcal{C} \mathbb{P}\left(\inf_{t \in [s, +\infty[} Z(t) - \epsilon t \leq 0\right) = \mathcal{C} \mathbb{P}\left(\sup_{t \in [s, +\infty[} -Z(t) + \epsilon t \geq 0\right) \\ &\leq \mathcal{C} \mathbb{P}\left(\sup_{t \in [s, +\infty[} -Z(t) + \epsilon t \geq 0, -Z(s) + \epsilon s \leq 0\right) + \mathcal{C} \mathbb{P}(-Z(s) + \epsilon s \geq 0) \\ &= \mathcal{C} \mathbb{P}\left(\sup_{t \in [0, +\infty[} -Z^s(t) + \epsilon t \geq -(-Z(s) + \epsilon s), -Z(s) + \epsilon s \leq 0\right) + \mathcal{C} \mathbb{P}(-Z(s) + \epsilon s > 0). \end{aligned} \tag{6.55}$$

From the stationarity and independence of increments, the process $-Z^s + \epsilon$ is equal in law to $-Z + \epsilon$ and independent from $-Z(s) + \epsilon s$. From Corollary VII.2 in [3], the supremum over $[0, +\infty[$ of the process $-Z^s + \epsilon$ follows an exponential distribution with parameter α , where α is the non-trivial zero of $\Psi_{-Z+\epsilon}$. From this, combined with the independence from $-Z(s) + \epsilon s$, (6.55) becomes

$$\begin{aligned} & \mathcal{C} \mathbb{E} \left(e^{\alpha(-Z(s)+\epsilon s)} \mathbf{1}_{\{-Z(s)+\epsilon s \leq 0\}} \right) + \mathcal{C} \mathbb{P}(-Z(s) + \epsilon s > 0) \\ & \leq \mathcal{C} \mathbb{E} \left(e^{\alpha(-Z(s)+\epsilon s)/2} \mathbf{1}_{\{-Z(s)+\epsilon s \leq 0\}} \right) + \mathcal{C} \mathbb{P} \left(e^{\alpha(-Z(s)+\epsilon s)/2} > 1 \right). \end{aligned}$$

We have used the decreases of the negative exponential for the first term and composed by the function $x \mapsto \exp(\frac{\alpha}{2}x)$ in the probability of the second term. The above is less than

$$\mathcal{C} \mathbb{E} \left(e^{\frac{\alpha}{2}(-Z(s)+\epsilon s)} \right) + \mathcal{C} \mathbb{P} \left(e^{\frac{\alpha}{2}(-Z(s)+\epsilon s)} > 1 \right) \leq 2\mathcal{C} \mathbb{E} \left(e^{\alpha/2(-Z(s)+\epsilon s)} \right),$$

where we have used Markov's inequality in the second term.

$$= 2\mathcal{C} e^{s\Psi_{-Z+\epsilon}(\alpha/2)}.$$

Since $\Psi_{-Z+\epsilon}$ is negative on $]0, \alpha[$, we get the result with $c_1 = 2\mathcal{C}$ and $c_2 = -\Psi_{-Z+\epsilon}(\alpha/2)$. \square

We can now prove Theorem 1.13.

Proof. of Theorem 1.13

We fix $y > 0$. In order to get Z^\uparrow from Z_y^\uparrow , we use the decomposition given by Theorem 25 in [8]. Let m_y be the point where the process Z_y^\uparrow reaches its infimum, $m_y := \sup\{s \geq 0, Z_y^\uparrow(s-) \wedge Z_y^\uparrow(s) = \inf_{[0, +\infty[} Z_y^\uparrow\}$. Note that from the absence of negative jumps the infimum is always reached at least at $m_y -$ so $Z_y^\uparrow(m_y -) = \inf_{[0, +\infty[} Z_y^\uparrow$. Moreover, since Z is regular for $]0, +\infty[$, Z_y^\uparrow is actually continuous at m_y (see the theorem in [8]). The decomposition given by Theorem 25 in [8] states that

- The two processes $(Z_y^\uparrow(m_y + s) - Z_y^\uparrow(m_y), s \geq 0)$ and $(Z_y^\uparrow(s), 0 \leq s < m_y)$ are independent,
- $(Z_y^\uparrow(m_y + s) - Z_y^\uparrow(m_y), s \geq 0)$ is equal in law to Z^\uparrow .

Now,

$$\begin{aligned} I(Z_y^\uparrow) &= \int_0^{m_y} e^{-Z_y^\uparrow(u)} du + \int_{m_y}^{+\infty} e^{-Z_y^\uparrow(u)} du \\ &= \int_0^{m_y} e^{-Z_y^\uparrow(u)} du + e^{-Z_y^\uparrow(m_y)} \int_0^{+\infty} e^{-(Z_y^\uparrow(m_y+u) - Z_y^\uparrow(m_y))} du \\ &\geq e^{-y} \int_0^{+\infty} e^{-(Z_y^\uparrow(m_y+u) - Z_y^\uparrow(m_y))} du, \end{aligned}$$

because almost surely $Z_y^\uparrow(m_y) \leq y$. Then, because of the above decomposition of the trajectory of Z_y^\uparrow , the latter is equal in law to $e^{-y}I(Z^\uparrow)$. We thus get

$$I(Z^\uparrow) \stackrel{sto}{\leq} e^y I(Z_y^\uparrow), \quad (6.56)$$

where $\overset{sto}{\leq}$ denotes a stochastic inequality. As a consequence we only need to prove the result for $I(Z_y^\uparrow)$. We now choose $\epsilon > 0$ as in Lemma 6.1. We have

$$\begin{aligned} 0 \leq I(Z_y^\uparrow) &= \int_0^{\mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0])} e^{-Z_y^\uparrow(t)} dt + \int_{\mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]}^{+\infty} e^{-Z_y^\uparrow(t)} dt \\ &\leq \mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]) + \int_{\mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]}^{+\infty} e^{-Z_y^\uparrow(t)} dt. \end{aligned}$$

Then, for $t \geq \mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0])$, we have $Z_y^\uparrow(t) \geq y + \epsilon t$, so

$$\begin{aligned} 0 \leq I(Z_y^\uparrow) &\leq \mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]) + \int_{\mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]}^{+\infty} e^{-y - \epsilon t} dt \\ &\leq \mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]) + \int_0^{+\infty} e^{-y - \epsilon t} dt \\ &= \mathcal{R}(Z_y^\uparrow(\cdot) - (y + \epsilon \cdot),] - \infty, 0]) + e^{-y}/\epsilon. \end{aligned}$$

According to Lemma 6.1, this is almost surely finite and admits some finite exponential moments. Thanks to (6.56) we have the same for $I(Z^\uparrow)$ which is the expected result. \square

Remark 6.2. *The arguments of the proof of Theorem 1.13 can be extended for a more general Lévy process Z if we assume that : Z drifts to $+\infty$, Z is regular for $]0, +\infty[$ and $] - \infty, 0[$, $\exists \lambda < 0$ s.t. $\mathbb{E}[\exp(\lambda Z(1))] < +\infty$ (which, according to Theorem 25.3 in [26], is equivalent to the fact that the restriction to $] - \infty, -1[$ of the Lévy measure of Z admits some finite exponential moments of negative order).*

We need an analogous of Lemma 2.6 in order to compare, as we did in subsection 4.3, the exponential functionals $I(Z^\uparrow)$ and $I(Z)$. We define m to be the point where the process Z reaches its infimum : $m := \sup\{s \geq 0, Z(s-) \wedge Z(s) = \inf_{[0, +\infty[} Z\}$. Here again, from the absence of negative jumps, the infimum is always reached at least at $m-$ so $Z(m-) = \inf_{[0, +\infty[} Z$. Also, since Z is regular for $]0, +\infty[$, Z is actually continuous at m (see the proof of Theorem 25 in [8]). The arguments in the proof of Theorem 25 of [8] yields

Lemma 6.3. $Z(m + \cdot) - Z(m)$ has the same law as Z^\uparrow .

We can now prove Proposition 1.15.

Proof. of Proposition 1.15

We have

$$\begin{aligned} I(Z) &= \int_0^m e^{-Z(t)} dt + \int_m^{+\infty} e^{-Z(t)} dt \\ &= \int_0^m e^{-Z(t)} dt + e^{-Z(m)} \int_m^{+\infty} e^{-(Z(m+t) - Z(m))} dt \\ &\geq \int_m^{+\infty} e^{-(Z(m+t) - Z(m))} dt. \end{aligned}$$

For the last inequality we have used the fact that almost surely $Z(m) \leq 0$. Lemma 6.3 tells us that the last term is equal in law to $I(Z^\uparrow)$. We thus get $I(Z^\uparrow) \overset{sto}{\leq} I(Z)$ where $\overset{sto}{\leq}$ denotes a stochastic inequality. Proposition 1.15 follows.

□

Remark 6.4. *The finiteness of $I(Z^\uparrow)$ can be proved directly using Proposition 1.15 combined with the well known finiteness of $I(Z)$ (see Theorem 1 in [4]). As in the spectrally negative case, this comparison with the functional of the non-conditioned process could not be used to derive the existence of exponential moments for $I(Z^\uparrow)$ (Theorem 1.13), since the right tail of $I(Z)$ is heavier (see Remark 1.14). However, similarly to the spectrally negative case, Proposition 1.15 leads to a relevant comparison of the left tails of $I(Z^\uparrow)$ and $I(Z)$.*

Remark 6.5. *Here again, the arguments of the proof of Proposition 1.15 can be extended for a more general Lévy process Z if we assume that Z drifts to $+\infty$ and Z is regular for $]0, +\infty[$ and $] -\infty, 0[$.*

7. ASYMPTOTIC TAIL AT 0 IN THE PRESENCE OF POSITIVE JUMPS

7.1. Proof of Theorem 1.16. As in the Introduction, Y is a Lévy process drifting to $+\infty$. Let us denote by (A, d, π) its Lévy triplet. π therefore denotes its Lévy measure and for $x > 0$, $\bar{\pi}(x) := \pi([x, +\infty[)$.

We first prove (1.9). It is trivially satisfied if the support of π is bounded from above. We thus assume that it is not. Let us fix $x \in]0, 1[$ and define

$$\pi^a := \pi(\cdot \cap [\log(1/x), +\infty[) \quad \text{and} \quad \pi^b := \pi(\cdot \cap] -\infty, \log(1/x)[).$$

Let Y^a and Y^b be two independent Lévy processes which generating triplets are respectively $(0, 0, \pi^a)$ and (A, d, π^b) . This yields $Y \stackrel{\mathcal{L}}{=} Y^a + Y^b$ according to the Lévy-Khintchine formula. Let J denote the random instant of the first jump of Y^a (note that J follows an exponential distribution with parameter $\bar{\pi}(\log(1/x))$) and $\tilde{Y} := Y(J + \cdot) - Y(J)$. Because of the Markov property \tilde{Y} is equal in law to Y and independent from $\sigma(Y^a(s), Y^b(s), 0 \leq s \leq J)$. We have

$$\begin{aligned} \mathbb{P}(I(Y) \leq x) &\geq \mathbb{P}\left(J \leq x/3, \int_0^J e^{-Y^b(t)} dt \leq 2x/3, e^{-Y(J)} I(\tilde{Y}) \leq x/3\right) \\ &\geq \mathbb{P}\left(J \leq x/3, \inf_{[0, J]} Y^b \geq -\log(2), e^{-Y(J)} I(\tilde{Y}) \leq x/3\right). \end{aligned}$$

Note that in the above event $Y(J) = Y^a(J) + Y^b(J) \geq \log(1/x) - \log(2) = \log(1/2x)$ so $e^{-Y(J)} \leq 2x$. We thus have

$$\begin{aligned} \mathbb{P}(I(Y) \leq x) &\geq \mathbb{P}\left(J \leq x/3, \inf_{[0, J]} Y^b \geq -\log(2), 2x I(\tilde{Y}) \leq x/3\right) \\ &= \mathbb{P}\left(J \leq x/3, \inf_{[0, J]} Y^b \geq -\log(2)\right) \times \mathbb{P}\left(I(\tilde{Y}) \leq 1/6\right). \end{aligned}$$

In the last equality we have used the independence between \tilde{Y} and $\sigma(Y^a(s), Y^b(s), 0 \leq s \leq J)$. Then,

$$\mathbb{P}(I(Y) \leq x) \geq \mathbb{P}\left(J \leq x/3, \inf_{[0, 1]} Y^b \geq -\log(2)\right) \times \mathbb{P}\left(I(\tilde{Y}) \leq 1/6\right),$$

where we have used the fact that, in the above event, $J < x/3 < 1$. Then, since Y^a and Y^b are independent we obtain

$$\begin{aligned} \mathbb{P}(I(Y) \leq x) &\geq \mathbb{P}(J \leq x/3) \times \mathbb{P}\left(\inf_{[0, 1]} Y^b \geq -\log(2)\right) \times \mathbb{P}\left(I(\tilde{Y}) \leq 1/6\right) \\ &= \left(1 - e^{-x\bar{\pi}(\log(1/x))/3}\right) \times \mathbb{P}\left(\inf_{[0, 1]} Y^b \geq -\log(2)\right) \times \mathbb{P}\left(I(\tilde{Y}) \leq 1/6\right). \end{aligned}$$

When x goes to 0, the first factor is equivalent to $x\bar{\pi}(\log(1/x))/3$ while the second factor converges to $\mathbb{P}(\inf_{[0,1]} Y \geq -\log(2))$ and the third factor remains constant. We thus get (1.9).

The idea to prove (1.10) is to first prove (1.11) and then to make appear a Poisson process in the trajectory of a general Y . We thus now determine the left tail of $I(\alpha N)$ where N is a standard Poisson process and $\alpha > 0$. Let us denote by p the parameter of N .

We proceed by studying the asymptotic behavior of the log-Laplace transform of $I(\alpha N)$ at infinity. By a property of standard Poisson processes, it is easy to see that $I(\alpha N)$ can be written as

$$\frac{1}{p} \sum_{k=0}^{+\infty} e^{-\alpha k} e_k,$$

where $(e_k)_{k \in \mathbb{N}}$ is a sequence of *iid* exponential random variable with parameter 1. This allows to compute the log-Laplace transform of $I(\alpha N)$:

$$\forall \lambda \geq 0, \quad -\log \left(\mathbb{E} \left[e^{-\lambda I(\alpha N)} \right] \right) = \sum_{k=0}^{+\infty} \log \left(1 + \frac{\lambda}{p} e^{-\alpha k} \right). \quad (7.57)$$

Then, note that for any integer $k \geq 0$ we have

$$\log \left(1 + \frac{\lambda}{p} e^{-\alpha(k+1)} \right) \leq \int_k^{k+1} \log \left(1 + \frac{\lambda}{p} e^{-\alpha y} \right) dy \leq \log \left(1 + \frac{\lambda}{p} e^{-\alpha k} \right).$$

Combining with (7.57) we get for any $\lambda \geq 0$,

$$J(\lambda) \leq -\log \left(\mathbb{E} \left[e^{-\lambda I(\alpha N)} \right] \right) \leq J(\lambda) + \log \left(1 + \frac{\lambda}{p} \right), \quad (7.58)$$

where $J(\lambda) := \int_0^{+\infty} \log(1 + \frac{\lambda}{p} e^{-\alpha y}) dy$. Making the change of variable $u = \lambda e^{-\alpha y}/p$ we get

$$J(\lambda) = \frac{1}{\alpha} \int_0^{\lambda/p} \frac{\log(1+u)}{u} du.$$

Now, λ only determines the intervalle of integration. Since the above integral converges to $+\infty$ when λ goes to infinity we have, for any $M > 0$,

$$J(\lambda) \underset{\lambda \rightarrow +\infty}{\sim} \frac{1}{\alpha} \int_M^{\lambda/p} \frac{\log(1+u)}{u} du.$$

In particular, we can fix $\epsilon > 0$ and choose M so large such that $\forall u \geq M$, $(1 - \epsilon) \log(u) \leq \log(1+u) \leq (1 + \epsilon) \log(u)$. This implies in particular that

$$J(\lambda) \underset{\lambda \rightarrow +\infty}{\sim} \frac{1}{\alpha} \int_1^{\lambda/p} \frac{\log(u)}{u} du.$$

Now making the change of variable $v = \log(u)$ we get that the right hand side equals

$$\frac{1}{\alpha} \int_0^{\log(\lambda/p)} v dv = \frac{1}{2\alpha} (\log(\lambda/p))^2.$$

As a consequence we have $J(\lambda) \sim (\log(\lambda))^2/2\alpha$, so putting into (7.58) we get

$$-\log \left(\mathbb{E} \left[e^{-\lambda I(\alpha N)} \right] \right) \underset{\lambda \rightarrow +\infty}{\sim} \frac{1}{2\alpha} (\log(\lambda))^2. \quad (7.59)$$

We now deduce the left tail of $I(\alpha N)$ by standard methods. From Markov's inequality we have

$$\forall x > 0, \quad \mathbb{P}(I(\alpha N) \leq x) \leq e \mathbb{E} \left[e^{-I(\alpha N)/x} \right]. \quad (7.60)$$

Let us fix $\epsilon > 0$. We have

$$\begin{aligned} \forall x > 0, \mathbb{E} \left[e^{-I(\alpha N)/x^{1+\epsilon}} \right] &\leq \mathbb{E} \left[e^{-I(\alpha N)/x^{1+\epsilon}} \mathbf{1}_{\{I(\alpha N) \leq x\}} \right] + \mathbb{E} \left[e^{-I(\alpha N)/x^{1+\epsilon}} \mathbf{1}_{\{I(\alpha N) > x\}} \right] \\ &\leq \mathbb{P}(I(\alpha N) \leq x) + e^{-1/x^\epsilon}. \end{aligned} \quad (7.61)$$

Combining (7.60) and (7.61) with (7.59) we get

$$1 \leq \liminf_{x \rightarrow 0} \frac{-\log(\mathbb{P}(I(\alpha N) \leq x))}{\frac{1}{2\alpha}(\log(x))^2} \leq \limsup_{x \rightarrow 0} \frac{-\log(\mathbb{P}(I(\alpha N) \leq x))}{\frac{1}{2\alpha}(\log(x))^2} \leq (1 + \epsilon)^2.$$

Letting ϵ go to 0 we get (1.11).

We now prove (1.10). Since, by assumption, $\pi(\cdot \cap]0, +\infty[)$ is non zero, there exist $0 < \gamma_1 < \gamma_2 < +\infty$ such that $\pi([\gamma_1, \gamma_2]) > 0$. Then, for $\eta \in]0, 1[$ we define

$$\pi^{\eta,1} := \eta\pi(\cdot \cap [\gamma_1, \gamma_2]) \quad \text{and} \quad \pi^{\eta,2} := \pi - \eta\pi(\cdot \cap [\gamma_1, \gamma_2]).$$

We set $Y^{\eta,1}$ and $Y^{\eta,2}$ to be two independent Lévy processes which generating triplets are respectively $(0, 0, \pi^{\eta,1})$ and $(A, d, \pi^{\eta,2})$. We have $\pi = \pi^{\eta,1} + \pi^{\eta,2}$ so according to the Lévy-Khintchine formula,

$$Y \stackrel{\mathcal{L}}{=} Y^{\eta,1} + Y^{\eta,2}.$$

Let us justify that η can be chosen in such a way so that $Y^{\eta,2}$ drifts to $+\infty$. First let us assume that $\mathbb{E}[(Y(1))^-] < +\infty$. Then $\mathbb{E}[Y(1)]$ is defined and belongs to $]0, +\infty[$ (otherwise Y would not drift to $+\infty$ according to the law of large number and Theorem 36.7 in [26]). Since $\mathbb{E}[Y(1)] = \mathbb{E}[Y^{\eta,1}(1)] + \mathbb{E}[Y^{\eta,2}(1)]$ and $\mathbb{E}[Y^{\eta,1}(1)] < \gamma_2\eta\pi([\gamma_1, \gamma_2])$, we have that $\mathbb{E}[Y^{\eta,2}(1)]$ is positive for η small enough. By the law of large numbers, this implies that $Y^{\eta,2}$ drifts to $+\infty$ for such small η . We now assume that $\mathbb{E}[(Y(1))^-] = +\infty$, then we also have $\mathbb{E}[(Y(1))^+] = +\infty$ (otherwise Y would not drift to $+\infty$ according to the law of large numbers). Since Y drifts to $+\infty$, Y is in the case (1) of Remark 37.13 of [26]. Note that $\pi^{\eta,2}$ and π have the same asymptotic tails at $+\infty$ and $-\infty$ so replacing π by $\pi^{\eta,2}$ does not change the finiteness of K^- (resp. the infiniteness of K^+) from Remark 37.13 of [26]. As a consequence for any $\eta \in]0, 1[$, $Y^{\eta,2}$ is also in the case (1) of Remark 37.13 of [26], and therefore drifts to $+\infty$. In any case we can chose η such that $Y^{\eta,2}$ drifts to $+\infty$. We thus choose such an $\eta_0 \in]0, 1[$ and denote by m_2 the point at which $Y^{\eta_0,2}$ reaches its minimum.

$Y^{\eta_0,1}$ is a compound Poisson process, we define N the counting process of its jumps : $N(t) := \#\{s \in [0, t], Y^{\eta_0,1}(s) - Y^{\eta_0,1}(s-) > 0\}$. N is thus a standard Poisson process with parameter $c_0 := \eta_0\pi([\gamma_1, \gamma_2])$ and it is independent of $Y^{\eta_0,2}$. We have

$$\begin{aligned} I(Y) &= \int_0^{+\infty} e^{-(Y^{\eta_0,1}(t) + Y^{\eta_0,2}(t))} dt \\ &\leq e^{-Y^{\eta_0,2}(m_2)} \int_0^{+\infty} e^{-Y^{\eta_0,1}(t)} dt \\ &\leq e^{-Y^{\eta_0,2}(m_2)} \int_0^{+\infty} e^{-\gamma_1 N(t)} dt \\ &= e^{-Y^{\eta_0,2}(m_2)} I(\gamma_1 N). \end{aligned}$$

Then, from the independence between the two factors

$$\mathbb{P}(I(\gamma_1 N) \leq x/2) \times \mathbb{P}\left(e^{-Y^{\eta_0,2}(m_2)} \leq 2\right) \leq \mathbb{P}(I(Y) \leq x). \quad (7.62)$$

Applying (1.11) to $I(\gamma_1 N)$ we get, for $\epsilon > 0$ and all x small enough,

$$e^{-(1+\epsilon)(\log(x))^2/2\gamma_1} \leq \mathbb{P}(I(\gamma_1 N) \leq x/2).$$

Combining with (7.62), we get the result with $c = (1 + \epsilon)/2\gamma_1$. Since ϵ can be chosen as close as we want to 0 and γ_1 can be chosen as close as we want to S , (1.10) follows.

7.2. Proof of Proposition 1.17. We now prove Proposition 1.17. The ingredients of the proof are : the factorization identity of $I(Y)$ from [14], Theorem 2.5 of [17] which gives the asymptotic tail of exponential functionals of subordinators, easy calculations on the ascending ladder height process of Y and on regularly varying functions. We give all the details in order to make the proof rigorous.

As in the statement of the proposition, we assume that Y is spectrally positive and π satisfies (A) for some $\alpha > 0$. Note that this implies that $\bar{\pi}(\log(\cdot))$ has $-\alpha$ -regular variation so $\bar{\pi}(x) = o(e^{-\eta x})$ for any $\eta \in]0, \alpha[$ and the Laplace exponent of Y , Ψ_Y , is defined on $] - \infty, \alpha[$, where

$$\forall t \geq 0, \lambda \in] - \infty, \alpha[, \mathbb{E} \left[e^{\lambda Y(t)} \right] = e^{t\Psi_Y(\lambda)}.$$

Let H be the the ascending ladder height process associated with Y and Φ_H denote its Laplace exponent :

$$\Phi_H(\lambda) := -\log \left(\mathbb{E} \left[e^{-\lambda H(1)} \right] \right).$$

H is a proper subordinator since Y drifts to $+\infty$. Let d_H and π_H denote respectively the drift and the Lévy measure of H and for any $x > 0$, $\bar{\pi}_H(x) := \pi_H([x, +\infty[)$. The Lévy-Kintchine formula reads

$$\forall \lambda \geq 0, \Phi_H(\lambda) = d_H \lambda + \int_0^{+\infty} (1 - e^{-\lambda x}) \pi_H(dx). \quad (7.63)$$

According to [3], Section VII.1 (applied to $-Y$), the descending ladder height process associated with Y has its Laplace exponent equal to $\lambda + \kappa_Y$. According to the Wiener-Hopf factorization for Laplace exponents we have

$$\forall \lambda \geq 0, \Psi_Y(-\lambda) = -(-\lambda + \kappa_Y)\Phi_H(\lambda) = (\lambda - \kappa_Y)\Phi_H(\lambda). \quad (7.64)$$

Note that the above equality is true up to a multiplicative constant but we can consider that the normalization of the local time (from which is defined the ascending ladder height process) has been chosen in such a way that this constant equals 1. Combining the above factorization with (7.63) we obtain for all $\lambda \geq 0$,

$$\begin{aligned} \Psi_Y(-\lambda) &= d_H \lambda^2 - \left(\int_0^{+\infty} x \pi_H(dx) + d_H \right) \kappa_Y \lambda + \int_0^{+\infty} \lambda (1 - e^{-\lambda x}) \pi_H(dx) \\ &\quad + \kappa_Y \int_0^{+\infty} (e^{-\lambda x} - 1 + \lambda x) \pi_H(dx). \end{aligned}$$

The fact that $\bar{\pi}(x) = o(e^{-\eta x})$ for any $\eta \in]0, \alpha[$ together with (7.64) implies that Φ_H can be analytically extended on $] - \alpha, +\infty[$. As a consequence $H(1)$ admits some finite exponential moments and, according to Theorem 25.3 in [26], so does the Lévy measure π_H . This justifies the existence of the term $\int_0^{+\infty} x \pi_H(dx)$ that we made appear in the previous expression. By integration by parts we obtain

$$\Psi_Y(-\lambda) = d_H \lambda^2 - \left(\int_0^{+\infty} x \pi_H(dx) + d_H \right) \kappa_Y \lambda + \int_0^{+\infty} \lambda (1 - e^{-\lambda x}) (\pi_H(dx) + \kappa_Y \bar{\pi}_H(x) dx)$$

Thanks to the Lévy-Kintchine formula applied to Y we can identify, in the above expression, the Lévy measure of Y . We obtain

$$\pi_H(dx) + \kappa_Y \bar{\pi}_H(x) dx = \bar{\pi}(x) dx.$$

This leads to the following differential equation in the sens of distributions : $\overline{\pi}_H' - \kappa_Y \overline{\pi}_H = -\overline{\pi}$. Solving this equation we get

$$\forall x \geq 0, \overline{\pi}_H(x) = K e^{\kappa_Y x} + e^{\kappa_Y x} \int_x^{+\infty} e^{-\kappa_Y y} \overline{\pi}(y) dy,$$

where K is a positive constant. Now, note that $\overline{\pi}_H(x)$ converges to 0 when x goes to $+\infty$ and it is no difficult to see that the second term in the above expression goes to 0 when x goes to $+\infty$. This imposes $K = 0$. Then,

$$\frac{\overline{\pi}_H(x)}{\overline{\pi}(x)} = e^{\kappa_Y x} \int_x^{+\infty} e^{-\kappa_Y y} \frac{\overline{\pi}(y)}{\overline{\pi}(x)} dy = \int_0^{+\infty} e^{-\kappa_Y z} \frac{\overline{\pi}(z+x)}{\overline{\pi}(x)} dz.$$

$\overline{\pi}$ is non-increasing so, for any $z \geq 0$, we have $\overline{\pi}(z+x)/\overline{\pi}(x) \leq 1$. By the assumption (A), $\overline{\pi}(z+x)/\overline{\pi}(x)$ converges, for any fixed $z \geq 0$, to $e^{-\alpha z}$ when x goes to $+\infty$. By dominated convergence, we conclude that $\overline{\pi}_H(x)/\overline{\pi}(x)$ converges to $1/(\alpha + \kappa_Y)$ when x goes to $+\infty$, so

$$\overline{\pi}_H(x) \underset{x \rightarrow +\infty}{\sim} \frac{1}{\alpha + \kappa_Y} \overline{\pi}(x).$$

As a consequence, π_H satisfies (A) with the same α . We can thus apply Theorem 2.5 of [17] and we get the following equivalent for m_H , the density of the exponential functional $I(H)$.

$$m_H(x) \underset{x \rightarrow 0}{\sim} \mathbb{E}[I(H)^{-\alpha}] \overline{\pi}_H(\log(1/x)) \underset{x \rightarrow 0}{\sim} \frac{\mathbb{E}[I(H)^{-\alpha}]}{\alpha + \kappa_Y} \overline{\pi}(\log(1/x)).$$

Now, let us fix $\epsilon > 0$. Thanks to the previous equivalent, there exists $y_0 \in]0, 1[$ such that

$$\forall y \in]0, y_0[, (1 - \epsilon) \frac{\mathbb{E}[I(H)^{-\alpha}]}{\alpha + \kappa_Y} \overline{\pi}(\log(1/y)) \leq m_H(y) \leq (1 + \epsilon) \frac{\mathbb{E}[I(H)^{-\alpha}]}{\alpha + \kappa_Y} \overline{\pi}(\log(1/y)). \quad (7.65)$$

According to Corollary 2.1(2) in [14], the density m_Y of $I(Y)$ is given by the following expression for $x > 0$.

$$\begin{aligned} m_Y(x) &= \frac{x^{-\kappa_Y - 1}}{\Gamma(\kappa_Y)} \int_0^{+\infty} e^{-y/x} y^{\kappa_Y} m_H(y) dy \geq \frac{x^{-\kappa_Y - 1}}{\Gamma(\kappa_Y)} \int_0^{y_0} e^{-y/x} y^{\kappa_Y} m_H(y) dy \\ &\geq (1 - \epsilon) \frac{\mathbb{E}[I(H)^{-\alpha}] x^{-\kappa_Y - 1}}{(\alpha + \kappa_Y) \Gamma(\kappa_Y)} \int_0^{y_0} e^{-y/x} y^{\kappa_Y} \overline{\pi}(\log(1/y)) dy \\ &= (1 - \epsilon) \frac{\mathbb{E}[I(H)^{-\alpha}] x^{-\kappa_Y - 1}}{(\alpha + \kappa_Y) \Gamma(\kappa_Y)} \overline{\pi}(\log(1/x)) \int_0^{y_0} e^{-y/x} y^{\kappa_Y} \frac{\overline{\pi}(\log(1/y))}{\overline{\pi}(\log(1/x))} dy \\ &= (1 - \epsilon) \frac{\mathbb{E}[I(H)^{-\alpha}]}{(\alpha + \kappa_Y) \Gamma(\kappa_Y)} \overline{\pi}(\log(1/x)) \int_0^{+\infty} e^{-u} u^{\kappa_Y} \frac{\overline{\pi}(\log(1/ux))}{\overline{\pi}(\log(1/x))} \mathbb{1}_{u < y_0/x} du, \end{aligned} \quad (7.66)$$

where we have used the lower bound of (7.65) and made the change of variable $u = y/x$. Since $\overline{\pi}(\log(\cdot))$ has $-\alpha$ -regular variation, we have, according to Karamata representation theorem,

$$\overline{\pi}(\log(z)) = z^{-\alpha} \exp \left(c(z) + \int_1^z \frac{r(t)}{t} dt \right). \quad (7.67)$$

In this representation c is a bounded measurable function that converges to a constant c at $+\infty$, and r is a bounded measurable function that converges to 0 at $+\infty$. By decreasing y_0 if necessary, we can assume that for all $z \geq 1/y_0$: $c - \epsilon \leq c(z) \leq c + \epsilon$ and $|r(z)| \leq \epsilon$. As a consequence, for all $x \leq y_0$ and $u \in]0, y_0/x[$ we have

$$\begin{aligned} \frac{\overline{\pi}(\log(1/ux))}{\overline{\pi}(\log(1/x))} &\geq u^\alpha \exp \left(c(1/ux) - c(1/x) + \int_{1/x}^{1/ux} \frac{r(t)}{t} dt \right) \\ &\geq u^\alpha \exp(-2\epsilon - \epsilon |\log(u)|) = u^\alpha (u \wedge 1/u)^\epsilon e^{-2\epsilon}. \end{aligned}$$

In the last inequality we have used the fact that, since $1/x$ and $1/xu$ are greater than $1/y_0$ we have $-2\epsilon \leq c(1/ux) - c(1/x) \leq 2\epsilon$ and $\epsilon |\log(u)| = \epsilon |\log(1/ux) - \log(1/x)| \geq \int_{1/x}^{1/ux} \frac{r(t)}{t} dt \geq -\epsilon |\log(1/ux) - \log(1/x)| = -\epsilon |\log(u)|$. Putting into (7.66) we get for all $x \leq y_0$,

$$m_Y(x) \geq (1 - \epsilon)e^{-2\epsilon} \frac{\mathbb{E}[I(H)^{-\alpha}]}{(\alpha + \kappa_Y)\Gamma(\kappa_Y)} \bar{\pi}(\log(1/x)) \int_0^{+\infty} e^{-u} u^{\alpha + \kappa_Y} (u \wedge 1/u)^\epsilon \mathbf{1}_{u < y_0/x} du.$$

By monotone convergence in the integral of the right hand side we obtain

$$\liminf_{x \rightarrow 0} \frac{m_Y(x)}{\bar{\pi}(\log(1/x))} \geq (1 - \epsilon)e^{-2\epsilon} \frac{\mathbb{E}[I(H)^{-\alpha}]}{(\alpha + \kappa_Y)\Gamma(\kappa_Y)} \int_0^{+\infty} e^{-u} u^{\alpha + \kappa_Y} (u \wedge 1/u)^\epsilon du.$$

Letting ϵ go to 0 we obtain, still by monotone convergence,

$$\begin{aligned} \liminf_{x \rightarrow 0} \frac{m_Y(x)}{\bar{\pi}(\log(1/x))} &\geq \frac{\mathbb{E}[I(H)^{-\alpha}]}{(\alpha + \kappa_Y)\Gamma(\kappa_Y)} \int_0^{+\infty} e^{-u} u^{\alpha + \kappa_Y} du \\ &= \frac{\mathbb{E}[I(H)^{-\alpha}]\Gamma(\alpha + \kappa_Y + 1)}{(\alpha + \kappa_Y)\Gamma(\kappa_Y)} = \frac{\mathbb{E}[I(H)^{-\alpha}]\Gamma(\alpha + \kappa_Y)}{\Gamma(\kappa_Y)}. \end{aligned} \quad (7.68)$$

For the upper bound, using again the expression of m_Y from Corollary 2.1(2) in [14] we have, for $x > 0$,

$$m_Y(x) = \frac{x^{-\kappa_Y - 1}}{\Gamma(\kappa_Y)} \int_0^{y_0} e^{-y/x} y^{\kappa_Y} m_H(y) dy + \frac{x^{-\kappa_Y - 1}}{\Gamma(\kappa_Y)} \int_{y_0}^{+\infty} e^{-y/x} y^{\kappa_Y} m_H(y) dy. \quad (7.69)$$

If $x < y_0/\kappa_Y$, then $(y \mapsto e^{-y/x} y^{\kappa_Y})$ is decreasing on $[y_0, +\infty[$. As a consequence, if x is small enough the second term of (7.69) is less than

$$\frac{x^{-\kappa_Y - 1}}{\Gamma(\kappa_Y)} e^{-y_0/x} y_0^{\kappa_Y} \int_{y_0}^{+\infty} m_H(y) dy.$$

For the first term of (7.69) we can proceed as we did for the lower bound of $m_Y(x)$ and get that it is less than

$$(1 + \epsilon)e^{2\epsilon} \frac{\mathbb{E}[I(H)^{-\alpha}]}{(\alpha + \kappa_Y)\Gamma(\kappa_Y)} \bar{\pi}(\log(1/x)) \int_0^{+\infty} e^{-u} u^{\alpha + \kappa_Y} (u \vee 1/u)^\epsilon du.$$

Then, since $\bar{\pi}(\log(\cdot))$ has $-\alpha$ -regular variation we have $x^\eta = o(\bar{\pi}(\log(1/x)))$ for any $\eta > \alpha$. As a consequence the upper bound of the second term of (7.69) is negligible compared to the upper bound of the first term. We thus get

$$\limsup_{x \rightarrow 0} \frac{m_Y(x)}{\bar{\pi}(\log(1/x))} \leq (1 + \epsilon)e^{2\epsilon} \frac{\mathbb{E}[I(H)^{-\alpha}]}{(\alpha + \kappa_Y)\Gamma(\kappa_Y)} \int_0^{+\infty} e^{-u} u^{\alpha + \kappa_Y} (u \vee 1/u)^\epsilon du.$$

Here again, we can let ϵ go to 0 and obtain the same value as in (7.68). Thus, combing with (7.68) we get the result.

Acknowledgements :

We are grateful to two anonymous referees for their helpful comments that allowed us to considerably improve the paper. In particular, we thank one of the referees for the simpler and nicer proof he or she suggested to show the existence of exponential moments for $I(V^\uparrow)$.

REFERENCES

- [1] P. Andreatti and A. Devulder. Localization and number of visited valleys for a transient diffusion in random environment. *Electronic Journal of Probability*, pages 1–59, 2015.
- [2] P. Andreatti, A. Devulder, and G. Véchambre. Renewal structure and local time for diffusions in random environment. *ALEA, Lat. Am. J. Probab. Math. Stat.*, **13**:863–923, 2016.
- [3] J. Bertoin. *Lévy Processes*. Tracts in Mathematics, Cambridge, 1996.

- [4] J. Bertoin and M. Yor. Exponential functionals of lévy processes. *Probab. Surveys*, 2:191–212, 2005.
- [5] Jean Bertoin and Maria-Emilia Caballero. Entrance from $0+$ for increasing semi-stable markov processes. *Bernoulli*, 8(2):195–205, 04 2002.
- [6] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular Variation*. Cambridge University Press, 1987. Cambridge Books Online.
- [7] T. Brox. A one-dimensional diffusion process in a Wiener medium. *Ann. Probab.*, 14(4): 1206–1218, 1986.
- [8] R. Doney. *Fluctuation Theory for Lévy Processes (Summer school, St-Floor France)*. Springer, 2005.
- [9] A. Lindner J. Bertoin and R. Maller. On continuity properties of the law of integrals of lévy processes.
- [10] A. Kuznetsov and J. C. Pardo. Fluctuations of stable processes and exponential functionals of hypergeometric lévy processes. *Acta Applicandae Mathematicae*, 123(1):113–139, 2013.
- [11] K. Maulik and B. Zwart. Tail asymptotics for exponential functionals of lévy processes. *Stochastic Processes and their Applications*, 116(2):156 – 177, 2006.
- [12] F. Petit P. Carmona and M. Yor. On the distribution and asymptotic results for exponential functionals of Lévy processes. In *Exponential functionals and principal values related to Brownian motion*, Bibl. Rev. Mat. Iberoamericana, pages 73–130. Rev. Mat. Iberoamericana, Madrid, 1997.
- [13] J. C. Pardo. The upper envelope of positive self-similar markov†processes. *Journal of Theoretical Probability*, 22(2):514–542, 2009.
- [14] J. C. Pardo, P. Patie, and M. Savov. A Wiener-Hopf type factorization for the exponential functional of Lévy processes. *Journal of the London Mathematical Society*, 86(3):930, 2012.
- [15] J.C. Pardo. On the future infimum of positive self-similar markov processes. *Stochastics*, 78(3):123–155, 2006.
- [16] J.C. Pardo and V. Rivero. Self-similar markov processes. *Boletín de la Sociedad Matemática Mexicana*.
- [17] J.C. Pardo, V. Rivero, and K. van Schaik. On the density of exponential functionals of lévy processes. *Bernoulli*, 19(5A):1938–1964, 11 2013.
- [18] P. Patie. Infinite divisibility of solutions to some self-similar integro-differential equations and exponential functionals of lévy processes. *Ann. Inst. H. Poincaré Probab. Statist.*, (3):667–684, 08.
- [19] P. Patie. Law of the absorption time of some positive self-similar Markov processes. *Ann. Probab.*, 40(2):765–787, 2012.
- [20] P. Patie and M. Savov. Extended factorizations of exponential functionals of Lévy processes. *Electron. J. Probab.*, 17:22, 2012.
- [21] P. Patie and M. Savov. Exponential functional of Lévy processes: Generalized Weierstrass products and wiener-hopf factorization. *Comptes Rendus Mathématique*, 351(9-10):393–396, 2013.
- [22] P. Patie and M. Savov. Spectral expansions of non self adjoint generalized Laguerre semigroups. *Preprint, arxiv 1506.01625v2*, 2015.
- [23] V. Rivero. Recurrent extensions of self-similar markov processes and Cramér’s condition. *Bernoulli*, 11(3):471–509, 2005.
- [24] V. Rivero. Recurrent extensions of self-similar markov processes and Cramér’s condition II. *Bernoulli*, 13(4):1053–1070, 2007.
- [25] V. Rivero. Tail asymptotics for exponential functionals of lévy processes: The convolution equivalent case. *Ann. Inst. H. Poincaré Probab. Statist.*, 48(4): 1081–1102, 2012.
- [26] K. Sato. *Lévy Processes and Infinitely Divisible Distributions*. Cambridge studies in advanced mathematics, 1999.
- [27] K. Sato and M. Yamazato. On distribution functions of class I. *Z. Wahrscheinlichkeitstheorie*, 43:273–308, 1978.
- [28] A. Singh. Rates of convergence of a transient diffusion in a spectrally negative lévy potential. *Ann. Probab.*, 36: 279–318, 2008.
- [29] G. Véchambre. Almost sure behavior for the local time of a diffusion in a spectrally negative Lévy environment. *Preprint*, 2016+.
- [30] G. Véchambre. Path decomposition of spectrally negative Lévy processes, and application to the local time of a diffusion in those environments. *Preprint, arxiv 1605.05084*, 2016.

UNIVERSITÉ D’ORLÉANS, LABORATOIRE MAPMO - FÉDÉRATION DENIS POISSON, BÂTIMENT DE MATHÉMATIQUES - RUE DE CHARTRES B.P. 6759 - 45067 ORLÉANS CEDEX 2, FRANCE

E-mail address: Gregoire.Vechambre@ens-rennes.fr