



**HAL**  
open science

## Supplement to ” Non-asymptotic convergence analysis for the Unadjusted Langevin Algorithm ”

Alain Durmus, Eric Moulines

► **To cite this version:**

Alain Durmus, Eric Moulines. Supplement to ” Non-asymptotic convergence analysis for the Unadjusted Langevin Algorithm ”. 2015. hal-01176084v1

**HAL Id: hal-01176084**

**<https://hal.science/hal-01176084v1>**

Preprint submitted on 14 Jul 2015 (v1), last revised 15 Jul 2018 (v5)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Supplement to “Non-asymptotic convergence analysis for the Unadjusted Langevin Algorithm”

Alain Durmus <sup>\*</sup> and Éric Moulines <sup>\*\*</sup>

*LTCI, Telecom ParisTech & CNRS, 46 rue Barrault, 75634 Paris Cedex 13, France,*  
*e-mail: <sup>\*</sup>alain.durmus@telecom-paristech.fr; <sup>\*\*</sup>eric.moulines@telecom-paristech.fr*

## 1. Discussion of Theorem 16

Note that

$$u_n^{(2)}(\gamma) \leq \sum_{i=1}^n \{A_0 \gamma_i^2 + A_1 \gamma_i^3\} \prod_{k=i+1}^n (1 - \kappa \gamma_k / 2), \quad (\text{S1})$$

where  $\kappa$  is given by (12), and

$$A_0 \stackrel{\text{def}}{=} 2L^2 \kappa^{-1} d, \quad (\text{S2})$$

$$A_1 \stackrel{\text{def}}{=} 2L^2 \kappa^{-1} d(m+L)^{-1} + dL^4 (\kappa^{-1} + (m+L)^{-1})(m^{-1} + 6^{-1}(m+L)^{-1}). \quad (\text{S3})$$

If  $(\gamma_k)_{k \geq 1}$  is a constant step size,  $\gamma_k = \gamma$  for all  $k \geq 1$ , then a straightforward consequence of Theorem 16 and (S1) is the following result, which gives the minimal number of iterations  $n$  and a step-size  $\gamma$  to get  $W_2(\delta_{x^*} Q_n^\gamma, \pi)$  smaller than  $\epsilon > 0$ .

**Corollary S1** (of Theorem 16). *Assume **H1** and **H3**. Let  $x^*$  be the unique minimizer of  $U$ . Let  $x \in \mathbb{R}^d$  and  $\epsilon > 0$ . Set for all  $k \in \mathbb{N}$ ,  $\gamma_k = \gamma$  with*

$$\gamma = \frac{-2A_0 \kappa^{-1} + (4\kappa^{-1}(A_0^2 \kappa^{-1} + \epsilon^2 A_1))^{1/2}}{4A_1 \kappa^{-1}} \wedge (m+L)^{-1}, \quad (\text{S4})$$

$$n = \lceil 2(\kappa\gamma)^{-1} \{-\log(\epsilon^2/4) + \log(2d/m)\} \rceil. \quad (\text{S5})$$

Then  $W_2(\delta_{x^*} Q_n^\gamma, \pi) \leq \epsilon$ .

Note that if  $\gamma$  is given by (S4), and is different from  $1/(m+L)$ , then  $\gamma \leq \epsilon(4A_1 \kappa^{-1})^{-1/2}$  and  $2\kappa^{-1}(A_0 \gamma + A_1 \gamma^2) = \epsilon^2/2$ . Therefore,

$$\gamma \geq (\epsilon^2 \kappa / 4) \left\{ A_0 + \epsilon (A_1 / (4\kappa))^{1/2} \right\}^{-1}.$$

It is shown in [S1, Corollary 1] that under **H3**, for constant step size for any  $\epsilon > 0$ , we can choose  $\gamma$  and  $n \geq 1$  such that if for all  $k \geq 1$ ,  $\gamma_k = \gamma$ , then  $\|\nu^* Q_n^\gamma - \pi\|_{\text{TV}} \leq \epsilon$  where  $\nu^*$  is the Gaussian measure on  $\mathbb{R}^d$  with mean  $x^*$  and covariance matrix  $L^{-1} \mathbf{I}_d$ . We stress that the results in [S1, corollary 1] hold only for a particular choice of the initial distribution  $\nu^*$ , (which might seem a rather artificial assumption) whereas Theorem 16 hold for any initial distribution in  $\mathcal{P}_2(\mathbb{R}^d)$ .

We compare the optimal value of  $\gamma$  and  $n$  obtained from Corollary S1 with those given in [S1, Corollary 1]. This comparison is summarized in Table 1 and Table 2; for simplicity, we provide only the dependencies of the optimal stepsize  $\gamma$  and minimal number of simulations  $n$  as a function of the dimension  $d$ , the precision  $\epsilon$  and the constants  $m, L$ . It can be seen that the dependency on the dimension is significantly better than those in [S1, Corollary 1].

Parameter	$d$	$\epsilon$	$L$	$m$
Theorem 16 and (30)	$d^{-1}$	$\epsilon^2$	$L^{-2}$	$m^2$
[S1, Corollary 1]	$d^{-2}$	$\epsilon^2$	$L^{-2}$	$m$

TABLE 1

Dependencies of  $\gamma$ 

Parameter	$d$	$\epsilon$	$L$	$m$
Theorem 16 and (30)	$d \log(d)$	$\epsilon^{-2}  \log(\epsilon) $	$L^2$	$ \log(m)  m^{-3}$
[S1, Corollary 1]	$d^3$	$\epsilon^{-2}  \log(\epsilon) $	$L^3$	$ \log(m)  m^{-2}$

TABLE 2

Dependencies of  $n$ 

### 1.1. Explicit bounds for $\gamma_k = \gamma_1 k^\alpha$ with $\alpha \in (0, 1]$

We give here a bound on the sequences  $(u_n^{(1)}(\gamma))_{n \geq 0}$  and  $(u_n^{(2)}(\gamma))_{n \geq 0}$  for  $(\gamma_k)_{k \geq 1}$  defined by  $\gamma_1 < 1/(m+L)$  and  $\gamma_k = \gamma_1 k^{-\alpha}$  for  $\alpha \in (0, 1]$ . Also for that purpose we introduce for  $t \in \mathbb{R}_+^*$ ,

$$\psi_\beta(t) = \begin{cases} (t^\beta - 1)/\beta & \text{for } \beta \neq 0 \\ \log(t) & \text{for } \beta = 0. \end{cases} \quad (\text{S6})$$

We easily get for  $a \geq 0$  that for all  $n, p \geq 1$ ,  $n \leq p$

$$\psi_{1-a}(p+1) - \psi_{1-a}(n) \leq \sum_{k=n}^p k^{-a} \leq \psi_{1-a}(p) - \psi_{1-a}(n) + 1, \quad (\text{S7})$$

and for  $a \in \mathbb{R}$

$$\sum_{k=n}^p k^{-a} \leq \psi_{1-a}(p+1) - \psi_{1-a}(n) + 1. \quad (\text{S8})$$

1. For  $\alpha = 1$ , using that for all  $t \in \mathbb{R}$ ,  $(1+t) \leq e^t$  and by (S7) and (S8), we have

$$u_n^{(1)}(\gamma) \leq (n+1)^{-\kappa\gamma_1/2}, \quad u_n^{(2)}(\gamma) \leq (n+1)^{-\kappa\gamma_1/2} \sum_{j=0}^1 \mathbf{A}_j(\psi_{\kappa\gamma_1/2-1-j}(n+1) + 1).$$

2. For  $\alpha \in (0, 1)$ , by (S7) and Lemma 39 applied with  $\ell = \lceil n/2 \rceil$ , where  $\lceil \cdot \rceil$  is the ceiling function, we have

$$\begin{aligned} u_n^{(1)}(\gamma) &\leq \exp(-\kappa\gamma_1\psi_{1-\alpha}(n+1)/2) \\ u_n^{(2)}(\gamma) &\leq \sum_{j=0}^1 \mathbf{A}_j \left( 2\kappa^{-1}\gamma_1^{j+1}(n/2)^{-\alpha(j+1)} + \gamma_1^{j+2} (\psi_{1-\alpha(j+2)}(\lceil n/2 \rceil) + 1) \right) \\ &\quad \times \exp\{-(\kappa\gamma_1/2)(\psi_{1-\alpha}(n+1) - \psi_{1-\alpha}(\lceil n/2 \rceil))\}. \end{aligned} \quad (\text{S9})$$

### 1.2. Optimal strategy with a fixed number of iterations

**Corollary S2.** Assume **H1**, **H3**, and  $(\gamma_k)_{k \geq 1}$  is a constant sequence,  $\gamma_k = \gamma$  for all  $k \geq 1$ . Let  $n \geq 1$ , and set

$$\begin{aligned}\gamma^+ &\stackrel{\text{def}}{=} 2(\kappa n)^{-1}(\log(\kappa n/2) + \log(2(\|x - x^*\|^2 + d/m)) - \log(2\kappa^{-1}\mathbf{A}_0)) \\ \gamma_- &\stackrel{\text{def}}{=} 2(\kappa n)^{-1}(\log(\kappa n/2) + \log(2(\|x - x^*\|^2 + d/m)) - \log(2\kappa^{-1}(\mathbf{A}_0 + 2\mathbf{A}_1(m+L)^{-1}))) .\end{aligned}$$

Assume  $\gamma^+ \in (0, (m+L)^{-1})$ . Then, the optimal choice of  $\gamma$  to minimize the bound on  $W_2(\delta_x Q_n^\gamma, \pi)$  given by Theorem 16 belongs to  $[\gamma_-, \gamma^+]$ . Moreover if  $\gamma = \gamma_+$ , then the bound on  $W_2^2(\delta_x Q_n^\gamma, \pi)$  is equivalent to  $4\mathbf{A}_0(\kappa^2 n)^{-1} \log(\kappa n/2)$  as  $n \rightarrow +\infty$ .

Similarly, we have the following result.

**Corollary S3.** Assume **H1** and **H3**. Let  $(\gamma_k)_{k \geq 1}$  be the decreasing sequence, defined by  $\gamma_k = \gamma_\alpha k^{-\alpha}$ , with  $\alpha \in (0, 1)$ . Let  $n \geq 1$  and set

$$\gamma_\alpha \stackrel{\text{def}}{=} 2(1-\alpha)\kappa^{-1}(2/n)^{1-\alpha} \log(\kappa n/(2(1-\alpha))) .$$

Assume  $\gamma_\alpha \in (0, (m+L)^{-1})$ . Then the bound on  $W_2^2(\delta_x Q_n^\gamma, \pi)$  given by Theorem 16 is smaller for large  $n$  than  $8(1-\alpha)\mathbf{A}_0(\kappa^2 n)^{-1} \log(\kappa n/(2(1-\alpha)))$ .

*Proof.* Follows from (S1), (S9) and the choice of  $\gamma_\alpha$ .  $\square$

## 2. Generalization of Theorem 16

In this section, we weaken the assumption  $\gamma_1 \leq 1/(m+L)$  of Theorem 16. We assume now:

**G1.** The sequence  $(\gamma_k)_{k \geq 1}$  is non-increasing, and there exists  $\rho > 0$  and  $n_1$  such that  $(1+\rho)\gamma_{n_1} \leq 2/(m+L)$ .

Under **G1**, we denote by

$$n_0 \stackrel{\text{def}}{=} \min \{k \in \mathbb{N} \mid \gamma_k \leq 2/(m+L)\} \tag{S10}$$

We first give an extension of Theorem 14. Denote in the sequel  $(\cdot)_+ = \max(\cdot, 0)$ . Recall that under **H3**,  $x^*$  is the unique minimizer of  $U$ , and  $\kappa$  is defined in (14)

**Theorem S4.** Assume **H1**, **H3** and **G1**. Then for all  $n, p \in \mathbb{N}^*$ ,  $n \leq p$ ,

$$\int_{\mathbb{R}^d} \|x - x^*\|^2 \mu_0 Q_n^p(dx) \leq E_{n,p}(\mu_0, \gamma) ,$$

where

$$\begin{aligned}E_{n,p}(\mu_0, \gamma) &\stackrel{\text{def}}{=} \exp\left(-\sum_{k=n}^p \gamma_k \kappa + \sum_{k=n}^{n_0-1} L^2 \gamma_k^2\right) \int_{\mathbb{R}^d} \|x - x^*\|^2 \mu_0(dx) \\ &+ 2d\kappa^{-1} + 2d \left\{ \prod_{k=n}^{n_0-1} (\gamma_{n_0-1} L^2)^{-1} (1 + L^2 \gamma_k^2) \right\} \exp\left(-\sum_{k=n}^p \kappa \gamma_k + \sum_{k=n}^{n_0-1} \gamma_k^2 m L\right) .\end{aligned} \tag{S11}$$

*Proof.* For any  $\gamma > 0$ , we have for all  $x \in \mathbb{R}^d$ :

$$\int_{\mathbb{R}^d} \|y - x^*\|^2 R_\gamma(x, dy) = \|x - \gamma \nabla U(x) - x^*\|^2 + 2\gamma d.$$

Using that  $\nabla U(x^*) = 0$ , (12) and **H1**, we get from the previous inequality:

$$\begin{aligned} \int_{\mathbb{R}^d} \|y - x^*\|^2 R_\gamma(x, dy) &\leq (1 - \kappa\gamma) \|x - x^*\|^2 + \gamma \left( \gamma - \frac{2}{m+L} \right) \|\nabla U(x) - \nabla U(x^*)\|^2 + 2\gamma d \\ &\leq \eta(\gamma) \|x - x^*\|^2 + 2\gamma d, \end{aligned}$$

where  $\eta(\gamma) = (1 - \kappa\gamma + \gamma L(\gamma - 2/(m+L)))_+$ . Denote for all  $k \geq 1$ ,  $\eta_k = \eta(\gamma_k)$ . By a straightforward induction, we have by definition of  $Q_n^p$  for  $p, n \in \mathbb{N}$ ,  $p \leq n$ ,

$$\int_{\mathbb{R}^d} \|x - x^*\|^2 \mu_0 Q_n^p(dx) \leq \prod_{k=n}^p \eta_k \int_{\mathbb{R}^d} \|x - x^*\| \mu_0(dx) + (2d) \sum_{i=n}^p \prod_{k=i+1}^p \eta_k \gamma_i, \quad (\text{S12})$$

with the convention that for  $n, p \in \mathbb{N}$ ,  $n < p$ ,  $\prod_p^n = 1$ . For the first term of the right hand side, we simply use the bound, for all  $x \in \mathbb{R}$ ,  $(1+x) \leq e^x$ , and we get by **G1**

$$\prod_{k=n}^p \eta_k \leq \exp \left( - \sum_{k=n}^p \kappa \gamma_k + \sum_{k=n}^{n_0-1} L^2 \gamma_k^2 \right), \quad (\text{S13})$$

where  $n_0$  is defined in (S10). Consider now the second term in the right hand side of (S12).

$$\begin{aligned} \sum_{i=n}^p \prod_{k=i+1}^p \eta_k \gamma_i &\leq \sum_{i=n_0}^p \prod_{k=i+1}^p (1 - \kappa \gamma_k) \gamma_i + \sum_{i=n}^{n_0-1} \prod_{k=i+1}^p \eta_k \gamma_i \\ &\leq \kappa^{-1} \sum_{i=n_0}^p \left\{ \prod_{k=i+1}^p (1 - \kappa \gamma_k) - \prod_{k=i}^p (1 - \kappa \gamma_k) \right\} + \left\{ \sum_{i=n}^{n_0-1} \prod_{k=i+1}^{n_0-1} (1 + L^2 \gamma_k^2) \gamma_i \right\} \prod_{k=n_0}^p (1 - \kappa \gamma_k) \end{aligned} \quad (\text{S14})$$

Since  $(\gamma_k)_{k \geq 1}$  is nonincreasing, we have

$$\begin{aligned} \sum_{i=n}^{n_0-1} \prod_{k=i+1}^{n_0-1} (1 + L^2 \gamma_k^2) \gamma_i &= \sum_{i=n}^{n_0-1} (\gamma_i L^2)^{-1} \left\{ \prod_{k=i}^{n_0-1} (1 + L^2 \gamma_k^2) - \prod_{k=i+1}^{n_0-1} (1 + L^2 \gamma_k^2) \right\} \\ &\leq \prod_{k=n}^{n_0-1} (\gamma_{n_0-1} L^2)^{-1} (1 + L^2 \gamma_k^2). \end{aligned}$$

Furthermore for  $k < n_0$   $\gamma_k > 2/(m+L)$ . This implies with the bound  $(1+x) \leq e^x$  on  $\mathbb{R}$ :

$$\prod_{k=n_0}^p (1 - \kappa \gamma_k) \leq \exp \left( - \sum_{k=n_0}^p \kappa \gamma_k \right) \exp \left( \sum_{k=n_0}^{n_0-1} \kappa \gamma_k \right) \leq \exp \left( - \sum_{k=n_0}^p \kappa \gamma_k \right) \exp \left( \sum_{k=n_0}^{n_0-1} \gamma_k^2 m L \right).$$

Using the two previous inequalities in (S14), we get

$$\sum_{i=n}^p \prod_{k=i+1}^p \eta_k \gamma_i \leq \kappa^{-1} + \left\{ \prod_{k=n}^{n_0-1} (\gamma_{n_0-1} L^2)^{-1} (1 + L^2 \gamma_k^2) \right\} \exp \left( - \sum_{k=n}^p \kappa \gamma_k + \sum_{k=n}^{n_0-1} \gamma_k^2 m L \right). \quad (\text{S15})$$

Combining (S13) and (S15) in (S12) concluded the proof.  $\square$

We now deal with bounds on  $W_2(\mu_0 Q_\gamma^n, \pi)$  under **G1**. But before we preface our result by some technical lemmas.

**Lemma S5.** *Assume **H1** and **H3**. Let  $\zeta_0 \in \mathcal{P}_2(\mathbb{R}^d \times \mathbb{R}^d)$ ,  $(Y_t, \bar{Y}_t)_{t \geq 0}$  such that  $(Y_0, \bar{Y}_0)$  is distributed according to  $\zeta_0$  and given by (21). Let  $(\mathcal{F}_t)_{t \geq 0}$  be the filtration associated with  $(B_t)_{t \geq 0}$  with  $\mathcal{F}_0$ , the  $\sigma$ -field generated by  $(Y_0, \bar{Y}_0)$ . Then for all  $n \geq 0$ ,  $\epsilon_1 > 0$  and  $\epsilon_2 > 0$ ,*

$$\begin{aligned} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \left\| Y_{\Gamma_{n+1}} - \bar{Y}_{\Gamma_{n+1}} \right\|^2 \right] &\leq \{1 - \gamma_{n+1}(\kappa - 2\epsilon_1) + \gamma_{n+1}L((1 + \epsilon_2)\gamma_{n+1} - 2/(m + L))_+\} \|Y_{\Gamma_n} - \bar{Y}_{\Gamma_n}\|^2 \\ &\quad + \gamma_{n+1}^2(1/(2\epsilon_1) + (1 + \epsilon_2^{-1})\gamma_{n+1}) \left( dL^2 + (L^4\gamma_{n+1}/2) \|Y_{\Gamma_n} - x^*\|^2 + dL^4\gamma_{n+1}^2/12 \right). \end{aligned}$$

*Proof.* Let  $n \geq 0$  and  $\epsilon_1 > 0$ , and set  $\Delta_n = Y_{\Gamma_n} - \bar{Y}_{\Gamma_n}$  by definition we have:

$$\begin{aligned} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \|\Delta_{n+1}\|^2 \right] &= \|\Delta_n\|^2 + \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \left\| \int_{\Gamma_n}^{\Gamma_{n+1}} \{ \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}) \} ds \right\|^2 \right] \\ &\quad - 2\gamma_{n+1} \langle \Delta_n, \nabla U(Y_{\Gamma_n}) - \nabla U(\bar{Y}_{\Gamma_n}) \rangle - 2 \int_{\Gamma_n}^{\Gamma_{n+1}} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} [\langle \Delta_n, \{ \nabla U(Y_s) - \nabla U(Y_{\Gamma_n}) \} \rangle ds]. \end{aligned}$$

Using the two inequalities  $|\langle a, b \rangle| \leq \epsilon_1 \|a\|^2 + (4\epsilon_1)^{-1} \|b\|^2$  and (12), we get

$$\begin{aligned} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \|\Delta_{n+1}\|^2 \right] &\leq \{1 - \gamma_{n+1}(\kappa - 2\epsilon_1)\} \|\Delta_n\|^2 - 2\gamma_{n+1}/(m + L) \|\nabla U(Y_{\Gamma_n}) - \nabla U(\bar{Y}_{\Gamma_n})\|^2 \\ &\quad + \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \left\| \int_{\Gamma_n}^{\Gamma_{n+1}} \{ \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}) \} ds \right\|^2 \right] + \frac{1}{2\epsilon_1} \int_{\Gamma_n}^{\Gamma_{n+1}} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \|\nabla U(Y_s) - \nabla U(Y_{\Gamma_n})\|^2 \right] ds. \end{aligned}$$

Using  $\|a + b\|^2 \leq (1 + \epsilon_2) \|a\|^2 + (1 + \epsilon_2^{-1}) \|b\|^2$  and the Jensen's inequality, we have

$$\begin{aligned} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \left\| \int_{\Gamma_n}^{\Gamma_{n+1}} \{ \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}) \} ds \right\|^2 \right] &\leq (1 + \epsilon_2)\gamma_{n+1}^2 \|\nabla U(Y_{\Gamma_n}) - \nabla U(\bar{Y}_{\Gamma_n})\|^2 \\ &\quad + \gamma_{n+1}(1 + \epsilon_2^{-1}) \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \int_{\Gamma_n}^{\Gamma_{n+1}} \|\nabla U(Y_s) - \nabla U(Y_{\Gamma_n})\|^2 ds \right]. \end{aligned}$$

This result and **H1** imply,

$$\begin{aligned} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \|\Delta_{n+1}\|^2 \right] &\leq \{1 - \gamma_{n+1}(\kappa - 2\epsilon_1) + \gamma_{n+1}L((1 + \epsilon_2)\gamma_{n+1} - 2/(m + L))_+\} \|\Delta_n\|^2 \\ &\quad + ((1 + \epsilon_2^{-1})\gamma_{n+1} + (2\epsilon_1)^{-1}) \int_{\Gamma_n}^{\Gamma_{n+1}} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \|\nabla U(Y_s) - \nabla U(Y_{\Gamma_n})\|^2 \right] ds. \quad (\text{S16}) \end{aligned}$$

By **H1**, the Markov property of  $(Y_t)_{t \geq 0}$  and (66), we have

$$\int_{\Gamma_n}^{\Gamma_{n+1}} \mathbb{E}^{\mathcal{F}_{\Gamma_n}} \left[ \|\nabla U(Y_s) - \nabla U(Y_{\Gamma_n})\|^2 \right] ds \leq L^2 \left( d\gamma_{n+1}^2 + dL^2\gamma_{n+1}^4/12 + (L^2\gamma_{n+1}^3/2) \|Y_{\Gamma_n} - x^*\|^2 \right).$$

Plugging this bound in (S16) concludes the proof.  $\square$

**Lemma S6.** Let  $(\gamma_k)_{k \geq 1}$  be a nonincreasing sequence of positive numbers. Let  $\varpi, \beta > 0$  be positive constants satisfying  $\varpi^2 \leq 4\beta$  and  $\tau > 0$ . Assume there exists  $N \geq 1$ ,  $\gamma_N \leq \tau$  and  $\gamma_N \varpi \leq 1$ . Then for all  $n \geq 0$ ,  $j \geq 2$

(i)

$$\begin{aligned} \sum_{i=1}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi + \gamma_k \beta (\gamma_k - \tau)_+) \gamma_i^j &\leq \sum_{i=N}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi) \gamma_i^j \\ &\quad + \left\{ \beta^{-1} \gamma_1^{j-2} \prod_{k=1}^{N-1} (1 + \gamma_k^2 \beta) \right\} \prod_{k=N}^{n+1} (1 - \varpi \gamma_k) . \end{aligned}$$

(ii) For all  $\ell \in \{N, \dots, n\}$ ,

$$\sum_{i=N}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi) \gamma_i^j \leq \exp \left( - \sum_{k=\ell}^{n+1} \varpi \gamma_k \right) \sum_{i=N}^{\ell-1} \gamma_i^j + \frac{\gamma_\ell^{j-1}}{\varpi} .$$

*Proof.* By definition of  $N$  we have

$$\begin{aligned} &\sum_{i=1}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi + \gamma_k \beta (\gamma_k - \tau)_+) \gamma_i^j \\ &\leq \sum_{i=N}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi) \gamma_i^j + \left\{ \sum_{i=1}^{N-1} \prod_{k=i+1}^{N-1} (1 + \gamma_k^2 \beta) \gamma_i^j \right\} \prod_{k=N}^{n+1} (1 - \gamma_k \varpi) . \end{aligned} \quad (\text{S17})$$

Using that  $(\gamma_k)_{k \geq 1}$  is nonincreasing, we have

$$\begin{aligned} \sum_{i=1}^{N-1} \prod_{k=i+1}^{N-1} (1 + \gamma_k^2 \beta) \gamma_i^j &\leq \sum_{i=1}^{N-1} \frac{\gamma_i^{j-2}}{\beta} \left\{ \prod_{k=i}^{N-1} (1 + \gamma_k^2 \beta) - \prod_{k=i+1}^{N-1} (1 + \gamma_k^2 \beta) \right\} \\ &\leq \beta^{-1} \gamma_1^{j-2} \prod_{k=1}^{N-1} (1 + \gamma_k^2 \beta) . \end{aligned}$$

Plugging this inequality in (S17) concludes the proof of (i). Let  $\ell \in \{N, \dots, n+1\}$ . Since  $(\gamma_k)_{k \geq 1}$  is nonincreasing and for every  $x \in \mathbb{R}$ ,  $(1+x) \leq e^x$ , we get

$$\begin{aligned} \sum_{i=N}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi) \gamma_i^j &= \sum_{i=N}^{\ell-1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi) \gamma_i^j + \sum_{i=\ell}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi) \gamma_i^j \\ &\leq \sum_{i=N}^{\ell-1} \exp \left( - \sum_{k=i+1}^{n+1} \varpi \gamma_k \right) \gamma_i^j + \gamma_\ell^{j-1} \sum_{i=\ell}^{n+1} \prod_{k=i+1}^{n+1} (1 - \gamma_k \varpi) \gamma_i \\ &\leq \exp \left( - \sum_{k=\ell}^{n+1} \varpi \gamma_k \right) \sum_{i=N}^{\ell-1} \gamma_i^j + \frac{\gamma_\ell^{j-1}}{\varpi} . \end{aligned}$$

□

**Lemma S7.** Let  $(\gamma_k)_{k \geq 1}$  be a nonincreasing sequence of positive numbers,  $\varpi, \beta, \tau > 0$  be positive real numbers, and  $N \geq 1$  satisfying the assumptions of Lemma S6. Let  $\mathbf{P} \in \mathbb{N}^*$ ,  $C_i \geq 0$ ,  $i = 0, \dots, \mathbf{P}$  be positive constants and  $(u_n)_{n \geq 0}$  be a sequence of real numbers with  $u_0 \geq 0$  satisfying for all  $n \geq 0$

$$u_{n+1} \leq (1 - \gamma_{n+1}\varpi + \beta\gamma_{n+1}(\gamma_{n+1} - \tau)_+)u_n + \sum_{i=0}^{\mathbf{P}} C_j \gamma_{n+1}^{j+2}.$$

Then for all  $n \geq 1$ ,

$$\begin{aligned} u_n \leq & \left\{ \prod_{k=1}^{N-1} (1 + \beta\gamma_k^2) \right\} \prod_{k=N}^n (1 - \gamma_k\varpi) u_0 + \sum_{j=0}^{\mathbf{P}} C_j \sum_{i=N}^n \prod_{k=i+1}^n (1 - \gamma_k\varpi) \gamma_i^{j+2} \\ & + \left\{ \sum_{j=0}^{\mathbf{P}} C_j \beta^{-1} \gamma_1^j \prod_{k=1}^{N-1} (1 + \gamma_k^2 \beta) \right\} \prod_{k=N}^n (1 - \varpi\gamma_k). \end{aligned}$$

*Proof.* This is a consequence of a straightforward induction and Lemma S6-(i).  $\square$

**Proposition S8.** Assume **H 1**, **H 3** and **G 1**. Let  $x^*$  be the unique minimizer of  $U$ . Let  $\zeta_0 \in \mathcal{P}_2(\mathbb{R}^d \times \mathbb{R}^d)$ ,  $(Y_t, \bar{Y}_t)_{t \geq 0}$  such that  $(Y_0, \bar{Y}_0)$  is distributed according to  $\zeta_0$  and given by (21). Then for all  $n \geq 0$  and  $t \in [\Gamma_n, \Gamma_{n+1}]$ :

$$\mathbb{E} \left[ \|Y_t - \bar{Y}_t\|^2 \right] \leq \tilde{u}_n^{(1)} \mathbb{E} \left[ \|Y_0 - \bar{Y}_0\|^2 \right] + \tilde{u}_n^{(4)} + \tilde{u}_{t,n}^{(5)},$$

where

$$\tilde{u}_n^{(1)}(\gamma) \stackrel{\text{def}}{=} \left\{ \prod_{k=1}^{n_1-1} (1 + L^2(1 + \rho)\gamma_k^2) \right\} \prod_{k=n_1}^n (1 - \kappa\gamma_k/2), \quad (\text{S18})$$

$$\begin{aligned} \tilde{u}_n^{(4)}(\gamma) & \stackrel{\text{def}}{=} \sum_{i=n_1}^n \gamma_i^2 f(\gamma_i) \prod_{k=i+1}^n (1 - \kappa\gamma_k/2) \\ & + f(\gamma_1)(L^2(1 + \rho))^{-1} \left\{ \prod_{k=1}^{n_1-1} (1 + \gamma_k^2(1 + \rho)L^2) \right\} \prod_{k=n_1}^n (1 - \kappa\gamma_k/2), \end{aligned}$$

where for  $\gamma > 0$ ,

$$\begin{aligned} f(\gamma) & = \{2\kappa^{-1} + (1 + \rho^{-1})\gamma\} (dL^2 + \delta L^4 \gamma/2 + dL^4 \gamma^2/12), \\ \delta & = \max_{i \geq 1} \left\{ e^{-2m\Gamma_{i-1}} \mathbb{E} \left[ \|Y_0 - x^*\|^2 \right] + (1 - e^{-2m\Gamma_{i-1}})(d/m) \right\}, \end{aligned}$$

and

$$\tilde{u}_{t,n}^{(5)}(\gamma) \stackrel{\text{def}}{=} \frac{m+L}{2} \left( \frac{(t - \Gamma_n)^3 L^2}{3} E_{1,n}(\mu_0, \gamma) + (t - \Gamma_n)^2 d \right),$$

where  $E_{1,n}(\mu_0, \gamma)$  is given by (S11) and  $\mu_0$  is the initial distribution of  $\bar{Y}_0$ .



*Proof.* Lemma S5 with  $\epsilon_1 = \kappa/4$  and  $\epsilon_2 = \rho$ , G1, Lemma S6, (65) imply for all  $n \geq 0$

$$\mathbb{E} \left[ \|Y_{\Gamma_n} - \bar{Y}_{\Gamma_n}\|^2 \right] \leq \tilde{u}_n^{(1)}(\gamma) \mathbb{E} \left[ \|Y_0 - \bar{Y}_0\|^2 \right] + \tilde{u}_n^{(4)}(\gamma). \quad (\text{S19})$$

Now let  $n \geq 0$  and  $t \in [\Gamma_n, \Gamma_{n+1}]$ . By (21),

$$\|Y_t - \bar{Y}_t\|^2 = \|Y_{\Gamma_n} - \bar{Y}_{\Gamma_n}\|^2 - 2 \int_{\Gamma_n}^t \langle \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}), Y_s - \bar{Y}_s \rangle ds. \quad (\text{S20})$$

Moreover for all  $s \in [\Gamma_n, \Gamma_{n+1}]$ , by (12) we get

$$\begin{aligned} \langle \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}), Y_s - \bar{Y}_s \rangle &= \langle \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}), Y_s - \bar{Y}_{\Gamma_n} + \bar{Y}_{\Gamma_n} - \bar{Y}_s \rangle \\ &\geq (m+L)^{-1} \|\nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n})\|^2 + \langle \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}), \bar{Y}_{\Gamma_n} - \bar{Y}_s \rangle. \end{aligned} \quad (\text{S21})$$

Since  $|\langle a, b \rangle| \leq (m+L)^{-1} \|a\|^2 + (m+L) \|b\|^2/4$ , we have

$$\begin{aligned} \langle \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}), \bar{Y}_{\Gamma_n} - \bar{Y}_s \rangle &\geq \\ &- (m+L)^{-1} \|\nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n})\|^2 - (m+L) \|\bar{Y}_s - \bar{Y}_{\Gamma_n}\|^2/4. \end{aligned}$$

Using this inequality in (S21), we get

$$\langle \nabla U(Y_s) - \nabla U(\bar{Y}_{\Gamma_n}), Y_s - \bar{Y}_s \rangle \geq -(m+L) \|\bar{Y}_s - \bar{Y}_{\Gamma_n}\|^2/4,$$

and (S20) becomes

$$\|Y_t - \bar{Y}_t\|^2 \leq \|Y_{\Gamma_n} - \bar{Y}_{\Gamma_n}\|^2 + ((m+L)/2) \int_{\Gamma_n}^t \|\bar{Y}_s - \bar{Y}_{\Gamma_n}\|^2 ds \quad (\text{S22})$$

Therefore, by the previous inequality, it remains to bound the expectation of  $\|\bar{Y}_s - \bar{Y}_{\Gamma_n}\|^2$ . By (21) and using  $\nabla U(x^*) = 0$ ,

$$\|\bar{Y}_s - \bar{Y}_{\Gamma_n}\|^2 = \left\| -(s - \Gamma_n)(\nabla U(\bar{Y}_{\Gamma_n}) - \nabla U(x^*)) + \sqrt{2}(B_s - B_{\Gamma_n}) \right\|^2.$$

Then taking the expectation, using the Markov property of  $(B_t)_{t \geq 0}$  and H1, we have

$$\mathbb{E} \left[ \|\bar{Y}_s - \bar{Y}_{\Gamma_n}\|^2 \right] \leq (s - \Gamma_n)^2 L^2 \mathbb{E} \left[ \|\bar{Y}_{\Gamma_n} - x^*\|^2 \right] + 2(s - \Gamma_n)d. \quad (\text{S23})$$

The proof follows from taking the expectation in (S22), combining (S19)-(S23), and using Theorem S4.  $\square$

**Theorem S9.** Assume H1, H3 and G1. Then for all  $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$  and  $n \geq 1$ ,

$$W_2^2(\mu_0 Q_\gamma^n, \pi) \leq \tilde{u}_n^{(1)}(\gamma) W_2^2(\mu_0, \pi) + \tilde{u}_n^{(2)}(\gamma), \quad (\text{S24})$$

where  $(\tilde{u}_n^{(1)})_{n \geq 0}$  is given by (S18) and

$$\begin{aligned} \tilde{u}_n^{(2)}(\gamma) &\stackrel{\text{def}}{=} \sum_{i=n_1}^n \gamma_i^2 \phi(\gamma_i) \prod_{k=i+1}^n (1 - \kappa \gamma_k/2) \\ &\quad + \phi(\gamma_1) (L^2(1 + \rho))^{-1} \left\{ \prod_{k=1}^{n_1-1} (1 + \gamma_k^2(1 + \rho)L^2) \right\} \prod_{k=n_1}^n (1 - \kappa \gamma_k/2), \end{aligned} \quad (\text{S25})$$

where

$$\phi(\gamma) = \{2\kappa^{-1} + (1 + \rho^{-1})\gamma\} (dL^2 + dL^4\gamma/(2m) + dL^4\gamma^2/12).$$

*Proof.* Let  $\zeta_0$  be an optimal transference plan of  $\mu_0$  and  $\pi$ . Let  $(Y_t, \bar{Y}_t)_{t \geq 0}$  with  $(Y_0, \bar{Y}_0)$  distributed according to  $\zeta_0$  and defined by (21). By definition of  $W_2$  and since for all  $t \geq 0$ ,  $\pi$  is invariant for  $P_t$ ,  $W_2^2(\mu_0 Q^n, \pi) \leq \mathbb{E}[\|Y_{\Gamma_n} - X_{\Gamma_n}\|^2]$ . Then the proof follows from Proposition S8 since  $Y_0$  is distributed according to  $\pi$  and by (16), which shows that  $\delta \leq d/m$ .  $\square$

### 2.1. Explicit bound based on Theorem S9 for $\gamma_k = \gamma_1 k^\alpha$ with $\alpha \in (0, 1]$

We give here a bound on the sequences  $(\tilde{u}_n^{(1)}(\gamma))_{n \geq 1}$  and  $(\tilde{u}_n^{(2)}(\gamma))_{n \geq 1}$  for  $(\gamma_k)_{k \geq 1}$  defined by  $\gamma_1 > 0$  and  $\gamma_k = \gamma_1 k^{-\alpha}$  for  $\alpha \in (0, 1]$ . Recall that  $\psi_\beta$  is given by (S6). First note, since  $(\gamma_k)_{k \geq 1}$  is nonincreasing, for all  $n \geq 1$ , we have

$$\begin{aligned} \tilde{u}_n^{(2)}(\gamma) &\leq \sum_{j=0}^1 \mathbf{B}_j \sum_{i=n_1}^n \gamma_i^{j+2} \prod_{k=i+1}^n (1 - \kappa\gamma_k/2) \\ &\quad + \sum_{j=0}^1 \mathbf{B}_j (L^2(1 + \rho))^{-1} \gamma_1^j \left\{ \prod_{k=1}^{n_1-1} (1 + \gamma_k^2(1 + \rho)L^2) \right\} \prod_{k=n_1}^n (1 - \kappa\gamma_k/2), \end{aligned} \quad (\text{S26})$$

where

$$\mathbf{B}_1 = \mathbf{b}dL^2, \mathbf{B}_2 = \mathbf{b}(dL^4/(2m) + \gamma_1 dL^4/12), \mathbf{b} = 2\kappa^{-1} + (1 + \rho^{-1})\gamma_1.$$

1. For  $\alpha = 1$  and  $n_1 = 1$ , by (S7) and (S8), we have

$$\begin{aligned} \tilde{u}_n^{(1)}(\gamma) &\leq (n+1)^{-\kappa\gamma_1/2} \\ \tilde{u}_n^{(2)}(\gamma) &\leq (n+1)^{-\kappa\gamma_1/2} \sum_{j=0}^1 \mathbf{B}_j \left\{ \gamma_1^{j+2} (\psi_{\kappa\gamma_1/2-1-j}(n+1) + 1) + (L^2(1 + \rho))^{-1} \gamma_1^j \right\}. \end{aligned}$$

For  $n_1 > 1$ , since  $(\gamma_k)_{k \geq 0}$  is non increasing, using again (S7), (S8), and the bound for  $t \in \mathbb{R}$ ,  $(1+t) \leq e^t$

$$\begin{aligned} \tilde{u}_n^{(1)}(\gamma) &\leq (n+1)^{-\kappa\gamma_1/2} \exp \left\{ \kappa\gamma_1 \psi_0(n_1)/2 + L^2(1 + \rho) \gamma_1^2 (\psi_{-1}(n_1 - 1) + 1) \right\} \\ \tilde{u}_n^{(2)}(\gamma) &\leq (n+1)^{-\kappa\gamma_1/2} \sum_{j=0}^1 \mathbf{B}_j \left( \gamma_1^{j+2} (\psi_{\kappa\gamma_1/2-1-j}(n+1) - \psi_{\kappa\gamma_1/2-1-j}(n_1) + 1) \right. \\ &\quad \left. + (\gamma_1^j / (L^2(1 + \rho))) \exp \left\{ \kappa\gamma_1 \psi_0(n_1)/2 + L^2(1 + \rho) \gamma_1^2 (\psi_{-1}(n_1 - 1) + 1) \right\} \right). \end{aligned}$$

Thus, for  $\gamma_1 > \kappa/2$ , we get a bound in  $\mathcal{O}(n^{-1})$ .

2. For  $\alpha \in (0, 1)$  and  $n_1 = 1$ , by (S7) and Lemma S6-(ii) applied with  $\ell = \lceil n/2 \rceil$ , we have

$$\begin{aligned} \tilde{u}_n^{(1)}(\gamma) &\leq \exp(-(\kappa\gamma_1/2)\psi_{\alpha-1}(n+1)) \\ \tilde{u}_n^{(2)}(\gamma) &\leq \sum_{j=0}^1 \mathbf{B}_j \left\{ \gamma_1^{j+2} (\psi_{1-\alpha(j+2)}(\lceil n/2 \rceil) + 1) \exp \left\{ -(\kappa\gamma_1/2)(\psi_{1-\alpha}(n+1) - \psi_{1-\alpha}(\lceil n/2 \rceil)) \right\} \right. \\ &\quad \left. + 2\kappa^{-1} \gamma_1^{j+1} (n/2)^{-\alpha(j+1)} + \gamma_1^j \exp \left\{ -(\kappa\gamma_1/2)\psi_{\alpha-1}(n+1) \right\} \right\}. \end{aligned}$$

For  $n_1 > 1$  and  $\lceil n/2 \rceil \geq n_1$ , since  $(\gamma_k)_{k \geq 0}$  is non increasing, using again (S7), and Lemma S6-(ii) applied with  $\ell = \lceil n/2 \rceil$ , and the bound for  $t \in \mathbb{R}$ ,  $(1+t) \leq e^t$ , we get

$$\begin{aligned} \tilde{u}_n^{(1)}(\gamma) &\leq \exp \left\{ -\kappa\gamma_1(\boldsymbol{\psi}_{\alpha-1}(n+1) - \boldsymbol{\psi}_{1-\alpha}(n_1))/2 + L^2(1+\rho)\gamma_1^2(\boldsymbol{\psi}_{1-2\alpha}(n_1-1) + 1) \right\} \\ \tilde{u}_n^{(2)}(\gamma) &\leq \sum_{j=0}^1 \mathbf{B}_j \left\{ 2\kappa^{-1}\gamma_1^{j+1}(n/2)^{-\alpha(j+1)} \right. \\ &\quad + \gamma_1^{j+2} (\boldsymbol{\psi}_{1-\alpha(j+2)}(\lceil n/2 \rceil) - \boldsymbol{\psi}_{1-\alpha(j+2)}(n_1) + 1) \exp \left\{ -(\kappa\gamma_1/2)(\boldsymbol{\psi}_{1-\alpha}(n+1) - \boldsymbol{\psi}_{1-\alpha}(\lceil n/2 \rceil)) \right\} \\ &\quad \left. + (\gamma_1^j/(L^2(1+\rho))) \exp \left\{ -\kappa\gamma_1(\boldsymbol{\psi}_{\alpha-1}(n+1) - \boldsymbol{\psi}_{1-\alpha}(n_1))/2 + L^2(1+\rho)\gamma_1^2(\boldsymbol{\psi}_{1-2\alpha}(n_1-1) + 1) \right\} \right\} . \end{aligned}$$

### 3. Explicit bound on the MSE under strongly convex assumptions for $\gamma_k = \gamma_1 k^{-\alpha}$ with $\alpha \in [0, 1]$

Without loss of generality, assume that  $\|f\|_{\text{Lip}} = 1$ . In the following, denote by  $\Omega(x) \stackrel{\text{def}}{=} \|x - x^*\|^2 + d/m$  and  $C$  a constant (which may take different values upon each appearance), which does not depend on  $m, L, \gamma_1, \alpha$  and  $\|x - x^*\|$ .

1. First for  $\alpha = 0$ , recall by (S1), (29) and (30) we have for all  $p \geq 1$ ,

$$u_p^{(1)}(\gamma)W_2^2(\delta_x, \pi) + u_p^{(2)}(\gamma) \leq 2\Omega(x)(1 - \kappa\gamma_1/2)^p + 2\kappa^{-1}(\mathbf{A}_0\gamma_1 + \mathbf{A}_1\gamma_1^2),$$

where  $\mathbf{A}_0$  and  $\mathbf{A}_1$  are given by (S2) and (S3) respectively. So by Proposition 18 and Lemma 39, we have the following bound for the bias

$$\left\{ \mathbb{E}_x[\hat{\pi}_n^N(f)] - \pi(f) \right\}^2 \leq C \left( \frac{\kappa^{-1} \exp(-\kappa N \gamma_1/2) \Omega(x)}{\gamma_1 n} + \kappa^{-1} \mathbf{A}_0 \gamma_1 \right).$$

Therefore plugging this inequality and the one given by Theorem 24 in (32) implies:

$$\text{MSE}_f(N, n) \leq C \left( \kappa^{-1} \mathbf{A}_0 \gamma_1 + \frac{\kappa^{-2} + \kappa^{-1} \exp(-\kappa N \gamma_1/2) \Omega(x)}{n \gamma_1} \right). \quad (\text{S27})$$

So with fixed  $\gamma_1$  this bound is of order  $\gamma_1$ . If we fix the number of iterations  $n$ , we can optimize the choice of  $\gamma_1$ . Set

$$\gamma_{*,0}(n) = (\kappa^{-1} \mathbf{A}_0)^{-1} (C_{\text{MSE},0}/n)^{1/2}, \text{ where } C_{\text{MSE},0} \stackrel{\text{def}}{=} \kappa^{-3} \mathbf{A}_0,$$

and (S27) becomes if  $\gamma_1 \leftarrow \gamma_{*,0}(n)$ ,

$$\text{MSE}_f(N, n) \leq C (C_{\text{MSE},0} n)^{-1/2} (\kappa^{-1} \exp(-\kappa N \gamma_{*,0}(n)/2) \Omega(x) + C_{\text{MSE},0}).$$

Setting  $N_0(n) = 2(\kappa \gamma_{*,0}(n))^{-1} \log(\Omega(x))$ , we end up with

$$\text{MSE}_f(N_0(n), n) \leq C (C_{\text{MSE},0}/n)^{1/2}.$$

Note that  $N_0(n)$  is of order  $n^{1/2}$ .

2. For  $\alpha \in (0, 1/2)$ , recall by (S1), (29) and (30) we have for all  $p \geq 1$ ,

$$u_p^{(1)}(\gamma)W_2^2(\delta_x, \pi) \leq 2\Omega(x) \prod_{i=1}^p (1 - \kappa\gamma_i/2). \quad (\text{S28})$$

So by Proposition 18, Lemma 39, (S7) and (S9), we have the following bound for the bias

$$\{\mathbb{E}_x[\hat{\pi}_n^N(f)] - \pi(f)\}^2 \leq C \left( \frac{\kappa^{-1}\mathbf{A}_0\gamma_1}{(1-2\alpha)n^\alpha} + \frac{\kappa^{-1} \exp\{-\kappa\gamma_1 N^{1-\alpha}/(2(1-\alpha))\} \Omega(x)}{\gamma_1 n^{1-\alpha}} \right).$$

Plugging this inequality and the one given by Theorem 24 in (32) implies:

$$\text{MSE}_f(N, n) \leq C \left( \frac{\kappa^{-1}\mathbf{A}_0\gamma_1}{(1-2\alpha)n^\alpha} + \frac{\kappa^{-1} \exp\{-\kappa\gamma_1 N^{1-\alpha}/(2(1-\alpha))\} \Omega(x) + \kappa^{-2}}{\gamma_1 n^{1-\alpha}} \right). \quad (\text{S29})$$

At fixed  $\gamma_1$ , this bound is of order  $n^{-\alpha}$ , and is better than (S27) for  $(\gamma_k)_{k \geq 1}$  constant. If we fix the number of iterations  $n$ , we can optimize the choice of  $\gamma_1$  again. Set

$$\gamma_{*,\alpha}(n) = (\kappa^{-1}\mathbf{A}_0/(1-2\alpha))^{-1} (C_{\text{MSE},\alpha}/n^{1-2\alpha})^{1/2}, \quad \text{where } C_{\text{MSE},\alpha} \stackrel{\text{def}}{=} \kappa^{-3}\mathbf{A}_0/(1-2\alpha),$$

(S29) becomes with  $\gamma_1 \leftarrow \gamma_{*,\alpha}(n)$ ,

$$\begin{aligned} \text{MSE}_f(N, n) \\ \leq C (C_{\text{MSE},\alpha} n)^{-1/2} \left( \kappa^{-1} \exp\{-\kappa N^{1-\alpha} \gamma_{*,\alpha}(n)/(2(1-\alpha))\} \Omega(x) + C_{\text{MSE},\alpha} \right). \end{aligned}$$

Setting  $N_\alpha(n) = \{2(1-\alpha)\kappa^{-1}\gamma_{*,\alpha}(n) \log(\Omega(x))\}^{1/(1-\alpha)}$ , we end up with

$$\text{MSE}_f(N_\alpha(n), n) \leq C (C_{\text{MSE},\alpha}/n)^{1/2}.$$

It is worthwhile to note that the order of  $N_\alpha(n)$  in  $n$  is  $n^{(1-2\alpha)/(2(1-\alpha))}$ , and  $C_{\text{MSE},\alpha}$  goes to infinity as  $\alpha \rightarrow 1/2$ .

3. If  $\alpha = 1/2$ , (S28) still holds. By Lemma 39, (S7) and (S9), we have the following bound for the bias

$$\{\mathbb{E}_x[\hat{\pi}_n^N(f)] - \pi(f)\}^2 \leq C \left( \frac{\kappa^{-1}\mathbf{A}_0\gamma_1 \log(n)}{n^{1/2}} + \frac{\kappa^{-1} \exp\{-\kappa\gamma_1 N^{1/2}/4\} \Omega(x)}{\gamma_1 n^{1/2}} \right).$$

Plugging this inequality and the one given by Theorem 24 in (32) implies:

$$\text{MSE}_f(N, n) \leq C \left( \frac{\kappa^{-1}\mathbf{A}_0\gamma_1 \log(n)}{n^{1/2}} + \frac{\kappa^{-1} \exp\{-\kappa\gamma_1 N^{1/2}/4\} \Omega(x) + \kappa^{-2}}{\gamma_1 n^{1/2}} \right). \quad (\text{S30})$$

At fixed  $\gamma_1$ , the order of this bound is  $\log(n)n^{-1/2}$ , and is the best bound for the MSE. Fix the number of iterations  $n$ , and we now optimize the choice of  $\gamma_1$ . Set

$$\gamma_{*,1/2}(n) = (\kappa^{-1}\mathbf{A}_0)^{-1} (C_{\text{MSE},1/2}/\log(n))^{1/2}, \quad \text{where } C_{\text{MSE},1/2} \stackrel{\text{def}}{=} \kappa^{-3}\mathbf{A}_0,$$

and (S30) becomes with  $\gamma_1 \leftarrow \gamma_{\star,1/2}(n)$ ,

$$\begin{aligned} \text{MSE}_f(N, n) &\leq C \left( \frac{\log(n)}{nC_{\text{MSE},1/2}} \right)^{1/2} \left( \kappa^{-1} \exp \left\{ -\kappa N^{1/2} \gamma_{\star,1/2}(n)/4 \right\} \Omega(x) + \frac{C_{\text{MSE},1/2}}{\log(n)} \right). \end{aligned}$$

Setting  $N_{1/2}(n) = (4\kappa^{-1}\gamma_{\star,1/2}(n) \log(\Omega(x)))^2$ , we end up with

$$\text{MSE}_f(N_{1/2}(n), n) \leq C \left( \frac{\log(n)C_{\text{MSE},1/2}}{n} \right)^{1/2}.$$

We can see that we obtain a worse bound than for  $\alpha = 0$  and  $\alpha \in (0, 1/2)$ .

4. For  $\alpha \in (1/2, 1]$ , (S28) still holds. By Lemma 39, (S7) and (S9), we have the following bound for the bias

$$\{\mathbb{E}_x[\hat{\pi}_n^N(f)] - \pi(f)\}^2 \leq C \left( \frac{\kappa^{-1}A_0\gamma_1}{n^{1-\alpha}} + \frac{\kappa^{-1} \exp \left\{ -\kappa\gamma_1 N^{1-\alpha}/(2(1-\alpha)) \right\} \Omega(x)}{\gamma_1 n^{1-\alpha}} \right).$$

Plugging this inequality and the one given by Theorem 24 in (32) implies:

$$\text{MSE}_f(N, n) \leq C \left( \frac{\kappa^{-1}A_0\gamma_1}{n^{1-\alpha}} + \frac{\kappa^{-1} \exp \left\{ -\kappa\gamma_1 N^{1-\alpha}/(2(1-\alpha)) \right\} \Omega(x) + \kappa^{-2}}{\gamma_1 n^{1-\alpha}} \right).$$

For fixed  $\gamma_1$ , the MSE is of order  $n^{1-\alpha}$ , and is worse than for  $\alpha = 1/2$ . For a fixed number of iteration  $n$ , optimizing  $\gamma_1$  would imply to choose  $\gamma_1 \rightarrow +\infty$  as  $n \rightarrow +\infty$ . Therefore, in that case, the best choice of  $\gamma_1$  is the largest possible value  $1/(m+L)$ .

5. For  $\alpha = 1$ , (S28) still holds. By Lemma 39, (S7) and (S9), the bias is upper bounded by

$$\{\mathbb{E}_x[\hat{\pi}_n^N(f)] - \pi(f)\}^2 \leq C \left( \frac{\kappa^{-1}A_0\gamma_1}{\log(n)} + \frac{\kappa^{-1}N^{-\kappa\gamma_1/2}\Omega(x)}{\gamma_1 \log(n)} \right).$$

Plugging this inequality and the one given by Theorem 24 in (32) implies:

$$\text{MSE}_f(N, n) \leq C \left( \frac{\kappa^{-1}A_0\gamma_1}{\log(n)} + \frac{\kappa^{-1}N^{-\kappa\gamma_1/2}\Omega(x) + \kappa^{-2}}{\gamma_1 \log(n)} \right).$$

For fixed  $\gamma_1$ , the order of the MSE is  $(\log(n))^{-1}$ . For a fixed number of iterations, the conclusions are the same than for  $\alpha \in (1/2, 1)$ .

## References

- [S1] A. Dalalyan. Theoretical guarantees for approximate sampling from a smooth and log-concave density. submitted 1412.7392, arXiv, December 2014.