



HAL
open science

A relaxation model for liquid-vapor phase change with metastability

Francois James, H el ene Mathis

► **To cite this version:**

Francois James, H el ene Mathis. A relaxation model for liquid-vapor phase change with metastability. 2015. hal-01178947v1

HAL Id: hal-01178947

<https://hal.science/hal-01178947v1>

Preprint submitted on 21 Jul 2015 (v1), last revised 14 Mar 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche franais ou  trangers, des laboratoires publics ou priv es.

A RELAXATION MODEL FOR LIQUID-VAPOR PHASE CHANGE WITH METASTABILITY

FRANÇOIS JAMES AND HÉLÈNE MATHIS

ABSTRACT. We propose a model that describes phase transition including metastable states present in the van der Waals Equation of State. From a convex optimization problem on the Helmholtz free energy of a mixture, we deduce a dynamical system that is able to depict the mass transfer between two phases, for which equilibrium states are either metastable states, stable states or a coexistent state. The dynamical system is then used as a relaxation source term in an isothermal 4×4 two-phase model. We use a Finite Volume scheme that treats the convective part and the source term in a fractional step way. Numerical results illustrate the ability of the model to capture phase transition and metastable states.

CONTENTS

1. Introduction	1
2. Thermodynamics and the van der Waals EoS	3
2.1. Thermodynamics of a single fluid	4
2.2. Example: the van der Waals EOS	5
2.3. Thermodynamics of phase transition	6
2.4. Equilibrium states	9
3. Dynamical system and phase transition	10
3.1. Construction of the dynamical system	10
3.2. Equilibria of the dynamical system	12
3.3. Numerical illustrations of the dynamical system behavior	13
4. The isothermal model	16
4.1. Definition of the model	17
4.2. Hyperbolicity and entropy	18
4.3. Structure of the waves	19
5. Numerical approximation	21
5.1. Definition of the splitting strategy	21
5.2. Numerical results	22
6. Conclusion	27
References	30

1. INTRODUCTION

In the last decades considerable research has been devoted to the simulation of liquid-vapor phase change, which are of major importance in several industrial applications. For instance liquid-vapor flows are present in water circuit of pressurized water reactors in which the water can be submitted to saturation pressure and temperature [11, 10]. The phenomena we are interested in are complex phase

2010 *Mathematics Subject Classification.* 35Q79, 35L40, 76T10, 37N10, 80A10.

Key words and phrases. thermodynamics of phase transition - metastable states - nonlinear hyperbolic system with relaxation - van der Waals equation of state.

changes including the possible appearance of metastable states. [An example is the metastable vapor which is](#) a gaseous state where the pressure is higher than the saturation pressure. Such states are very unstable and a very small perturbation brings out a droplet of liquid inside the gas [23]. [This](#) phenomenon can appear at saturated pressure (or at saturated temperature for metastable liquid). It is the case in pressurized water reactor during a loss of coolant accident when sudden vaporization occurs due to the drop of pressure inside the superheated liquid [19].

We focus in this paper on situations where the heterogeneity of the fluid and the thermodynamical conditions allow the diphasic flow to be described by a compressible averaged model using Euler type equations. Other models can be considered, which account for very small scale by means of Korteweg type tensors including dispersive and dissipative effects. Such models allow to preserve metastable states but give only a microscopic description of the flow, see [30, 31, 1, 6, 24, 20] and the references cited herein. In the averaged models framework, one can distinguish between one-fluid and two-fluid models.

The one-fluid model approach consists in describing the fluid flow as a single substance that can be present in its vapor or its liquid phase. Assuming that the thermodynamical equilibrium is reached instantaneously (quasi-static process), then the Euler system has to be closed by an Equation of State (EoS) able to depict either the pure phases (liquid or vapor) and the phase transition. A typical example is the van der Waals EoS, which is well-known to depict stable and metastable states below the critical temperature. However this EoS is not valid in the so-called spinodal zone where the pressure is a decreasing function of the density. This forbids the use of instantaneous kinetic exchanges, since the pressure is always given by the EoS, and a decreasing pressure leads to a loss of hyperbolicity in Euler equations, hence to instabilities and computational failure.

To overcome this defect, and recover that the phase transition happens at constant pressure, temperature and chemical potential, the van der Waals pressure is commonly corrected by the Maxwell equal area rule [8]. This construction leads to a correct constant pressure in the spinodal zone but removes the metastable regions.

Another way to provide a unique EoS able to depict pure phases and phase transition has been studied in [22, 2, 13, 16, 25]. It consists in considering that each phase is depicted by its own convex energy (that is its own monotone pressure law). According to the second principle of thermodynamics, the mixture equilibrium energy corresponds to the inf-convolution of the partial energies. As a result the mixture equilibrium energy coincides with the convex hull of the minimum of the two partial energies. The resulting pressure law of the mixture turns out to be composed of the monotone branches of the liquid and vapor pressure laws and is constant in the phase transition zone. Hence it is clear that such a construction prevents the appearance of metastable phases.

Still in the context of one-fluid models, it is also possible to drop out the assumption of instantaneous equilibrium. The model involves then relaxation terms, which can be of various forms, but are the expression of a pulling back force to the equilibrium. We have to consider extended versions of the Euler system, which is supplemented by partial differential equations on additional quantities such as the volume fraction of the vapor phase or partial masses. This approach has been used in [5, 17] and in [9, 14] in the isothermal case. In the later references the question of preservation of metastable states is addressed.

We mention briefly another way to describe diphasic flows, which consists in considering a two-fluid approach to model liquid-vapor phase change. Initially developed to depict the motion of multicomponent flows [4], such a modeling assumes

that the fluid can locally be present under both phase. Hence the model admits two pressures, two velocities and two temperatures and is supplemented by additional equations on the volume fraction. Phase transition is achieved by chemical and mechanical relaxation processes, in the limit where the kinetics is considered infinitely fast, see for instance [29, 32, 27].

One of the present drawbacks of the averaged models (one or two-fluid) with relaxation is that there is no global agreement on the equations satisfied by the fractions and on the transfer terms [12]. Moreover the preservation of the metastable states seems to be out of reach in this framework.

We intend in this paper to provide a model able to depict liquid-vapor phase change and metastable states of a single component, say some liquid in interaction with its own vapor. We focus on a one-fluid description of the motion while the phase transition is driven by transfer terms that will be coupled to fluid equations through a finite relaxation speed.

The modelling of the phase transition is the core of this work. For the sake of simplicity we assume the system to be isothermal. We propose transfer terms obtained through the minimization of the Helmholtz free energy of the two-phase system. We use for both phases the same equation of state which has to be non monotone, typically the reduced van der Waals equation. This choice allows us to recover all possible equilibria: pure phases (liquid or vapor), metastable states and coexistence states characterized by the equality of pressures and chemical potentials. These are physically admissible states, but the set of equilibrium states also contains the spinodal zone, which is irrelevant. Thus the key point now is to characterize the physical stability, and hence admissibility, of these states. It turns out that this has to be done in terms of their dynamical behaviour. More precisely, we design a dynamical system which is able to depict all the *stable* equilibria of the system as attraction points. In particular we show that metastable states and mixtures have different basins of attraction, so that they can be differentiated only by their long-time behaviour with respect to initial conditions. Hence there is no hope to recover both metastable states and coexistent states under the assumption of instantaneous equilibrium, which amounts precisely to choose *a priori* this long-time behaviour.

This dynamical system is used as a transfer term in an isothermal two-phase model in the spirit of [26] and [3]. The extended Euler system we obtain in this way provides a regularization of the isothermal Euler equation with van der Waals EoS, which takes the form of a mixture zone surrounding the physical interfaces, see Section 5 below.

The paper is organized as follows. Section 2 is devoted to the thermodynamics of a two-phase fluid. We provide the definitions of the common potentials and give some details on the reduced van der Waals model. Assuming that both phases follow the same non monotone EoS, we describe the thermodynamical equilibrium as the result of a minimization process on the Helmholtz free energy of the two-phase fluid. The section ends with the study of the equilibria of the optimality system. Section 3 is the core of this work. It is devoted to the construction of the dynamical system based on the results of the previous section. A few numerical simulations illustrate the ability of the system to catch both the Maxwell line and the metastable states in the van der Waals EoS. The dynamical system is then plugged as a source term in a 4×4 isothermal model, see section 4. In section 5 we present several numerical illustrations which assess the ability of the model to deal with metastable states. They are obtained using a classical finite volume schemes that treats the convective part and the source terms with a time-splitting technique.

2. THERMODYNAMICS AND THE VAN DER WAALS EOS

2.1. Thermodynamics of a single fluid. Consider a fluid of mass $M \geq 0$ occupying a volume $V \geq 0$, assumed to be at constant temperature T . If the fluid is homogeneous and at rest, its behavior is entirely described by its mass, its volume and its Helmholtz free energy F . According to Gibbs' formalism [15], the fluid is at equilibrium if its Helmholtz free energy is a function, also denoted by F , of its mass M and volume V :

$$(1) \quad F : (M, V) \rightarrow F(M, V).$$

Notice that we do not address yet the stability of equilibrium states. At this level the involved quantities are extensive, which means that they share the same scalings as the volume V . This corresponds to the notion of homogeneous sample: for twice the volume, the mass is double, and the energy as well. The mathematical consequence of this notion is that extensive variables have to be related through positively homogeneous functions of degree 1 (PH1), namely

$$(2) \quad \forall \lambda > 0, \quad F(\lambda M, \lambda V) = \lambda F(M, V).$$

We assume in addition, and without loss of generality, that the energy function F belongs to $\mathcal{C}^2(\mathbb{R}^+ \times \mathbb{R}^+)$.

Remark 2.1. *The regularity of F seems to preclude phase transitions, but this is not the case because no convexity assumption is made at this stage. We shall come back to this in more details in the next section.*

We introduce now intensive quantities, by which we mean that they do not depend on the volume scaling. A typical example is the density $\rho = M/V$, but such quantities also appear as derivatives of the the equilibrium relation (2), which are homogeneous of degree 0 by construction. In this way two fundamental quantities are to be considered, namely the pressure p and the chemical potential μ , defined by

$$(3) \quad p = -\frac{\partial F}{\partial V}(M, V), \quad \mu = \frac{\partial F}{\partial M}(M, V).$$

Notice that these quantities are defined only when the system is at equilibrium, and we recover the classical thermodynamic relation for isothermal flows

$$(4) \quad dF = -pdV + \mu dM.$$

Another classical property of thermodynamical potentials is the so-called Gibbs relation, which results from the Euler relation for PH1 functions:

$$F(M, V) = \nabla F(M, V) \cdot \begin{pmatrix} M \\ V \end{pmatrix}.$$

Using (3), we obtain

$$(5) \quad F(M, V) = \mu M - pV.$$

For most of the following computations it is useful to rewrite the preceding relations using intensive variables. For a fixed volume V , we denote f the Helmholtz free energy per unit volume, which is a function of the density $\rho = M/V$:

$$(6) \quad f(\rho) = f\left(\frac{M}{V}\right) = \frac{1}{V}F(M, V) = F\left(\frac{M}{V}, 1\right).$$

We keep the notations p and μ the pressure and the chemical potential as functions of the density ρ :

$$(7) \quad p(\rho) = p\left(\frac{M}{V}\right) = -\frac{\partial E}{\partial V}\left(\frac{M}{V}, 1\right), \quad \mu(\rho) = \mu\left(\frac{M}{V}\right) = \frac{\partial E}{\partial M}\left(\frac{M}{V}, 1\right).$$

Thus we obtain an intensive form of the Gibbs relation (5)

$$(8) \quad f(\rho) = \rho\mu(\rho) - p(\rho).$$

On the other hand, from the definitions of p , μ and f we obtain easily the following relations

$$(9) \quad \mu(\rho) = f'(\rho), \quad p(\rho) = \rho f'(\rho) - f(\rho), \quad \rho\mu'(\rho) = p'(\rho).$$

2.2. Example: the van der Waals EOS. The extensive energy of a van der Waals (monoatomic) fluid is

$$(10) \quad F(M, V) = -\frac{aM^2}{V} + RT \left(M \log \frac{M}{V - Mb} - M \right),$$

where R stands for the perfect gases constant and a and b are positive constants ($a = b = 0$ leads to the perfect gases law). Since we consider the isothermal model, the temperature T is a parameter here. We refer to [23, Ch. 7] for a justification of this law from statistical thermodynamics. The constant b is proportional to the proper volume of a particule such that $V > Mb$, and the potential $\frac{a}{V}$ depicts the interaction between particules.

The pressure and the chemical potential associated read

$$(11) \quad \begin{aligned} p\left(\frac{M}{V}\right) &= -\frac{aM^2}{V^2} + RT \frac{M}{V - Mb}, \\ \mu\left(\frac{M}{V}\right) &= -2\frac{aM}{V} + RT \log \frac{M}{V - Mb} + RT \frac{Mb}{V - Mb}. \end{aligned}$$

The intensive quantities are

$$(12) \quad \begin{aligned} f(\rho) &= -a\rho^2 + RT\rho \left(\log \left(\frac{\rho}{1 - b\rho} \right) - 1 \right), \\ p(\rho) &= -a\rho^2 + \frac{\rho RT}{1 - b\rho}, \\ \mu(\rho) &= -2a\rho + RT \log \frac{\rho}{1 - b\rho} + RT \frac{b\rho}{1 - b\rho}. \end{aligned}$$

The behavior of the isotherm curves in the pressure-density plan is depicted in Figure 1. The critical temperature T_c is the lower limit of temperatures such that the pressure is an increasing function of the density. At $T = T_c$ the pressure curve admits an unique horizontal inflection point, called the critical point, denoted C on Figure 1.


In the sequel we will consider the reduced form of the van der Waals equation (see [23, Ch. 8]). Denoting (ρ_c, p_c) the coordinates of the critical point one has

$$\begin{aligned} p'(\rho_c) &= -2a\rho_c + \frac{RT_c}{(1 - b\rho_c)^2} = 0, \\ p''(\rho_c) &= -2a + \frac{2bRT_c}{(1 - b\rho_c)^3} = 0. \end{aligned}$$

Considering normalized critical quantities, that is setting $\rho_c = 1$, $p_c = 1$ and $T_c = 1$, one obtains the reduced van der Waals EOS with $R = 8/3$, $a = 3$, and $b = 1/3$ and

$$(13) \quad \begin{aligned} f(\rho) &= -3\rho^2 + \frac{8}{3}\rho (\log(3\rho/(3 - \rho)) - 1), \\ p(\rho) &= -3\rho^2 + \frac{8\rho}{3 - \rho}, \\ \mu(\rho) &= -6\rho + \frac{8}{3} \log(3\rho/(3 - \rho)) + \frac{8}{3}(3\rho/(3 - \rho)) \end{aligned}$$

FIGURE 1. Phase diagram for the van der Waals EoS in the (p, ρ) plan. Below the critical temperature T_C , the isotherm curve decreases in the spinodal zone which is delimited by the densities $\rho^- < \rho^+$. In that area the isotherm is commonly replaced by an horizontal segment (dashed line) which coincides with the isobaric line at constant pressure $p = p^*$. Such a construction defines the two densities ρ_1^* and ρ_2^* .



for $0 < \rho < 3$.

Below the critical temperature T_c the pressure is not monotone with respect to the density (see the red curve on Figure 1): in a region called spinodal zone, delimited for a given temperature by the densities $\rho^- < \rho^+$, the pressure decreases with respect to the density and thus leading to unstable states. According to (9) the energy f is non-convex in the spinodal zone.

In that region the isotherm is commonly replaced by the Maxwell area rule in order to recover that phase transition happens at constant pressure and constant chemical potential. The Maxwell construction is commonly applied on the pressure (see [8] for instance) in such a way that the two zones delimited by the van der Waals isotherm and the Maxwell line (above and below the Maxwell line respectively) have the same area. This is not the case in Figure 1 because in our context, the equal area rule is obtained on the chemical potential, see Section 2.4 below (Proposition 2.1). In any case, the idea is that the isotherm curve is replaced locally by the horizontal segment, the so-called Maxwell line, that coincides with some isobaric line $p = p^*$. Such a construction defines two densities, denoted ρ_1^* and ρ_2^* , as well as the value of p^* . An equivalent way to compute this correction is to build the convex hull of the function f , see [2] and [16].

However this construction removes the admissible regions delimited by the densities where the pressure law is still nondecreasing. Such regions are called metastable regions. At a given temperature these regions correspond to densities belonging to the range $[\rho_1^*, \rho^-]$ and $[\rho^+, \rho_2^*]$.

2.3. Thermodynamics of phase transition. The van der Waals model depicts the thermodynamical behavior of a pure substance under its liquid state, gaseous state and the coexistence state. The non convexity of the EoS allows to capture metastable states but does not give a relevant representation of the coexistence phase. A convenient way to cope with this problem is to represent the fluid under consideration as a set of several copies of itself under different pure phases (liquid

or gaseous phases). Such a representation is used in [22, 5, 17, 2, 16, 13] assuming that the fluid is described by two copies, each one satisfying a convex EoS that differs from the one of another copy. See also [21] where such a representation is used in the context of adsorption-desorption of a mixture.

We adopt here a slightly modified approach. Let us consider I copies of the pure substance, $I \geq 1$ being some integer *a priori* arbitrary. Each copy is depicted by its mass $M_i \geq 0$ and its volume $V_i > 0$, and is assumed to be at equilibrium. Hence a copy can occupy a volume with zero mass. We suppose that each copy follows the same non-convex energy function $F(M_i, V_i)$, typically the van der Waals extensive energy given by (10). This assumption contrasts with the aforementioned references where different convex energy functions are considered, and the number of copies is prescribed.

Thanks to the mass conservation, the complete system has a total mass $M = \sum_{i=1}^I M_i$. Assuming that the copies are immiscible, the total volume is $V = \sum_{i=1}^I V_i$ (for a mixture of gas, one has $V_i = V$, $\forall i = 1, \dots, I$, and this condition implies Dalton's law, see [8]). Out of thermodynamical equilibrium the free Helmholtz energy of the system reads

$$\mathcal{F}((M_i, V_i)_i) = \sum_{i=1}^I F(M_i, V_i).$$

Let us fix the total mass M and volume V of the system. According to the second principle of thermodynamics (see [8] for instance), the thermodynamical equilibrium corresponds to the solutions of the constrained optimization problem

$$\inf_{I \geq 1, M_i \geq 0, V_i \geq 0} \left\{ \sum_{i=1}^I F(M_i, V_i); \sum_{i=1}^I V_i = V, \sum_{i=1}^I M_i = M \right\}.$$

We stress the fact that the total number I of possible copies is not fixed *a priori* here. However, as a consequence of Carathéodory's theorem in dimension 2 (see e.g. [28, Ch. 17]), we can state the Gibbs phase rule (see [23, Ch. 9]), which gives the expected result in this context.

Lemma 2.1 (Gibbs phase rule). *We have $I \leq 2$.*

In the sequel we use just as well the term *phase* as the term *copy*.

Taking into account Lemma 2.1 and the above intensive formulations, we rewrite the constrained optimization problem for a fixed mass M and volume V , hence for a fixed ρ , as

$$(14) \quad \inf_{\substack{\rho_1 \geq 0 \\ \rho_2 \geq 0}} \{ \alpha_1 f(\rho_1) + \alpha_2 f(\rho_2) \},$$

under the constraints

$$(15) \quad \alpha_1 + \alpha_2 = 1,$$

$$(16) \quad \alpha_1 \rho_1 + \alpha_2 \rho_2 = \rho.$$

Here $\alpha_i = \frac{V_i}{V} \in [0, 1]$ denotes the volume fraction and $\rho_i = \frac{M_i}{V_i} \geq 0$ the density of the phase $i = 1, 2$. Notice that we removed the optimization on the phase number I , since single phases can be recovered by $\rho_1 = \rho_2$ (and any α_i) or one $\alpha_i = 0$ (with undetermined ρ_j , $j \neq i$).

We rule out the equality case by noticing that, provided $\rho_1 \neq \rho_2$, we can rewrite the constraints (15)-(16) as

$$(17) \quad \begin{aligned} \alpha_1 : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ &\rightarrow [0, 1] & \alpha_2 : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ &\rightarrow [0, 1] \\ (\rho, \rho_1, \rho_2) &\mapsto \frac{\rho - \rho_2}{\rho_1 - \rho_2}, & (\rho, \rho_1, \rho_2) &\mapsto -\frac{\rho - \rho_1}{\rho_1 - \rho_2}. \end{aligned}$$

We have $\alpha_1 \geq 0$ if and only if $\rho_1 \leq \rho \leq \rho_2$ or $\rho_1 \geq \rho \geq \rho_2$. Therefore, accounting on the reduced form of the van der Waals model (13), we shall assume in the sequel that the densities ρ , ρ_1 and ρ_2 satisfy

$$(H_1) \quad 0 < \rho_1 \leq \rho \leq \rho_2 < 3 \quad \text{and} \quad \rho_1 < \rho_2.$$

This is not a restriction, as we shall see below (see Proposition 3.1).

One can also introduce the mass fraction $\varphi_i = \frac{M_i}{M}$, $i = 1, 2$ defined by

$$(18) \quad \begin{aligned} \varphi_1 : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ &\rightarrow [0, 1] & \varphi_2 : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ &\rightarrow [0, 1] \\ (\rho, \rho_1, \rho_2) &\mapsto \frac{\frac{1}{\rho} - \frac{1}{\rho_2}}{\frac{1}{\rho_1} - \frac{1}{\rho_2}}, & (\rho, \rho_1, \rho_2) &\mapsto \frac{\frac{1}{\rho} - \frac{1}{\rho_1}}{\frac{1}{\rho_2} - \frac{1}{\rho_1}}, \end{aligned}$$

that satisfy $\varphi_1 + \varphi_2 = 1$ and $\varphi_1 \varphi_2 \geq 0$ if and only if the assumption (H_1) is satisfied. Such quantities will be useful in the mathematical study of the isothermal two-phase flow model introduced in Section 4.

Remark 2.2. *In the aforementioned references [22, 5, 17, 2, 16, 13] the method consists in describing the two-phase fluid by a coexistence of two copies of the same substance. The description can be made either on the extensive variables or the intensive one. Unlike our present approach the two copies do not follow the same EoS: each copy is described by its own strictly convex extensive energy F_i , $i = 1, 2$, which is a function of the mass M_i and the volume V_i of the phase. Following the second principle of thermodynamics [8], at equilibrium, the extensive energy of the two-phase fluid is given by*

$$(19) \quad F((M, V)) := F_1 \square F_2(M, V) = \min_{V_1 \geq 0, M_1 \geq 0} F_1(M_1, V_1) + F_2(M - M_1, V - V_1),$$

under the constraints of mass conservation $M = M_1 + M_2$ and immiscibility (without vacuum) $V = V_1 + V_2$. This operation \square is called *inf-convolution operation* in the convex analysis framework [18]. In [16] the authors investigate the link between the inf-convolution, the Legendre transform and the $(\max, +)$ algebra. The Legendre transform of a energy F is a convex function $(M^*, V^*) \rightarrow F^*(M^*, V^*)$ defined by

$$F^*(M^*, V^*) = \inf_{M \geq 0, V \geq 0} \{M^* M + V^* V - F(M, V)\}.$$

The inf-convolution is transformed into an addition by the Legendre transform which implies that

$$(F_1 \square F_2)^* = F_1^* + F_2^*.$$

Moreover in the case of convex lower semi continuous (slc) functions F_i , $i = 1, 2$, one has

$$F_1 \square F_2 = (F_1 \square F_2)^{**} = (F_1^* + F_2^*)^*.$$

It means that the energy F_{eq} of the two-phase fluid at equilibrium is given by

$$(20) \quad F_{eq} = F^{**},$$

where F^{**} is the convex hull of the energy F . As it is proved in [2] the construction of the convex hull of the energy F is equivalent to the Maxwell construction. Hence the operation (20) removes the metastable regions.

2.4. Equilibrium states. This section is devoted to the characterization of the equilibrium states of the thermodynamical system, that is states that realize the infimum in the optimization problem (14)–(15)–(16). The function we minimize is defined by:

$$\begin{aligned} \tilde{\mathcal{F}} : [0, 1] \times [0, 1] \times \mathbb{R}^+ \times \mathbb{R}^+ &\rightarrow \mathbb{R} \\ (\alpha_1, \alpha_2, \rho_1, \rho_2) &\mapsto \alpha_1 f(\rho_1) + \alpha_2 f(\rho_2), \end{aligned}$$

and is C^1 on the space $\{\rho_i \geq 0, i = 1, 2\}$. The constraints (15)–(16) are affine, so they are also C^1 . We are thus in position to use the Lagrange multipliers characterization of the infimum (see [28, Ch. 28]): $\lambda_\alpha \in \mathbb{R}$ and $\lambda_\rho \in \mathbb{R}$ respectively correspond to the constraints (15) and (16).

Using the definition (8) of the free energy f and the pressure and chemical potential definitions (9), one deduces the following optimality system of equations

$$(21) \quad f(\rho_1) + \lambda_\alpha + \lambda_\rho \rho_1 = 0,$$

$$(22) \quad f(\rho_2) + \lambda_\alpha + \lambda_\rho \rho_2 = 0,$$

$$(23) \quad \alpha_1 \mu(\rho_1) + \lambda_\rho \alpha_1 = 0,$$

$$(24) \quad \alpha_2 \mu(\rho_2) + \lambda_\rho \alpha_2 = 0.$$

From this optimality system we deduce immediately that there are two different kinds of equilibria.

Lemma 2.2. *Under hypothesis (H₁), the equilibrium states are*

- (1) **Pure liquid or gaseous states:** $\alpha_1 = 0$ (resp. $\alpha_2 = 0$), with $\rho_2 = \rho$, $\rho_1 < \rho$ arbitrary (resp. $\rho_1 = \rho$, $\rho < \rho_2$ arbitrary)
- (2) **Coexistence states:** $\alpha_1 \alpha_2 \neq 0$, with (ρ_1, ρ_2) satisfying $\mu(\rho_1) = \mu(\rho_2)$ and $p(\rho_1) = p(\rho_2)$.

Proof. The case $\alpha_1 = 0$ corresponds to the saturation of the constraint $\alpha_1 + \alpha_2 = 1$ see (15). It leads to $\alpha_2 = 1$ and thus $\rho_2 = \rho$ that is only the phase 2 is present. Conversely if $\alpha_2 = 1$ only the phase 1 remains.

On the other hand let us assume $\alpha_1 \alpha_2 \neq 0$. Then (23) and (24) lead to the equality of the chemical potentials

$$\mu(\rho_1) = \mu(\rho_2) = -\lambda_\rho.$$

Then the intensive Gibbs relation (8) allows to rewrite the conditions (21) and (22) as

$$\rho_i \mu(\rho_i) - p(\rho_i) + \lambda_\alpha + \lambda_\rho \rho_i = 0, \quad i = 1, 2.$$

Since $-\lambda_\rho = \mu(\rho_i)$, this leads to the pressures equality

$$p(\rho_1) = p(\rho_2) = \lambda_\alpha.$$

□

To proceed further we need the following result for coexistence states.

Proposition 2.1. *Under hypothesis (H₁) and if $\alpha_1 \alpha_2 \neq 0$, the following propositions are equivalent and uniquely define the couple of densities $\rho_1^* < \rho_2^*$.*

- (1) *The chemical potentials and the pressures are equal*

$$(25) \quad \mu(\rho_1^*) = \mu(\rho_2^*),$$

$$(26) \quad p(\rho_1^*) = p(\rho_2^*).$$

- (2) *The Maxwell's area rule on the chemical potential holds*

$$(27) \quad \int_0^1 \mu(\rho_2 + t(\rho_1 - \rho_2)) dt = \mu(\rho_1^*) = \mu(\rho_2^*).$$

(3) *The difference of the Helmholtz free energies reads*

$$(28) \quad f(\rho_2^*) - f(\rho_1^*) = \mu(\rho_1^*)(\rho_2^* - \rho_1^*) = \mu(\rho_2^*)(\rho_2^* - \rho_1^*).$$

Proof. The identities (27) and (28) are equivalent since $f'(\rho) = \mu(\rho)$, see (9). Now assume (25)-(26) hold. Then the intensive Gibbs relation (8) gives (28). Conversely (28) implies the chemical potentials equality and thus the pressures equality. Now the uniqueness of (ρ_1^*, ρ_2^*) follows easily from a geometrical argument using the Maxwell's area rule. \square

The most famous characterizations of diphasic equilibria are (25)-(25) and (27), although the latter is usually written in terms of pressure. We can recover this form by writing the intensive relations with the specific volume $\tau = V/M$ instead of ρ . The third relation (28) is not so classic but will be useful in the sequel.

The density ρ_1^* (resp. ρ_2^*) separates the range of pure gaseous state and the range of metastable gas (resp. the range of pure liquid state and the range of metastable liquid), see Figure 1. These densities define the coexistence pressure p^* and the coexistence chemical potential μ^* :

$$(29) \quad p^* = p(\rho_1^*) = p(\rho_2^*), \quad \mu^* = \mu(\rho_1^*) = \mu(\rho_2^*).$$

We emphasize that the necessary conditions for equilibrium contain both unstable (spinodal zone), metastable and stable states. Nothing at this stage can make the difference, which turns out to be of dynamical nature, in a way we make precise now.

3. DYNAMICAL SYSTEM AND PHASE TRANSITION

This section is devoted to the construction and the analysis of a dynamical system deduced from the optimality conditions given in Lemma 2.2 and Proposition 2.1. We will prove that the equilibria of this dynamical system are either pure liquid/vapor states, pure metastable states or coexistence states in the spinodal zone (that is states satisfying the properties (25)-(27) of Proposition 2.1). We emphasize that the difference between metastable states and coexistent states actually relies on the long-time dynamical behaviour of the solutions to the dynamical system. No static characterization can be given.

The section ends with numerical illustrations that assess the ability of the dynamical system to preserve metastable states.

3.1. Construction of the dynamical system. We want to construct a dynamical system which equilibria coincide with the equilibria of the optimization problem depicted in Lemma 2.2. To do so, we force the dynamical system to dissipate the Helmholtz free energy defined by

$$(30) \quad \begin{aligned} \mathcal{F} : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ &\rightarrow \mathbb{R} \\ (\rho, \rho_1, \rho_2) &\mapsto \alpha_1(\rho, \rho_1, \rho_2)f(\rho_1) + \alpha_2(\rho, \rho_1, \rho_2)f(\rho_2), \end{aligned}$$

under the optimization constrains (15) and (16).

Assuming that ρ , ρ_1 and ρ_2 are now time-dependent functions, we can compute the derivative of the total Helmholtz free energy \mathcal{F} with respect to time and deduce appropriate time derivatives of ρ , ρ_1 and ρ_2 such that \mathcal{F} is dissipated in time. Moreover we want pure states (either liquid, vapor or metastable) to be equilibria of the dynamical system. Hence we have forced the time derivatives of ρ , ρ_1 and ρ_2 to vanish in case of pure state (that is when $\alpha_1\alpha_2 = 0$).

For sake of readability we denote

$$(31) \quad \mathbf{r} = \begin{pmatrix} \rho \\ \rho_1 \\ \rho_2 \end{pmatrix}$$

the vector of admissible densities satisfying the assumption (H_1) . By the definition (17) of the volume fractions α_i , $i = 1, 2$, one has

$$\nabla_{\mathbf{r}}\alpha_1(\mathbf{r}) = -\nabla_{\mathbf{r}}\alpha_2(\mathbf{r}) = \frac{1}{\rho_1 - \rho_2} \begin{pmatrix} 1 \\ -\alpha_1(\mathbf{r}) \\ -\alpha_2(\mathbf{r}) \end{pmatrix}.$$

From this and the definition (30) of \mathcal{F} , we easily get the gradient of the free energy

$$(32) \quad \nabla_{\mathbf{r}}\mathcal{F}(\mathbf{r}) = \frac{1}{\rho_1 - \rho_2} \begin{pmatrix} f(\rho_1) - f(\rho_2) \\ \alpha_1(\mathbf{r})(\rho_2(\mu(\rho_2) - \mu(\rho_1)) + p(\rho_1) - p(\rho_2)) \\ \alpha_2(\mathbf{r})(\rho_1(\mu(\rho_2) - \mu(\rho_1)) + p(\rho_1) - p(\rho_2)) \end{pmatrix}$$

Note at once that it can be expressed in terms of relative free energy

$$(33) \quad \nabla_{\mathbf{r}}\mathcal{F}(\mathbf{r}) = \frac{1}{\rho_1 - \rho_2} \begin{pmatrix} f(\rho_1) - f(\rho_2) \\ \alpha_1(\mathbf{r})f(\rho_2|\rho_1) \\ -\alpha_2(\mathbf{r})f(\rho_1|\rho_2) \end{pmatrix},$$

where the relative free energy of ρ_2 with respect to ρ_1 is defined by

$$(34) \quad f(\rho_2|\rho_1) := f(\rho_2) - f(\rho_1) - \mu(\rho_1)(\rho_2 - \rho_1).$$

We turn now to the definition of our dynamical system. Because of the mass conservation, we obviously impose that

$$\dot{\rho} = 0.$$

The main idea to proceed further is that we want the system to dissipate the total Helmholtz free energy \mathcal{F} . Combining the definition (30) of \mathcal{F} and the expression of $\nabla_{\mathbf{r}}\mathcal{F}(\mathbf{r})$, one computes the time derivative of \mathcal{F} along some trajectory, that is

$$(35) \quad \dot{\mathcal{F}}(\mathbf{r}) = \frac{1}{\rho_1 - \rho_2} (\alpha_1(\mathbf{r})f(\rho_2|\rho_1)\dot{\rho}_1 - \alpha_2(\mathbf{r})f(\rho_1|\rho_2)\dot{\rho}_2).$$

We now propose the following dynamical system

$$(36) \quad \begin{cases} \dot{\rho} = 0, \\ \dot{\rho}_1 = -(\rho - \rho_1)(\rho - \rho_2)f(\rho_2|\rho_1), \\ \dot{\rho}_2 = +(\rho - \rho_1)(\rho - \rho_2)f(\rho_1|\rho_2). \end{cases}$$

An easy computation shows that this system dissipates \mathcal{F} along its trajectories (under the assumption (H_1)). Indeed using the expression (35) of $\dot{\mathcal{F}}$ and the equations of $\dot{\rho}_1$ and $\dot{\rho}_2$, one gets

$$\dot{\mathcal{F}}(\mathbf{r}) = -(\rho - \rho_1) [\alpha_1(\mathbf{r})f(\rho_2|\rho_1)]^2 + (\rho - \rho_2) [\alpha_2(\mathbf{r})f(\rho_1|\rho_2)]^2.$$

The multiplicative term $(\rho - \rho_1)(\rho - \rho_2)$ in the equations on $\dot{\rho}_1$ and $\dot{\rho}_2$ ensures that the right hand side of the system vanishes in case of pure states (either pure liquid, vapor or metastable states).

Remark 3.1. *We emphasize that the choice of the right hand side of (36) is somewhat arbitrary. Other terms might be more efficient, but this one was chosen for its simplicity and its interpretation in terms of relative free energy.*

One can check easily that $\overbrace{\alpha_1\rho_1 + \alpha_2\rho_2} = 0$ so that it is consistent with the total mass conservation equation $\dot{\rho} = 0$.

An equivalent dynamical system can be written in terms of the time derivatives of the volume fractions α_i and of the partial masses $\alpha_i\rho_i$, $i = 1, 2$. Accounting for the

constraints (15)-(16) and for the system (36), some straightforward computations lead to

$$(37) \quad \begin{cases} \dot{\alpha}_1 &= -\dot{\alpha}_2 \\ &= \alpha_1^2(\rho - \rho_1)f(\rho_2|\rho_1) + \alpha_2^2(\rho - \rho_2)f(\rho_1|\rho_2), \\ \dot{\alpha}_1\rho_1 &= -\dot{\alpha}_2\rho_2 \\ &= \alpha_1^2\rho_2(\rho - \rho_1)f(\rho_2|\rho_1) + \alpha_2^2\rho_1(\rho - \rho_2)f(\rho_1|\rho_2). \end{cases}$$

This formulation does not allow to compute single phase systems, since the determination of the partial densities ρ_i is impossible when $\alpha_i = 0$. Thus we rather use the dynamical system (36).

3.2. Equilibria of the dynamical system. The major result of this paragraph is that the attractors of the system are either pure liquid/vapor states, including metastable states, or the coexistence state defined by (27)-(26), see Proposition 2.1.

Proposition 3.1. *The dynamical system (36) satisfies the following properties.*

- (1) *If the assumption (H_1) is satisfied at $t = 0$, then it is preserved in time (in particular, the assumption $\rho_1 < \rho_2$ is preserved).*
- (2) *If $\alpha_1(0) = 0$ (resp. $\alpha_1(0) = 1$) then $\alpha_1(t) = 0$ (resp. $\alpha_1(t) = 1$), for all time.*

Proof. First we address the preservation of the hypothesis (H_1). Some straightforward computations lead to

$$(38) \quad \dot{\rho}_1 - \dot{\rho}_2 = (\rho - \rho_1)(\rho - \rho_2)(\rho_1 - \rho_2)(\mu(\rho_2) - \mu(\rho_1)).$$

Thus if $\rho_1 - \rho_2$ is zero at $t = 0$ then this property is preserved for all time and the sign of the difference $\rho_1 - \rho_2$ is also preserved. The property on the volume fraction α_1 is proved in the same way noting that

$$(39) \quad \dot{\alpha}_1(\mathbf{r}) = -\alpha_1(\mathbf{r})(\rho - \rho_1)(\rho_1(\mu(\rho_1) - \mu(\rho_2)) - p(\rho_1) + p(\rho_2)).$$

□

We turn now to the characterization of the equilibria of the dynamical system. Since the total mass ρ is constant in time, they are parametrized by ρ . Moreover it arises that their thermodynamical characterization can be given in terms of attraction basins. We refer the reader to figures 2 and 3 for graphical references.

Theorem 3.1. *Assume that the initial data $(\rho(0), \rho_1(0), \rho_2(0))$ of the system (36) satisfy (H_1). Then the equilibria are given by*

- (1) **spinodal zone:** $\rho_1^* < \rho(0) < \rho_2^*$. *The unique equilibrium is (ρ_1^*, ρ_2^*) characterized by Proposition 2.1, with α_i given by (17). The attraction basin is $(0, \rho) \times (\rho, 3)$.*
- (2) **pure gaseous states:** $\rho(0) < \rho_1^*$ (resp. **pure liquid states:** $\rho(0) > \rho_2^*$). *The set of equilibria is $\{\rho\} \times (\rho, 3)$, with $\alpha_1 = 1$ (resp. $(0, \rho) \times \{\rho\}$ with $\alpha_2 = 1$). The attraction basin is $(0, \rho) \times (\rho, 3)$.*
- (3) **metastable states:** $\rho_1^* \leq \rho(0) \leq \rho_-$ (resp. $\rho_+ \leq \rho(0) \leq \rho_2^*$). *There are two sets of equilibria*
 - (a) **perturbation within the phase:** $\rho_2(0) \leq \rho_-$ (resp. $\rho_1(0) \geq \rho_+$). *The set of equilibria is $\{\rho\} \times (\rho, \rho_-)$, with $\alpha_1 = 1$ (resp. $(\rho_+, \rho) \times \{\rho\}$ with $\alpha_2 = 1$). The attraction basin is then $(0, \rho) \times (\rho, \rho_-)$ (resp. $(\rho_+, \rho) \times (\rho, 3)$).*
 - (b) **outside perturbation:** $\rho_- \leq \rho_2(0) \leq \rho_2^*$ (resp. $\rho_+ \geq \rho_1(0) \geq \rho_1^*$). *There is a unique equilibrium (ρ_1^*, ρ_2^*) , characterized by Proposition 2.1, with α_i given by (17). The attraction basin is $(0, \rho) \times (\rho_-, 3)$ (resp. $(0, \rho_+) \times (\rho, 3)$).*

Proof. We look for Lyapunov functions for each case.

Spinodal zone. Let us define

$$(40) \quad G(\mathbf{r}) = \alpha_1 f(\rho_1) + \alpha_2 f(\rho_2) - \mu^*(\alpha_1 \rho_1 + \alpha_2 \rho_2) + p^*(\alpha_1 + \alpha_2),$$

where μ^* and p^* are defined by the Maxwell area rule (29) on the chemical potential μ , and α_i are the functions of ρ, ρ_1, ρ_2 given by (17). The intensive version of the Gibbs relation (8) implies that $G(\mathbf{r}^*) = 0$ where $\mathbf{r}^* = (\rho, \rho_1^*, \rho_2^*)^T$. Straightforward computations show that $\nabla_{\mathbf{r}} G = \nabla_{\mathbf{r}} \mathcal{F} - (\mu^*, 0, 0)^T$, this implies that $\nabla_{\mathbf{r}} G(\mathbf{r}^*) = 0$, and that $\dot{G}(\mathbf{r}(t)) \leq 0$ since the free energy is dissipated. Hence the function G is a Lyapunov function on the admissible domain defined by (H_1) , which contains the unique equilibrium \mathbf{r}^* , see Figure 3.

Pure stable states The expected equilibrium states are now $\bar{\mathbf{r}} = (\rho, \rho, \rho_2)$ for any $\rho_2 \geq \rho$, see Figure 2-left. We introduce the function

$$(41) \quad G^{(1)}(\mathbf{r}) = f(\rho_1) - \bar{\mu}\rho_1 + \bar{p}, \quad \text{where } \bar{\mu} = \mu(\rho), \bar{p} = p(\rho).$$

Once again, the Gibbs relation leads to $G^{(1)}(\bar{\mathbf{r}}) = 0$ for any equilibrium $\bar{\mathbf{r}}$, and we easily have $\nabla_{\mathbf{r}} G^{(1)}(\bar{\mathbf{r}}) = 0$. Now we have

$$\frac{d}{dt} G^{(1)}(\mathbf{r}(t)) = -(\mu(\rho_1) - \mu(\rho))(\rho - \rho_1)(\rho - \rho_2)f(\rho_2|\rho_1).$$

Since $\rho_1 \leq \rho \leq \rho_2$ and μ is nondecreasing on $]0, \rho_1^*]$ the right-hand side in the preceding relation is nonpositive if $f(\rho_2|\rho_1) \geq 0$. But using again the Gibbs relation and (34), this can be rewritten

$$(42) \quad f(\rho_2) - f(\rho_1) \geq \mu(\rho_1)(\rho_2 - \rho_1).$$

This convexity inequality holds true for all (ρ_1, ρ_2) such that $\rho_1 \leq \rho_1^*$, by the very definition of ρ_1^* (see the definition (20) of F^{**}). Since we consider pure liquid states, one has $\rho_1 \leq \rho \leq \rho_1^*$. Hence $G^{(1)}$ is dissipated in time, leading to the conclusion.

Metastable states Two types of equilibria are encountered in this situation, with two distinct attraction basins, see Figure 2-right.

- the metastable basin, which appears in yellow in the figure, corresponds to perturbation of a given state within the same phase. It is actually an attraction basin, because the function $G^{(1)}$ defined above (41) is also a Lyapunov function in this domain. Indeed the convexity argument still holds true for any (ρ_1, ρ_2) such that $\rho_2 \leq \rho_-$, since f is convex below ρ_- .
- the unstable basin, corresponding to perturbation of the state by the other phase. This basin is governed by the same Lyapunov function as for the spinodal zone (40).

□

Remark 3.2. *This is a formalization of the remark in Landau & Lifshitz [23, §21] “[...]we must distinguish between metastable and stable equilibrium states. A body in a metastable state may not return to it after a sufficient deviation”*

3.3. Numerical illustrations of the dynamical system behavior. We provide in this paragraph some numerical examples to illustrate the behavior of the dynamical system (36). First we give the characteristic values of the reduced van der Waals EoS at a fixed nondimensionalized temperature $T = 0.85$. Figure 4 presents the corresponding isothermal curve in the (ρ, μ) plan.

The densities ρ^- and ρ^+ correspond to the extrema of the chemical potential. The densities ρ_1^* et ρ_2^* are obtained by the Maxwell’s equal area rule construction on the chemical potential (27) or equivalently by solving (25)-(26).

Table 1 contains the values of the characteristic densities and the corresponding values of the pressure and chemical potential at $T = 0.85$.




FIGURE 2. Attraction basin for pure states. The left figure represents the attraction basin for a pure gaseous state (an equivalent figure may be sketched for a pure liquid state). The attraction points are in red, the non-attainable states being blue. The right figure represent the attraction basin for a metastable vapor state, with the same color coding as the left figure. The attraction basin is pointed out in yellow.





FIGURE 3. Attraction basin for spinodal states: the attraction point is given in red, the blue area refers to non-attainable states.

Using a Backward Differentiation Formula (provided by Scilab for stiff problem), we provide numerical illustrations of the attraction basins depicted in Theorem 3.1. To do so, we fix 50 initial values of $\rho(0)$ in a given interval and set the partial densities $\rho_i(0)$ as perturbations of $\rho(0)$. Then the dynamical system is solved for a final time $T_f = 10^3$ with a time step set to 10^{-3} s. We then provide the graphs of the quantities $\alpha_1(T_f)$, $\rho_i(T_f)$ and $\alpha_1(T_f)p(\rho_1(T_f)) + \alpha_2(T_f)p(\rho_2(T_f))$, plotted with respect to the density $\rho(0)$, for each one of the 50 initial conditions $(\rho(0), \rho_1(0), \rho_2(0))$. The mixture pressure profile is compared with the classical van der Waals pressure curve and the Maxwell line.


FIGURE 4. Isotherm curve in the (ρ, μ) plan at $T = 0.85$.

ρ	$p(\rho)$	$\mu(\rho)$
$\rho^- = 0.5810799446067$	0.62055388470356498	-3.68447967881140137
$\rho^+ = 1.4888047089018$	0.04962960899844759	-4.24339302065563029
$\rho_1^* = 0.31972996451885$	0.504491649787487	-3.97717851100986
$\rho_2^* = 1.8071403273364$		

TABLE 1. Characteristic densities at $T = 0.85$ and corresponding pressure and chemical potential values

Metastable states: The initial density $\rho(0)$ takes on 50 values in $[\rho_1^*, \rho^-]$. For this computation, we set $\rho_1(0) = \rho(0) - 0.1$ and $\rho_2(0) = \rho(0) + 0.2$ so that we can observe the perturbation within and outside the metastable vapor zone. According to Figure 5-top left, the mixture pressure presents two different parts: the left part (for $\rho < 0.45$) remains on the van der Waals pressure curve and a second part coincides with the Maxwell line. The first part corresponds to the perturbation of the metastable vapor state within the phase, while the right part corresponds to the perturbation outside the metastable vapor, that is when $\rho^- \leq \rho_2(0) \leq \rho_2^*$. One can check on the volume fraction curve (see Figure 5-top right) that $\alpha_1(T_f) = 1$ for $\rho(0) \leq 0.45$, that is only the phase 1 is present. Then $0 < \alpha_1 < 1$ and the corresponding final partial densities $\rho_1, 2(T_f)$ coincide with the densities ρ_1^* and ρ_2^* respectively, which explains that the pressure matches with the Maxwell line.

Perturbation of the density ρ in the whole domain: Figure 6 corresponds to an initial density $\rho(0)$ which takes on 50 values between 0.2 and 1.8, while $\rho_1(0) = \rho(0) - 0.1$ and $\rho_2(0) = \rho(0) + 0.1$. One observes that an initial perturbation of the density $\rho(0)$ leads to final states which belong to either pure vapor/liquid states, including metastable states, or the coexistent state. Hence the mixture




FIGURE 5. Numerical illustration of a perturbation within the metastable vapor zone. The initial density $\rho(0)$ takes on 50 values in $[\rho_1^*, \rho^-]$ and $\rho_i(0)$ are perturbations of $\rho(0)$ under the assumption (H_1) . The top-left figure corresponds to the mixture pressure at final time $T_f = 10^3$. For densities $\rho(0) < 0.45$, the pressure coincides with the reduced van der Waals pressure, while for $\rho(0) > 0.45$ it matches with the Maxwell line. One notices that the volume fraction $\alpha_1(T_f)$ is either constant equal to 1 (for $\rho(0) < 0.45$) that is only the phase 1 is present, or takes on values in $]0,1[$ that is the system has reached the coexistent state.

pressure coincides with the admissible branches of the van der Waals pressure curve or with the Maxwell line.

However the convergence is not obvious for $\rho(0)$ close to ρ_- on the left (resp. to ρ_+ on the right). This can be observed in Figure 6 top-right (plot of the volume fraction), where the parts on the left and on the right should be straight lines. Actually, since for ρ close to ρ_- the perturbation chosen is in the spinodal zone, we expect the equilibrium to be on the Maxwell line, which is not the case. We suppose that the final time is not large enough to ensure the actual convergence. As a matter of fact, we observed that the requested time to reach convergence is larger in the metastable zone than in the spinodal one.

4. THE ISOTHERMAL MODEL

This section is devoted to the definition of a 4×4 van der Waals isothermal two-phase flow model of partial differential equations with a relaxation source term accounting for mass transfer.

After defining the model, we state that the homogeneous system is hyperbolic except for densities belonging to the spinodal zone. An interesting feature is that the system boils down to the classical p -system in pure phases that is when $\alpha_1\alpha_2 =$




FIGURE 6. Numerical illustration of a perturbation of $\rho(0)$ in the whole domain. The mixture pressure $\alpha_1(T_f)p(\rho_1(T_f)) + \alpha_2(T_f)p(\rho_2(T_f))$ coincides with the admissible branches of the reduced van der Waals pressure in the pure liquid/vapor states, including metastable state. In the spinodal zone it corresponds to the Maxwell line.

0. We prove that the system admits an energy function that decreases in time. However because of the non convexity of the van der Waals EoS, this energy is non convex.

4.1. Definition of the model. Since we are interested in the modeling of metastable phase transition, the liquid-vapor flows that we consider are submitted to strong thermodynamical perturbations. Hence we propose to depict the dynamic of the flows by a compressible averaged model, namely Euler type equations. In order to model the phase transition, the dynamic part of the model is classically coupled with a source term which carries the mass transfer.

Because we want to model also metastable states, the equilibria of the source term have to be either pure liquid/vapor states, metastable states or the coexistence state given by (27)-(26). Hence we propose to couple the dynamical system (36) introduced in the previous section with a modified version of the isothermal two-phase model proposed in [3].

The system contains the balance equations accounting for the conservation of the total mass and the total momentum of the two-phase flow. Two others equations depict the evolution of the partial densities. Now the unknowns ρ, ρ_1 and ρ_2 are functions of time t and space x . The two phases evolve with the same velocity u . The momentum equation contains a pressure term that reads $\alpha_1 p(\rho_1) + \alpha_2 p(\rho_2)$. For sake of readability we skip the dependence of α_i in ρ, ρ_1 and ρ_2 . The system

reads

$$(43) \quad \begin{cases} \partial_t \rho + \partial_x(\rho u) = 0, \\ \partial_t \rho_1 + \partial_x(\rho_1 u) = -\frac{1}{\varepsilon}(\rho - \rho_1)(\rho - \rho_2)f(\rho_2|\rho_1), \\ \partial_t \rho_2 + \partial_x(\rho_2 u) = \frac{1}{\varepsilon}(\rho - \rho_1)(\rho - \rho_2)f(\rho_1|\rho_2), \\ \partial_t(\rho u) + \partial_x(\rho u^2 + \alpha_1 p(\rho_1) + \alpha_2 p(\rho_2)) = 0, \end{cases}$$

where ε is a relaxation parameter that determines the rate at which the chemical potentials and pressures of the two phases reach equilibrium. The chemical potential μ and the pressure p follow the van der Waals model and are given by (11). The system (43) will be supplemented with initial conditions on the velocity u and on the densities ρ and ρ_i , $i = 1, 2$ satisfying the assumption (H_1).

From (43) one can recover the two equations on the partial masses $\alpha_i \rho_i$ and deal with a system of the classical form

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0, \\ \partial_t(\alpha_1 \rho_1) + \partial_x(\alpha_1 \rho_1 u) = RHS, \\ \partial_t(\alpha_2 \rho_2) + \partial_x(\alpha_2 \rho_2 u) = -RHS, \\ \partial_t(\rho u) + \partial_x(\rho u^2 + \alpha_1 p(\rho_1) + \alpha_2 p(\rho_2)) = 0. \end{cases}$$

In the present paper we choose to focus on the system (43), which allows us to define the densities ρ_i even when $\alpha_i = 0$, which is not the case in the aforementioned system.

4.2. Hyperbolicity and entropy.

Theorem 4.1. *The total Helmholtz free energy \mathcal{F} defined by (30) is convex for $\rho \in]0, 3[$, $u \in \mathbb{R}$ and*

- $(\rho_1, \rho_2) \in (0, \rho) \times ((\rho, \rho_-) \cup (\rho_+, 3))$, if $\rho \leq \rho_-$
- $(\rho_1, \rho_2) \in (0, \rho_-) \times (\rho_+, 3)$, if $\rho \in (\rho_-, \rho_+)$
- $(\rho_1, \rho_2) \in ((0, \rho_-) \times (\rho_+, \rho)) \times (\rho, 3)$, if $\rho \geq \rho_+$

Proof. The function \mathcal{F} is a convex combination of $f(\rho_1)$ and $f(\rho_2)$ where f is the intensive Helmholtz free energy (8). By definition of ρ_- and ρ_+ (see Figure 1), f is convex on $(0, \rho_-] \cup [\rho_+, 3)$, so the result follows. \square

Let us introduce the function

$$(44) \quad \mathcal{E}(\rho, \rho_1, \rho_2, u) = \frac{\rho u^2}{2} + \alpha_1 f(\rho_1) + \alpha_2 f(\rho_2).$$

Using the relations (32) one deduces

Proposition 4.1. *The function (44) satisfies the following equation*

$$(45) \quad \partial_t \mathcal{E} + \partial_x(u(\mathcal{E} + \alpha_1 p(\rho_1) + \alpha_2 p(\rho_2))) = (\partial_{\rho_1} f) \dot{\rho}_1 + (\partial_{\rho_2} f) \dot{\rho}_2 \leq 0.$$

Note that \mathcal{E} is convex where \mathcal{F} is, that is in the domains given in Theorem 4.1. However it is not an entropy for the system since \mathcal{F} is non-convex in the complements of the aforementioned domains.

We turn now to the determination of the eigenvalues of the homogeneous system (43). If we set $\mathbf{Y} = (\rho, \rho_1, \rho_2, u)$, for smooth solutions, the homogeneous system can be written as

$$(46) \quad \partial_t \mathbf{Y} + A(\mathbf{Y}) \partial_X \mathbf{Y} = 0,$$

where the matrix $A(\mathbf{Y})$ is defined by

$$(47) \quad A(\mathbf{Y}) = \begin{pmatrix} u & 0 & 0 & \rho \\ 0 & u & 0 & \rho_1 \\ 0 & 0 & u & \rho_2 \\ A_1(\mathbf{Y}) & A_2(\mathbf{Y}) & A_3(\mathbf{Y}) & u \end{pmatrix},$$

and

$$(48) \quad \begin{aligned} A_1(\mathbf{Y}) &= \frac{p(\rho_1) - p(\rho_2)}{\rho(\rho_1 - \rho_2)}, \\ A_2(\mathbf{Y}) &= \frac{\alpha_1}{\rho(\rho_1 - \rho_2)}(p(\rho_2) - p(\rho_1)) + \frac{\alpha_1}{\rho}p'(\rho_1), \\ A_3(\mathbf{Y}) &= \frac{\alpha_2}{\rho(\rho_1 - \rho_2)}(p(\rho_2) - p(\rho_1)) + \frac{\alpha_2}{\rho}p'(\rho_2). \end{aligned}$$

The characteristic equation of $A(\mathbf{Y})$ is given by

$$(u - \lambda)^2(u - c - \lambda)(u + c - \lambda),$$

with the speed of sound

$$(49) \quad c := c(\mathbf{r}) = \sqrt{\frac{1}{\rho}(\alpha_1(\rho, \rho_1, \rho_2)\rho_1 p'(\rho_1) + \alpha_2(\rho, \rho_1, \rho_2)\rho_2 p'(\rho_2))}.$$

Thus we obtain three distinct eigenvalues for the matrix $A(\mathbf{Y})$:

$$(50) \quad \lambda_1(\mathbf{Y}) = u - c, \quad \lambda_2(\mathbf{Y}) = \lambda_3(\mathbf{Y}) = u, \quad \lambda_4(\mathbf{Y}) = u + c.$$

Note that the eigenvalues are real if \mathbf{r} satisfies

$$\alpha_1(\mathbf{r})\rho_1 p'(\rho_1) + \alpha_2(\mathbf{r})\rho_2 p'(\rho_2) \geq 0.$$

Accounting on relations (9), it is equivalent to the following hyperbolicity condition

$$(51) \quad \alpha_1(\mathbf{r})\rho_1^2 \mu'(\rho_1) + \alpha_2(\mathbf{r})\rho_2^2 \mu'(\rho_2) \geq 0.$$

The right eigenvectors $r_i(\mathbf{Y})$, $i = 1, \dots, 4$, that satisfy $A(\mathbf{Y})r_i(\mathbf{Y}) = \lambda_i(\mathbf{Y})r_i(\mathbf{Y})$ can be chosen as

$$(52) \quad r_1(\mathbf{Y}) = \begin{pmatrix} -\frac{\rho}{c} \\ -\frac{\rho_1}{c} \\ -\frac{\rho_2}{c} \\ 1 \end{pmatrix}, \quad r_2(\mathbf{Y}) = \begin{pmatrix} -\frac{A_3}{A_1} \\ 0 \\ 1 \\ 0 \end{pmatrix}, \quad r_3(\mathbf{Y}) = \begin{pmatrix} -\frac{A_2}{A_1} \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad r_4(\mathbf{Y}) = \begin{pmatrix} \frac{\rho}{c} \\ \frac{\rho_1}{c} \\ \frac{\rho_2}{c} \\ 1 \end{pmatrix},$$

where the quantities A_1 , A_2 and A_3 are defined in (48).

If the densities ρ, ρ_1 and ρ_2 satisfy (H₁) and (51), the matrix $A(\mathbf{Y})$ is diagonalizable in \mathbb{R} and its eigenvectors span the whole space \mathbb{R}^4 so that the system is hyperbolic.

4.3. Structure of the waves. In this paragraph we study the structure of the waves. Assuming that the densities ρ, ρ_1 and ρ_2 satisfy (H₁) and (51), one can observe that the waves are either genuinely non linear or linearly degenerate.

Straightforward computations lead to the following property which will be useful in the sequel.

Proposition 4.2. *The speed of sound c , function of state \mathbf{r} , satisfies the following properties relations*

$$(53) \nabla p(\mathbf{r}) \cdot \mathbf{r} = \rho c^2(\mathbf{r}),$$

$$(54) \quad \nabla c(\mathbf{r}) = \frac{1}{2c(\mathbf{r})} \begin{pmatrix} -\frac{c(\mathbf{r})^2}{\rho} + \frac{1}{\rho(\rho_1 - \rho_2)}(\rho_1 p'(\rho_1) - \rho_2 p'(\rho_2)) \\ \frac{\alpha_1(\mathbf{r})}{\rho} \left(\frac{\rho_2 p'(\rho_2) - \rho_1 p'(\rho_1)}{\rho_1 - \rho_2} + p'(\rho_1) + \rho_1 p''(\rho_1) \right) \\ \frac{\alpha_2(\mathbf{r})}{\rho} \left(\frac{\rho_2 p'(\rho_2) - \rho_1 p'(\rho_1)}{\rho_1 - \rho_2} + p'(\rho_2) + \rho_2 p''(\rho_2) \right) \end{pmatrix}.$$

Let us start with the waves associated to the wave speed $u - c$ and $u + c$.

Proposition 4.3. *The characteristic fields associated to the waves speed $\lambda_1(\mathbf{Y}) = u - c$ and $\lambda_4(\mathbf{Y}) = u + c$ are genuinely non linear i.e. $\nabla_{\mathbf{Y}} \lambda_1(\mathbf{Y}) \cdot r_1(\mathbf{Y}) \neq 0$ and $\nabla_{\mathbf{Y}} \lambda_4(\mathbf{Y}) \cdot r_4(\mathbf{Y}) \neq 0$ for admissible state vector \mathbf{Y} that is for densities (ρ, ρ_1, ρ_2) satisfying (H_1) and (51).*

Proof. We introduce the notation $D(\mathbf{r}) = (c(\mathbf{r}))^2$. We consider the wave associated to the eigenvalue $\lambda_1(\mathbf{Y})$. One has

$$(55) \quad \begin{aligned} \nabla_{\mathbf{Y}} \lambda_1(\mathbf{Y}) \cdot r_1(\mathbf{Y}) &= \begin{pmatrix} \nabla_{\mathbf{r}} c(\mathbf{r}) \\ 1 \end{pmatrix} \cdot r_1(\mathbf{Y}) \\ &= -\frac{1}{2c(\mathbf{r})^2} \left(\rho \frac{\partial D}{\partial \rho} + \rho_1 \frac{\partial D}{\partial \rho_1} + \rho_2 \frac{\partial D}{\partial \rho_2} \right) + 1 \\ &= -\frac{1}{2\rho c(\mathbf{r})^2} (\alpha_1(\mathbf{r}) \rho_1^2 p''(\rho_1) + \alpha_2(\mathbf{r}) \rho_2^2 p''(\rho_2)) + 1. \end{aligned}$$

The densities are assumed to be strictly positive. Under the hypothesis (H_1) and (51) the mass fractions α_i are positive and the second derivative of the van der Waals pressure (11) is a strictly negative function of the density. Thus $\nabla_{\mathbf{Y}} \lambda_1(\mathbf{Y}) \cdot r_1(\mathbf{Y}) \neq 0$. Similarly we can state that $\nabla_{\mathbf{Y}} \lambda_4(\mathbf{Y}) \cdot r_4(\mathbf{Y}) \neq 0$ that conclude the proof. \square

We now study the wave associated to the speed u .

Proposition 4.4. *The characteristic fields associated to the waves $\lambda_2(\mathbf{Y}) = \lambda_3(\mathbf{Y}) = u$ linearly degenerate i.e. $\nabla_{\mathbf{Y}} \lambda_2(\mathbf{Y}) \cdot r_2(\mathbf{Y}) = 0$ and $\nabla_{\mathbf{Y}} \lambda_3(\mathbf{Y}) \cdot r_3(\mathbf{Y}) = 0$ for admissible state vector \mathbf{Y} that is for densities (ρ, ρ_1, ρ_2) satisfying (51).*

Proof. We deduce from the eigenvalues (50) the relation

$$(56) \quad \nabla_{\mathbf{Y}} \lambda_i(\mathbf{Y}) \cdot r_i(\mathbf{Y}) = (0, 0, 0, 1)^T \cdot r_i(\mathbf{Y}),$$

for $i = \{1, 2\}$. Then introducing the right eigenvectors (52) in (56), it is easily checked that $\nabla_{\mathbf{Y}} \lambda_i(\mathbf{Y}) \cdot r_i(\mathbf{Y}) = 0$ for $i = \{1, 2\}$ and this complete the proof. \square

We now address the determination of the Riemann invariants of the system. These computations are made easier using the following property.

Proposition 4.5. *The mass and volume fractions α_1 and φ_1 , defined by (17) and (18), satisfy the following non conservative equations*

$$(57) \quad \begin{aligned} \partial_t \alpha_1 + u \partial_x \alpha_1 &= 0, \\ \partial_t \varphi_1 + u \partial_x \varphi_1 &= 0. \end{aligned}$$

Proposition 4.6. *The Riemann invariants associated to the wave of speed u are*

$$(58) \quad \{u, \bar{p}\},$$

with $\bar{p}(\mathbf{r}) = \alpha_1(\mathbf{r})p(\rho_1) + \alpha_2(\mathbf{r})p(\rho_2)$. The volume and mass fractions are Riemann invariants associated to the wave of speed $u \pm c$:

$$(59) \quad \{\alpha_1, \varphi_1\}.$$

Proof. Because the field associated to the speed u is linearly degenerate, u is clearly a Riemann invariant for this wave. Using the gradient of \bar{p} with respect to \mathbf{r} , a straightforward computation gives

$$(60) \quad \nabla_{\mathbf{Y}} \bar{p}(\mathbf{Y}) \cdot r_2(\mathbf{Y}) = 0.$$

On the other hand the volume fractions α_i and the mass fractions φ_i satisfy the equations (57). Thus the fractions are Riemann invariants for the waves of speed $u \pm c$. \square

The characterization of the third Riemann invariant for the waves of speed $u \pm c$ is more intricate and is not addressed here.

5. NUMERICAL APPROXIMATION

This section is devoted to numerical experiments. We do not wish to elaborate here on efficient numerical schemes for this problem, but merely to illustrate some typical behaviours of the model. Hence we limit ourselves to a simple finite volume scheme, coupled to a time-splitting method for the source terms. Considering the stiffness of the problem, a complete numerical study is mandatory but far beyond the aim of this paper.

5.1. Definition of the splitting strategy. We rewrite the system (43) in a more compact form, considering the following Cauchy problem

$$(61) \quad \begin{aligned} \partial_t W + \partial_x F(W) &= S(W), \\ W(t=0, x) &= W_0(x), \quad \forall x \in \mathbb{R}, \end{aligned}$$

where

$$\begin{aligned} W &= (\rho, \rho_1, \rho_2, \rho u)^T, \\ F(W) &= (\rho u, \rho_1 u, \rho_2 u, \rho u^2 + \alpha_1 p(\rho_1) + \alpha_2 p(\rho_2))^T, \\ S(W) &= \frac{1}{\varepsilon} (0, -(\rho - \rho_1)(\rho - \rho_2) f(\rho_2 | \rho_1), (\rho - \rho_1)(\rho - \rho_2) f(\rho_1 | \rho_2), 0), \end{aligned}$$

and ε is the relaxation parameter. Note that we exclude pure phase initial data, so that the equations on the partial densities are not multiplied by α_i anymore.

Convective terms and source terms are taken into account by a fractional step approach. We denote Δt the time step and Δx the length of the cell $(x_{i-1/2}, x_{i+1/2})$ on the regular mesh. Let W^n be the Finite Volume approximation at time $t^n = n\Delta t$, $n \in \mathbb{N}$. The approximated solution W^{n+1} of the Cauchy problem

$$(62) \quad \begin{cases} \partial_t W + \partial_x F(W) = S(W), & t \in (t^n, t^{n+1}), x \in \mathbb{R}, \\ W(t^n, x) = W^n(x), & \forall x \in \mathbb{R}, \end{cases}$$

is approximated by splitting the problem in two steps. The first one corresponds to the convective part

$$(63) \quad \begin{cases} \partial_t W + \partial_x F(W) = 0, & t \in (t^n, t^{n+1}), x \in \mathbb{R} \\ W(t^n, x) = W^n(x), & \forall x \in \mathbb{R}, \end{cases}$$

which provides $W^{n,-}$. The second step corresponds to the relaxation process

$$(64) \quad \begin{cases} \partial_t W = S(W), & t \in (t^n, t^{n+1}), x \in \mathbb{R} \\ W(t^n, x) = W^{n,-}(x), & \forall x \in \mathbb{R}, \end{cases}$$

which finally gives W^{n+1} .

Numerical scheme for the convective part. We consider a classical HLL numerical flux. We adopt the following classical notation

$$(65) \quad W_i^n = \frac{1}{\Delta x} \int_{x_{i-1/2}}^{x_{i+1/2}} W(t^n, x) dx, \quad n \geq 0, \quad i \in \mathbb{R}.$$

The scheme is the following

$$(66) \quad \Delta x (W_i^{n,-} - W_i^n) + \Delta t (F_{i+1/2}^n - F_{i-1/2}^n) = 0,$$

together with

$$F_{i+1/2}^n = \begin{cases} F(W_i^n), & \text{if } 0 \leq s_L, \\ \frac{s_R F(W_i^n) - s_L F(W_{i+1}^n) + s_L s_R (W_{i+1}^n - W_i^n)}{s_R - s_L}, & \text{if } s_L \leq 0 \leq s_R, \\ F(W_{i+1}^n), & \text{if } 0 \geq s_R, \end{cases}$$

where $s_R = \max(u_i^n + c_i^n, u_{i+1}^n + c_{i+1}^n)$ and $s_L = \min(u_i^n - c_i^n, u_{i+1}^n - c_{i+1}^n)$. The time step is subjected to the classical CFL condition

$$(67) \quad \frac{\Delta t}{\Delta x} |\lambda_{\max}| \leq 1,$$

where λ_{\max} is the maximal speed of wave computed on each cell of the mesh.

Numerical treatment of the source terms. The initial condition for this step is $W^{n,-}$ which is assumed to be admissible, that is $(\rho^{n,-}, \rho_1^{n,-}, \rho_2^{n,-}) \in \mathcal{R}$. The total density ρ and the momentum ρu remain unchanged during this step. Only the densities ρ_1 and ρ_2 may vary which leads to the following system

$$(68) \quad \begin{aligned} \partial_t \rho &= \partial_t (\rho u) = 0, \\ \partial_t \rho_1 &= -\frac{1}{\varepsilon} (\rho - \rho_1) (\rho - \rho_2) f(\rho_2 | \rho_1), \\ \partial_t \rho_2 &= \frac{1}{\varepsilon} (\rho - \rho_1) (\rho - \rho_2) f(\rho_1 | \rho_2). \end{aligned}$$

At this stage we merely use a classic explicit order 4 Runge-Kutta method to integrate the source term.

Such a treatment enforces tough constraints on the time step: the computations were performed with 1000 iterations using a time step of 10^{-6} . We emphasize that this does not ensure the actual convergence to the equilibrium state. This pleads for a more efficient method, for instance a semi-implicit scheme in the spirit of [7].

5.2. Numerical results. We present here numerical results that assess the ability of the model (43) to capture phase transition and metastable states.

We consider the van der Waals pressure in its reduced form (13) at a constant subcritical temperature $T = 0.85$. At this fixed temperature the extrema ρ^- and ρ^+ of the pressure and the values ρ_1^* and ρ_2^* defined by the Maxwell construction on the chemical potential are given in Table 1.

We propose test cases with Riemann initial conditions that is

$$(69) \quad W(t = 0, x) = w_0(x) = \begin{cases} W_L, & \text{if } x \leq 0, \\ W_R, & \text{if } x > 0. \end{cases}$$

The following test cases are set on the domain $[0, 1]$, with an uniform mesh of 10000 cells and Neumann boundary conditions at $x = 0$ and $x = 1$.




FIGURE 7. Riemann problem with phase transition: density ρ , volume fraction α_1 and pressure p .

5.2.1. *Riemann problem with phase transition.* The initial state W_L and W_R are

$$\begin{aligned} \rho_L = \rho_{1,L} = 0.3, \quad \rho_{2,L} = \rho_2^*, \quad u_L = 0, \\ \rho_R = \rho_{2,R} = 1.9, \quad \rho_{1,R} = \rho_1^*, \quad u_R = 0. \end{aligned}$$

The left state is a pure stable gas and the right state is a pure stable liquid. Various value of ε are considered. The solution is at time $t = 0.1s$. One can observe the appearance of a mixture zone on both sides of the interface, see in particular the pressure profile Figure 7-bottom. The results with $\varepsilon = 10^{-4}$ or 10^{-6} are similar expect on the velocity profile (see Figure 8-bottom) where the intermediate state on the left of the interface is slightly modified for $\varepsilon = 10^{-4}$.




FIGURE 8. Riemann problem with phase transition: partial densities ρ_1 and ρ_2 (top) and velocity u .

5.2.2. *Cavitation by double rarefaction.* The test consists in a liquid state submitted to a double rarefaction wave. The initial state is given by $\rho = \rho_2^* - 10^{-3}$, $\rho_2 = \rho_2^*$, $\rho_1 = \rho_1^*$ and the velocities are $u_R = 0.3 = -u_L$. The total density corresponds to a metastable liquid, and the initial volume fraction is $\alpha_1 \simeq 0.000672$, which means that phase 2 is predominant. The solution is computed at time $t = 0.1$ s. We observe on the plot of the volume fraction (Figure 9-top) that a bubble of stable vapor appears around the interface $x = 0.5$. The value of ϵ does not modify the profile of the bubble. However the pressure profile is sharper for $\epsilon = 10^{-6}$ than for 10^{-4} , see Figure 9-bottom.

5.2.3. *Nucleation by double shock.* The test consists in a pure stable gaseous state submitted to a double shock wave. The initial state is given by $\rho = \rho_1 = 0.3$,




FIGURE 9. Cavitation by double rarefaction: density ρ and volume fraction α_1 (top) and pressure p .

$\rho_2 = 0.1$ and the velocities are $u_R = -0.3 = -u_L$. The solution is computed at time $t = 0.1s$. Note that phase 2 is not present initially but is fixed in the spinodal zone. We observe two different behaviours of the solution depending on the value of ε . For $\varepsilon = 10^{-6}$, the profile of the volume fraction (see Figure 11-top) shows that a droplet of liquid appears around $x = 0.5$ which is not the case for $\varepsilon = 10^{-4}$ where there is no droplet. For $\varepsilon = 10^{-6}$, one observes on the density plot (see Figure 11-top) that the liquid state inside the droplet admits a density ρ close to ρ_2^* with small oscillations. The droplet is surrounded by two mixture areas with $\rho = \rho_1^*$. The pressure curve inside the droplet presents oscillations (see Figure 11-bottom) which might be due to a loss of hyperbolicity.




FIGURE 10. Cavitation by double rarefaction: partial densities ρ_1 and ρ_2 (top) and velocity u .

5.2.4. *Acoustic perturbation of a metastable state.* This example consists in a constant metastable vapor state, with a perturbation in the velocity. The initial state is $\rho = 0.42$, $\rho_1 = 0.32$, $\rho_2 = 0.52$ and the velocities are $u_L = 0.4$, $u_R = 0$. Both densities ρ_1 and ρ_2 are in the metastable state, and we impose a compression from the left with velocity 0.4.

The compression induces the appearance of droplet of pure liquid which moves from the left to the right, see the time evolution on Figure 13. With a smaller velocity perturbation the structure of the waves is similar, but there is no creation of a droplet at the interface. One can check on Figure 13 that the density ρ inside the droplet is larger than ρ_2^* . The droplet is surrounded by two areas with a mixture state with $\rho = \rho_1^*$. The velocity and pressure profiles exhibit spikes on both sides




FIGURE 11. Nucleation by double shock: density ρ and volume fraction α_1 (top) and pressure p .

of the droplet, which amplitude decreases when ϵ decreases, see Figures 14 and 15. Notice also that the pressure in the mixture zone is not at the value p^* . It seems that when ϵ decreases the value is closer, this may indicate that the source term has not reached the equilibrium state yet.

6. CONCLUSION

The core of this work is the formalization in terms of a dynamical system of the thermodynamics of phase transition, using the van der Waals EoS. It leads to a mathematical characterization of metastable states, compared to stable coexistent two-phase states. The dynamical behaviour of the solutions is crucial here, and seems to preclude the use of instantaneous exchange kinetic. When coupled to a




FIGURE 12. Nucleation by double shock: partial densities ρ_1 and ρ_2 (top) and velocity u .

simple hydrodynamic model, namely the isothermal Euler equations, it evidences abilities to cope with metastable states as well as bubble or droplet generation. This preliminary study gives rise to a wide bunch of open questions and problems, and we believe that the methodology can be used in a much larger context.

First, in the same isothermal context, the construction of the dynamical system (the right-hand side in the extended Euler equations) can be addressed. We deliberately used a simple and readable function, which possibly could be improved. In any case, the behaviour of the coexistence zone around the interface has to be investigated in more details, as well as the role of ε .

Next, an obvious mandatory issue is the numerical treatment of the coupled system. We have chosen here the simplest numerical strategy that allowed us to




FIGURE 13. Velocity perturbation of a metastable state: density ρ at times 0.1, 0.2, 0.4




FIGURE 14. Velocity perturbation of a metastable state: velocity u at times 0.1, 0.2, 0.4

illustrate our purpose. The explicit treatment of the stiff relaxation term enforces tough constraints on the time step, and prevents the simulation of more realistic metastable cases.




FIGURE 15. Velocity perturbation of a metastable state: pressure p at times 0.1, 0.2, 0.4

Finally, we attend to include temperature dependance to obtain a fully heat, mass and mechanical transfer model in order to compare our results to those of [29] and [32].

REFERENCES

- [1] R. Abeyaratne and J. K. Knowles. Kinetic relations and the propagation of phase boundaries in solids. *Arch. Rational Mech. Anal.*, 114(2):119–154, 1991.
- [2] G. Allaire, G. Faccanoni, and S. Kokh. A strictly hyperbolic equilibrium phase transition model. *C. R. Math. Acad. Sci. Paris*, 344(2):135–140, 2007.
- [3] A. Ambroso, C. Chalons, F. Coquel, and T. Galié. Relaxation and numerical approximation of a two-fluid two-pressure diphasic model. *M2AN Math. Model. Numer. Anal.*, 43(6):1063–1097, 2009.
- [4] M. R. Baer and J. W. Nunziato. A two phase mixture theory for the deflagration to detonation (ddt) transition in reactive granular materials. *Int. J. Multiphase Flow*, 12(6):861–889, 1986.
- [5] T. Barberon and P. Helluy. Finite volume simulation of cavitating flows. *Computers and Fluids*, 34(7):832–858, 2005.
- [6] N. Bedjaoui, C. Chalons, F. Coquel, and P. G. LeFloch. Non-monotonic traveling waves in van der Waals fluids. *Anal. Appl. (Singap.)*, 3(4):419–446, 2005.
- [7] S. Boscarino and G. Russo. On a class of uniformly accurate IMEX Runge-Kutta schemes and applications to hyperbolic systems with relaxation. *SIAM J. Sci. Comput.*, 31(3):1926–1945, 2009.
- [8] H. B. Callen. *Thermodynamics and an introduction to thermostatistics, second edition*. Wiley and Sons, 1985.
- [9] A. Corli and H. Fan. The Riemann problem for reversible reactive flows with metastability. *SIAM J. Appl. Math.*, 65(2):426–457 (electronic), 2004/05.
- [10] J.-M. Delhayé, M. Giot, L. Mahias, P. Raymond, and C. Renault. *Thermohydraulique des racteurs*. EDP Sciences, 1998.
- [11] J.-M. Delhayé, M. Giot, and M. I. Riethmüller. *Thermohydraulics of two-phase systems for industrial design and nuclear engineering*. Hemisphere Publishing Corporation, 1981.
- [12] D.A. Drew. Mathematical modeling of two-phase flow. *Ann. Rev. Fluid Mech.*, 15:261–291, 1983.

- [13] G. Faccanoni, S. Kokh, and G. Allaire. Modelling and simulation of liquid-vapor phase transition in compressible flows based on thermodynamical equilibrium. *ESAIM Math. Model. Numer. Anal.*, 46(5):1029–1054, 2012.
- [14] H. Fan and X.-B. Lin. A dynamical systems approach to traveling wave solutions for liquid/vapor phase transition. In *Infinite dimensional dynamical systems*, volume 64 of *Fields Inst. Commun.*, pages 101–117. Springer, New York, 2013.
- [15] J.W. Gibbs. *The Collected Works of J. Willard Gibbs, vol I: Thermodynamics*. Yale University Press, 1948.
- [16] P. Helluy and H. Mathis. Pressure laws and fast Legendre transform. *Math. Models Methods Appl. Sci.*, 21(4):745–775, 2011.
- [17] P. Helluy and N. Seguin. Relaxation models of phase transition flows. *M2AN Math. Model. Numer. Anal.*, 40(2):331–352, 2006.
- [18] J.-B. Hiriart-Urruty and C. Lemaréchal. *Fundamentals of convex analysis*. Grundlehren Text Editions. Springer-Verlag, Berlin, 2001.
- [19] A.R. Imre, I.F. Barna, G. zsl, G. Hzi, and T. Kraska. Theoretical study of flashing and water hammer in a supercritical water cycle during pressure drop. *Nuclear Engineering and Design*, 240(6):1569 – 1574, 2010.
- [20] F. Jaegle, C. Rohde, and C. Zeiler. A multiscale method for compressible liquid-vapor flow with surface tension. In *CEMRACS’11: Multiscale coupling of complex models in scientific computing*, volume 38 of *ESAIM Proc.*, pages 387–408. EDP Sci., Les Ulis, 2012.
- [21] F. James, M. Sepúlveda, and P. Valentin. Statistical thermodynamics models for multicomponent isothermal diphasic equilibria. *Math. Models Methods Appl. Sci.*, 7(1):1–29, 1997.
- [22] S. Jaouen. *Etude mathématique et numérique de stabilité pour des modèles hydrodynamiques avec transition de phase*. PhD thesis, Université Paris VI, November 2001.
- [23] L. Landau and E. Lifschitz. *A Course of theoretical physics, vol 5, Statistical Physics*. Pergamon Press, 1959-1969.
- [24] P. G. LeFloch. Kinetic relations for undercompressive shock waves. Physical, mathematical, and numerical issues. In *Nonlinear partial differential equations and hyperbolic wave phenomena*, volume 526 of *Contemp. Math.*, pages 237–272. Amer. Math. Soc., Providence, RI, 2010.
- [25] H. Mathis. *Étude théorique et numérique des écoulements avec transition de phase*. PhD thesis, Université de Strasbourg, 2010.
- [26] A. Murrone and H. Guillard. A five equation reduced model for compressible two phase flow problems. *J. Comput. Phys.*, 202(2):664–698, 2005.
- [27] M. Pelanti and K.-M. Shyue. A mixture-energy-consistent six-equation two-phase numerical model for fluids with interfaces, cavitation and evaporation waves. *J. Comput. Phys.*, 259:331–357, 2014.
- [28] R.T. Rockafellar. *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. Reprint of the 1970 original, Princeton Paperbacks.
- [29] R. Saurel, F. Petitpas, and R. Abgrall. Modelling phase transition in metastable liquids: application to cavitating and flashing flows. *J.Fluid Mech.*, 607:313–350, 2008.
- [30] M. Slemrod. Admissibility criteria for propagating phase boundaries in a van der Waals fluid. *Arch. Rational Mech. Anal.*, 81(4):301–315, 1983.
- [31] M. Slemrod. Dynamic phase transitions in a van der Waals fluid. *J. Differential Equations*, 52(1):1–23, 1984.
- [32] A. Zein, M. Hantke, and G. Warnecke. Modeling phase transition for compressible two-phase flows applied to metastable liquids. *J. Comp. Phys.*, 229:1964–2998, 2010.

MATHÉMATIQUES – ANALYSE, PROBABILITÉS, MODÉLISATION – ORLÉANS (MAPMO), UNIVERSITÉ D’ORLÉANS & CNRS UMR 7349, BP 6759, F-45067 ORLÉANS CEDEX 2, FRANCE
E-mail address: francois.james@univ-orleans.fr

LABORATOIRE JEAN LERAY, UNIVERSITÉ DE NANTES & CNRS UMR 6629, BP 92208, F-44322 NANTES CEDEX 3, FRANCE
E-mail address: helene.mathis@univ-nantes.fr