



HAL
open science

Compact Brownian surfaces I. Brownian disks

Jérémie Bettinelli, Gregory Miermont

► **To cite this version:**

Jérémie Bettinelli, Gregory Miermont. Compact Brownian surfaces I. Brownian disks. 2015. hal-01182224v2

HAL Id: hal-01182224

<https://hal.science/hal-01182224v2>

Preprint submitted on 20 Oct 2015 (v2), last revised 12 Feb 2016 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Compact Brownian surfaces I. Brownian disks

Jérémie Bettinelli* Grégory Miermont†

October 20, 2015

Abstract

We show that, under certain natural assumptions, large random plane bipartite maps with a boundary converge after rescaling to a one-parameter family $(BD_L, 0 < L < \infty)$ of random metric spaces homeomorphic to the closed unit disk of \mathbb{R}^2 , the space BD_L being called the *Brownian disk of perimeter L* and unit area. These results can be seen as an extension of the convergence of uniform plane quadrangulations to the Brownian map, which intuitively corresponds to the limit case where $L = 0$. Similar results are obtained for maps following a Boltzmann distribution, in which the perimeter is fixed but the area is random.

Contents

1	Introduction	2
1.1	Motivation	2
1.2	Maps	3
1.3	The case of quadrangulations	4
1.4	Universal aspects of the limit	5
1.5	Convergence of Boltzmann maps	10
1.6	Further comments and organization of the paper	11
2	Definition of Brownian disks	12
2.1	First-passage bridges and random continuum forests	12
2.2	Snakes	14
2.3	Brownian disks	15
3	The Schaeffer bijection and two variants	16
3.1	The original Cori–Vauquelin–Schaeffer bijection	16
3.2	Slices	18
3.3	Plane quadrangulations with a boundary	18

*CNRS & Institut Élie Cartan de Lorraine

†ENS de Lyon & Institut Universitaire de France

1	INTRODUCTION	2
4	Proof of Theorem 1	20
4.1	Subsequential convergence	20
4.2	Identification of the limit	23
5	Boltzmann random maps and well-labeled mobiles	28
5.1	The Bouttier–Di Francesco–Guitter bijection	28
5.2	Random mobiles	29
6	Convergence of the encoding processes	31
6.1	Convergence for an infinite forest	32
6.2	Convergence for a conditioned forest	34
6.3	Convergence of the white contour and label processes	38
7	Proof of the invariance principle	39
7.1	Convergence of conditioned pointed maps	40
7.2	De-pointing	43
7.3	Proof of the convergence of Boltzmann maps	44

1 Introduction

1.1 Motivation

Random maps are a natural discrete version of random surfaces. It has been shown in recent years that their scaling limits can provide “canonical” models of random metric spaces homeomorphic to a surface of a given topology. More precisely, given a random planar map M , one can consider it as a random finite metric space by endowing its vertex set with the usual graph metric, and multiply this graph metric by a suitable renormalizing factor that converges to 0 as the size of the map M is sent to infinity. One is then interested in the convergence in distribution of the resulting sequence of rescaled maps, in the Gromov–Hausdorff topology [22] (or pointed Gromov–Hausdorff topology if one is interested in non-compact topologies), to some limiting random metric space.

Until now, the topology for which this program has been carried out completely is that of the sphere, for a large (and still growing) family of different random maps models, see [26, 34, 6, 2, 11, 1], including for instance the case of uniform triangulations of the sphere with n faces, or uniform random maps of the sphere with n edges. The limiting metric space, called the Brownian map, turns out not only to have the topology of the sphere [28, 33], as can be expected, but also to be independent (up to a scale constant) of the model of random maps that one chooses, provided it is, in some sense, “reasonable.” See however [3, 27] for natural models of random maps that converge to qualitatively different metric spaces. These two facts indeed qualify the Brownian map as being a canonical random geometry on the sphere. Note that a non-compact variant of the Brownian map, called the *Brownian plane*, has been introduced in [20] and shown to be the scaling limit of some natural models of random quadrangulations.

However, for other topologies allowing higher genera and boundary components, only partial results are known [7, 8, 10, 9]. Although subsequential convergence results have been obtained for rescaled random maps in general topologies, it has not been shown that the limit is uniquely defined and independent of the choice of the extraction. The goal of this paper and its companion [12] is to fill in this gap by showing convergence of a natural model of random maps on a given compact surface S to a random metric space with same topology, which one naturally can call the “Brownian S .”

This paper will focus exclusively on the particular case of the disk topology, which requires quite specific arguments, and indeed serves as a building block to construct the boundaries of general compact Brownian surfaces in [12].

1.2 Maps

To state our results, let us recall some important definitions and set some notation. We first define the objects that will serve as discrete models for a metric space with the disk topology.

A *plane map* is an embedding of a finite connected multigraph into the 2-dimensional sphere, and considered up to orientation-preserving homeomorphisms of the latter. The *faces* of the map are the connected components of the complement of edges, and can be then shown to be homeomorphic to 2-dimensional open disks. For every oriented edge e , with origin vertex v , we can consider the oriented edge e' that follows e in counterclockwise order around v , and define the *corner* incident to e as a small open angular sector between e and e' . It does not matter how we choose these regions as long as they are pairwise disjoint. The number of corners contained in a given face f is called the *degree* of that face; equivalently, it is the number of oriented edges to the left of which f lies — we say that f is *incident* to these oriented edges, or to the corresponding corners. We let $\mathbf{V}(\mathbf{m})$, $\mathbf{E}(\mathbf{m})$, $\mathbf{F}(\mathbf{m})$ denote the sets of vertices, edges and faces of a map \mathbf{m} , or simply \mathbf{V} , \mathbf{E} , \mathbf{F} when the mention of \mathbf{m} is clear from the context.

If \mathbf{m} is a map, we can view it as a metric space $(\mathbf{V}(\mathbf{m}), d_{\mathbf{m}})$, where $d_{\mathbf{m}}$ is the graph metric on the set $\mathbf{V}(\mathbf{m})$ of vertices of \mathbf{m} . For simplicity, we will sometimes denote this metric space by \mathbf{m} as well and, if $a > 0$, we denote by $a\mathbf{m}$ the metric space $(\mathbf{V}(\mathbf{m}), ad_{\mathbf{m}})$.

For technical reasons, the maps we consider will always implicitly be *rooted*, which means that one of the corners (equivalently, one of the oriented edges) is distinguished and called the *root*. The face f_* incident to the root is called the *root face*. Since we want to consider objects with the topology of a disk, we insist that the root face is an *external face* to the map, whose incident edges forms the *boundary* of the map, and call its degree the *perimeter* of the map. By contrast, the non-root faces are called *internal faces*. Note that the boundary of the external face is in general not a simple curve (see Figure 1). As a result, the topological space obtained by removing the external face from the surface in which the map is embedded is not necessarily a surface with a boundary, in the sense that every point does not have a neighborhood homeomorphic to some open set of $\mathbb{R} \times \mathbb{R}_+$. However, removing any Jordan domain from the external face does of course result in a surface with a boundary, which is homeomorphic to the 2-dimensional disk.

1.3 The case of quadrangulations

The first part of the paper is concerned exclusively with a particular family of maps, for which the results are the simplest to obtain and to state. A *quadrangulation with a boundary* is a rooted plane map whose internal faces all have degree 4. It is a simple exercise to see that this implies in fact that the perimeter is necessarily an even number. For $l, n \in \mathbb{N}$, we let $\mathbf{Q}_{l,n}$ be the set of quadrangulations with a boundary having n internal faces and perimeter $2l$.




Figure 1: A quadrangulation from $\mathbf{Q}_{23,19}$. The root is the corner indicated by the red arrow.

Our main result in the context of random quadrangulations is the following.

Theorem 1. *Let $L \in [0, \infty)$ be fixed, and $(l_n, n \geq 1)$ be a sequence of integers such that $l_n/\sqrt{2n} \rightarrow L$ as $n \rightarrow \infty$. Let Q_n be uniformly distributed over $\mathbf{Q}_{l_n,n}$. There exists a random compact metric space BD_L such that*

$$\left(\frac{9}{8n}\right)^{1/4} Q_n \xrightarrow[n \rightarrow \infty]{(d)} \text{BD}_L$$

where the convergence holds in distribution for the Gromov–Hausdorff topology.

The random metric space BD_L is called the *Brownian disk with perimeter L* and unit area. We will give in Section 2 an explicit description of BD_L (as well as versions with general areas, see also Section 1.5) in terms of certain stochastic processes, and the convention for the scaling constant $(9/8)^{1/4}$ is here to make the description of these processes simpler. The main properties of BD_L are the following; they follow from [10, Theorems 1–3].

Proposition 2. *Let $L > 0$ be fixed. Almost surely, the space BD_L is homeomorphic to the closed unit disk of \mathbb{R}^2 . Moreover, almost surely, the Hausdorff dimension of BD_L is 4, while that of its boundary ∂BD_L is 2.*

We stress that the case $L = 0$, corresponding to the situation where $l_n = o(\sqrt{n})$, is the statement of [10, Theorem 4], which says that BD_0 is the so-called Brownian map. Since the Brownian map is a.s. homeomorphic to the sphere [28], this means that the boundaries of the approximating random maps are too small to be seen in the limit. This particular case generalizes the convergence of uniform random quadrangulations, obtained in [26, 34], corresponding to the case where $l_n = 2$ for every $n \geq 1$.

The case where $l_n/\sqrt{n} \rightarrow \infty$ is also of interest, and is the object of [10, Theorem 5], showing that, in this case, $(2l_n)^{-1/2}Q_n$ converges to the so-called Brownian Continuum Random Tree [4, 5]. This means that the boundary takes over the planar geometry and folds the map into a tree-shaped object.

We will prove our result by using the already studied case of plane maps without boundary, together with some surgical methods. Heuristically, we will cut Q_n along certain geodesics into elementary pieces of planar topology, to which we can apply a variant of the convergence of random spherical quadrangulations to the Brownian map. The idea of cutting into *slices* quadrangulations with a boundary along geodesics appears in Bouttier and Guitter [15, 16]. The use of these slices (also called *maps with a piecewise geodesic boundary*) plays an important role in Le Gall’s approach [26] to the uniqueness of the Brownian map in the planar case, which requires to introduce the scaling limits of these slices. The previously cited works are influential to our approach. It however requires to glue an infinite number of metric spaces along geodesic boundaries, which could create potential problems when passing to the limit.

1.4 Universal aspects of the limit

Another important aspect is that of *universality* of the spaces BD_L . Indeed, we expect these spaces to be the scaling limit of many other models of random maps with a boundary, as in the case of the Brownian map, which corresponds to $L = 0$. In the latter case, it has indeed been proved, starting in Le Gall’s work [26], that the Brownian map is the unique scaling limit for a large family of natural models of discrete random maps, see [6, 2, 11, 1]. The now classical approach to universality developed in [26] can be generalized to our context, as we illustrate in the case of *critical bipartite Boltzmann maps*.

1.4.1 Boltzmann random maps

Let \mathbf{B} be the set of bipartite rooted plane maps, that is, the set of rooted plane maps with faces all having even degrees (equivalently, this is the set of maps whose internal faces all have even degrees). For $l \in \mathbb{Z}_+$, let \mathbf{B}_l be the set of bipartite maps $\mathbf{m} \in \mathbf{B}$ with perimeter¹ $2l$. Note that when $l = 1$, meaning that the root face has degree 2, there is a natural bijection between \mathbf{B}_1 and $\mathbf{B} \setminus \mathbf{B}_0$, consisting in gluing together the two edges of the root face into one edge.

Let $q = (q_1, q_2, \dots)$ be a sequence of non-negative *weights*. We assume throughout that $q_i > 0$ for at least one index $i \geq 2$. The Boltzmann measure associated with the sequence q is the measure $W(q; \cdot)$ on \mathbf{B} defined by

$$W(q; \mathbf{m}) = \prod_{f \in \mathbf{F}(\mathbf{m}) \setminus \{f_*\}} q_{\deg(f)/2}.$$

¹By convention, the vertex map \circ consisting of no edges and only one vertex, “bounding” a face of degree 0, is considered as an element of \mathbf{B} , so that $\mathbf{B}_0 = \{\circ\}$. It will only appear incidentally in the analysis.

This defines a non-negative, σ -finite measure, and by convention the vertex-map receives a weight $W(q, \circ) = 1$. In what follows, the weight sequence q is considered fixed and its mention will be implicit, so that we denote for example $W(\mathbf{m}) = W(q; \mathbf{m})$, and likewise for the variants of W to be defined below.

We aim at understanding various probability measures obtained by conditioning W with respect to certain specific subsets of \mathbf{B} . It is a simple exercise to check that $W(\mathbf{B}_l)$ is non-zero for every $l \in \mathbb{N}$, and that $W(\mathbf{B}_l)$ is finite for one value of $l > 0$ if and only if it is finite for all values of $l > 0$. In this case, it makes sense to define the Boltzmann probability measures

$$\mathbb{W}_l = W(\cdot | \mathbf{B}_l) = \frac{W(\cdot \cap \mathbf{B}_l)}{W(\mathbf{B}_l)}, \quad l \geq 0.$$

A random map with distribution \mathbb{W}_l has a root face of fixed degree $2l$, but a random number of vertices, edges and faces.

Likewise, we can consider conditioned versions of W given both the perimeter and the “size” of the map, where the size can be alternatively the number of vertices, edges or internal faces². We let $\mathbf{B}_{l,n}^{\mathbf{V}}$, $\mathbf{B}_{l,n}^{\mathbf{E}}$, $\mathbf{B}_{l,n}^{\mathbf{F}}$ be the subsets of \mathbf{B}_l consisting of maps with respectively $n + 1$ vertices, n edges and n internal faces. (The choice of $n + 1$ vertices instead of a more natural choice of n vertices is technical and will make the statements simpler.)

In all the statements involving a given weight sequence q and a symbol $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$ (for “size”), it will always be tacitly imposed that (l, n) belongs to the set

$$\mathcal{E}^{\mathbf{S}}(q) = \{(l, n) \in \mathbb{Z}_+^2 : W(\mathbf{B}_{l,n}^{\mathbf{S}}) > 0\}.$$

Note that for $(l, n) \in \mathcal{E}^{\mathbf{S}}(q)$, it holds that $W(\mathbf{B}_{l,n}^{\mathbf{S}}) < \infty$ since $W(\mathbf{B}_l) < \infty$. In this way, we can define the distribution

$$\mathbb{W}_{l,n}^{\mathbf{S}}(\cdot) = W(\cdot | \mathbf{B}_{l,n}^{\mathbf{S}}).$$

It will be useful in the following to know what the set $\mathcal{E}^{\mathbf{S}}(q)$ looks like. More precisely, let

$$\mathcal{E}_l^{\mathbf{S}}(q) = \{n \geq 0 : (l, n) \in \mathcal{E}^{\mathbf{S}}(q)\}. \quad (1)$$

As above, when the weight sequence q is unequivocally fixed, we will drop the mention of it from the notation and write $\mathcal{E}^{\mathbf{S}}$ and $\mathcal{E}_l^{\mathbf{S}}$.

Define three numbers $h^{\mathbf{V}}$, $h^{\mathbf{E}}$, $h^{\mathbf{F}}$ by

$$h^{\mathbf{V}} = \gcd(\{n \geq 2 : q_{2n} > 0\}), \quad h^{\mathbf{E}} = \gcd(\{n \geq 1 : q_{2n} > 0\}), \quad h^{\mathbf{F}} = 1. \quad (2)$$

Then we have the following lemma, which is a slight generalization of [38, Section 6.3.1].

²We could also consider other ways to measure the size of a map \mathbf{m} , e.g. considering combinations of the form $x_{\mathbf{V}}|\mathbf{V}(\mathbf{m})| + x_{\mathbf{E}}|\mathbf{E}(\mathbf{m})| + x_{\mathbf{F}}|\mathbf{F}(\mathbf{m})|$ for some $x_{\mathbf{V}}, x_{\mathbf{E}}, x_{\mathbf{F}} \geq 0$ with sum 1 as is done for instance in [38] (in fact, due to the Euler formula, there is really only one degree of freedom rather than two). We will not address this here but we expect our results to hold in this context as well.

Lemma 3. *Let q be a weight sequence, and let \mathbf{S} be one of the three symbols \mathbf{V} , \mathbf{E} , \mathbf{F} . There exists an integer $\beta^{\mathbf{S}} \geq 0$ such that for every $l \geq 1$, there exists a set $R_l^{\mathbf{S}} \subset \{0, 1, \dots, l\beta^{\mathbf{S}} - 1\}$ such that*

$$\mathcal{E}_l^{\mathbf{S}}(q) = R_l^{\mathbf{S}} \cup (l\beta^{\mathbf{S}} + h^{\mathbf{S}}\mathbb{Z}_+).$$

In fact, note that $\mathcal{E}_l^{\mathbf{F}}(q) = \mathbb{Z}_+$, which amounts to the fact that, for any q and any $n \geq 0$, $l \geq 1$, there is at least one map \mathbf{m} with n internal faces and perimeter $2l$ such that $W(q; \mathbf{m}) > 0$. As a consequence, we can always take $\beta^{\mathbf{F}} = 0$.

1.4.2 Admissible, regular critical weight sequences

Let us introduce some terminology taken from [30]. Let

$$f_q(x) = \sum_{k \geq 0} x^k \binom{2k+1}{k} q_{k+1}, \quad x \geq 0.$$

This defines a totally monotone function with values in $[0, \infty]$.

Definition 4. *We say that q is admissible if the equation*

$$f_q(z) = 1 - \frac{1}{z} \tag{3}$$

admits a solution $z > 1$. We also say that q is regular critical if moreover this solution satisfies

$$z^2 f_q'(z) = 1$$

and if there exists $\varepsilon > 0$ such that $f_q(z + \varepsilon) < \infty$.

Note that q being regular critical means that the graphs of f_q and of $x \mapsto 1 - 1/x$ are tangent at the point of abscissa z , and in particular, by convexity of f_q , the solution z to (3) is unique. We denote by

$$\mathcal{Z}_q = z$$

this solution, which will play an important role in the discussion to come.

To give a little more insight into this definition, let us introduce at this point a measure on maps that looks less natural at first sight than the Boltzmann measure \mathbb{W}_l , but which will turn out to be better-behaved from the bijective point of view on which this work relies. Let \mathbf{B}^\bullet be the set of pairs (\mathbf{m}, v_*) where $\mathbf{m} \in \mathbf{B}$ is a rooted bipartite map and $v_* \in \mathbf{V}(\mathbf{m})$ is a distinguished vertex. We also let \mathbf{B}_l^\bullet be the subset of \mathbf{B}^\bullet consisting of the maps having perimeter $2l$. We let $W^\bullet(q; \cdot)$ be the measure on \mathbf{B}^\bullet defined by

$$W^\bullet(q; \{(\mathbf{m}, v_*)\}) = W(\mathbf{m}), \quad (\mathbf{m}, v_*) \in \mathbf{B}^\bullet, \tag{4}$$

as well as the probability measures \mathbb{W}^\bullet and \mathbb{W}_l^\bullet , defined by conditioning W^\bullet respectively on \mathbf{B}^\bullet and \mathbf{B}_l^\bullet . Note that, if $\phi(\mathbf{m}, v_*) = \mathbf{m}$ denotes the map from \mathbf{B}^\bullet to \mathbf{B} that forgets

the marked point, then \mathbb{W}_l is absolutely continuous with respect to $\phi_*\mathbb{W}_l^\bullet$, with density function given by

$$d\mathbb{W}_l(\mathbf{m}) = \frac{K_l}{|\mathbf{V}|} d(\phi_*\mathbb{W}_l^\bullet)(\mathbf{m}), \quad (5)$$

where $|\mathbf{V}|$ should be understood as the random variable $\mathbf{m} \mapsto |\mathbf{V}(\mathbf{m})|$ giving the number of vertices of the map, and $K_l = \mathbb{W}_l^\bullet[1/|\mathbf{V}|]^{-1}$. This fact will be useful later.

Proposition 1 in [30] shows that the sequence q of non-negative weights is admissible if and only if $W^\bullet(q; \mathbf{B}_1^\bullet) < \infty$ (this is in fact the defining condition of admissibility in [30]). We see that this clearly implies that $W(q; \mathbf{B}_1) < \infty$, and even that $W(q; \mathbf{B}_l) < \infty$ for every $l \geq 1$. Moreover, in this case, the constant \mathcal{Z}_q has a nice interpretation in terms of the pointed measures. Namely, it holds that

$$\mathcal{Z}_q = 1 + W^\bullet(\mathbf{B}_1^\bullet)/2. \quad (6)$$

From now on, our attention will be exclusively focused on regular critical weight sequences. It is not obvious at this point how to interpret the definition, which will become clearer when we see how to code maps with decorated trees. However, let us explain now in which context this property typically intervenes, and refer the reader to the upcoming Subsection 1.4.3 for two applications. For instance, if one wants to study uniform random quadrangulations with a boundary and with n faces as we did in the first part of this paper, it is natural to consider the sequence $q^\circ = \delta_2 = (0, 1, 0, 0, \dots)$ and to note that $\mathbb{W}_{l,n}^\mathbf{F}(q^\circ; \cdot)$ is the uniform distribution on $\mathbf{Q}_{l,n}$. Here, note that the sequence q° is not admissible, but the probability measure $\mathbb{W}_{l,n}^\mathbf{F}(q^\circ; \cdot)$ does make sense because $0 < W(\mathbf{B}_{l,n}^\mathbf{F}) < \infty$, due to the fact that there are finitely many quadrangulations with a boundary of perimeter $2l$, and with n internal faces. Now, it can be checked that $q = 12^{-1}q^\circ$ is admissible and regular critical, and that $\mathbb{W}_{l,n}^\mathbf{F}(q; \cdot) = \mathbb{W}_{l,n}^\mathbf{F}(q^\circ; \cdot)$ is still the uniform distribution on $\mathbf{Q}_{l,n}$. This way of transforming a “naturally given” weight sequence q° into a regular weight sequence while leaving $\mathbb{W}_{l,n}^\mathbf{S}$ invariant is common and very useful.

The main result is the following. Let q be a regular critical weight sequence. Define $\rho_q = 2 + \mathcal{Z}_q^3 f_q''(\mathcal{Z}_q)$ and let $\sigma_{\mathbf{E}}, \sigma_{\mathbf{V}}, \sigma_{\mathbf{F}}$ be the non-negative numbers with squares

$$\sigma_{\mathbf{E}}^2 = \frac{\rho_q}{\mathcal{Z}_q}, \quad \sigma_{\mathbf{V}}^2 = \rho_q, \quad \sigma_{\mathbf{F}}^2 = \frac{\rho_q}{\mathcal{Z}_q - 1}. \quad (7)$$

For $L > 0$, we denote by $\mathcal{S}_L^\mathbf{S}$ the set of sequences $(l_k, n_k)_{k \geq 0} \in (\mathcal{E}^\mathbf{S})^\mathbb{N}$ such that $l_k, n_k \rightarrow \infty$ with $l_k \sim L\sigma_{\mathbf{S}}\sqrt{n_k}$ as $k \rightarrow \infty$.

Theorem 5. *Let \mathbf{S} denote one of the symbols $\mathbf{V}, \mathbf{E}, \mathbf{F}$, and $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^\mathbf{S}$ for some $L > 0$. For $k \geq 0$, denote by M_k a random map with distribution $\mathbb{W}_{l_k, n_k}^\mathbf{S}$. Then*

$$\left(\frac{4\sigma_{\mathbf{S}}^2}{9} n_k \right)^{-1/4} M_k \xrightarrow[k \rightarrow \infty]{(d)} \text{BD}_L$$

in distribution for the Gromov–Hausdorff topology.

Remark 1. The intuitive meaning for these renormalization constants is the following: in a large random map with Boltzmann distribution, it can be checked that the numbers $|\mathbf{V}|$ and $|\mathbf{F}|$ of vertices and faces are of order $|\mathbf{E}|/\mathcal{Z}_q$ and $|\mathbf{E}|(1 - 1/\mathcal{Z}_q)$ respectively, where $|\mathbf{E}|$ is the number of edges, and that conditioning on having n edges is asymptotically the same as conditioning on having (approximately) n/\mathcal{Z}_q vertices, or $n(1 - 1/\mathcal{Z}_q)$ faces.

Remark 2. In fact, the above result is also valid in the case where $L = 0$, with the interpretation that BD_0 is the Brownian map. The proof of this claim can be obtained by following ideas similar to [10, Section 6.1]. However, a full proof requires the convergence of a map with law $\mathbb{W}_{1,n}^{\mathbf{S}}$, rescaled by $(4\sigma_{\mathbf{S}}^2 n/9)^{1/4}$, to the Brownian map, and this has been explicitly done only in the case where $\mathbf{S} = \mathbf{V}$ in [26, Section 9]. In fact, building on the existing literature [30, 32], it is easy to adapt the argument to work for $\mathbf{S} = \mathbf{F}$ in the same way, while the case $\mathbf{S} = \mathbf{E}$, which is slightly different, can be tackled by the methods of [1]. Writing all the details would add a consequent number of pages to this already lengthy paper, so we will omit the proof.

1.4.3 Applications

Let us give two interesting specializations of Theorem 5. If $p \geq 2$ is an integer, a $2p$ -angulation with a boundary is a map whose internal faces all have degree $2p$. The computations of the various constants appearing in the statement of Theorem 5 have been performed in Section 1.5.1 of [30]. These show that the weight sequence

$$q = \frac{(p-1)^{p-1}}{p^p \binom{2p-1}{p}} \delta_p$$

is regular critical, that $\mathbb{W}_{l,n}^{\mathbf{F}}$ is the uniform law on the set of $2p$ -angulations with n faces and perimeter $2l$ in this case, and that the constants are

$$\mathcal{Z}_q = \frac{p}{p-1}, \quad \rho_q = p, \quad \sigma_{\mathbf{E}}^2 = p-1, \quad \sigma_{\mathbf{V}}^2 = p, \quad \sigma_{\mathbf{F}}^2 = p(p-1).$$

Therefore, in this situation, Theorem 5 for $\mathbf{S} = \mathbf{F}$ gives the following result, that clearly generalizes Theorem 1.

Corollary 6. *Let $L \in (0, \infty)$ be fixed, $(l_n, n \geq 1)$ be a sequence of integers such that $l_n \sim L\sqrt{p(p-1)n}$ as $n \rightarrow \infty$, and M_n be uniformly distributed over the set of $2p$ -angulations with n internal faces and with perimeter $2l_n$. Then the following convergence holds in distribution for the Gromov–Hausdorff topology:*

$$\left(\frac{9}{4p(p-1)n} \right)^{1/4} M_n \xrightarrow[n \rightarrow \infty]{(d)} \text{BD}_L.$$

Next, consider the case where $q_k = a^{-k}$, $k \geq 1$ for some $a > 0$. In this case, for every $\mathbf{m} \in \mathbf{B}$, a simple computation shows that

$$W(\mathbf{m}) = a^{-|\mathbf{E}(\mathbf{m})|+l}$$

so that $\mathbb{W}_{l,n}^{\mathbf{E}}$ is the uniform distribution over bipartite maps with n edges and a perimeter $2l$. It was shown in [30, Section 1.5.2] (and implicitly recovered in [1, Proposition 2]) that choosing $a = 1/8$ makes q regular critical and that, in this case,

$$\mathcal{Z}_q = \frac{3}{2}, \quad \rho_q = \frac{27}{4}, \quad \sigma_{\mathbf{E}}^2 = \frac{9}{2}.$$

Thus, one deduces the following statement, that should be compared to [1, Theorem 1].

Corollary 7. *Let M_n be a uniform random bipartite map with n edges and with perimeter $2l_n$, where $l_n \sim 3L\sqrt{n/2}$ for some $L > 0$. Then the following convergence holds in distribution for the Gromov–Hausdorff topology:*

$$(2n)^{-1/4} M_n \xrightarrow[n \rightarrow \infty]{(d)} \text{BD}_L.$$

1.5 Convergence of Boltzmann maps

The models we have presented so far consist in taking a random map with a fixed size and perimeter and letting both these quantities go to infinity in an appropriate regime. However, it is legitimate to ask about the behavior of a typical random map with law \mathbb{W}_l or \mathbb{W}_l^\bullet when $l \rightarrow \infty$, so that the perimeter is fixed and large, while the total size is left free.

For every $L \geq 0$ and $A > 0$, we define a random metric space $\text{BD}_{L,A}$, which we interpret as the Brownian disk with area A and perimeter L . For concreteness, the space $\text{BD}_{L,A}$ has same distribution as $A^{1/4} \text{BD}_{A^{-1/2}L}$. To motivate the definition, note that $\text{BD}_{L,1}$ has same distribution as BD_L and that if Q_n is a uniform random element in $\mathbf{Q}_{[L\sqrt{2n}], [An]}$, then $(8n/9)^{-1/4} Q_n$ converges in distribution for the Gromov–Hausdorff topology to $\text{BD}_{L,A}$ by virtue of Theorem 1. See also Remark 3 in Section 2.3 below.

Let \mathcal{A}^\bullet be a stable random variable with index $1/2$, with distribution given by

$$\mathbb{P}(\mathcal{A}^\bullet \in dA) = \frac{1}{\sqrt{2\pi}A^3} \exp\left(-\frac{1}{2A}\right) dA \mathbf{1}_{\{A>0\}}.$$

Note that $\mathbb{E}[1/\mathcal{A}^\bullet] = 1$, so that the formula

$$\frac{\mathbb{P}(\mathcal{A}^\bullet \in dA)}{A} = \frac{1}{\sqrt{2\pi}A^5} \exp\left(-\frac{1}{2A}\right) dA \mathbf{1}_{\{A>0\}}$$

also defines a probability distribution, and we let \mathcal{A} be a random variable with this distribution. We define the *free Brownian disk* with perimeter 1 to be a space with same law as $\text{BD}_{1,\mathcal{A}}$, where this notation means that conditionally given $\mathcal{A} = A$, it has same distribution as $\text{BD}_{1,A}$. Likewise, the *free pointed Brownian disk* with perimeter 1 has same distribution as $\text{BD}_{1,\mathcal{A}^\bullet}$.

For future reference, for $L > 0$, it is natural to define the law of the free Brownian disk (resp. free pointed Brownian disk) with perimeter L by scaling, setting it to be the law of $\sqrt{L} \text{BD}_{1,\mathcal{A}}$ or equivalently of $\text{BD}_{L,L^2\mathcal{A}}$ (resp. $\sqrt{L} \text{BD}_{1,\mathcal{A}^\bullet} \stackrel{(d)}{=} \text{BD}_{L,L^2\mathcal{A}^\bullet}$). We let FBD_L (resp. FBD_L^\bullet) stand for the free Brownian disk (resp. free pointed Brownian disk) with perimeter L .

Theorem 8. *Let q be a regular critical weight sequence. For $l \in \mathbb{N}$, let B_l (resp. B_l^\bullet) be distributed according to $\mathbb{W}_l(q; \cdot)$ (resp. $\mathbb{W}_l^\bullet(q; \cdot)$). Then*

$$\left(\frac{2l}{3}\right)^{-1/2} B_l \xrightarrow[l \rightarrow \infty]{(d)} \text{FBD}_1$$

and respectively

$$\left(\frac{2l}{3}\right)^{-1/2} B_l^\bullet \xrightarrow[l \rightarrow \infty]{(d)} \text{FBD}_1^\bullet$$

in distribution for the the Gromov–Hausdorff topology.

It is remarkable that the renormalization in this theorem does not involve q whatsoever!

1.6 Further comments and organization of the paper

The very recent preprint [35] by Miller and Sheffield aims at providing an axiomatic characterization of the Brownian map in terms of elementary properties. In this work, certain measures on random disks play a central role. We expect that these measures, denoted by $\mu_{\text{DISK}}^{k,L}$ for $k \in \{0, 1\}$ and $L > 0$, are respectively the laws of the free Brownian disk ($k = 0$) and the pointed free Brownian disk ($k = 1$) with perimeter $L > 0$. Miller and Sheffield define these measures directly in terms of the metric balls in certain versions of the Brownian map, and it is not immediate, though it is arguably very likely, that this definition matches the one given in the present paper. Establishing such a connection would be interesting from the perspective of [35] since, for example, it is not established that $\mu_{\text{DISK}}^{k,L}$ is supported on compact metric spaces, due to the possibly wild behavior of the boundary from a metric point of view. We hope to address such questions in future work.

Note also that [35] introduces another measure on metric spaces, called μ_{MET}^L , which intuitively corresponds to the law of a variant of a metric ball in the Brownian map, with a given boundary length. A description of this measure in terms of slices is given in [35], which is very much similar to the one we describe in the current work. However, there is a fundamental difference, which is that μ_{MET}^L does not satisfy the invariance under re-rooting that is essential to our study of random disks. In a few words, in a random disk with distribution μ_{MET}^L , all points of the boundary are equidistant from some special point (the center of the ball), while it is very likely that no such point exists a.s. in $\text{BD}_{L,A}$, or under the law $\mu_{\text{DISK}}^{k,L}$.

It would be natural to consider the operation that consists in gluing Brownian disks, say with same perimeter, along their boundaries, hence constructing what should intuitively be a random sphere with a self-avoiding loop. However, this operation is in general badly behaved from a metric point of view (in the sense of [17, Chapter 3] say), and it is not clear that the resulting space has the same topology as the topological gluing. The reason for this difficulty is that we require to glue along curves that are not Lipschitz, since the boundaries of the spaces BD_L have Hausdorff dimension 2 (by contrast, the

gluings considered in Section 4 of the present paper are all along geodesics.) At present, such questions remain to be investigated.

The rest of the paper is organized as follows. In Section 2, we give a self-standing definition of the limiting objects. As in many papers on random maps, we rely on bijective tools, and Section 3 introduces these tools. Section 4 is dedicated to the proof of Theorem 1. In Sections 5–7, we address the question of universality and prove Theorems 5 and 8.

Acknowledgments. This work is partly supported by the GRAAL grant ANR-14-CE25-0014. We also acknowledge partial support from the Isaac Newton Institute for Mathematical Sciences where part of this work was conducted, and where G.M. benefited from a Rothschild Visiting Professor position during January 2015.

We thank Erich Baur, Timothy Budd, Guillaume Chapuy, Nicolas Curien, Igor Kortchemski, Jean-François Le Gall, Jason Miller, Gourab Ray and Scott Sheffield, for useful remarks and conversations during the elaboration of this work.

2 Definition of Brownian disks

Recall that the Brownian map BD_0 is defined ([24], see also [31]) in terms of a certain stochastic process called the normalized Brownian snake. Likewise, the spaces BD_L , $L > 0$ of Theorem 1 are defined in terms of stochastic processes, as we now discuss.

2.1 First-passage bridges and random continuum forests

The first building blocks of the Brownian disks are first-passage bridges of Brownian motion. Informally, given A , $L > 0$, the first-passage bridge at level $-L$ and time A is a Brownian motion conditioned to first hit $-L$ at time A . To be more precise, let us introduce some notation. We let X be the canonical continuous process, and $\mathcal{G}_s = \sigma(X_u, u \leq s)$ be the associated canonical filtration. Denote by \mathbb{P} the law of standard Brownian motion, and by \mathbb{P}^A the law of standard Brownian motion killed at time $A > 0$. For $L \geq 0$, let $T_L = \inf\{s \geq 0 : X_s = -L\}$ be the first hitting time of $-L$. We denote the density function of its law by

$$j_L(A) = \frac{\mathbb{P}(T_L \in dA)}{dA} = \frac{L}{\sqrt{2\pi A^3}} \exp\left(-\frac{L^2}{2A}\right). \quad (8)$$

With this notation, the law \mathbb{F}_L^A of the first-passage bridge at level $-L$ and at time A can informally be seen as $\mathbb{P}^A(\cdot | T_L = A)$. It is best defined by an absolute continuity relation with respect to \mathbb{P}^A . Namely, for every $s \in (0, A)$ and every non-negative random variable G that is measurable with respect to \mathcal{G}_s , we let

$$\mathbb{F}_L^A(G) = \mathbb{P}^A \left[G \mathbf{1}_{\{T_L > s\}} \frac{j_{L+X_s}(A-s)}{j_L(A)} \right]. \quad (9)$$

It can be seen [18] that this definition is consistent and uniquely extends to a law \mathbb{F}_L^A on \mathcal{G}_A , supported on continuous processes, and for which $\mathbb{F}_L^A(T_L = A) = 1$.

An alternative description of first-passage bridges, which will be useful to us later, is the following.

Proposition 9. *Let $A, L > 0$. Then for every $y \in (0, L)$ and for every non-negative random variable G that is measurable with respect to \mathcal{G}_{T_y} , we have*

$$\mathbb{F}_L^A[G] = \mathbb{P}^A \left[G \mathbf{1}_{\{T_y < A\}} \frac{j_{L-y}(A - T_y)}{j_L(A)} \right]. \quad (10)$$

Moreover, this property characterizes \mathbb{F}_L^A among all measures on \mathcal{G}_A supported on continuous functions.

Proof. The definition of \mathbb{F}_L^A implies that the process $\mathcal{M} = (\mathbf{1}_{\{T_L > s\}} j_{L+X_s}(A-s)/j_L(A), 0 \leq s < A)$ is a $(\mathcal{G}_s, 0 \leq s < A)$ -martingale. Therefore, for every stopping time T such that $T < A$ a.s. under \mathbb{F}_L^A , and for every $E \in \mathcal{G}_T$, we have

$$\mathbb{F}_L^A(E) = \lim_{s \uparrow A} \mathbb{P}^A[\mathbf{1}_{E \cap \{T \leq s\}} \mathcal{M}_s] = \lim_{s \uparrow A} \mathbb{P}^A[\mathbf{1}_{E \cap \{T \leq s\}} \mathbb{P}^A[\mathcal{M}_s | \mathcal{G}_T]] = \lim_{s \uparrow A} \mathbb{P}^A[\mathbf{1}_{E \cap \{T \leq s\}} \mathcal{M}_T],$$

and this is equal to $\mathbb{P}^A[\mathbf{1}_E \mathcal{M}_T]$. The formula is obtained by applying this result to $T = T_y$, and by a standard approximation procedure of a general measurable function by weighted sums of indicator functions.

The fact that \mathbb{F}_L^A is characterized by these formulas comes from the following observation. Define $\tilde{\mathbb{F}}_L^A$ on \mathcal{G}_{T_y} as being absolutely continuous with respect to $\mathbb{P}^A|_{\mathcal{G}_{T_y}}$, with density \mathcal{M}_{T_y} . Then for every $s < A$, $\tilde{\mathbb{F}}_L^A(T_y < s) = \mathbb{P}^A[\mathbf{1}_{\{T_y < s\}} j_{L-y}(A - T_y)/j_L(A)]$, and this clearly converges to 0 as $y \uparrow L$. Therefore, T_y converges $\tilde{\mathbb{F}}_L^A$ -a.s. to A as $y \uparrow L$. Then for every $s < A$ and $E \in \mathcal{G}_s$, similar manipulations to the above ones show that

$$\tilde{\mathbb{F}}_L^A(E) = \lim_{y \uparrow L} \tilde{\mathbb{F}}_L^A(E \cap \{T_y > s\}) = \lim_{y \uparrow L} \mathbb{P}^A[\mathbf{1}_{E \cap \{T_y > s\}} \mathcal{M}_{T_y}] = \lim_{y \uparrow L} \mathbb{P}^A[\mathbf{1}_{E \cap \{T_y > s\}} \mathcal{M}_s]$$

and this is $\lim_{y \uparrow L} \mathbb{F}_L^A(E \cap \{T_y > s\}) = \mathbb{F}_L^A(E)$. \square

It is convenient to view a first-passage bridge as encoding a random continuum forest. This is a classical construction that can be summarized as follows, see for instance [36]. Here we work under \mathbb{F}_L^A . For $0 \leq s \leq s' \leq A$, define $\underline{X}_{s,s'} = \inf\{X_u : s \leq u \leq s'\}$ and let

$$d_X(s, s') = X_s + X_{s'} - 2\underline{X}_{s \wedge s', s \vee s'} \quad s, s' \in [0, A]. \quad (11)$$

The function d_X on $[0, A]^2$ is a pseudo-distance, to which one can associate a random metric space $\mathcal{F}_L^A = [0, A]/\{d_X = 0\}$, endowed with the quotient metric induced from d_X . This metric space is a.s. a compact \mathbb{R} -tree, that is, a compact geodesic metric space into which \mathbb{S}^1 cannot be embedded. It comes with a distinguished geodesic of length L , which is the image of the first hitting times $\{T_y, 0 \leq y \leq L\}$ under the canonical projection $p_X : [0, A] \rightarrow \mathcal{F}_L^A$. It is convenient to view this segment as the *floor* of a forest of \mathbb{R} -trees, these trees being exactly of the form $\mathcal{T}_y = p_X((T_{y-}, T_y))$, corresponding to the excursions of X above its past infimum. One should imagine that the \mathbb{R} -tree \mathcal{T}_y is grafted at the point $p_X(T_y)$ of the floor lying at distance y from $p_X(0)$.

2.2 Snakes

We now enrich the random “real forest” described above by assigning labels to it. Informally speaking, the trees of the forest are labeled by independent Brownian snakes [23, 21], while the floor of the forest is labeled by a Brownian bridge with variance factor 3.

More precisely, let X be a first-passage bridge with law \mathbb{F}_L^A . Conditionally given X , we let $(Z_s^0, 0 \leq s \leq A)$ be a centered Gaussian process with covariance function

$$\text{Cov}(Z_s^0, Z_{s'}^0) = \inf_{u \in [s \wedge s', s \vee s']} (X_u - \underline{X}_u) \quad s, s' \in [0, A],$$

where $\underline{X}_u = \inf_{0 \leq v \leq u} X_v$ is the past infimum of X . Note in particular that Z_s^0 and $Z_{s'}^0$ are independent if s, s' belong to two different excursion intervals of X above \underline{X} . It is classical [23] that Z^0 admits a continuous modification, see also [7] for a discussion in the current context. For this modification, we a.s. have $Z_{T_y}^0 = 0$ for every $y \in [0, L]$ (for a given y , this comes directly from the variance formula). The process Z^0 is sometimes called the *head of the Brownian snake* driven by the process $X - \underline{X}$, the reason being that it can be obtained as a specialization of a path-valued Markov process called the Brownian snake [23] driven by $X - \underline{X}$. The process Z^0 itself is not Markov.

Let also b be a standard Brownian bridge of duration L , so that

$$\text{Cov}(b_y, b_{y'}) = \frac{y(L - y')}{L}, \quad 0 \leq y \leq y' \leq L.$$

We define the process Z to be

$$Z_s = Z_s^0 + \sqrt{3} b_{T^{-1}(s)}, \quad 0 \leq s \leq A, \quad (12)$$

where $T^{-1}(s) = \sup\{y \geq 0 : T_y \leq s\}$. We abuse notation and still denote by \mathbb{F}_L^A the law of the pair (X, Z) thus defined, so that \mathbb{F}_L^A is seen as a probability distribution on the space $\mathcal{C}([0, A], \mathbb{R})^2$. In the same spirit, we will still denote by \mathcal{G}_t the natural filtration $\sigma(\{(X_s, Z_s), 0 \leq s \leq t\})$. Note that the absolute continuity relations (9) and (10) are still valid *verbatim* with these extended notation and, in particular, the density function involves only X and not Z .

It is classical that a.s. under \mathbb{F}_L^A , Z is a class function on $[0, A]$ for the equivalence relation $\{d_X = 0\}$, so that Z can also be seen as a function on the forest \mathcal{F}_L^A . Note that $T^{-1}(T_y) = y$ for every $0 \leq y \leq L$, which corresponds to the fact that, in the above depiction of the random forest, the point $p_X(T_y)$ receives label $\sqrt{3} b_y$.

It is a simple exercise to check that the above definition of Z is equivalent to the following quicker (but more obscure) one. Conditionally given X , we have that Z is Gaussian, centered, with covariance function

$$\text{Cov}(Z_s, Z_{s'}) = \underline{X}_{s, s'} - \underline{X}_{s'} - 3\underline{X}_s(L + \underline{X}_{s'})/L \quad s, s' \in [0, A].$$

Similarly as (11), we define a pseudo-distance function using the process Z instead of X , but with an extra twist. As above, let $\underline{Z}_{s, s'} = \inf\{Z_u : u \in [s, s']\}$ for $0 \leq s \leq s' \leq A$,

and this time we extend the definition to $0 \leq s' < s \leq A$ by setting

$$\underline{Z}_{s,s'} = \inf\{Z_u : u \in [s, A] \cup [0, s']\} = \underline{Z}_{s,A} \wedge \underline{Z}_{0,s'},$$

so if we see $[0, A]$ as a circle by identifying 0 with A , $\underline{Z}_{s,s'}$ is the minimum of Z on the directed arc from s to s' . We let

$$d_Z(s, s') = Z_s + Z_{s'} - 2 \max(\underline{Z}_{s,s'}, \underline{Z}_{s',s}) \quad s, s' \in [0, A]. \quad (13)$$

2.3 Brownian disks

We are now ready to give the definition of Brownian disks. Consider the set \mathcal{D} of all pseudo-distances d on $[0, A]$ satisfying the two properties

$$\left\{ \begin{array}{l} \{d_X = 0\} \subseteq \{d = 0\} \\ d \leq d_Z. \end{array} \right.$$

The set \mathcal{D} is nonempty (it contains the zero pseudo-distance) and contains a maximal element D^* defined by

$$D^*(s, s') = \inf \left\{ \sum_{i=1}^k d_Z(s_i, t_i) : \begin{array}{l} k \geq 1, \quad t_1, s_2, \dots, s_k \in [0, A], \quad s_1 = s, \quad t_k = s', \\ d_X(t_i, s_{i+1}) = 0 \text{ for every } i \in \{1, \dots, k-1\} \end{array} \right\}, \quad (14)$$

see [17, Chapter 3]. The Brownian disk $\text{BD}_{L,A}$ with area A and perimeter L is the quotient set $[0, A]/\{D^* = 0\}$, endowed with the quotient metric induced from D^* (which we still denote by D^* for simplicity), and considered under the law \mathbb{F}_L^A . In the case $A = 1$, we drop the second subscript and write $\text{BD}_L = \text{BD}_{L,1}$.

Remark 3. Observe that, by usual scaling properties of Gaussian random variables, under the law \mathbb{F}_L^A , the scaled pair $((\lambda^{1/2} X_{s/\lambda}, 0 \leq s \leq \lambda A), (\lambda^{1/4} Z_{s/\lambda}, 0 \leq s \leq \lambda A))$ has law $\mathbb{F}_{\lambda^{1/2}L}^{\lambda A}$, from which we deduce that the random metric space $\lambda^{-1/4} \text{BD}_{\lambda^{1/2}L, \lambda A}$ has the same distribution as $\text{BD}_{L,A}$.

The reason why we say that $\text{BD}_{L,A}$ has “area” A is that it naturally comes with a non-negative measure of total mass A , which is the image of the Lebesgue measure on $[0, A]$ by the canonical projection $\mathbf{p} : [0, A] \rightarrow \text{BD}_{L,A}$. It will be justified later that $\text{BD}_{L,A}$ is a.s. homeomorphic to the closed unit disk, so that the term *area* makes more sense in this context. Furthermore, the boundary $\partial \text{BD}_{L,A}$ will be shown to be equal to $\mathbf{p}(\{T_y : 0 \leq y \leq L\})$, so that it can be endowed with a natural non-negative measure with total mass L , which is the image of the Lebesgue measure on $[0, L]$ by $y \mapsto \mathbf{p}(T_y)$. This justifies the term “perimeter”.

3 The Schaeffer bijection and two variants

This work strongly relies on powerful encodings of discrete maps by trees and related objects. In this section we present the encodings we will need: the original Cori–Vauquelin–Schaeffer bijection [19, 37], a variant for so-called *slices* [26] and a variant for plane quadrangulations with a boundary (particular case of [14]). We only give the constructions from the encoding objects to the considered maps and refer the reader to the aforementioned works for converse constructions and proofs.

3.1 The original Cori–Vauquelin–Schaeffer bijection

Let (\mathbf{t}, ℓ) be a *well-labeled tree* with n edges. Recall that this means that \mathbf{t} is a rooted plane tree with n edges, and $\ell : \mathbf{V}(\mathbf{t}) \rightarrow \mathbb{Z}$ is a labeling function such that $\ell(u) - \ell(v) \in \{-1, 0, 1\}$ whenever u and v are neighboring vertices in \mathbf{t} . It is usual to “normalize” ℓ in such a way that the root vertex of \mathbf{t} gets label 0, but we will also consider different conventions: in fact, all our discussion really deals with the function ℓ up to addition of a constant. For simplicity, in the following, we let $\ell_* = \min\{\ell(v) : v \in \mathbf{V}(\mathbf{t})\} - 1$.

Note. Throughout this paper, whenever a function f is defined at a vertex v , we extend its definition to any corner c incident to v by setting $f(c) = f(v)$. In particular, the label $\ell(c)$ of a corner is understood as the label of the incident vertex.

Let $c_0, c_1, \dots, c_{2n-1}$ be the sequence of corners of \mathbf{t} in contour order, starting from the root corner. We extend the list of corners by periodicity, setting $c_{2n+i} = c_i$ for every $i \geq 0$, and adding one corner c_∞ incident to a vertex v_* not belonging to \mathbf{t} , with label $\ell(c_\infty) = \ell(v_*) = \ell_*$. Once this is done, we define the *successor* functions by setting

$$s(i) = \inf\{j > i : \ell(c_j) = \ell(c_i) - 1\} \in \mathbb{Z}_+ \cup \{\infty\}, \quad i \in \{0, 1, \dots, 2n-1\},$$

and $s(c_i) = c_{s(i)}$. The Cori–Vauquelin–Schaeffer construction consists in linking c_i with $s(c_i)$ by an arc, in a non-crossing fashion, for every $i \in \{0, 1, \dots, 2n-1\}$. The embedded graph \mathbf{q} with vertex set $\mathbf{V}(\mathbf{t}) \cup \{v_*\}$ and edge set the set of arcs (excluding the edges of \mathbf{t}) is then a quadrangulation, which is rooted according to some convention (we omit details here as this point is not important for our purposes), and is naturally pointed at v_* . Moreover, the labels on $\mathbf{V}(\mathbf{q})$ inherited from those on \mathbf{t} (and still denoted by ℓ) are exactly the relative distances to v_* in \mathbf{q} :

$$d_{\mathbf{q}}(v, v_*) = \ell(v) - \ell_*, \quad v \in \mathbf{V}(\mathbf{q}).$$

(This entirely determines ℓ as soon as the value $\ell(v_0)$ is known for some specific v_0 , but recall that in general we do not want to fix the normalization of ℓ .) See Figure 2 for an example of the construction.




Figure 2: The Cori-Vauquelin-Schaeffer bijection. There are two possible rootings of q : they are indicated with question marks.




Figure 3: A map with geodesic boundary is associated with a well-labeled tree via the modified Schaeffer bijection. The maximal geodesic is represented in red on the left and the shuttle is the green chain on the right.

3.2 Slices

We now follow [26] and describe a modification of the previous construction that goes as follows. See Figure 3 for an example, and compare with Figure 2.

Rather than appending to \mathbf{t} a single corner c_∞ incident to a vertex v_* , we add a sequence of corners $c'_1, c'_2, \dots, c'_{\ell(c_0)-\ell_*-1}, c'_{\ell(c_0)-\ell_*} = c_\infty$, and set labels $\ell(c'_i) = \ell(c_0) - i$, so in particular this is consistent with the label we already set for c_∞ . Also, instead of extending the sequence $c_0, c_1, \dots, c_{2n-1}$ by periodicity, we add an extra corner c_{2n} to the right of c_0 and we let $c_{2n+i} = c'_i$ for $i \in \{1, 2, \dots, \ell(c_0) - \ell_*\}$. The definition of the successor

$$s(i) = \inf\{j > i : \ell(c_j) = \ell(c_i) - 1\}, \quad s(c_i) = c_{s(i)} \quad (15)$$

then makes sense for $i \in \{0, 1, \dots, 2n + \ell(c_0) - \ell_* - 1\}$, and we can draw the arcs from c_i to $s(c_i)$ for every $i \in \{0, 1, \dots, 2n + \ell(c_0) - \ell_* - 1\}$. In particular, note that the arcs link c_{2n} with $c'_1, c'_2, \dots, c'_{\ell(c_0)-\ell_*-1}, c_\infty$ into a chain, which we call *shuttle*, and to which are connected the arcs $c_i \rightarrow s(c_i)$ with $i \leq 2n - 1$ and $s(i) > 2n - 1$. Let \mathbf{q}' be the map obtained by this construction. This map contains two distinguished geodesic chains, which are, on the one hand, the so-called *maximal geodesic* made of arcs between consecutive successors $c_0, s(c_0), s(s(c_0)), \dots, c_\infty$ and, on the other hand, the shuttle linking $c_{2n}, c'_1, c'_2, \dots, c'_{\ell(c_0)-\ell_*-1}, c_\infty$. Note that both chains indeed have the same length (number of edges), equal to $\ell(c_0) - \ell_*$. In particular, we have $d_{\mathbf{q}'}(c_0, c_\infty) = d_{\mathbf{q}}(c_0, c_\infty) = \ell(c_0) - \ell_*$, where \mathbf{q} is the quadrangulation from the previous section, constructed from the same well-labeled tree (\mathbf{t}, ℓ) . These two chains are incident to a face of \mathbf{q}' of degree $2d_{\mathbf{q}}(c_0, c_\infty)$, and all other faces have degree 4. Observe that the maximal geodesic and the shuttle only intersect at the root vertex of the tree and v_* ; as a result, the boundary of the degree $2d_{\mathbf{q}}(c_0, c_\infty)$ -face is a simple curve. Finally, the quadrangulation \mathbf{q} can then be obtained from \mathbf{q}' by identifying one by one the edges of the maximal geodesic with the edges of the shuttle, in the same order.

3.3 Plane quadrangulations with a boundary

We now present the variant for plane quadrangulation with a boundary, which is a particular case of the Bouttier–Di Francesco–Guitter bijection [14]. We rather use the presentation of [9], better fitted to our situation.

The encoding object of a plane quadrangulation with a boundary having n internal faces and perimeter $2l$ is a forest $\mathbf{f} = (\mathbf{t}_1, \dots, \mathbf{t}_l)$ of l trees with n edges in total, together with a labeling function $\ell : \mathbf{V}(\mathbf{f}) = \bigsqcup_i \mathbf{V}(\mathbf{t}_i) \rightarrow \mathbb{Z}$ satisfying the following:

- for $1 \leq i \leq l$, the tree \mathbf{t}_i equipped with the restriction of ℓ to $V(\mathbf{t}_i)$ is a well-labeled tree;
- for $1 \leq i \leq l$, we have $\ell(\rho_{i+1}) \geq \ell(\rho_i) - 1$, where ρ_i denotes the root vertex of \mathbf{t}_i and setting $\ell(\rho_{l+1}) = \ell(\rho_1)$ by convention.

Note that the condition on the labels of the root vertices is different from the condition on the labels of neighboring vertices of a given tree. The reader familiar with the Bouttier–Di Francesco–Guitter bijection may recognize the label condition for faces of even degree more than 4. We will come back to this during Section 5.

Here and later, it will be convenient to normalize ℓ by asking that $\ell(\rho_1) = 0$. As before, we define $\ell_* = \min\{\ell(v) : v \in \mathbf{V}(\mathbf{f})\} - 1$. We identify \mathbf{f} with the map obtained by adding l edges linking the roots $\rho_1, \rho_2, \dots, \rho_l$ of the successive trees in a cycle. This map has two faces, one of degree $2n + l$ (the bounded one on Figure 4) and one of degree l (the unbounded one on Figure 4). We then follow a procedure similar to that of Section 3.1. We let $c_0, c_1, \dots, c_{2n+l-1}$ be the sequence of corners of the face of degree $2n + l$ in contour order, starting from the root corner of \mathbf{t}_1 . We extend this list by periodicity and add one corner c_∞ incident to a vertex v_* lying inside the face of degree $2n + l$, with label $\ell(c_\infty) = \ell(v_*) = \ell_*$. We define the successor functions by (15) and draw an arc from c_i to $s(c_i)$ for every $i \in \{0, 1, \dots, 2n + l - 1\}$, in such a way that this arc does not cross the edges of \mathbf{f} , or other arcs.

The embedded graph \mathbf{q}'' with vertex set $\mathbf{V}(\mathbf{f}) \cup \{v_*\}$ and edge set given by the added arcs is a plane quadrangulation with a boundary, whose external face is the degree- $2l$ face corresponding to the face of degree l . It is rooted at the corner of the unbounded face that is incident to the root vertex of \mathbf{t}_1 , and it is naturally pointed at v_* . See Figure 4.




Figure 4: The bijection for a plane quadrangulation with a boundary. The l edges we added to \mathbf{f} are represented by dotted lines and the root edge of \mathbf{q}'' is represented in red. Note that we used for ℓ the normalization given by $\ell(\rho_1) = 0$.

The above mapping is a bijection between previously described labeled forests and the set of pointed plane quadrangulations (\mathbf{q}, v_*) with a boundary having n internal faces and

perimeter $2l$ that further satisfy the property that $d_{\mathbf{q}}(e_*^+, v_*) = d_{\mathbf{q}}(e_*^-, v_*) + 1$, where e_* denotes the root edge of \mathbf{q} , that is, the oriented edge incident to the root face that directly precedes the root corner in the contour order (see Figure 4). In words, the pointed quadrangulations that are in the image of the above mapping are those whose root edge *points away from* the distinguished vertex v_* .

The requirement that the root edge is directed away from the distinguished vertex is not a serious issue, as we can dispose of this constraint simply by re-rooting along the boundary:

Lemma 10. *Let (Q, v_*) be uniformly distributed in the set $\mathbf{Q}_{l,n}^{\bullet,+}$ of rooted and pointed quadrangulations such that $Q \in \mathbf{Q}_{l,n}$ and such that the root edge e_* points away from v_* . Let c' be a uniformly chosen random corner incident to the root face of Q , and let Q' be the map Q re-rooted at c' . Then Q' is a uniform random element of $\mathbf{Q}_{l,n}$.*

Proof. The probability that Q' is a given rooted map \mathbf{q}' is equal

$$\mathbb{P}(Q' = \mathbf{q}') = \frac{1}{2l} \sum_{v \in \mathbf{V}(\mathbf{q}')} \sum_c^+ \mathbb{P}((Q, v_*) = (\mathbf{q}, v)) = \frac{1}{2l} \sum_{v \in \mathbf{V}(\mathbf{q}')} \sum_c^+ \frac{1}{|\mathbf{Q}_{l,n}^{\bullet,+}|},$$

where the factor $1/2l$ is the probability that c' is chosen to be the root corner of \mathbf{q}' , the symbol \sum_c^+ stands for the sum over all corners incident to the root face of \mathbf{q}' that point away from v , and \mathbf{q} is the map \mathbf{q}' re-rooted at the corner c .

Now fix the vertex $v \in \mathbf{V}(\mathbf{q}')$. Due to the bipartite nature of \mathbf{q}' , among the $2l$ oriented edges incident to the root face, l are pointing away from v , and l are pointing toward v . Indeed, let \tilde{c}_0 be the root corner of Q , and $\tilde{c}_1, \tilde{c}_2, \dots, \tilde{c}_{2l-1}, \tilde{c}_{2l} = \tilde{c}_0$ be the corners incident to the root face in cyclic order. The sequence $(d_{\mathbf{q}'}(\tilde{c}_i, v_*), 0 \leq i \leq 2l)$ takes integer values, varies by ± 1 at every step as \mathbf{q}' is bipartite, and takes the same value at times 0 and $2l$. This means that l of its increments are equal to $+1$ and l are equal to -1 , respectively corresponding to edges that point away from v and toward v .

Therefore, the sum \sum_c^+ contains exactly l elements. Noting that every map in $\mathbf{Q}_{l,n}$ has $n + l + 1$ vertices, by the Euler characteristic formula, this gives

$$\mathbb{P}(Q' = \mathbf{q}') = \frac{n + l + 1}{2|\mathbf{Q}_{l,n}^{\bullet,+}|},$$

which depends only on n, l and not on the particular choice of \mathbf{q}' . □

4 Proof of Theorem 1

4.1 Subsequential convergence

Recall the construction of Section 3.3 and consider an encoding labeled forest (\mathbf{f}, ℓ) . We will further encode it by a pair of real-valued functions. Before we proceed, it will be convenient to add an extra vertex-tree ρ_{l+1} with label $\ell(\rho_{l+1}) = \ell(\rho_1)$ to the forest. This

extra vertex does not really play a part but its introduction will make the presentation simpler. We also add l edges between ρ_i and ρ_{i+1} , for $1 \leq i \leq l$. See Figure 5.

We let $c_0, c_1, \dots, c_{2n+l-1}$ be as in the previous section and we add to this list the corner c_{2n+l} incident to the extra vertex-tree ρ_{l+1} . We define the contour and label processes on $[0, 2n + l]$ by

$$C(j) = d_{\mathbf{f}}(c_j, \rho_{l+1}) - l \quad \text{and} \quad \ell(j) = \ell(c_j), \quad 0 \leq j \leq 2n + l$$

and by linear interpolation between integer values.




Figure 5: The contour and label processes associated with the labeled forest of Figure 4. The extra vertex-tree ρ_7 and the edges linking the roots are represented with a dashed line. Note that the normalization we chose for the labels is equivalent to imposing $\ell(0) = 0$.

Let us fix $L \in (0, \infty)$ and a sequence $(l_n, n \geq 1)$ such that $l_n \sim L\sqrt{2n}$ as $n \rightarrow \infty$. We let (F_n, ℓ_n) be uniformly distributed over the set of labeled forests of l_n trees with n edges in total, and let (Q_n, v_*) be the random pointed quadrangulation associated with F_n via the bijection of Section 3.3. Note that up to re-rooting Q_n at a uniform corner incident to the root face, we may assume that Q_n is uniform in $\mathbf{Q}_{l_n, n}$ by Lemma 10.

We let C_n, ℓ_n be the associated contour and label processes, and we define their renormalized versions

$$C_{(n)}(s) = \frac{C_n((2n + l_n)s)}{\sqrt{2n}}, \quad \ell_{(n)}(s) = \left(\frac{9}{8n}\right)^{1/4} \ell_n((2n + l_n)s), \quad 0 \leq s \leq 1.$$

We let $D_n(i, j)$ be the distance in Q_n between the vertices incident to the i -th and j -th corner of F_n , for $i, j \in \{0, 1, \dots, 2n + l_n\}$. We extend D_n to a continuous function on $[0, 2n + l_n]^2$ by “bilinear interpolation,” writing $\{s\} = s - \lfloor s \rfloor$ for the fractional part of s , and then setting

$$D_n(s, t) = (1 - \{s\})(1 - \{t\})D_n(\lfloor s \rfloor, \lfloor t \rfloor) + \{s\}(1 - \{t\})D_n(\lfloor s \rfloor + 1, \lfloor t \rfloor) + (1 - \{s\})\{t\}D_n(\lfloor s \rfloor, \lfloor t \rfloor + 1) + \{s\}\{t\}D_n(\lfloor s \rfloor + 1, \lfloor t \rfloor + 1). \quad (16)$$

We finally define its renormalized version

$$D_{(n)}(s, t) = \left(\frac{9}{8n}\right)^{1/4} D_n((2n + l_n)s, (2n + l_n)t), \quad 0 \leq s, t \leq 1. \quad (17)$$

It is shown in [10] that, from every increasing family of positive numbers, one can extract a further subsequence along which

$$(C_{(n)}, \ell_{(n)}, D_{(n)}) \xrightarrow[n \rightarrow \infty]{(d)} (X, Z, D)$$

in distribution in the space $\mathcal{C}([0, 1]) \times \mathcal{C}([0, 1]) \times \mathcal{C}([0, 1]^2)$. (At this moment, the need of extracting a subsequence is caused by the last coordinate $D_{(n)}$ and the convergence without extraction holds if one drops this coordinate.) Here, D is a random pseudo-distance on $[0, 1]$ and (X, Z) has the distribution³ \mathbb{F}_L^1 defined in Section 2. This means that X is a first-passage bridge, attaining level $-L$ for the first time at time 1, and Z is the associated snake process.

Moreover, the pointed random metric space $(\mathbf{V}(Q_n), (9/8n)^{1/4}d_{Q_n}, v_*)$ converges in distribution, still along the same subsequence, to the random metric space $([0, 1]/\{D = 0\}, D, x_*)$, in the sense of the pointed Gromov–Hausdorff topology. Here, we let $x_* = \mathbf{p}(s_*)$, where $\mathbf{p} : [0, 1] \rightarrow [0, 1]/\{D = 0\}$ is the canonical projection, and s_* is the (a.s. unique [10, Lemma 11]) point in $[0, 1]$ at which Z reaches its global minimum.

Proposition 11 ([10]). *Almost surely, the space $\mathbf{D} = [0, 1]/\{D = 0\}$ is a topological disk whose boundary $\partial\mathbf{D}$ satisfies*

$$\mathbf{p}^{-1}(\partial\mathbf{D}) = \{s \in [0, 1] : X_s = \underline{X}_s\}. \quad (18)$$

Almost surely, the Hausdorff dimension of \mathbf{D} is 4, and that of $\partial\mathbf{D}$ is 2.

Recall that v_* is a uniform random vertex in Q_n , conditionally given the latter. From this observation, we obtain an invariance under re-rooting property of (\mathbf{D}, D, x_*) , along the same lines as [25].

Lemma 12. *Let U be a uniform random variable in $[0, 1]$, independent of (X, Z, D) . Then the two pointed spaces $(\mathbf{D}, D, \mathbf{p}(U))$ and (\mathbf{D}, D, x_*) have the same distribution.*

The following lemma is an easy consequence of the study of geodesics done in [9].

Lemma 13. *Almost surely, for every $x \in \mathbf{D} \setminus \partial\mathbf{D}$, there exists a geodesic from x to x_* that does not intersect $\partial\mathbf{D}$. Moreover, this is the only geodesic from x to x_* for μ -almost every $x \in \mathbf{D}$, where $\mu = \mathbf{p}_*(\text{Leb}_{[0,1]})$.*

Proof. For $s \in [0, 1]$, we define the path $\Phi_s : [0, D(s_*, s)] \rightarrow \mathbf{D}$ by

$$\Phi_s(w) = \mathbf{p} \left(\sup \{r : \underline{Z}_{r,s} = Z_{s_*} + w\} \right), \quad 0 \leq w \leq D(s_*, s) = Z_s - Z_{s_*}.$$

It is shown in [9, Proposition 23] that the path Φ_s is a geodesic from x_* to $\mathbf{p}(s)$ in \mathbf{D} and that a.s. all the geodesics from x_* are of this form. We call *increase point* of a function a point t such that the function is greater than its value at t on a small interval

³This is of course an abuse of notation since (X, Z) previously denoted the canonical process, however we did not want to introduce a further specific notation at this point.

of the form $[t - \varepsilon, t]$ or $[t, t + \varepsilon]$ for some $\varepsilon > 0$. Clearly, for $0 \leq w < D(s_*, s)$, the point $\sup \{r : \underline{Z}_{r,s} = Z_{s_*} + w\}$ is an increase point of the process Z , which is furthermore different from 0. On the other hand, the expression (18) shows that $\mathbf{p}^{-1}(\partial\mathbf{D})$ is made only of increase points of X , together with the point 0. Moreover, [9, Lemma 18] states that, a.s., the processes X and Z do not share any increase points. As a consequence, Φ_s may only intersect $\partial\mathbf{D}$ at its endpoint $\mathbf{p}(s)$ and the first statement follows.

In addition, [9, Proposition 17] entails that, for $0 \leq s \leq t \leq 1$, $D(s, t) = 0$ if and only if one of the following occurs:

$$(19a) \quad X_s = X_t = \underline{X}_{s,t};$$

$$(19b) \quad Z_s = Z_t = \underline{Z}_{s,t} \text{ or } Z_s = Z_t = \underline{Z}_{t,s}.$$

Moreover, for $s \neq t$, only one of the previous situations can happen. In some sense, this can be thought of as a continuous version of the bijection from Section 3.3: point (19a) constructs the continuous random forest and drawing an arc between a corner and its successor becomes, in the limit, identifying points with the same label and such that the labels visited in between in the contour order are all larger (point (19b)). Standard properties of the process Z then allow us to conclude that $\text{Leb}_{[0,1]}(\{s : \exists t \neq s : D(s, t) = 0\}) = 0$, so that, for μ -almost every $x \in \mathbf{D}$, the set $\mathbf{p}^{-1}(x)$ is a singleton and the only geodesic from x_* to x is thus $\Phi_{\mathbf{p}^{-1}(x)}$. \square

Combining Lemmas 12 and 13, we see that the conclusion of the latter is still valid if x_* is replaced by a uniformly chosen point in \mathbf{D} , that is, a random point of the form $\mathbf{p}(U)$ as in the first lemma. Finally, we will use the following result.

Lemma 14 ([10]). *The following properties hold almost surely.*

- $D \in \mathcal{D}$.
- $D(s, s_*) = Z_s - Z_{s_*}$ for every $s \in [0, 1]$.

4.2 Identification of the limit

Recall the notation D^* from Section 2.3. In this section, we show the following result.

Theorem 15. *Almost surely, it holds that $D = D^*$.*

Theorem 1 is an immediate consequence of this. Indeed, since D^* is a measurable function of (X, Z) , this shows that D^* is the only possible subsequential limit of $D_{(n)}$. This, combined with the tightness of the sequence $(D_{(n)}, n \geq 1)$ that we alluded to above, implies that $D_{(n)}$ converges in distribution to D^* .

In turn, this convergence implies that of $(9/8n)^{1/4}Q_n$ to $\text{BD}_L = (\mathbf{D}, D^*)$ in the Gromov–Hausdorff sense and even that of the pointed space $((9/8n)^{1/4}Q_n, v_*)$ to (BD_L, x_*) , where we recall that $x_* = \mathbf{p}(s_*)$. Let us recall how to prove this fact. First, one can assume

that the convergence of $(C_{(n)}, \ell_{(n)}, D_{(n)})$ to (X, Z, D^*) is almost-sure, by using Skorokhod's representation theorem. Then we define a correspondence \mathcal{R}_n between Q_n and BD_L by

$$\mathcal{R}_n = \{(v_{\lfloor (2n+l_n)s \rfloor}, \mathbf{p}(s)) : s \in [0, 1]\} \cup \{(v_*, \mathbf{p}(s_*))\},$$

where v_i is the vertex of Q_n incident to the i -th corner c_i . It is elementary to see from the uniform convergence of $D_{(n)}$ to D^* that the distortion of \mathcal{R}_n with respect to the metrics $(9/8n)^{1/4}d_{Q_n}$ and D^* converges to 0 as $n \rightarrow \infty$.

Recall that $[a, b]$ is an excursion interval of X above \underline{X} if $a < b$ and $X_a = X_b = \underline{X}_b$. Let us arrange the excursion intervals of X above \underline{X} as $[a_i, b_i]$, $i \geq 1$ in decreasing order of length. For a given i , the excursion interval $[a_i, b_i]$ encodes a Brownian map with a geodesic boundary in the sense of [26]. Namely, let $d_Z^i(s, t) = Z_s + Z_t - 2Z_{s \wedge t, s \vee t}$ for $s, t \in [a_i, b_i]$, and let \tilde{D}^i be the maximal pseudo-distance on $[a_i, b_i]$ that is less than d_Z^i and satisfies $d_X(s, t) = 0 \implies \tilde{D}^i(s, t) = 0$. More explicitly,

$$\tilde{D}^i(s, t) = \inf \left\{ \sum_{j=1}^k d_Z^i(s_j, t_j) : \begin{array}{l} k \geq 1, \quad s_1, t_1, s_2, t_2, \dots, s_k, t_k \in [a_i, b_i], \quad s_1 = s, \quad t_k = t, \\ d_X(t_j, s_{j+1}) = 0 \text{ for every } j \in \{1, \dots, k-1\} \end{array} \right\}.$$

The next key lemma states that the distance D can be identified as a metric gluing of the maps with geodesic boundaries $([a_i, b_i], \tilde{D}^i)/\{\tilde{D}^i = 0\}$, $i \geq 1$. This guides the intuition of its proof, which will partly consist in going back to the discrete slices that compose the quadrangulations with boundaries we took the limit of.

Lemma 16. *Let \tilde{D} be the pseudo-distance on $\bigcup_{i \geq 1} [a_i, b_i]$ defined by $\tilde{D}(s, t) = \tilde{D}^i(s, t)$ if $s, t \in [a_i, b_i]$ for some $i \geq 1$, and $\tilde{D}(s, t) = \infty$ otherwise. Then for almost every $(s, t) \in [0, 1]^2$ with respect to the Lebesgue measure, it holds that*

$$D(s, t) = \inf \left\{ \sum_{j=1}^k \tilde{D}(s_j, t_j) : \begin{array}{l} k \geq 1, \quad s_1, t_1, s_2, t_2, \dots, s_k, t_k \in [0, 1], \quad s_1 = s, \quad t_k = t, \\ d_Z(t_j, s_{j+1}) = 0 \text{ for every } j \in \{1, \dots, k-1\} \end{array} \right\}.$$

Moreover, the above infimum is attained.

Proof. Clearly, $D(s, t) \leq d_Z(s, t) \leq d_Z^i(s, t)$ whenever $s, t \in [a_i, b_i]$, so that $D(s, t) \leq \tilde{D}(s, t)$ for $s, t \in [0, 1]$ and, as a consequence, the left-hand side is smaller than the right-hand side. We then only need to prove the converse inequality.

Let us first define the discrete analogs to the functions \tilde{D}^i . We consider the i -th largest tree \mathbf{t} of F_n and we suppose that it is visited between times a_i^n and b_i^n in the contour order of F_n . For $j, k \in \{a_i^n, \dots, b_i^n\}$, we let $\tilde{D}_n(j, k)$ be the distance in the slice corresponding to \mathbf{t} between the vertices v_j and v_k incident to the j -th and k -th corner of F_n . In other words, $\tilde{D}_n(j, k)$ is the length of a shortest path linking v_j to v_k and that do not "traverse" the images in Q_n of the maximal geodesic and shuttle of the aforementioned slice. We then extend \tilde{D}_n to a continuous function on $[a_i^n, b_i^n]^2$ by bilinear interpolation, and define its renormalized version $\tilde{D}_{(n)}^i$ on a subsquare of $[0, 1]^2$ by the analog of (17). We define $\tilde{D}_{(n)}^i$ arbitrarily for $i > l_n$.

As an easy consequence of [26, Proposition 9.2], we obtain that

$$\left(C_{(n)}, \ell_{(n)}, (\tilde{D}_{(n)}^i)_{i \geq 1} \right) \xrightarrow[n \rightarrow \infty]{} \left(X, Z, (\tilde{D}^i)_{i \geq 1} \right) \quad (20)$$

in distribution in the space $\mathcal{C}([0, 1]) \times \mathcal{C}([0, 1]) \times \mathcal{C}([0, 1]^2)^N$. Applying Skorokhod's representation theorem, we furthermore assume from now on that the previous convergence holds a.s.

It suffices to prove the claimed formula where s, t are replaced by two independent uniform random variables U, V , independent of (X, Z, D) . Let $\gamma : [0, D(U, V)] \rightarrow \mathbf{D}$ be the geodesic in (\mathbf{D}, D) from $\mathbf{p}(U)$ to $\mathbf{p}(V)$, which by Lemmas 12 and 13 is unique and does not intersect $\partial \mathbf{D}$, a.s. Let also $\text{Im}(\gamma) = \gamma([0, D(U, V)])$ be the image of γ and define

$$I(U, V) = \{i \geq 1 : \mathbf{p}^{-1}(\text{Im}(\gamma)) \cap [a_i, b_i] \neq \emptyset\}.$$

Claim 1. *The set $I(U, V)$ is finite almost surely.*

Proof. Let us argue by contradiction, assuming that $I(U, V)$ is infinite with positive probability. Then it holds that, still with positive probability, there is an increasing integer sequence $(i_n)_{n \geq 1}$ and a sequence $(r_n)_{n \geq 1}$ with values in $[0, D(U, V)]$ such that $\gamma(r_n) \in \mathbf{p}([a_{i_n}, b_{i_n}])$. Then, up to extraction, the sequence (r_n) converges to some limit r , and if $s_n \in [a_{i_n}, b_{i_n}]$ is a choice of a given element in $\mathbf{p}^{-1}(\gamma(r_n))$, then, again up to possibly further extraction, (s_n) converges to a limit s with $\mathbf{p}(s) = \gamma(r)$. By construction, s is not in $\bigcup_{i \geq 1} (a_i, b_i)$, since the intervals in this union are pairwise disjoint. This implies that $X_s = \underline{X}_s$, meaning that $\gamma(r) = \mathbf{p}(s) \in \partial \mathbf{D}$, which is the contradiction we were looking for. \diamond

Let γ_l^i, γ_r^i be the left and right “geodesic boundaries” of the space $([a_i, b_i]/d_Z^i, d_Z^i)$, defined by

$$\gamma_l^i(t) = \mathbf{p}(\inf\{s \in [a_i, b_i] : Z_s = Z_{a_i} - t\}), \quad \gamma_r^i(t) = \mathbf{p}(\sup\{s \in [a_i, b_i] : Z_s = Z_{a_i} - t\}),$$

where t ranges over $[0, Z_{a_i} - \underline{Z}_{a_i, b_i}]$ (recall that $Z_{a_i} = Z_{b_i}$). Those are geodesic paths in (\mathbf{D}, D) from $\mathbf{p}(a_i)$ to $\mathbf{p}(s_i^*)$, where s_i^* is the (a.s. unique [29]) time in $[a_i, b_i]$ at which Z attains its infimum on that same interval. Alternatively, these paths are parts of the geodesics Φ_{a_i} and Φ_{b_i} introduced earlier. Note also that $\text{Im}(\gamma_l^i) \cap \text{Im}(\gamma_r^i)$ is not necessarily reduced to $\{\mathbf{p}(s_i^*)\}$.

Claim 2. *It is not possible to find $r_1 < r_2 < r_3$ such that $\gamma(r_1), \gamma(r_3) \in \text{Im}(\gamma_l^i)$ and $\gamma(r_2) \notin \text{Im}(\gamma_l^i)$. The same statement is valid for γ_r^i instead of γ_l^i .*

Proof. Indeed, such a situation would clearly violate the uniqueness of the geodesic γ , since we could replace it between times r_1 and r_3 by the arc of γ_l^i from $\gamma(r_1)$ to $\gamma(r_3)$, and still obtain a geodesic from $\mathbf{p}(U)$ to $\mathbf{p}(V)$, distinct from γ . \diamond

Claim 3. *Almost surely, for every $i \geq 1$, the topological boundary of $\mathbf{p}([a_i, b_i])$ in (\mathbf{D}, D) is included in $\text{Im}(\gamma_l^i) \cup \text{Im}(\gamma_r^i)$.*

Proof. This claim is relatively obvious with the interpretation that $([a_i, b_i]/\{\tilde{D}^i = 0\}, \tilde{D}^i)$ is a space with geodesic boundaries given by γ_l^i, γ_r^i , but since we are not referring explicitly to these spaces, let us give a complete proof for this. In fact, the topological boundary of $\mathbf{p}([s, t])$ for any $a_i \leq s \leq t \leq b_i$ is given by [9, Lemma 21] but, as the proof is quite short, we restate the arguments here. Note that $\mathbf{p}([a_i, b_i])$ is closed so that every point in $\partial\mathbf{p}([a_i, b_i])$ is of the form $\mathbf{p}(s')$ for some $s' \in [a_i, b_i]$ and is a limit of a sequence of points of the form $\mathbf{p}(s_n)$, $n \geq 1$, where $s_n \notin [a_i, b_i]$ for every $n \geq 1$. Up to extraction, (s_n) converges to a limit $s \notin (a_i, b_i)$ such that $D(s, s') = 0$. If $s \in \{a_i, b_i\}$ then the claim follows immediately. Otherwise, $s \neq s'$ and, as mentioned during the proof of Lemma 13, this implies $d_X(s, s') = 0$ (19a) or $d_Z(s, s') = 0$ (19b). It cannot hold that $d_X(s, s') = 0$ because $s' \in [a_i, b_i]$ while $s \notin [a_i, b_i]$, so necessarily $d_Z(s, s') = 0$. Assuming for instance that $Z_s = Z_{s'} = \underline{Z}_{s, s'}$, so that $Z_u \geq Z_{s'}$ for every $u \in [a_i, s']$, this implies that $d_Z(s, \inf\{u \in [a_i, b_i] : Z_u = Z_s\}) = 0$. Finally, we get that $\mathbf{p}(s) = \gamma_l^i(Z_{a_i} - Z_s) \in \text{Im}(\gamma_l^i)$. Similarly, if $Z_s = Z_{s'} = \underline{Z}_{s', s}$, we obtain that $\mathbf{p}(s) \in \text{Im}(\gamma_r^i)$. \diamond

From the three claims above, we obtain that there exists a finite number of points $x_1, x_2, \dots, x_{k+1} \in \mathbf{D}$ and integers i_1, \dots, i_k with $x_1 = \mathbf{p}(U)$, $x_{k+1} = \mathbf{p}(V)$, such that γ visits the points x_1, x_2, \dots, x_{k+1} in this order, and such that the segment of γ between x_j and x_{j+1} is

(21a) either included in $\text{Im}(\gamma_l^{i_j})$ or included in $\text{Im}(\gamma_r^{i_j})$

(21b) or included in $\mathbf{p}([a_{i_j}, b_{i_j}])$ and such that its intersection with $\text{Im}(\gamma_l^{i_j}) \cup \text{Im}(\gamma_r^{i_j})$ is a subset of $\{x_j, x_{j+1}\}$.

Indeed, Claims 1 and 2 entail that $\text{Im}(\gamma) \cap (\bigcup_i \text{Im}(\gamma_l^i) \cup \text{Im}(\gamma_r^i))$ is a finite union of segments satisfying (21a) and the parts of γ linking two successive such segments satisfy (21b), by Claim 3. Since the segment of γ between x_j and x_{j+1} is included in $\mathbf{p}([a_{i_j}, b_{i_j}])$ in both cases, we may choose $s_j, t_j \in [a_{i_j}, b_{i_j}]$ such that $x_j = \mathbf{p}(s_j)$ and $x_{j+1} = \mathbf{p}(t_j)$. For any such choice,

$$D(U, V) = \sum_{j=1}^k D(s_j, t_j).$$

We will soon justify that we can choose s_j, t_j satisfying the extra property that $D(s_j, t_j) = \tilde{D}(s_j, t_j)$ on the event $\{\max(I(U, V)) \leq N\}$. Since, by definition, $\mathbf{p}(t_j) = \mathbf{p}(s_{j+1})$, one has $d_X(t_j, s_{j+1}) = 0$ or $d_Z(t_j, s_{j+1}) = 0$ so that $\tilde{D}(t_j, s_{j+1}) = 0$ or $d_Z(t_j, s_{j+1}) = 0$. Similarly, $\tilde{D}(U, s_1) = 0$ or $d_Z(U, s_1) = 0$ and $\tilde{D}(t_k, V) = 0$ or $d_Z(t_k, V) = 0$. As a result, up to potentially doubling some s_j 's and t_j 's, we wrote $D(U, V)$ in the desired form and we conclude the proof by letting $N \rightarrow \infty$, as $\max(I(U, V)) < \infty$ almost surely by Claim 1.

Let us work from now on on the event $\{\max(I(U, V)) \leq N\}$ and justify the possibility of choosing s_j, t_j as previously claimed. If the segment of γ between x_j and x_{j+1} satisfies (21a), then the claim readily follows from Lemma 14, as

$$D(s_j, t_j) \geq |D(s_j, s_*) - D(t_j, s_*)| = |Z_{s_j} - Z_{t_j}| \geq d_Z^i(s_j, t_j) \geq \tilde{D}(s_j, t_j)$$

(Recall that the converse inequality always holds.) We now suppose that the segment of γ between x_j and x_{j+1} satisfies (21b) and we go back to the discrete setting. For $u \in \{0, 1/(2n + l_n), \dots, 1\}$, we denote by $c_n(u)$ the $(2n + l_n)u$ -th corner of F_n . We let $a_{i_j}^n$ and $b_{i_j}^n$ be such that $c_n(a_{i_j}^n)$ and $c_n(b_{i_j}^n)$ are the first and last corners of the i_j -th largest tree of F_n . Standard properties of Brownian motion and the convergence $C_{(n)} \rightarrow X$ entail that $a_{i_j}^n \rightarrow a_{i_j}$ and $b_{i_j}^n \rightarrow b_{i_j}$. Choose two sequences $s_j^n, t_j^n \in [a_{i_j}^n, b_{i_j}^n]$ indexed by n such that $s_j^n \rightarrow s_j$ and $t_j^n \rightarrow t_j$. We denote by u_j^n and v_j^n the vertices incident respectively to $c_n(s_j^n)$ and $c_n(t_j^n)$ and we let γ_j^n be a geodesic in Q_n from u_j^n to v_j^n .

We also let \mathbf{V}_j be the set of vertices of i_j -th largest tree of F_n that do not belong to the maximal geodesic of the slice corresponding to this tree, seen as a subset of $\mathbf{V}(Q_n)$. We will see that $\text{Im}(\gamma_j^n) \setminus \mathbf{V}_j$ is only constituted of vertices “close” to the extremities of γ_j^n in the scale $n^{-1/4}$. Notice first that the middle point $\gamma_j^n(\lfloor D_{Q_n}(u_j^n, v_j^n)/2 \rfloor)$ of γ_j^n necessarily belongs to \mathbf{V}_j for large n . Indeed, let us assume otherwise. Then, for infinitely many values of n , we can find real numbers $u_n \notin [a_{i_j}^n, b_{i_j}^n]$ such that $c_n(u_n)$ is incident to the middle point of γ_j^n . Up to further extraction, we may suppose that $u_n \rightarrow u \notin (a_{i_j}, b_{i_j})$, so that $\mathbf{p}(u)$ does not belong to the interior of $\mathbf{p}([a_{i_j}, b_{i_j}])$. As $\mathbf{p}(u)$ is at mid-distance between x_j and x_{j+1} , we obtain a contradiction with (21b).

We then let $\tilde{s}_j^n \in [a_{i_j}^n, b_{i_j}^n]$ be such that $c_n(\tilde{s}_j^n)$ is incident to

$$\gamma_j^n(\min \{ \alpha \leq D_{Q_n}(u_j^n, v_j^n)/2 : \gamma_j^n(\beta) \in \mathbf{V}_j \text{ for all } \beta \in [\alpha, D_{Q_n}(u_j^n, v_j^n)/2] \})$$

and, symmetrically, $\tilde{t}_j^n \in [a_{i_j}^n, b_{i_j}^n]$ be such that $c_n(\tilde{t}_j^n)$ is incident to

$$\gamma_j^n(\max \{ \alpha \geq D_{Q_n}(u_j^n, v_j^n)/2 : \gamma_j^n(\beta) \in \mathbf{V}_j \text{ for all } \beta \in [D_{Q_n}(u_j^n, v_j^n)/2, \alpha] \}).$$

Up to further extraction, we may suppose that $\tilde{s}_j^n \rightarrow \tilde{s}_j$ and $\tilde{t}_j^n \rightarrow \tilde{t}_j$. We necessarily have $\mathbf{p}(s_j) = \mathbf{p}(\tilde{s}_j)$. Indeed, let us argue by contradiction and suppose that $\mathbf{p}(s_j) \neq \mathbf{p}(\tilde{s}_j)$. The definition immediately entails that $\mathbf{p}(\tilde{s}_j) \in \text{Im}(\gamma_l^{i_j}) \cup \text{Im}(\gamma_r^{i_j})$. But, as $\mathbf{p}(\tilde{s}_j) \in \mathbf{p}([a_{i_j}, b_{i_j}])$, the condition (21b) yields $\mathbf{p}(s_j) = \mathbf{p}(\tilde{s}_j)$, a contradiction. This implies $D(s_j, \tilde{s}_j) = 0$, which also implies $d_X(s_j, \tilde{s}_j) = 0$ or $d_Z(s_j, \tilde{s}_j) = 0$, so that $\tilde{D}(s_j, \tilde{s}_j) = 0$ as s_j and \tilde{s}_j both belong to the same excursion interval $[a_{i_j}, b_{i_j}]$. The same argument shows that $D(t_j, \tilde{t}_j) = \tilde{D}(t_j, \tilde{t}_j) = 0$. Finally, $D_{(n)}(\tilde{s}_j^n, \tilde{t}_j^n) = \tilde{D}_{(n)}^{i_j}(\tilde{s}_j^n, \tilde{t}_j^n)$ by construction and we obtain $D(\tilde{s}_j, \tilde{t}_j) = \tilde{D}(\tilde{s}_j, \tilde{t}_j)$ by (20), and then $D(s_j, t_j) = \tilde{D}(s_j, t_j)$ by the previous discussion. \square

Note that from the formula for $D(s, t)$ given in the statement of Lemma 16 and the definition of D^* , it holds that $D(s, t) \geq D^*(s, t)$ for Lebesgue-almost every $s, t \in [0, 1]$, so that equality holds since $D \leq D^*$ by Lemma 14. Since $D^* \leq d_Z$, which is continuous on $[0, 1]^2$ and null on the diagonal, we get immediately that the pseudo-distances D, D^* are continuous when seen as functions on $[0, 1]^2$, and by density we get that $D = D^*$. This proves Theorem 15.

5 Boltzmann random maps and well-labeled mobiles

5.1 The Bouttier–Di Francesco–Guitter bijection

There is a well-known extension of the Cori–Vauquelin–Schaeffer bijection to general maps. This extension, due to Bouttier, Di Francesco and Guitter [14], can roughly be described in the following way. Any *bipartite* map can be coded by an object called a *well-labeled mobile*. Namely, a mobile is a rooted plane tree \mathbf{t} (we usually call e_0 its root edge) together with a bicolouration of its vertices into “white vertices” and “black vertices.” We denote by $\mathbf{V}_\circ(\mathbf{t})$, $\mathbf{V}_\bullet(\mathbf{t})$ the corresponding sets of vertices, and ask that any two neighboring vertices carry different colors, and that $e_0^- \in \mathbf{V}_\circ(\mathbf{t})$, meaning that mobiles are rooted at a white vertex.

Moreover, the set $\mathbf{V}_\circ(\mathbf{t})$ carries a label function $\ell : \mathbf{V}_\circ(\mathbf{t}) \rightarrow \mathbb{Z}$, that satisfies the following property: if $v' \in \mathbf{V}_\bullet(\mathbf{t})$ is a black vertex, and if $v'_0, v'_1, \dots, v'_{k-1} \in \mathbf{V}_\circ(\mathbf{t})$ denote the neighbors of v' arranged in clockwise order around v' induced by the planar structure of \mathbf{t} (so that $k = \deg_{\mathbf{t}}(v')$), it holds that

$$\ell(v'_{i+1}) - \ell(v'_i) \geq -1, \quad \forall i \in \{0, 1, \dots, k-1\},$$

with the convention that $v'_k = v'_0$. A simple counting argument shows that, as soon as one of the labels, say that of $\ell(v'_0)$, is fixed, there are exactly $\binom{2k-1}{k}$ possible choices for the other labels $\ell(v'_1), \dots, \ell(v'_{k-1})$. At this point of the discussion, we do not insist that the label of any given vertex is fixed, so we really view ℓ as a function defined up an additive constant, as we did in Section 3. We will fix a normalization in the next section.

In our context of maps with a boundary, we use the following conventions. The objects encoding the bipartite maps with perimeter $2l$ (maps of \mathbf{B}_l) are forests $\mathbf{f} = (\mathbf{t}_1, \dots, \mathbf{t}_l)$ of l mobiles, together with a labeling function $\ell : \mathbf{V}_\circ(\mathbf{f}) = \bigsqcup_i \mathbf{V}_\circ(\mathbf{t}_i) \rightarrow \mathbb{Z}$ satisfying the following:

- for $1 \leq i \leq l$, the mobile \mathbf{t}_i equipped with the restriction of ℓ to $\mathbf{V}_\circ(\mathbf{t}_i)$ is a well-labeled mobile;
- for $1 \leq i \leq l$, we have $\ell(\rho_{i+1}) \geq \ell(\rho_i) - 1$, where ρ_i denotes the root vertex of \mathbf{t}_i and $\ell(\rho_{l+1}) = \ell(\rho_1)$.

Remark 4. These forests are in simple bijection with the set of mobiles rooted (unusually) at a black vertex of degree l . But since the external face really plays a different role from the other faces, we prefer indeed to view those as forests of individual mobiles, rather than one single mobile.

The BDG bijection is very similar to the construction presented in Section 3.3. We consider a forest $\mathbf{f} = (\mathbf{t}_1, \dots, \mathbf{t}_l)$ of l mobiles, labeled by ℓ as above and we set $\ell_* = \min\{\ell(v) : v \in \mathbf{V}_\circ(\mathbf{f})\} - 1$. We let $N^{\mathbf{V}}$ be its number of white vertices, $N^{\mathbf{F}}$ be its number of black vertices, and $N^{\mathbf{E}} = N^{\mathbf{V}} + N^{\mathbf{F}}$ be its total number of vertices. (The reason for this notation will become clear in a short moment.)

We identify \mathbf{f} with the map obtained by adding l edges linking the roots $\rho_1, \rho_2, \dots, \rho_l$ of the successive trees in a cycle. This map has one face of degree l incident to the l added edges and another face of degree $2N^{\mathbf{E}} - l$, incident to the l added edges as well as all the mobiles. In the latter face, we let $c_0, c_1, \dots, c_{2N^{\mathbf{V}}-l-1}$ be the sequence of corners incident to white vertices, listed in contour order, starting from the root corner of \mathbf{t}_1 . We extend this list by periodicity and add one corner c_∞ incident to a vertex v_* lying inside the face of degree $2N^{\mathbf{E}} - l$, with label $\ell(c_\infty) = \ell(v_*) = \ell_*$. We define the successor functions by (15) and draw arcs in a non-crossing fashion from c_i to $s(c_i)$ for every $i \in \{0, 1, \dots, 2N^{\mathbf{V}} - l - 1\}$. We root the resulting map at the corner of the degree $2l$ -face that is incident to the root vertex of \mathbf{t}_1 . We obtain a rooted bipartite map \mathbf{m} with perimeter $2l$, with vertex set $\mathbf{V}_\circ(\mathbf{t}) \cup \{v_*\}$, which is naturally pointed at v_* , and such that the root edge points away from v_* .

As in Section 3.3, the fact that the root edge necessarily points away from v_* is a bit unfortunate and we use the same trick in order to overcome this technicality. More precisely, we consider the map obtained from \mathbf{m} by forgetting its root and re-rooting it at a corner chosen uniformly at random among the $2l$ corners of the root face.

A noticeable fact about the BDG bijection is that the black vertices of the forest are in bijection with the internal faces of the map. More precisely, if $v \in \mathbf{V}_\bullet(\mathbf{f})$ corresponds to the face f of \mathbf{m} , then $\deg_{\mathbf{m}}(f) = 2 \deg_{\mathbf{f}}(v)$. Furthermore, the white vertices are bijectively associated with $\mathbf{V}(\mathbf{m}) \setminus \{v_*\}$ (so that we can naturally identify these two sets), in such a way that the label function ℓ gives distances to v_* via the formula

$$d_{\mathbf{m}}(v, v_*) = \ell(v) - \min_{\mathbf{V}_\circ(\mathbf{t})} \ell + 1. \quad (22)$$

As a result (and with the help of the Euler characteristic formula), note that $N^{\mathbf{V}} + 1$, $N^{\mathbf{F}}$ and $N^{\mathbf{E}}$ respectively correspond to the number of vertices, internal faces, and edges of \mathbf{m} — this explains the notation.

5.2 Random mobiles

We now show how to represent the pointed Boltzmann measures \mathbb{W}_l^\bullet of Section 1.4.2 in terms of random trees, via the BDG bijection. Let μ_\circ be the geometric distribution with parameter $1/\mathcal{Z}_q$, given by

$$\mu_\circ(k) = \frac{1}{\mathcal{Z}_q} \left(1 - \frac{1}{\mathcal{Z}_q}\right)^k, \quad k \geq 0.$$

Let also

$$\mu_\bullet(k) = \frac{\mathcal{Z}_q^k \binom{2k+1}{k} q_{k+1}}{f_q(\mathcal{Z}_q)} \quad k \geq 0.$$

Let \mathbb{M}_l be the law of a two-type Bienaymé–Galton–Watson forest, with l independent tree components, and in which even generations (white vertices) use the offspring distribution μ_\circ , while odd generations (black vertices) use the offspring distribution μ_\bullet .

Formally, we let $\mathbb{M}_l = (\mathbb{M}_1)^{\otimes l}$ where \mathbb{M}_1 is defined by

$$\mathbb{M}_1(\{\mathbf{t}\}) = \prod_{u \in \mathbf{V}_\circ(\mathbf{t})} \mu_\circ(k_u(\mathbf{t})) \prod_{u \in \mathbf{V}_\bullet(\mathbf{t})} \mu_\bullet(k_u(\mathbf{t}))$$

for every tree \mathbf{t} , where $k_u(\mathbf{t})$ is the number of children of u in \mathbf{t} . Finally, given a forest with law \mathbb{M}_l , the white vertices carry random integer labels with the following law. Let ξ_1, ξ_2, \dots be a sequence of i.i.d. random variables with shifted geometric(1/2) distributions

$$\mathbb{P}(\xi_i = l) = 2^{-l-2}, \quad l \geq -1, \quad i \geq 1,$$

and let (Y_1, \dots, Y_k) be distributed as the partial sums $(\xi_1, \xi_1 + \xi_2, \dots, \xi_1 + \dots + \xi_k)$ conditionally given $\xi_1 + \dots + \xi_{k+1} = 0$. We say that (Y_1, \dots, Y_k) is a *discrete bridge with shifted geometric steps*, and we let ν_k be the law of this random vector. It is simple to see that, if ν_k^0 is the uniform distribution on

$$\left\{ (x_1, \dots, x_k, x_{k+1}) \in \{-1, 0, 1, 2, \dots\}^{k+1} : \sum_{i=1}^{k+1} x_i = 0 \right\},$$

then ν_k is the image measure of ν_k^0 under $(x_1, \dots, x_{k+1}) \mapsto (\sum_{i=1}^j x_i, 1 \leq j \leq k)$.

Conditionally given the tree, if u is a black vertex with parent u_0 and children u_1, u_2, \dots, u_k , then the law ν_k of the label differences $(\ell(u_i) - \ell(u_0), 1 \leq i \leq k)$ is given by ν_k , while those label differences are independent as u ranges over all black vertices. Finally, the labels of the roots ρ_1, \dots, ρ_l of the forest have same law as $(0, Y_1, \dots, Y_l)$, where (Y_1, \dots, Y_l) has law ν_l . These specify entirely the law of the labels, and in fact, one sees that labels are uniform among all admissible labelings of the forest, in which the root ρ_1 of the first tree carries label 0. For simplicity, we still denote by \mathbb{M}_l the law of forest of well-labeled mobiles thus obtained.

For $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$, we let $\mathbf{B}_{l,n}^{\bullet, \mathbf{S}}$ be the set of pointed maps $(\mathbf{m}, v_*) \in \mathbf{B}^\bullet$ such that $\mathbf{m} \in \mathbf{B}_{l,n}^{\mathbf{S}}$ and we define

$$\mathbb{W}_{l,n}^{\bullet, \mathbf{S}} = W^\bullet(\cdot \mid \mathbf{B}_{l,n}^{\bullet, \mathbf{S}}), \quad (23)$$

where W^\bullet was defined by (4).

Proposition 17. *Let q be an admissible sequence, and $l \geq 1$. Then the image of \mathbb{M}_l under the Bouttier–Di Francesco–Guitter bijection is, after uniform re-rooting on the boundary, the probability measure \mathbb{W}_l^\bullet .*

For $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$, the same statement holds if we replace both \mathbb{M}_l with $\mathbb{M}_l(\cdot \mid N^{\mathbf{S}} = n)$ and \mathbb{W}_l^\bullet with $\mathbb{W}_{l,n}^{\bullet, \mathbf{S}}$.

This is proved by following the same steps as in [30, Proposition 7] and by applying a straightforward analog of Lemma 10; we omit the details. At this point, we can prove Lemma 3, which describes the set $\mathcal{E}^{\mathbf{S}}(q)$ of pairs (l, n) such that $\mathbb{W}(\mathbf{B}_{l,n}^{\mathbf{S}}) > 0$ or, equivalently, such that $\mathbb{M}_l(N^{\mathbf{S}} = n) > 0$.

Proof of Lemma 3. Let us fix the symbol \mathbf{S} . By Proposition 2.2 in [38], under the law \mathbb{M}_1 , there exist two constants α, h such that the support of $N^{\mathbf{S}}$ is included in $\alpha + h\mathbb{Z}_+$, and moreover, for every m large enough, $\mathbb{M}_1(N^{\mathbf{S}} = \alpha + hm) > 0$. In particular, there exists β such that $\mathbb{M}_1(N^{\mathbf{S}} = \beta + hm) > 0$ for every $m \geq 0$. This means that the support of the law of $N^{\mathbf{S}}$ under \mathbb{M}_1 is equal to $R \cup (\beta + h\mathbb{Z}_+)$, for some $R \subset \{0, 1, \dots, \beta - 1\}$. From this, we immediately deduce the similar result for forests under the distribution \mathbb{M}_l . Namely, the support of $N^{\mathbf{S}}$ under \mathbb{M}_l is equal to $R_l \cup (\beta l + h\mathbb{Z}_+)$, for some $R_l \subset \{0, 1, \dots, \beta l - 1\}$. From this observation and Proposition 17, using the remark at the end of the preceding section that the image of $N^{\mathbf{S}}$ under the BDG bijection is $|\mathbf{S}| - \mathbf{1}_{\{\mathbf{S}=\mathbf{V}\}}$, we obtain that the support of the law of $|\mathbf{S}(\mathbf{m})| - \mathbf{1}_{\{\mathbf{S}=\mathbf{V}\}}$ under \mathbb{W}_l^\bullet (or under \mathbb{W}_l by the absolute continuity relation (5)) is equal to

$$R_l \cup (\beta l + h\mathbb{Z}_+).$$

The result follows immediately from this, since the explicit form of h was computed in Section 6.3.1 of [38]. \square

Again, in all the following, when considering pairs (l, n) where l corresponds to the boundary length of a map, and n to its size (measured with respect to the symbol \mathbf{S}), it will always be implicitly assumed that $(l, n) \in \mathcal{E}^{\mathbf{S}}(q)$, which by Lemma 3 means that, up to finitely many exceptions,

$$n = \beta^{\mathbf{S}} l \pmod{h^{\mathbf{S}}}.$$

6 Convergence of the encoding processes

Let us now consider an infinite forest F with distribution $\mathbb{M}_\infty = (\mathbb{M}_1)^{\otimes \mathbb{N}}$. With it, we associate several exploration processes. Let v_0, v_1, v_2, \dots denote the vertices of F (black or white), listed in depth-first order, tree by tree. Let H be the so-called *height process* associated with F , that is, $H(i)$ denotes the distance between the vertex v_i and the root of the tree to which it belongs. For $i \geq 0$, we denote by $\hat{\ell}(i)$ the label of v_i , as well as $\hat{\ell}^0(i) = \hat{\ell}(i) - \ell(\rho_{(i)})$, where $\rho_{(i)}$ is the root of the tree to which v_i belongs. Note that this notion of label process differs from the one introduced during Section 4; we use the notation with a hat in order to avoid confusion. Recall also that, under \mathbb{M}_∞ , the labels are normalized in such a way that the root of the first tree gets label 0, so that the process $\hat{\ell}$ is defined without ambiguity. Finally, let $\Upsilon(i)$ be the number of fully explored trees at time i , that is, $\Upsilon(i) + 1 = p$ whenever v_i belongs to the p -th tree of F , that is $\rho_{(i)} = \rho_p$. We also let

$$\tau_l = \inf\{i \geq 0 : \Upsilon(i) = l\}$$

be the number of (black or white) vertices in the first l trees of the forest. Note for instance that, with the notation of Section 5.1, one has $N^{\mathbf{E}} = \tau_l$ under the law \mathbb{M}_l .

6.1 Convergence for an infinite forest

A key result is the following. Recall that \mathcal{Z}_q is given by (6) and $\rho_q = 2 + \mathcal{Z}_q^3 f_q''(\mathcal{Z}_q)$. Define

$$\sigma_q^2 = \frac{\mathcal{Z}_q \rho_q}{4}, \quad \sigma_{\mathbf{E}}^2 = \frac{\rho_q}{\mathcal{Z}_q}.$$

Proposition 18. *The following joint convergence holds in distribution in $\mathcal{C}(\mathbb{R}_+, \mathbb{R})$ under \mathbb{M}_∞ :*

$$\left(\frac{H(m \cdot)}{\sqrt{m}}, \frac{\Upsilon(m \cdot)}{\sqrt{m}}, \frac{\hat{\ell}^0(m \cdot)}{m^{1/4}} \right) \xrightarrow{m \rightarrow \infty} \left(\frac{2}{\sigma_q} (X - \underline{X}), -\sigma_{\mathbf{E}} \underline{X}, \sqrt{\frac{2\sigma_{\mathbf{E}}}{3}} Z^0 \right)$$

where $(X_t, t \geq 0)$ is a standard Brownian motion, $\underline{X}_t = \inf_{0 \leq s \leq t} X_s$ and Z^0 is the Brownian snake with driving process $X - \underline{X}$, introduced in Section 2.2.

Proof. We note that the two-type branching process with offspring distributions μ_\circ, μ_\bullet and alternating types is a critical branching process, in which the offspring distributions have small exponential moments (this is the place where we use the fact that q is regular critical), as discussed in Proposition 7 of [30]. Furthermore, the spatial displacements with distribution ν_k are centered and carried by $[-k, k]^k$ respectively. In particular, they have moments of all orders, which grow at most polynomially, in the sense that for every $a > 0$,

$$\langle \nu_k, |\cdot|^a \rangle = O(k^a),$$

where $|\cdot|$ is the Euclidean norm in \mathbb{R}^k . This is exactly what is needed to apply Theorems 1 and 3 in [32], which in our particular context stipulate that

$$\left(\frac{H(m \cdot)}{\sqrt{m}}, \frac{\Upsilon(m \cdot)}{\sqrt{m}}, \frac{\hat{\ell}^0(m \cdot)}{m^{1/4}} \right) \xrightarrow{m \rightarrow \infty} \left(\frac{2}{\sigma} (X - \underline{X}), -\frac{\sigma}{b_\circ} \underline{X}, \Sigma \sqrt{\frac{2}{\sigma}} Z^0 \right),$$

where the constants σ, b_\circ and Σ are defined in the following way. The mean matrix of the two-type Galton–Watson process under consideration is given by

$$\begin{pmatrix} 0 & m_\circ \\ m_\bullet & 0 \end{pmatrix},$$

where m_\circ is the mean of μ_\circ , and m_\bullet is the mean of μ_\bullet . Note that $m_\bullet = m_\circ^{-1}$ as an immediate consequence of the fact that q is regular critical. This matrix admits a left invariant vector $\mathbf{a} = (a_\circ, a_\bullet)$ normalized to be a probability, namely $a_\circ = (1 + m_\circ)^{-1}$ and $a_\bullet = (1 + m_\bullet)^{-1}$, and a right invariant $\mathbf{b} = (b_\circ, b_\bullet)$ normalized in such a way that the scalar product $\mathbf{a} \cdot \mathbf{b} = 1$, namely $b_\circ = (1 + m_\circ)/2$ and $b_\bullet = (1 + m_\bullet)/2$. Finally, with (μ_\circ, μ_\bullet) , one can associate a quadratic function $\mathbf{Q} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by

$$\mathbf{Q}(x_\circ, x_\bullet) = ((\sigma_\circ^2 + m_\circ(m_\circ - 1))x_\bullet^2, (\sigma_\bullet^2 + m_\bullet(m_\bullet - 1))x_\circ^2),$$

where σ_\circ^2 and σ_\bullet^2 are the variances of μ_\circ and μ_\bullet . Then σ^2 is given by the scalar product

$$\sigma^2 = \mathbf{a} \cdot \mathbf{Q}(\mathbf{b}).$$

Finally, Σ^2 is given by the formula

$$\Sigma^2 = \frac{1}{2} \sum_{k \geq 1} \frac{\mu_{\bullet}(k)}{m_{\bullet}} (\Sigma_{\bullet}^k)^2$$

where $(\Sigma_{\bullet}^k)^2 = \langle \nu_k, |\cdot|^2 \rangle = k(k+1)/3$, as can be checked in [30]. After computations, which have been performed in Section 3.2 of [30], one obtains in particular

$$m_{\circ} = \mathcal{Z}_q - 1, \quad b_{\circ} = \frac{\mathcal{Z}_q}{2}, \quad \sigma^2 = \frac{\mathcal{Z}_q \rho_q}{4}, \quad \Sigma^2 = \frac{\rho_q}{6}.$$

The conclusion follows. \square

We are also going to need the following fact. For every $m \geq 1$, let

$$\begin{aligned} \Lambda^{\mathbf{V}}(m) &= |\{i \in \{0, 1, \dots, m-1\} : v_i \in \mathbf{V}_{\circ}(\mathbf{f})\}|, \\ \Lambda^{\mathbf{F}}(m) &= |\{i \in \{0, 1, \dots, m-1\} : v_i \in \mathbf{V}_{\bullet}(\mathbf{f})\}|, \end{aligned} \quad (24)$$

be respectively the number of white vertices and the number of black vertices among the first m vertices of F in depth-first order. For convenience, we also let $\Lambda^{\mathbf{V}}(m) = m$ (the number of vertices of either type), so that $\Lambda^{\mathbf{S}}$ makes sense for every $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$. Define

$$a_{\mathbf{V}} = \mathcal{Z}_q^{-1}, \quad a_{\mathbf{F}} = 1 - \mathcal{Z}_q^{-1}, \quad a_{\mathbf{E}} = 1. \quad (25)$$

The first two quantities are the ones that appeared in the proof of Proposition 18, under the notation $a_{\mathbf{V}} = a_{\circ}$ and $a_{\mathbf{F}} = a_{\bullet}$. (Recall that, through the BDG bijection, \mathbf{V} correspond essentially to white vertices, \mathbf{F} to black vertices and \mathbf{E} to edges of the mobile, which are in direct bijection with the set of vertices of both colors.)

In the following statement and later, the notation $\text{oe}(n)$ stands for a quantity that is bounded from above by $c \exp(-c'n^{c''})$ for three positive constants c, c', c'' , uniformly in n .

Proposition 19. *Fix $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$. Then it holds that*

$$\left(\frac{\Lambda^{\mathbf{S}}(m \cdot)}{m} \right) \xrightarrow{m \rightarrow \infty} (a_{\mathbf{S}} t, t \geq 0)$$

in probability under \mathbb{M}_{∞} for the uniform topology over compact subintervals of \mathbb{R}_+ . More precisely, for every $K > 0$, one has the concentration result

$$\mathbb{M}_{\infty} \left(\max_{0 \leq k \leq Km} |\Lambda^{\mathbf{S}}(k) - a_{\mathbf{S}} k| > m^{3/4} \right) = \text{oe}(m).$$

Proof. The result is obvious for $\mathbf{S} = \mathbf{E}$, so that we suppose $\mathbf{S} \in \{\mathbf{V}, \mathbf{F}\}$. We first note that, since $\Lambda^{\mathbf{S}}(k) \leq k$, it suffices to prove the same bound with the maximum restricted over indices $k \in [m^{1/2}, Km]$. Now [32, Proposition 6 (ii)] shows that if $G_k^{\mathbf{S}}$ is the number of vertices in depth-first order (of either type) that have been visited before the k -th vertex of type \mathbf{S} (white if $\mathbf{S} = \mathbf{V}$, black if $\mathbf{S} = \mathbf{F}$), then $\mathbb{M}_{\infty}(|G_k^{\mathbf{S}} - a_{\mathbf{S}}^{-1} k| > k^{3/4}) = \text{oe}(k)$. Now

$|\Lambda^{\mathbf{S}}(k) - a_{\mathbf{S}} k| > m^{3/4}$ implies that $G_{a_{\mathbf{S}}k+m^{3/4}}^{\mathbf{S}} \leq k$ or $G_{(a_{\mathbf{S}}k-m^{3/4})_+}^{\mathbf{S}} \geq k$, the probability of which is bounded from above by

$$\begin{aligned} \mathbb{M}_{\infty}(|G_{a_{\mathbf{S}}k+m^{3/4}}^{\mathbf{S}} - a_{\mathbf{S}}^{-1}(a_{\mathbf{S}}k + m^{3/4})| \geq a_{\mathbf{S}}^{-1}m^{3/4}) \\ + \mathbb{M}_{\infty}(|G_{(a_{\mathbf{S}}k-m^{3/4})_+}^{\mathbf{S}} - a_{\mathbf{S}}^{-1}(a_{\mathbf{S}}k - m^{3/4})_+| \geq a_{\mathbf{S}}^{-1}m^{3/4}). \end{aligned}$$

Taking the maximum over all $k \in [m^{1/2}, Km] \cap \mathbb{Z}$, we see that this quantity is $o(m)$, as claimed. \square

6.2 Convergence for a conditioned forest

We now want a conditioned version of Proposition 19. We are going to need the following estimates. Recall the definition (2) of $h^{\mathbf{S}}$, the definition (8) of $j_L(A)$, and define $Q^{\mathbf{S}}(l, n) = \mathbb{M}_l(N^{\mathbf{S}} = n)$. We will also need the notation

$$\tau_l^{\mathbf{S}} = \Lambda^{\mathbf{S}}(\tau_l).$$

In words, $\tau_l^{\mathbf{E}} = \tau_l$ is the number of vertices in the l first trees of the forest, while $\tau_l^{\mathbf{V}}$ (resp. $\tau_l^{\mathbf{F}}$) is the number of white (resp. black) vertices in these trees.

Lemma 20. *Let $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$. Then*

$$\sup_{n \in \mathcal{E}_l^{\mathbf{S}}} \left| l^2 Q^{\mathbf{S}}(l, n) - h^{\mathbf{S}} j_{1/\sigma_{\mathbf{S}}} \left(\frac{n}{l^2} \right) \right| \xrightarrow{n \rightarrow \infty} 0.$$

Proof. Suppose first that $\mathbf{S} = \mathbf{E}$. In this case, a consequence of the convergence of the second component in Proposition 18 is that

$$\frac{\tau_l}{l^2} \xrightarrow[l \rightarrow \infty]{(d)} T_{1/\sigma_{\mathbf{E}}},$$

where we recall that $T_{1/\sigma_{\mathbf{E}}} = \inf\{t \geq 0 : \underline{X}_t = -1/\sigma_{\mathbf{E}}\}$ is a.s. a continuous function of X under the Wiener measure, due to the fact that $\underline{X}_{(T_{1/\sigma_{\mathbf{E}}}-\varepsilon)_+} > -1/\sigma_{\mathbf{E}} > \underline{X}_{T_{1/\sigma_{\mathbf{E}}}+\varepsilon}$ a.s. for every $\varepsilon > 0$. Moreover, τ_l under \mathbb{M}_{∞} is the sum of l i.i.d. random variables with same law as τ_1 : these are given by the number of vertices of the first l trees in the infinite forest of independent random mobiles. Since it is well known that $T_{1/\sigma_{\mathbf{E}}}$ follows a stable distribution with index $1/2$, with a density given by $j_{1/\sigma_{\mathbf{E}}}$, we conclude that τ_1 under \mathbb{M}_{∞} is in the domain of attraction of this law. The statement is then a consequence of the local limit theorem for stable random variables [13, Theorem 8.4.1].

The remaining two cases $\mathbf{S} \in \{\mathbf{V}, \mathbf{F}\}$ are now direct consequences of the case $\mathbf{S} = \mathbf{E}$ and of Proposition 19, which together imply that we have

$$\frac{\tau_l^{\mathbf{S}}}{l^2} = \frac{\Lambda^{\mathbf{S}}(\tau_l)}{l^2} = \frac{a_{\mathbf{S}} \tau_l (1 + o_{\mathbb{P}}(1))}{l^2}, \quad (26)$$

where $o_{\mathbb{P}}(1)$ denotes a quantity that converges to 0 in probability. This yields that

$$\frac{\tau_l^{\mathbf{S}}}{l^2} \xrightarrow[l \rightarrow \infty]{(d)} a_{\mathbf{S}} T_{1/\sigma_{\mathbf{E}}} \stackrel{(d)}{=} T_{\sqrt{a_{\mathbf{S}}}/\sigma_{\mathbf{E}}} = T_{1/\sigma_{\mathbf{S}}},$$

as can be checked using (7) and (25). The conclusion follows by the same arguments as in the case $\mathbf{S} = \mathbf{E}$. \square

In this section, it is convenient to consider processes whose total duration is not fixed. We let \mathcal{W} be the set of real-valued continuous functions f defined on an interval of the form $[0, \zeta]$ for some $\zeta = \zeta(f) \in [0, \infty)$. This set is endowed with the distance

$$\text{dist}(f, g) = \|f(\cdot \wedge \zeta(f)) - g(\cdot \wedge \zeta(g))\|_{\infty} + |\zeta(f) - \zeta(g)|$$

which makes it a complete separable metric space. For instance, the height process H under the law \mathbb{M}_l is a function with duration $\zeta(H) = N^{\mathbf{E}}$.

Recall the definition of $\mathcal{S}_L^{\mathbf{S}}$ given right after (7).

Proposition 21. *Let $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$ and $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^{\mathbf{S}}$ for some $L > 0$. Then, under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$, it holds that*

$$\left(\frac{H(a_{\mathbf{S}}^{-1} n_k \cdot)}{\sqrt{n_k}}, \frac{\Upsilon(a_{\mathbf{S}}^{-1} n_k \cdot)}{\sqrt{n_k}}, \frac{\hat{\ell}^{\mathbf{0}}(a_{\mathbf{S}}^{-1} n_k \cdot)}{n_k^{1/4}} \right) \xrightarrow[k \rightarrow \infty]{(d)} \left(\frac{2}{\sqrt{a_{\mathbf{S}}}} (X - \underline{X}), -\sigma_{\mathbf{S}} \underline{X}, \sqrt{\frac{2\sigma_{\mathbf{S}}}{3}} Z^{\mathbf{0}} \right),$$

in distribution in the space \mathcal{W}^3 where, in the limit, $X, Z^{\mathbf{0}}$ are understood under the law \mathbb{F}_L^1 defined in Section 2.2.

Proof. For simplicity, let Ξ_k denote the triple appearing in the left-hand side of the convergence. Denote by \mathcal{F}_p the σ -field generated by the p first trees of the generic (canonical process) infinite forest $\mathbf{f} = (\mathbf{t}_1, \mathbf{t}_2, \dots)$, together with their labels. Let G be measurable with respect to $\mathcal{F}_{l'_k}$, with $l'_k < l_k$. Then we have

$$\mathbb{M}_{l_k} [G | N^{\mathbf{S}} = n_k] = \mathbb{M}_{\infty} \left[G \frac{Q^{\mathbf{S}}(l_k - l'_k, n_k - \tau_{l'_k}^{\mathbf{S}})}{Q^{\mathbf{S}}(l_k, n_k)} \right], \quad (27)$$

where it should be understood that the quantity in the expectation is 0 whenever $\tau_{l'_k}^{\mathbf{S}} > n_k$. Now, we impose that $G = \Phi(\Xi'_k)$ is a continuous, bounded function of the triple of processes

$$\Xi'_k = \left(\frac{H(a_{\mathbf{S}}^{-1} n_k \cdot \wedge \tau_{l'_k}^{\mathbf{S}})}{\sqrt{n_k}}, \frac{\Upsilon(a_{\mathbf{S}}^{-1} n_k \cdot \wedge \tau_{l'_k}^{\mathbf{S}})}{\sqrt{n_k}}, \frac{\hat{\ell}^{\mathbf{0}}(a_{\mathbf{S}}^{-1} n_k \cdot \wedge \tau_{l'_k}^{\mathbf{S}})}{n_k^{1/4}} \right),$$

where we assume that $l'_k \sim L' \sigma_{\mathbf{S}} \sqrt{n_k}$ for some $0 < L' < L$. Proposition 18 shows the convergence in distribution

$$\left(\frac{H(a_{\mathbf{S}}^{-1} n_k \cdot)}{\sqrt{a_{\mathbf{S}}^{-1} n_k}}, \frac{\Upsilon(a_{\mathbf{S}}^{-1} n_k \cdot)}{\sqrt{a_{\mathbf{S}}^{-1} n_k}}, \frac{\hat{\ell}^{\mathbf{0}}(a_{\mathbf{S}}^{-1} n_k \cdot)}{(a_{\mathbf{S}}^{-1} n_k)^{1/4}} \right) \xrightarrow[k \rightarrow \infty]{(d)} \left(\frac{2}{\sigma_q} (X - \underline{X}), -\sigma_{\mathbf{E}} \underline{X}, \sqrt{\frac{2\sigma_{\mathbf{E}}}{3}} Z^{\mathbf{0}} \right),$$

where the limit is understood under the law \mathbb{P} . Using the convergence of the second component, the asymptotic behavior of l'_k and the fact that $a_{\mathbf{S}}\sigma_{\mathbf{S}}^2 = \sigma_{\mathbf{E}}^2$, it follows that

$$\frac{\tau_{l'_k}}{a_{\mathbf{S}}^{-1}n_k} \xrightarrow[k \rightarrow \infty]{(d)} T_{L'},$$

and that this convergence holds jointly with the previous one. From this, it follows that Ξ'_k converges in distribution under \mathbb{M}_{∞} to the triple

$$\Xi'_{\infty} = \left(\frac{2}{\sqrt{a_{\mathbf{S}}}\sigma_q}(X - \underline{X})_{\cdot \wedge T_{L'}}, -\sigma_{\mathbf{S}}\underline{X}_{\cdot \wedge T_{L'}}, \sqrt{\frac{2\sigma_{\mathbf{S}}}{3}}Z^{\mathbf{0}}_{\cdot \wedge T_{L'}} \right)$$

and an application of (26) implies that $\tau_{l'_k}^{\mathbf{S}}/n_k \rightarrow T_{L'}$ jointly with the above convergence. By the Skorokhod representation theorem, we may assume that the probability space is chosen in such a way that these convergences hold in the almost-sure sense, and then (27) together with Lemma 20 implies that

$$\mathbb{M}_{l_k} [\Phi(\Xi'_k) | N^{\mathbf{S}} = n_k] \xrightarrow[k \rightarrow \infty]{(d)} \mathbb{E} \left[\Phi(\Xi'_{\infty}) \frac{L^2}{(L-L')^2} \frac{j_{1/\sigma_{\mathbf{S}}}\left(\frac{(1-T_{L'})}{\sigma_{\mathbf{S}}^2(L-L')^2}\right)}{j_{1/\sigma_{\mathbf{S}}}\left(\frac{1}{\sigma_{\mathbf{S}}^2 L^2}\right)} \right]$$

and the limit can be re-expressed as

$$\mathbb{E} \left[\Phi(\Xi'_{\infty}) \frac{j_{L-L'}(1-T_{L'})}{j_L(1)} \right] = \mathbb{F}_L^1[\Phi(\Xi'_{\infty})].$$

By definition, $\Phi(\Xi'_{\infty})$ is $\mathcal{G}_{T_{L'}}$ -measurable and, by Galmarino's test, we have $\mathcal{G}_{T_{L'}} = \sigma(X_{\cdot \wedge T_{L'}}, Z^{\mathbf{0}}_{\cdot \wedge T_{L'}})$, so that, if it exists, the limit of the triple considered in the statement of the proposition necessarily has the claimed law, by virtue of Proposition 9.

To conclude the proof, it remains to prove that the laws of the processes under consideration are tight in \mathcal{W}^3 . We can argue as follows. Let f be a continuous function defined on an interval I , and $J \subseteq I$ be a subinterval of I . Denote by

$$\omega(f, \delta, J) = \sup_{s, t \in J, |t-s| \leq \delta} |f(t) - f(s)|$$

the modulus of continuity of f restricted to J , and let $\omega(f, \delta) = \omega(f, \delta, I)$.

Here, let Y_k denote either of the components of Ξ_k . Then, under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$,

$$\omega(Y_k, \delta) \leq \omega(Y_k, \delta, [0, a_{\mathbf{S}}\tau_{l'_k}/n_k]) + \omega(Y_k, \delta, [a_{\mathbf{S}}\tau_{l'_k}/n_k, a_{\mathbf{S}}\tau_{l_k}/n_k])$$

while the second component has same distribution as

$$\omega(Y_k, \delta, [0, a_{\mathbf{S}}\tau_{l_k-l'_k}/n_k])$$

by a symmetry argument (the l_k trees of the labeled forest are exchangeable). Choosing $l'_k \sim l_k/2$, we obtain from the convergence of Ξ'_k (for $L' = L/2$) that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \mathbb{M}_{l_k}(\omega(Y_k, \delta) \geq \varepsilon) \\ & \leq \limsup_{n \rightarrow \infty} \mathbb{M}_{l_k}(\omega(Y_k, \delta, [0, a_{\mathbf{S}} \tau_{l'_k}/n_k]) \geq \varepsilon/2) + \mathbb{M}_{l_k}(\omega(Y_k, \delta, [0, a_{\mathbf{S}} \tau_{l_k - l'_k}/n_k]) \geq \varepsilon/2) \\ & \leq 2 \mathbb{F}_L^1(\omega(Y_\infty, \delta, [0, T_{L/2}]) \geq \varepsilon/2), \end{aligned}$$

where Y_∞ is the limit of Y_k (for instance $Y_\infty = 2(X - \underline{X})/\sqrt{a_{\mathbf{S}}} \sigma_q$ if Y_k is the first component of Ξ_k). This quantity converges to 0 as $\delta \rightarrow 0$, for any fixed $\varepsilon > 0$. From this, it is an immediate consequence of the Ascoli–Arzela theorem that the laws of Ξ_k , $k \geq 1$ are relatively compact in \mathcal{W}^3 . \square

Corollary 22. *Let $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$ and $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^{\mathbf{S}}$ for some $L > 0$. Then, under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$, it holds that*

$$\left(\frac{H(a_{\mathbf{S}}^{-1} n_k \cdot)}{\sqrt{n_k}}, \frac{\Upsilon(a_{\mathbf{S}}^{-1} n_k \cdot)}{\sqrt{n_k}}, \frac{\hat{\ell}(a_{\mathbf{S}}^{-1} n_k \cdot)}{n_k^{1/4}} \right) \xrightarrow[k \rightarrow \infty]{(d)} \left(\frac{2}{\sqrt{a_{\mathbf{S}}}} (X - \underline{X}), -\sigma_{\mathbf{S}} \underline{X}, \sqrt{\frac{2\sigma_{\mathbf{S}}}{3}} Z \right),$$

in distribution in the space \mathcal{W}^3 where, in the limit, X, Z are understood under the law \mathbb{F}_L^1 , defined in Section 2.2.

Proof. It suffices to apply the preceding proposition, noting that one can get the following representation for the label process $\hat{\ell}$ in terms of $\hat{\ell}^0$:

$$\hat{\ell}(i) = \hat{\ell}^0(i) + \mathbf{B}(\Upsilon(i)),$$

where \mathbf{B} is a discrete bridge with shifted geometric step, with law ν_{l_k} defined in Section 5.2. It holds that, under our hypotheses, $\mathbf{B}(\sigma_{\mathbf{S}} \sqrt{n_k} \cdot) / \sqrt{2\sigma_{\mathbf{S}}} n_k^{1/4}$ converges in distribution to a Brownian bridge \mathbf{b} with duration L (see [10, Proposition 7]). Putting things together, we obtain that

$$\frac{\hat{\ell}(a_{\mathbf{S}}^{-1} n_k \cdot)}{n_k^{1/4}} = \frac{\hat{\ell}^0(a_{\mathbf{S}}^{-1} n_k \cdot)}{n_k^{1/4}} + \frac{\mathbf{B}(\sigma_{\mathbf{S}} \sqrt{n_k} (\Upsilon(a_{\mathbf{S}}^{-1} n_k \cdot) / \sigma_{\mathbf{S}} \sqrt{n_k}))}{n_k^{1/4}},$$

which converges in distribution to $s \mapsto \sqrt{2\sigma_{\mathbf{S}}/3} Z_s^0 + \sqrt{2\sigma_{\mathbf{S}}} \mathbf{b}_{-\underline{X}_s}$, jointly with the rescaled processes H and Υ . By definition (12), this yields the result. \square

Finally, we note that the convergence of $\Lambda^{\mathbf{S}}$ stated in Proposition 19 still holds under conditioned forests. Indeed, since the conditioning event $\{N^{\mathbf{S}} = n_k\}$ has a probability $Q^{\mathbf{S}}(l_k, n_k) = \Theta(l_k^{-2}) = \Theta(n_k^{-1})$ by Lemma 20, we obtain that for any $c' > 0$,

$$\begin{aligned} & \mathbb{M}_{l_k} \left(\max_{0 \leq i \leq N^{\mathbf{E}}} |\Lambda^{\mathbf{S}}(i) - a_{\mathbf{S}} i| > n_k^{3/4} \mid N^{\mathbf{S}} = n_k \right) \\ & \leq cn_k \mathbb{M}_\infty \left(\max_{0 \leq i \leq c'n_k} |\Lambda^{\mathbf{S}}(i) - a_{\mathbf{S}} i| > n_k^{3/4} \right) + \mathbb{M}_{l_k}(N^{\mathbf{E}} > c'n_k \mid N^{\mathbf{S}} = n_k). \end{aligned}$$

for some constant $c > 0$. The first term is $oe(n_k)$ by Proposition 19. The second term is equal to 0 if $\mathbf{S} = \mathbf{E}$ and $c' > 1$. If $\mathbf{S} = \mathbf{V}$, it can be bounded by

$$cn_k \mathbb{M}_\infty(\Lambda^{\mathbf{V}}(c'n_k) \leq n_k) = oe(n_k),$$

as soon as c' is chosen strictly larger than $a_{\mathbf{V}}^{-1}$, again by Proposition 19. The argument is the same if $\mathbf{S} = \mathbf{F}$. In particular, as $\Lambda^{\mathbf{S}}(N^{\mathbf{E}}) = N^{\mathbf{S}}$, this implies that, under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$, one has

$$\frac{N^{\mathbf{S}}}{N^{\mathbf{E}}} \longrightarrow a_{\mathbf{S}}$$

in probability as $k \rightarrow \infty$. This implies the following reformulation and refinement of the preceding corollary.

Corollary 23. *Let $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$ and $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^{\mathbf{S}}$ for some $L > 0$. Then, under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$, it holds that*

$$\left(\frac{H(N^{\mathbf{E}} \cdot)}{\sqrt{n_k}}, \frac{\Upsilon(N^{\mathbf{E}} \cdot)}{\sqrt{n_k}}, \frac{\hat{\ell}(N^{\mathbf{E}} \cdot)}{n_k^{1/4}} \right) \xrightarrow[k \rightarrow \infty]{(d)} \left(\frac{2}{\sqrt{a_{\mathbf{S}}} \sigma_q} (X - \underline{X}), -\sigma_{\mathbf{S}} \underline{X}, \sqrt{\frac{2\sigma_{\mathbf{S}}}{3}} Z \right),$$

in distribution in the space \mathcal{W}^3 , where, in the limit, X, Z are understood under the law \mathbb{F}_L^1 . Moreover, one has, still under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$,

$$\left(\frac{\Lambda^{\mathbf{S}}(N^{\mathbf{E}} t)}{N^{\mathbf{S}}}, 0 \leq t \leq 1 \right) \xrightarrow[k \rightarrow \infty]{} \text{Id}_{[0,1]}$$

in probability.

6.3 Convergence of the white contour and label processes

Finally, we consider a variant of the latest corollary where the height process is replaced by the slightly more convenient contour processes. First, given the forest F with possibly infinitely many trees, we let \bar{C} be its contour process, defined as follows. We first add edges linking the roots of consecutive trees as we did before (see for instance the left part of Figure 5). We then let c_0, c_1, \dots be the list of corners of the trees, arranged in contour order. The purpose of the added edges linking the roots is to “split in two” the root corners of the trees: a tree with p edges will thus have $2p+1$ corners. Finally, we let $\bar{C}(i)$ be the distance between c_i and the root of the tree to which it belongs. Note that there is slight difference with the contour function C defined in Section 4.1, where a downstep was separating the contours of successive trees, instead of the horizontal step we have here. It is a standard fact, proved in [21, Chapter 2.4], that the contour and height process of a forest are asymptotically similar in the following sense. First, let $f(i) + 1$ be the number of distinct vertices incident to the corners c_0, c_1, \dots, c_i (these vertices being $v_0, \dots, v_{f(i)}$). Then it holds that $f(i) \leq i$ for every $i \geq 0$, and for every $m \geq 0$, one has

$$\sup_{0 \leq i \leq m} |\bar{C}(i) - H(f(i))| \leq 1 + \sup_{0 \leq i \leq m} |H(i+1) - H(i)|$$

and

$$\max_{0 \leq i \leq m} \left| f(i) - \frac{i}{2} \right| \leq 1 + \frac{1}{2} \max_{0 \leq i \leq m} H_i. \quad (28)$$

From this, and the convergence of the rescaled height process stated in Corollary 23, it follows easily that under the same hypotheses,

$$\frac{\bar{C}(2N^{\mathbf{E}} \cdot)}{\sqrt{n_k}} \xrightarrow[k \rightarrow \infty]{(d)} \frac{2}{\sqrt{a_{\mathbf{S}}} \sigma_q} (X - \underline{X}),$$

where X is understood under \mathbb{F}_L^1 . Now, we want to consider the *white contour process* \bar{C}° , defined as follows. We let $c_0^\circ, c_1^\circ, \dots, c_{N^{\mathbf{E}}-1}^\circ$ be the list of corners that are incident to white vertices, arranged in contour order as above. Then $2\bar{C}^\circ(i)$ is the distance between c_i° and the root of the tree to which it belongs (note that this number is even). In the contour process, white vertices are visited once in every pair of steps, except at times when one of the trees has been fully explored. The number of such exceptions is $l_k = O(\sqrt{n_k})$, so clearly the preceding convergence implies

$$\frac{\bar{C}^\circ(N^{\mathbf{E}} \cdot)}{\sqrt{n_k}} \xrightarrow[k \rightarrow \infty]{(d)} \frac{1}{\sqrt{a_{\mathbf{S}}} \sigma_q} (X - \underline{X}),$$

where the limit is understood under \mathbb{F}_L^1 . Define $\Upsilon^\circ(i)$ to be the number of completely explored trees when visiting c_i° , as well as $\ell^\circ(i)$ to be the label of c_i° . Beware that the definitions of \bar{C}° , Υ° and ℓ° involve corners listed in contour order, instead of vertices listed in depth-first order, as in the definitions of H , Υ and $\hat{\ell}$. Similar arguments entail the following joint convergence.

Corollary 24. *Let $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$ and $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^{\mathbf{S}}$ for some $L > 0$. Then, under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$, it holds that*

$$\left(\frac{\bar{C}^\circ(N^{\mathbf{E}} \cdot)}{\sqrt{n_k}}, \frac{\Upsilon^\circ(N^{\mathbf{E}} \cdot)}{\sqrt{n_k}}, \frac{\ell^\circ(N^{\mathbf{E}} \cdot)}{n_k^{1/4}} \right) \xrightarrow[k \rightarrow \infty]{(d)} \left(\frac{1}{\sqrt{a_{\mathbf{S}}} \sigma_q} (X - \underline{X}), -\sigma_{\mathbf{S}} \underline{X}, \sqrt{\frac{2\sigma_{\mathbf{S}}}{3}} Z \right),$$

in distribution in the space \mathcal{W}^3 , where, in the limit, X, Z are understood under the law \mathbb{F}_L^1 .

7 Proof of the invariance principle

In this section, we prove Theorem 5 and Theorem 8. The arguments, originating in [26, Section 8], are now very standard, and have been applied successfully in [6, 2, 11, 1] in particular. Our approach is an easy adaptation of the arguments that can be found in either of these papers, so here we will be a bit sketchy. Let q be a regular critical sequence as in the previous section.

7.1 Convergence of conditioned pointed maps

The goal of this subsection is to prove the following analog of Theorem 5 under the pointed laws $\mathbb{W}_{l,n}^{\bullet,\mathbf{S}}$, defined by (23).

Theorem 25. *Let \mathbf{S} denote one of the symbols \mathbf{V} , \mathbf{E} , \mathbf{F} , and $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^{\mathbf{S}}$ for some $L > 0$. For $k \geq 0$, denote by M_k^\bullet a random map with distribution $\mathbb{W}_{l_k, n_k}^{\bullet,\mathbf{S}}$. Then*

$$\left(\frac{4\sigma_{\mathbf{S}}^2}{9} n_k \right)^{-1/4} M_k^\bullet \xrightarrow[k \rightarrow \infty]{(d)} \text{BD}_L$$

in distribution for the Gromov–Hausdorff topology.

Fix $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$ and $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^{\mathbf{S}}$ for some $L > 0$. For every $k \geq 0$, consider a forest (F, ℓ) of labeled mobiles with law $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$, and let M_k^\bullet be the random map with distribution $\mathbb{W}_{l_k, n_k}^{\bullet,\mathbf{S}}$ obtained by the process of Proposition 17.

The proof of the convergence of M_k^\bullet to the Brownian disk follows closely in spirit that of [26, Section 8]. Let

$$D'_k(i, j) = d_{M_k^\bullet}(c_i^\circ, c_j^\circ), \quad 0 \leq i, j \leq N^{\mathbf{E}}$$

(with the convention that $c_{N^{\mathbf{E}}}^\circ = c_0^\circ$) and extend D'_k to a continuous function on $[0, N^{\mathbf{E}}]^2$ by a formula similar to (16). In this way, D'_k satisfies the triangle inequality, and one has

$$D'_k(i, j) \leq \ell^\circ(i) + \ell^\circ(j) - 2 \max \left(\min_{[i \wedge j, i \vee j]} \ell^\circ, \min_{[i \vee j, N^{\mathbf{E}}] \cup [0, i \wedge j]} \ell^\circ \right) + 2, \quad (29)$$

see for instance [24, Lemma 3.1] for the special case of p -mobiles (in which black vertices all have degree p , which corresponds to $2p$ -angulations via the BDG bijection), but the proof in this general context is the same. Clearly, it also holds that, if c_i° and c_j° are incident to the same vertex, which means that $\bar{C}^\circ(i) = \bar{C}^\circ(j) = \min_{r \in [i \wedge j, i \vee j]} \bar{C}^\circ(r)$ and $\Upsilon^\circ(i) = \Upsilon^\circ(j)$ then $D'_k(i, j) = 0$. This generalizes to all $s, t \in [0, N^{\mathbf{E}}]$ rather than just integer values.

Now for $s, t \in [0, 1]$ let $D'_{(k)}(s, t) = (4\sigma_{\mathbf{S}}^2 n_k / 9)^{-1/4} D'_k(N^{\mathbf{E}} s, N^{\mathbf{E}} t)$. The same proof as [24, Proposition 3.2] (the key ingredients being (29) and the convergence of the rescaled labeled process ℓ° , established in Corollary 24) shows that, under \mathbb{M}_{l_k} , the laws of $D'_{(k)}$ are tight in the space $\mathcal{C}([0, 1]^2, \mathbb{R})$. Therefore, from any extraction, one can further extract a subsequence along which one has the following joint convergence in distribution under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$,

$$\left(\frac{\bar{C}^\circ(N^{\mathbf{E}} \cdot)}{\sqrt{n_k} / (\sqrt{a_{\mathbf{S}}} \sigma_q)}, \frac{\Upsilon^\circ(N^{\mathbf{E}} \cdot)}{\sigma_{\mathbf{S}} \sqrt{n_k}}, \frac{\ell^\circ(N^{\mathbf{E}} \cdot)}{(4\sigma_{\mathbf{S}}^2 n_k / 9)^{1/4}}, D'_{(k)} \right) \xrightarrow[k \rightarrow \infty]{(d)} (X - \underline{X}, -\underline{X}, Z, D') \quad (30)$$

where D' is some random continuous function on $[0, 1]^2$, and (X, Z) is the snake process under the law \mathbb{F}_L^1 introduced in Section 2.2. Recall the definition of the set \mathcal{D} in Section 2.3, as well as the definition of s_* , the a.s. unique time in $[0, 1]$ such that $Z_{s_*} = \inf Z$. We have the following result, which follows from a simple limiting argument, based on (29) and the discussion below, as well as (22) for the last point.

Lemma 26. *Almost-surely, it holds that*

- the random function D' is a pseudo-distance on $[0, 1]$, such that $D' \in \mathcal{D}$, and
- for every $s \in [0, 1]$, $D'(s, s_*) = Z_s - Z_{s_*}$.

Let D^* be the Brownian disk distance defined from the process (X, Z) by (14). By definition, it holds that $D' \leq D^*$. The conclusion will follow from the following re-rooting property.

Lemma 27. *Let U, V be two independent uniform random variables in $[0, 1]$, independent of the other random variables under consideration. Then $D'(U, V)$ and $D'(s_*, U)$ have the same distribution.*

Proof. This is again obtained by a limiting argument. The idea is to couple the random variables U and V with two uniformly chosen vertices of M_k^\bullet .

For $i \in \{1, 2, \dots, N^{\mathbf{V}}\}$, let $g(i)$ be the first time j such that c_j° is the i -th white vertex in depth-first order. We also let $g(0) = 0$ and extend by linear interpolation g into a continuous increasing function on $[0, N^{\mathbf{V}}]$. Recall the definition (24) of $\Lambda^{\mathbf{V}}(i)$ (for $0 \leq i \leq N^{\mathbf{E}}$), the number of white vertices among v_0, \dots, v_i , where v_0, v_1, \dots are listed in depth-first order. For $i \in \{1, 2, \dots, N^{\mathbf{V}}\}$, let

$$K(i) = \inf\{j \in \{0, 1, \dots, N^{\mathbf{E}}\} : \Lambda^{\mathbf{V}}(j) = i\},$$

so by definition, $v_{K(i)}$ is the i -th visited white vertex in depth-first order. Recall also from Section 6.3 that $f(j) + 1$ is the number of distinct vertices incident to the corners c_0, \dots, c_j , so if we let

$$K'(i) = \inf\{j \geq 0 : f(j) = i\},$$

then $c_{K'(i)}$ is the first time of visit of the vertex v_i in contour order. Consequently, $K'(K(i))$ is the first time of visit, in the contour sequence c_0, c_1, \dots , of $v_{K(i)}$. Finally, since $g(i)$ is the first index j such that c_j° is incident to $v_{K(i)}$, we have that

$$g(i) = K'(K(i))/2 + R_k(i) \tag{31}$$

where the error term satisfies $\sup_{1 \leq i \leq N^{\mathbf{V}}} |R_k(i)| \leq l_k = O(\sqrt{n_k})$, recalling the discussion of Section 6.3.

It follows from the last part of Corollary 23 and from (28) respectively that

$$\frac{K(N^{\mathbf{V}} \cdot)}{N^{\mathbf{E}}} \xrightarrow[k \rightarrow \infty]{} \text{Id}_{[0,1]}, \quad \frac{K'(N^{\mathbf{E}} \cdot)}{2N^{\mathbf{E}}} \xrightarrow[k \rightarrow \infty]{} \text{Id}_{[0,1]},$$

in probability under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$. From this and (31), we conclude that

$$\frac{g(N^{\mathbf{V}} \cdot)}{N^{\mathbf{E}}} \xrightarrow[k \rightarrow \infty]{} \text{Id}_{[0,1]}, \tag{32}$$

still in the same sense.

Now, if I is a uniform random variable on $\{1, 2, \dots, N^{\mathbf{V}}\}$ independent of the rest, then $v_{K(I)}$ is uniformly distributed among the white vertices of the forest, that is, among the vertices of the map M_k^\bullet distinct from the distinguished vertex v_* . Therefore, if we let $v_{(1)} = v_{K(\lceil N^{\mathbf{V}}U \rceil)}$ and $v_{(2)} = v_{K(\lceil N^{\mathbf{V}}V \rceil)}$, then $v_{(1)}, v_{(2)}$ can be coupled with two independent uniform vertices $v'_{(1)}, v'_{(2)}$ of M_k^\bullet in such a way that the conditional probability given M_k^\bullet that $v_{(i)} \neq v'_{(i)}$, $i \in \{1, 2\}$, is at most $1/|\mathbf{V}(M_k^\bullet)|$. The latter quantity, also equal to $1/(N^{\mathbf{V}} + 1)$, converges to 0 in probability under $\mathbb{M}_{l_k}(\cdot | N^{\mathbf{S}} = n_k)$ as $k \rightarrow \infty$.

Since v_* is a uniform random vertex of M_k^\bullet , we obtain that

$$d_{M_k^\bullet}(v_*, v'_{(1)}) \stackrel{(d)}{=} d_{M_k^\bullet}(v'_{(1)}, v'_{(2)}).$$

Due to the above discussion, outside a set of vanishing probability, we may assume that $v'_{(1)} = v_{(1)}$ and $v'_{(2)} = v_{(2)}$.

Now note that, by (22),

$$d_{M_k^\bullet}(v_*, v_{(1)}) = \ell^\circ(g(\lceil N^{\mathbf{V}}U \rceil)) - \inf \ell^\circ + 1,$$

and, by definition of D'_k ,

$$d_{M_k^\bullet}(v_{(1)}, v_{(2)}) = D'_k(g(\lceil N^{\mathbf{V}}U \rceil), g(\lceil N^{\mathbf{V}}V \rceil)).$$

Using (32), we conclude that

$$\left(\frac{4\sigma_{\mathbf{S}}^2 n_k}{9}\right)^{-1/4} d_{M_k^\bullet}(v_*, v_{(1)}) \xrightarrow[k \rightarrow \infty]{(d)} Z_U - \inf Z = D'(s_*, U),$$

while

$$\left(\frac{4\sigma_{\mathbf{S}}^2 n_k}{9}\right)^{-1/4} d_{M_k^\bullet}(v_{(1)}, v_{(2)}) \xrightarrow[k \rightarrow \infty]{(d)} D'(U, V).$$

It follows that $D'(U, V)$ and $D'(s_*, U)$ have same distribution, as claimed. \square

To conclude the proof of Theorem 25, we note that, by Lemma 27,

$$E[D'(U, V)] = E[D'(s_*, U)] = E[Z_U - \inf Z] = E[D^*(s_*, U)] = E[D^*(U, V)],$$

whence it follows that $D'(U, V) = D^*(U, V)$ a.s. since $D' \leq D^*$. Note that we have used the fact that $D^*(s_*, U)$ and $D^*(U, V)$ have same distribution, a fact that follows from Theorem 15 and Lemma 12. This is in fact the only place where we use the specific study of Sections 3 and 4.

This implies, by Fubini's theorem, that a.s. $D'(s, t) = D^*(s, t)$ for a.e. $s, t \in [0, 1]$, so that $D' = D^*$ a.s. by a density argument. This identifies D' uniquely, and shows that the convergence of $D'_{(k)}$ to $D' = D^*$ holds without having to pass to a subsequence. From there, showing the Gromov–Hausdorff convergence of $(4\sigma_{\mathbf{S}}^2 n_k/9)^{-1/4} M_k^\bullet$ under $\mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}}$ to BD_L is routine, see e.g. [10, Section 3.2].

7.2 De-pointing

Here we show how to dispose of the pointing that intervenes in Theorem 25. The argument closely follows the last section of [1], see also [11] for a similar situation.

Similarly to the absolute continuity relation (5), for $\mathbf{S} \in \{\mathbf{V}, \mathbf{E}, \mathbf{F}\}$, we have

$$d\mathbb{W}_{l,n}^{\mathbf{S}}(\mathbf{m}) = \frac{K_{l,n}^{\mathbf{S}}}{|\mathbf{V}|} d(\phi_* \mathbb{W}_{l,n}^{\bullet, \mathbf{S}})(\mathbf{m}) \quad (33)$$

where $K_{l,n}^{\mathbf{S}} = \mathbb{W}_{l,n}^{\bullet, \mathbf{S}}[1/|\mathbf{V}|]^{-1}$. In particular, $\mathbb{W}_{l,n}^{\mathbf{V}} = \phi_* \mathbb{W}_{l,n}^{\bullet, \mathbf{V}}$ for every l, n , so there is nothing more to prove for $\mathbf{S} = \mathbf{V}$; Theorem 5 is equivalent to Theorem 25 in this case.

Now suppose that $\mathbf{S} \in \{\mathbf{E}, \mathbf{F}\}$. Then, by Propositions 17 and 19, it holds that, for every $\varepsilon > 0$,

$$\mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \left(\left| |\mathbf{V}| - \frac{a_{\mathbf{V}}}{a_{\mathbf{S}}} n_k \right| > \varepsilon n_k \right) = \mathbb{M}_{l_k} \left(\left| N^{\mathbf{V}} + 1 - \frac{a_{\mathbf{V}}}{a_{\mathbf{S}}} n_k \right| > \varepsilon n_k \mid N^{\mathbf{S}} = n_k \right) = o_{\varepsilon}(n_k).$$

(Note that a bound of the form $o(1/n_k)$ would suffice for the argument to work.) From this and the fact that $1/|\mathbf{V}| \leq 1$, we obtain that, for every $\delta > 0$,

$$\mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \left[\left| \frac{a_{\mathbf{V}} n_k}{a_{\mathbf{S}} |\mathbf{V}|} - 1 \right| \leq \delta + \left(\frac{a_{\mathbf{V}}}{a_{\mathbf{S}}} n_k + 1 \right) \mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \left(\left| \frac{a_{\mathbf{V}}}{a_{\mathbf{S}}} n_k - |\mathbf{V}| \right| > \delta |\mathbf{V}| \right) \right] = \delta + o_{\varepsilon}(n_k),$$

since

$$\begin{aligned} \mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \left(\left| \frac{a_{\mathbf{V}}}{a_{\mathbf{S}}} n_k - |\mathbf{V}| \right| > \delta |\mathbf{V}| \right) &\leq \mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \left(|\mathbf{V}| \leq \frac{a_{\mathbf{V}} n_k}{a_{\mathbf{S}} 2} \right) \\ &\quad + \mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \left(\left| \frac{a_{\mathbf{V}}}{a_{\mathbf{S}}} n_k - |\mathbf{V}| \right| > \delta \frac{a_{\mathbf{V}} n_k}{a_{\mathbf{S}} 2} \right) = o_{\varepsilon}(n_k). \end{aligned}$$

From this it follows that $K_{l_k, n_k}^{\mathbf{S}} = \frac{a_{\mathbf{V}}}{a_{\mathbf{S}}} n_k (1 + o(1))$ as $k \rightarrow \infty$, and then that

$$\mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \left[\left| \frac{K_{l_k, n_k}^{\mathbf{S}}}{|\mathbf{V}|} - 1 \right| \right] \xrightarrow[k \rightarrow \infty]{} 0.$$

From this and (33), we obtain the following result.

Lemma 28. *For every $\mathbf{S} \in \{\mathbf{E}, \mathbf{F}\}$, and every $(l_k, n_k)_{k \geq 0} \in \mathcal{S}_L^{\mathbf{S}}$, one has*

$$\left\| \mathbb{W}_{l_k, n_k}^{\mathbf{S}} - \phi_* \mathbb{W}_{l_k, n_k}^{\bullet, \mathbf{S}} \right\| \xrightarrow[k \rightarrow \infty]{} 0,$$

where $\|\cdot\|$ is the total variation norm.

Theorem 5 is now a direct consequence of this statement combined with Theorem 25.

7.3 Proof of the convergence of Boltzmann maps

This section is dedicated to the proof of Theorem 8. As in the proof of Theorem 5, we first focus on random maps with distribution \mathbb{W}_l^\bullet , which are easier to handle, since they are directly related by Proposition 17 to random labeled forests with law \mathbb{M}_l .

The result follows from the fact that, for any measurable and bounded function Φ ,

$$\begin{aligned} \mathbb{W}_l^\bullet[\Phi] &= \sum_{n \in \mathcal{E}_l^\mathbf{V}} \mathbb{W}_l^\bullet(|\mathbf{V}| = n + 1) \mathbb{W}_{l,n}^{\bullet,\mathbf{V}}[\Phi] \\ &= \sum_{n \in \mathcal{E}_l^\mathbf{V}} \mathbb{M}_l(N^\mathbf{V} = n) \mathbb{W}_{l,n}^{\bullet,\mathbf{V}}[\Phi]. \end{aligned} \quad (34)$$

At this point, recall from Lemma 3 that $\mathcal{E}_l^\mathbf{V} = R_l^\mathbf{V} \cup (\beta^\mathbf{V}l + h^\mathbf{V}\mathbb{Z}_+)$, where $\beta^\mathbf{V} \geq 1$ and $R_l^\mathbf{V} \subset \{0, 1, \dots, \beta^\mathbf{V}l - 1\}$. For simplicity, let us use the notation $\beta = \beta^\mathbf{V}$ and $h = h^\mathbf{V}$. Therefore,

$$\begin{aligned} \mathbb{W}_l^\bullet[\Phi] &= \sum_{n \geq 0} \mathbb{M}_l(N^\mathbf{V} = \beta l + hn) \mathbb{W}_{l,\beta l + hn}^{\bullet,\mathbf{V}}[\Phi] + \mathcal{R}_l \\ &= \int_{\mathbb{R}_+} l^2 Q^\mathbf{V}(l, \beta l + h[l^2 A]) \mathbb{W}_{l,\beta l + h[l^2 A]}^{\bullet,\mathbf{V}}[\Phi] dA + \mathcal{R}_l \end{aligned}$$

where $|\mathcal{R}_l| \leq \|\Phi\|_\infty \mathbb{W}_l^\bullet(|\mathbf{V}| \leq \beta l)$. Recall that, under \mathbb{W}_l^\bullet , the random variable $|\mathbf{V}| - 1$ has same distribution as a sum of l i.i.d. random variables with distribution $Q^\mathbf{V}(1, \cdot)$. The proof of Lemma 20 yields that, under \mathbb{W}_l^\bullet

$$\frac{|\mathbf{V}|}{l^2} \xrightarrow[l \rightarrow \infty]{(d)} \frac{1}{\sigma_\mathbf{V}^2} \mathcal{A}^\bullet$$

where \mathcal{A}^\bullet is a stable random variable with density j_1 . Clearly, this implies that $\mathcal{R}_l \rightarrow 0$ as $l \rightarrow \infty$. Now assume that $\Phi = \varphi((2l/3)^{-1/2}M)$ where φ is a continuous and bounded function on the Gromov–Hausdorff space. Then one has, by Theorem 25,

$$\mathbb{W}_{l,\beta l + h[l^2 A]}^{\bullet,\mathbf{V}}[\varphi((2l/3)^{-1/2}M)] \xrightarrow[l \rightarrow \infty]{} \mathbb{E}\left[\varphi((h\sigma_\mathbf{V}^2 A)^{1/4} \text{BD}_{(h\sigma_\mathbf{V}^2 A)^{-1/2}})\right] = \mathbb{E}[\varphi(\text{BD}_{1,h\sigma_\mathbf{V}^2 A})],$$

where the last equality follows from Remark 3. At this point, we apply Lemma 20, which implies that for every $A > 0$,

$$l^2 Q^\mathbf{V}(l, \beta l + h[l^2 A]) \xrightarrow[l \rightarrow \infty]{} h j_{1/\sigma_\mathbf{V}}(hA) = h\sigma_\mathbf{V}^2 j_1(h\sigma_\mathbf{V}^2 A).$$

Since we are dealing with probability densities, the Scheffé Lemma implies that the latter convergence holds in fact in $L^1(dA)$, and we conclude that

$$\begin{aligned} \lim_{l \rightarrow \infty} \mathbb{W}_l^\bullet[\varphi((2l/3)^{-1/2}M)] &= \int_{\mathbb{R}_+} h\sigma_\mathbf{V}^2 j_1(h\sigma_\mathbf{V}^2 A) \mathbb{E}[\varphi(\text{BD}_{1,h\sigma_\mathbf{V}^2 A})] dA \\ &= \int_{\mathbb{R}_+} j_1(A) \mathbb{E}[\varphi(\text{BD}_{1,A})] dA \end{aligned}$$

and this is equal to $\mathbb{E}[\varphi(\text{FBD}_1^\bullet)]$. The second part of Theorem 8 follows.

To obtain the result under \mathbb{W}_l instead of \mathbb{W}_l^\bullet , note that (5) implies

$$\mathbb{W}_l(|\mathbf{V}| = n + 1) = K_l \frac{\mathbb{W}_l^\bullet(|\mathbf{V}| = n + 1)}{n + 1} \quad (35)$$

where $K_l = \mathbb{W}_l^\bullet[1/|\mathbf{V}|]^{-1}$. We then use the following lemma, which is certainly known, but for which we did not find a proper reference.

Lemma 29. *Let X_1, X_2, \dots be a sequence of i.i.d. r.v.s with values in $\{1, 2, 3, \dots\}$, and such that*

$$\mathbb{P}(X_1 > k) \underset{k \rightarrow \infty}{\sim} \frac{c}{k^\alpha}$$

for some constants $c \in (0, \infty)$ and $\alpha \in (0, 1)$. Then

$$\mathbb{E} \left[\frac{l^{1/\alpha}}{X_1 + \dots + X_l} \right] \xrightarrow{l \rightarrow \infty} \mathbb{E} \left[\frac{1}{S} \right]$$

where S is the limit in distribution of $(X_1 + \dots + X_l)/l^{1/\alpha}$ as $l \rightarrow \infty$ (so that S is a stable distribution of index α).

Proof. By hypothesis and standard facts on stable domains of attractions [13, Chapter 8], our hypotheses imply that as $s \uparrow 1$,

$$\mathbb{E}[s^{X_1}] = 1 - c'(1 - s)^\alpha(1 + o(1))$$

for some constant $c' \in (0, \infty)$ depending only on c and α . Applying this to $s = \exp(-\lambda)$ for $\lambda \geq 0$ implies that

$$\mathbb{E}[\exp(-\lambda X_1)] = 1 - c'\lambda^\alpha(1 + o(1))$$

as $\lambda \downarrow 0$, so that there exists $c'' \in (0, \infty)$ such that for every $\lambda \in [0, 1]$, one has

$$\begin{aligned} \mathbb{E}[\exp(-\lambda X_1)] &\leq 1 - c''\lambda^\alpha \\ &\leq \exp(-c''\lambda^\alpha). \end{aligned} \quad (36)$$

On the other hand, the assumption that $X_1 \geq 1$ a.s. implies that $\mathbb{E}[\exp(-\lambda X_1)] \leq \exp(-\lambda)$ for every $\lambda \geq 0$. This implies that, possibly by choosing c'' smaller, one can assume that (36) is valid for every $\lambda \geq 0$, as we supposed $\alpha \leq 1$.

Now note that, for every $x > 0$, one has (using the inequality $\mathbf{1}_{[0,1]}(u) \leq e \exp(-u)$ in the first step)

$$\begin{aligned} \mathbb{P}(X_1 + \dots + X_l \leq xl^{1/\alpha}) &\leq e \mathbb{E} \left[\exp \left(- (X_1 + \dots + X_l) / (xl^{1/\alpha}) \right) \right] \\ &\leq e \mathbb{E} \left[\exp \left(- X_1 / (xl^{1/\alpha}) \right) \right]^l \\ &\leq e \exp(-c''x^{-\alpha}), \end{aligned}$$

where we used the version of (36) valid for all λ at the last step. This stretched-exponential tail bound is uniform in l and clearly implies the convergence of all negative moments. \square

Since, as we observed, $|\mathbf{V}| - 1$ under \mathbb{W}_l^\bullet is distributed as a sum of l i.i.d. random variables satisfying the hypotheses of Lemma 29 with $\alpha = 1/2$ (by Lemma 20), this entails that

$$\lim_{l \rightarrow \infty} \frac{K_l}{l^2} = \mathbb{E} \left[\frac{\sigma_{\mathbf{V}^2}}{\mathcal{A}^\bullet} \right]^{-1} = \frac{1}{\sigma_{\mathbf{V}^2}}.$$

Repeating the previous argument, only changing $\mathbb{W}_l^\bullet(|\mathbf{V}| = n + 1)$ by $\mathbb{W}_l(|\mathbf{V}| = n + 1)$ in (34) and applying (35), and then performing the same steps using the equivalent we obtained for K_l , we obtain

$$\lim_{l \rightarrow \infty} \mathbb{W}_l [\varphi((2l/3)^{-1/2}M)] = \int_{\mathbb{R}_+} \frac{h\sigma_{\mathbf{V}^2}}{h\sigma_{\mathbf{V}^2}A} j_1(h\sigma_{\mathbf{V}^2}A) \mathbb{E}[\varphi(\text{BD}_{1,h\sigma_{\mathbf{V}^2}A})] dA$$

and this is $\mathbb{E}[\varphi(\text{FBD}_1)]$, as wanted.

References

- [1] C. Abraham. Rescaled bipartite planar maps converge to the Brownian map. *Preprint*, [arXiv:1312.5959](https://arxiv.org/abs/1312.5959), 2013.
- [2] L. Addario-Berry and M. Albenque. The scaling limit of random simple triangulations and random simple quadrangulations. *Preprint*, [arXiv:1306.5227](https://arxiv.org/abs/1306.5227), 2013.
- [3] M. Albenque and J.-F. Marckert. Some families of increasing planar maps. *Electron. J. Probab.*, 13:no. 56, 1624–1671, 2008.
- [4] D. J. Aldous. The continuum random tree. I. *Ann. Probab.*, 19(1):1–28, 1991.
- [5] D. J. Aldous. The continuum random tree. III. *Ann. Probab.*, 21(1):248–289, 1993.
- [6] J. Beltran and J.-F. Le Gall. Quadrangulations with no pendant vertices. *Bernoulli*, 19(4):1150–1175, 2013.
- [7] J. Bettinelli. Scaling limits for random quadrangulations of positive genus. *Electron. J. Probab.*, 15:no. 52, 1594–1644, 2010.
- [8] J. Bettinelli. The topology of scaling limits of positive genus random quadrangulations. *Ann. Probab.*, 40:no. 5, 1897–1944, 2012.
- [9] J. Bettinelli. Geodesics in Brownian surfaces (Brownian maps). [arXiv:1401.3602](https://arxiv.org/abs/1401.3602), to appear in *Ann. Inst. Henri Poincaré Probab. Stat.*, 2014.
- [10] J. Bettinelli. Scaling limit of random planar quadrangulations with a boundary. *Ann. Inst. Henri Poincaré Probab. Stat.*, 51(2):432–477, 2015.
- [11] J. Bettinelli, E. Jacob, and G. Miermont. The scaling limit of uniform random plane maps, *via* the Ambjørn–Budd bijection. *Electron. J. Probab.*, 19:no. 74, 1–16, 2014.

- [12] J. Bettinelli and G. Miermont. Compact Brownian surfaces II. The general case. *in preparation*, 2015.
- [13] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [14] J. Bouttier, P. Di Francesco, and E. Guitter. Planar maps as labeled mobiles. *Electron. J. Combin.*, 11:Research Paper 69, 27 pp. (electronic), 2004.
- [15] J. Bouttier and E. Guitter. Distance statistics in quadrangulations with a boundary, or with a self-avoiding loop. *J. Phys. A*, 42(46):465208, 44, 2009.
- [16] J. Bouttier and E. Guitter. Planar maps and continued fractions. *Comm. Math. Phys.*, 309(3):623–662, 2012.
- [17] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
- [18] L. Chaumont and J. C. Pardo. On the genealogy of conditioned stable Lévy forests. *ALEA Lat. Am. J. Probab. Math. Stat.*, 6:261–279, 2009.
- [19] R. Cori and B. Vauquelin. Planar maps are well labeled trees. *Canad. J. Math.*, 33(5):1023–1042, 1981.
- [20] N. Curien and J.-F. Le Gall. The Brownian plane. *Journal of Theoretical Probability*, pages 1–43, 2012.
- [21] T. Duquesne and J.-F. Le Gall. Random trees, Lévy processes and spatial branching processes. *Astérisque*, 281:vi+147, 2002.
- [22] M. Gromov. *Metric structures for Riemannian and non-Riemannian spaces*, volume 152 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1999.
- [23] J.-F. Le Gall. *Spatial branching processes, random snakes and partial differential equations*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1999.
- [24] J.-F. Le Gall. The topological structure of scaling limits of large planar maps. *Invent. Math.*, 169(3):621–670, 2007.
- [25] J.-F. Le Gall. Geodesics in large planar maps and in the Brownian map. *Acta Math.*, 205(2):287–360, 2010.
- [26] J.-F. Le Gall. Uniqueness and universality of the Brownian map. *Ann. Probab.*, 41(4):2880–2960, 2013.

- [27] J.-F. Le Gall and G. Miermont. Scaling limits of random planar maps with large faces. *Ann. Probab.*, 39(1):1–69, 2011.
- [28] J.-F. Le Gall and F. Paulin. Scaling limits of bipartite planar maps are homeomorphic to the 2-sphere. *Geom. Funct. Anal.*, 18(3):893–918, 2008.
- [29] J.-F. Le Gall and M. Weill. Conditioned Brownian trees. *Ann. Inst. H. Poincaré Probab. Statist.*, 42(4):455–489, 2006.
- [30] J.-F. Marckert and G. Miermont. Invariance principles for random bipartite planar maps. *Ann. Probab.*, 35(5):1642–1705, 2007.
- [31] J.-F. Marckert and A. Mokkadem. Limit of normalized random quadrangulations: the Brownian map. *Ann. Probab.*, 34(6):2144–2202, 2006.
- [32] G. Miermont. Invariance principles for spatial multitype Galton–Watson trees. *Ann. Inst. H. Poincaré Probab. Statist.*, 44(6):1128–1161, 2008.
- [33] G. Miermont. On the sphericity of scaling limits of random planar quadrangulations. *Electron. Commun. Probab.*, 13:248–257, 2008.
- [34] G. Miermont. The Brownian map is the scaling limit of uniform random plane quadrangulations. *Acta Math.*, 210(2):319–401, 2013.
- [35] J. Miller and S. Sheffield. An axiomatic characterization of the Brownian map. *Preprint*, [arXiv:1506.03806](https://arxiv.org/abs/1506.03806), 2015.
- [36] J. Pitman. *Combinatorial stochastic processes*, volume 1875 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2006. Lectures from the 32nd Summer School on Probability Theory held in Saint-Flour, July 7–24, 2002, With a foreword by Jean Picard.
- [37] G. Schaeffer. *Conjugaison d’arbres et cartes combinatoires aléatoires*. PhD thesis, Université Bordeaux I, 1998.
- [38] R. Stephenson. Local convergence of large critical multi-type Galton–Watson trees and applications to random maps. *Preprint*, [arXiv:1412.6911](https://arxiv.org/abs/1412.6911), 2014.