



HAL
open science

A Hybrid High-Order method for Leray–Lions elliptic equations on general meshes

Daniele Antonio Di Pietro, Jérôme Droniou

► **To cite this version:**

Daniele Antonio Di Pietro, Jérôme Droniou. A Hybrid High-Order method for Leray–Lions elliptic equations on general meshes. 2015. hal-01183484v1

HAL Id: hal-01183484

<https://hal.science/hal-01183484v1>

Preprint submitted on 8 Aug 2015 (v1), last revised 7 Dec 2016 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Hybrid High-Order method for Leray–Lions elliptic equations on general meshes

Daniele A. Di Pietro^{*1} and Jérôme Droniou^{†2}

¹University of Montpellier, Institut Montpellierain Alexander Grothendieck, 34095 Montpellier, France

²School of Mathematical Sciences, Monash University, Clayton, Victoria 3800, Australia

August 8, 2015

Abstract

In this work, we develop and analyze a Hybrid High-Order (HHO) method for steady non-linear Leray–Lions problems. The proposed method has several assets, including the support for arbitrary approximation orders and general polytopal meshes. This is achieved by combining two key ingredients devised at the local level: a gradient reconstruction and a high-order stabilization term that generalizes the one originally introduced in the linear case. The convergence analysis is carried out using a compactness technique. Extending this technique to HHO methods has prompted us to develop a set of discrete functional analysis tools whose interest goes beyond the specific problem and method addressed in this work: (direct and) reverse Lebesgue and Sobolev embeddings for local polynomial spaces, L^p -stability and $W^{s,p}$ -approximation properties for L^2 -projectors on such spaces, and Sobolev embeddings for hybrid polynomial spaces. Numerical tests are presented to validate the theoretical results for the original method and variants thereof.

2010 Mathematics Subject Classification: 65N08, 65N30, 65N12

Keywords: Hybrid High-Order methods, nonlinear elliptic equations, p -Laplacian, discrete functional analysis, convergence analysis, $W^{s,p}$ -stability of L^2 -projection on polynomials

1 Introduction

We are interested here in the numerical approximation of the steady Leray–Lions equation

$$-\operatorname{div}(\mathbf{a}(\cdot, u, \nabla u)) = f \quad \text{in } \Omega, \quad (1.1a)$$

$$u = 0 \quad \text{on } \partial\Omega, \quad (1.1b)$$

where $\Omega \subset \mathbb{R}^d$, $d \geq 1$, is a polytopal bounded connected domain of boundary $\partial\Omega$, while $\mathbf{a} : \Omega \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a (possibly nonlinear) function of its arguments, for which detailed assumptions are discussed in the following section. This model, which contains the p -Laplace equation, appears in the modelling of glacier motion [42], of incompressible turbulent flows in porous media [31] and in airfoil design [41]. Our goal is to design and analyze a discretization method for problem (1.1) inspired by the Hybrid High-Order (HHO) method introduced in [29] in the context of a linear diffusion model problem (see also [26] for degenerate advection–diffusion–reaction models). The proposed method offers several assets: (i) the construction is dimension-independent; (ii) fairly general meshes including polytopal elements and nonmatching interfaces are supported; (iii) arbitrary polynomial orders can be considered (including the case $k = 0$); (iv) it is efficiently

*daniele.di-pietro@umontpellier.fr

†jerome.droniou@monash.edu

parallelisable (the local stencil only connects a mesh element with its faces), and it has reduced computational cost (when solving by a first-order algorithm, the element-based unknowns can be eliminated by static condensation).

Numerical methods allowing for arbitrary-order discretizations and general meshes have received increasing attention over the last few years. Supporting general polytopal meshes is required, e.g., in the modelling of underground flows, where degenerate elements and nonconforming interfaces account for complex geometric features resulting from compaction, erosion, and the onset of fractures or faults. Another relevant application of polyhedral meshes is adaptive mesh coarsening [6, 11]. The literature on arbitrary-order polytopal methods for linear diffusion problems is vast. In this context, methods that have similarities (and differences) with the HHO method include, e.g., the Hybridizable Discontinuous Galerkin method of [23] (cf. also [22] for a precise study of its relation with the HHO method), the Virtual Element Method of [12, 13, 17], the High-Order Mimetic method of [47], the Weak Galerkin method of [50, 51], and the Multiscale Hybrid-Mixed method of [7].

The finite element approximation of nonlinear diffusion problems of Leray–Lions type on standard meshes has been studied in several papers; cf., e.g., [10, 42, 48]. The literature on polytopal meshes is, however, much more scarce, and is mainly restricted to the lowest-order case. We cite here, in particular, the two-dimensional Discrete Duality Finite Volume schemes studied in [4] (cf. also the precursor papers [1–3]), the Mixed Finite Volume scheme of [32] (inspired by [33]) valid in arbitrary space dimension, and the Mimetic Finite Difference method of [5] for $p \in (1, 2)$ and under more restrictive assumptions than (2.2). High-order discontinuous Galerkin approximations have also been considered in [20].

The starting point for the present work is the HHO method of [29]. In the lowest-order case, it has been shown in [29, Section 2.5] that this method belongs to the Hybrid Mixed Mimetic family [36], which includes the hybrid Mimetic Finite Differences [18], the Hybrid Finite Volume [39] and the Mixed Finite Volume [33]. The HHO method can therefore be seen as a higher order version of these schemes. The (hybrid) degrees of freedom (DOFs) for the HHO method are fully discontinuous polynomials of degree $k \geq 0$ at mesh elements and faces. The construction hinges on two key ingredients built element-wise: (i) a discrete gradient defined from element- and face-based DOFs; (ii) a high-order penalty term which vanishes whenever one of its arguments is a polynomial of degree $\leq (k + 1)$ inside the element. These ingredients are combined to build a local contribution, which is then assembled element-wise. A key feature reducing the computational cost is that only face-based DOFs are globally coupled, whereas element-based DOFs can be locally eliminated by a standard static condensation procedure.

The design of a HHO method for the nonlinear problem (1.1) entails several new ideas. A first difference with respect to the linear case is that a more natural choice is to seek the gradient reconstruction in the full space of vector-valued polynomials of degree $\leq k$ (as opposed to the space spanned by gradients of scalar-valued polynomials of degree $\leq (k + 1)$). The main consequence of this choice is that, when applied to the interpolates of smooth functions, the discrete gradient operator commutes with the L^2 -projector, and therefore enjoys L^p -stability properties (see below). A second important point is the design of a high-order stabilization term with appropriate scaling. Here, we propose a generalization of the stabilization term of [29] which preserves the property of vanishing whenever one of its arguments is a polynomial of degree $\leq (k + 1)$. As in the linear case, the construction hinges on the solution of small local linear problems inside each elements, and the possibility of statically condense element-based DOFs remains available.

The convergence analysis is carried out using a compactness argument in the spirit of [49], and under minimal regularity assumptions on the exact solution. Adapting the compactness argument has prompted us to develop discrete functional analysis tools whose interest goes beyond the specific method and problem considered in this work.

A first notable set of result are (direct and) reverse Lebesgue and Sobolev embeddings on local polynomial spaces (e.g., on mesh elements and faces, but curved geometries are also allowed). The term reverse refers to the fact that the *largest* exponent is bounded above by the term corresponding to the *lowest* exponent. Direct Sobolev embedding for broken spaces on fairly

general polytopal meshes are proved in [19, 27]; specific instances had already been established in [8, 15, 40, 44, 45]. Reverse embeddings, on the other hand, are established in [16, Theorem 4.5.11], but under the assumption that all mesh elements are affine-equivalent to one (or a finite number of) given fixed reference elements. This limitation is due to the very generic local finite element spaces considered therein. Exploiting the fact that we deal with *polynomial* local spaces, we can establish a more general version of reverse inequalities, that does not require to specify any particular geometry of the elements (only their non-degeneracy). Reverse Lebesgue embeddings are a crucial ingredient to prove the stability of the HHO method.

A second set of results concerns the stability and approximation properties of the L^2 -projector on local polynomial spaces. More specifically, we prove under very general geometric assumptions that the L^2 -projector is L^p -stable for any index $p \in [1, +\infty]$, and that it has optimal approximation properties in local polynomial spaces. Stability results for (global) projectors onto finite element spaces can be found in [9, 14, 21, 24]. However, these references mostly consider H^1 -stability, and assume quite restrictive (and sometimes difficult to check) geometrical assumptions on the meshes. These limitations are a consequence of dealing with projectors on global finite element spaces, that include some form of continuity property between the mesh elements. On discontinuous polynomial spaces such as the ones used in HHO methods, we can establish more general L^p - and $W^{s,p}$ -stability and approximation properties of local L^2 -projectors. The approximation results extend to the $W^{s,p}$ -setting the ones in [28, Section 1.4.4], based in turn on the ideas of [38].

Finally, a third set of discrete functional analysis tools are specific to polynomial spaces with a hybrid structure, i.e., using as DOFs polynomials at elements and faces. In this case, building on the results of [27] for discontinuous Galerkin methods (inspired by the low-order discrete functional analysis results of [33, 39]), we introduce a suitable discrete $W^{1,p}$ -like norm and prove a discrete counterpart of Sobolev embeddings and a compactness result for the discrete gradient reconstruction upon which the HHO method hinges.

The material is organized as follows: in Section 2 we recall a set of standard assumptions to write a weak formulation for problem (1.1); in Section 3 we detail the discrete setting by specifying the assumptions on the mesh and recalling the basic results on local polynomial spaces; in Section 4 we formulate the HHO method, state (without proof) the main stability and convergence results, and provide a few numerical examples; Section 5 collects the discrete functional analysis tools on hybrid polynomial spaces, which are used in Section 6 to prove the stability and convergence of the HHO method; in Section 7 we briefly address the treatment of other boundary conditions and hint at the modifications required in the analysis; a conclusion is given in Section 8 and, finally, in Appendix A we provide the proofs of the discrete functional analysis results on local polynomial spaces.

2 Continuous setting

In this section we detail the assumptions on the function \mathbf{a} and write a weak formulation for problem (1.1). Let $p \in (1, +\infty)$ be given, and denote by $p' := \frac{p}{p-1}$ the dual exponent of p , and by p^* the Sobolev exponent of p such that

$$p^* = \begin{cases} \frac{dp}{d-p} & \text{if } p < d, \\ +\infty & \text{if } p \geq d. \end{cases} \quad (2.1)$$

We assume that

$$\mathbf{a} : \Omega \times \mathbb{R} \times \mathbb{R}^d \mapsto \mathbb{R}^d \text{ is a Caratheodory function,} \quad (2.2a)$$

$$\begin{aligned} \exists \bar{a} \in L^{p'}(\Omega), \exists \beta_{\mathbf{a}} \in (0, +\infty), \exists r < \frac{p^*}{p'} : |\mathbf{a}(\mathbf{x}, s, \boldsymbol{\xi})| \leq \bar{a}(\mathbf{x}) + \beta_{\mathbf{a}}|s|^r + \beta_{\mathbf{a}}|\boldsymbol{\xi}|^{p-1} \\ \text{for a.e. } \mathbf{x} \in \Omega, \text{ for all } (s, \boldsymbol{\xi}) \in \mathbb{R} \times \mathbb{R}^d, \end{aligned} \quad (2.2b)$$

$$[\mathbf{a}(\mathbf{x}, s, \boldsymbol{\xi}) - \mathbf{a}(\mathbf{x}, s, \boldsymbol{\eta})] \cdot [\boldsymbol{\xi} - \boldsymbol{\eta}] \geq 0 \text{ for a.e. } \mathbf{x} \in \Omega, \text{ for all } (s, \boldsymbol{\xi}, \boldsymbol{\eta}) \in \mathbb{R} \times \mathbb{R}^d \times \mathbb{R}^d, \quad (2.2c)$$

$$\exists \lambda_{\mathbf{a}} \in (0, +\infty) : \mathbf{a}(\mathbf{x}, s, \boldsymbol{\xi}) \cdot \boldsymbol{\xi} \geq \lambda_{\mathbf{a}}|\boldsymbol{\xi}|^p \text{ for a.e. } \mathbf{x} \in \Omega, \text{ for all } (s, \boldsymbol{\xi}) \in \mathbb{R} \times \mathbb{R}^d, \quad (2.2d)$$

$$f \in L^{p'}(\Omega). \quad (2.2e)$$

Here, Carathéodory function means that $\mathbf{a}(\mathbf{x}, \cdot, \cdot)$ is continuous on $\mathbb{R} \times \mathbb{R}^d$ for a.e. $\mathbf{x} \in \Omega$, and $\mathbf{a}(\cdot, s, \boldsymbol{\xi})$ is measurable on Ω for all $(s, \boldsymbol{\xi}) \in \mathbb{R} \times \mathbb{R}^d$. The Euclidean dot product and norm in \mathbb{R}^d are denoted by $\mathbf{x} \cdot \mathbf{y}$ and $|\mathbf{x}|$, respectively. Classically [46], the weak formulation for (1.1) is

$$\begin{aligned} &\text{Find } u \in W_0^{1,p}(\Omega) \text{ such that, for all } v \in W_0^{1,p}(\Omega), \\ &\int_{\Omega} \mathbf{a}(\mathbf{x}, u(\mathbf{x}), \nabla u(\mathbf{x})) \cdot \nabla v(\mathbf{x}) \, d\mathbf{x} = \int_{\Omega} f(\mathbf{x})v(\mathbf{x}) \, d\mathbf{x}. \end{aligned} \quad (2.3)$$

3 Discrete setting

This section presents the discrete setting: admissible mesh sequences, analysis tools on such meshes, DOFs, reduction maps, and reconstruction operators.

3.1 Assumptions on the mesh

Denote by $\mathcal{H} \subset \mathbb{R}_*^+$ a countable set of meshsizes having 0 as its unique accumulation point. Following [28, Chapter 4], we consider h -refined mesh sequences $(\mathcal{T}_h)_{h \in \mathcal{H}}$ where, for all $h \in \mathcal{H}$, \mathcal{T}_h is a finite collection of nonempty disjoint open polyhedral elements T such that $\bar{\Omega} = \bigcup_{T \in \mathcal{T}_h} \bar{T}$ and $h = \max_{T \in \mathcal{T}_h} h_T$ with h_T standing for the diameter of the element T . A face F is defined as a hyperplanar closed connected subset of $\bar{\Omega}$ with positive $(d-1)$ -dimensional Hausdorff measure and such that (i) either there exist $T_1, T_2 \in \mathcal{T}_h$ such that $F \subset \partial T_1 \cap \partial T_2$ and F is called an interface or (ii) there exists $T \in \mathcal{T}_h$ such that $F \subset \partial T \cap \partial \Omega$ and F is called a boundary face. Interfaces are collected in the set \mathcal{F}_h^i , boundary faces in \mathcal{F}_h^b , and we let $\mathcal{F}_h := \mathcal{F}_h^i \cup \mathcal{F}_h^b$. The diameter of a face $F \in \mathcal{F}_h$ is denoted by h_F . For all $T \in \mathcal{T}_h$, $\mathcal{F}_T := \{F \in \mathcal{F}_h \mid F \subset \partial T\}$ denotes the set of faces contained in ∂T (with ∂T denoting the boundary of T) and, for all $F \in \mathcal{F}_T$, \mathbf{n}_{TF} is the unit normal to F pointing out of T . Symmetrically, for all $F \in \mathcal{F}_h$, we let $\mathcal{T}_F := \{T \in \mathcal{T}_h \mid F \subset \partial T\}$ the set of elements having F as a face.

Our analysis hinges on the following assumption on the mesh sequence.

Assumption 3.1 (Admissible mesh sequence). *For all $h \in \mathcal{H}$, \mathcal{T}_h admits a matching simplicial submesh \mathfrak{T}_h and there exists a real number $\varrho > 0$ such that, for all $h \in \mathcal{H}$: (i) for all simplex $S \in \mathfrak{T}_h$ of diameter h_S and inradius r_S , $\varrho h_S \leq r_S$, and (ii) for all $T \in \mathcal{T}_h$, and all $S \in \mathfrak{T}_h$ such that $S \subset T$, $\varrho h_T \leq h_S$.*

The simplicial submesh in this assumption is just a theoretical tool, and it is not used in the actual construction of the discretization method. Given an admissible mesh sequence, for all $h \in \mathcal{H}$, all $T \in \mathcal{T}_h$, and all $F \in \mathcal{F}_T$, h_F is uniformly comparable to h_T in the sense that (cf. [28, Lemma 1.42]):

$$\varrho^2 h_T \leq h_F \leq h_T. \quad (3.1)$$

Moreover, [28, Lemma 1.41] shows that there exists an integer N_ϱ depending on ϱ such that

$$\forall h \in \mathcal{H} : \max_{T \in \mathcal{T}_h} \text{card}(\mathcal{F}_T) \leq N_\varrho. \quad (3.2)$$

Finally, by [28, Lemma 1.40], there is an integer N_s depending on ϱ such that

$$\forall h \in \mathcal{H} : \max_{T \in \mathcal{T}_h} \text{card}(\{S \in \mathfrak{T}_h \mid S \subset T\}) \leq N_s. \quad (3.3)$$

3.2 Basic results on local polynomial spaces

The building blocks for the HHO method are local polynomial spaces on elements and faces. Let an integer $l \geq 0$ be fixed. Let U be a subset of \mathbb{R}^N (for some $N \geq 1$), H_U the affine space spanned by U , d_U its dimension, and assume that U has a non-empty interior in H_U . We denote by $\mathbb{P}^l(U)$ the space spanned by d_U -variate polynomials on H_U of total degree $\leq l$. In the following sections,

we will typically have $N = d$ and the set U will represent a mesh element or face. We note, in passing, that a subset U with curved boundaries is also allowed except in Lemma 3.6, which is why we use the different notation T instead of U in this lemma.

A key element in the construction are L^2 -projectors onto local polynomial spaces on bounded subsets $U \subset \mathbb{R}^N$. The L^2 -projector $\pi_U^l : L^1(U) \mapsto \mathbb{P}^l(U)$ is defined as follows: For any $w \in L^1(U)$, $\pi_U^l w$ is the unique element of $\mathbb{P}^l(U)$ such that

$$\forall v \in \mathbb{P}^l(U) : \int_U \pi_U^l w(\mathbf{x})v(\mathbf{x})d\mathbf{x} = \int_U w(\mathbf{x})v(\mathbf{x})d\mathbf{x}. \quad (3.4)$$

Note that the regularity $w \in L^1(U)$ suffices to integrate w against polynomials on U (which are bounded functions). In what follows, we state some stability and approximation properties for the L^2 -projector. The proofs are postponed to Appendix A.2.

Lemma 3.2 (L^p -stability of L^2 -projectors on polynomial spaces). *Let U be a measurable subset of \mathbb{R}^N , with inradius r_U and diameter h_U , such that*

$$\frac{r_U}{h_U} \geq \delta > 0. \quad (3.5)$$

Let $k \in \mathbb{N}$ and $p \in [1, +\infty]$. Then, there exists C only depending on N , δ , k and p such that

$$\forall g \in L^p(U) : \|\pi_U^k g\|_{L^p(U)} \leq C \|g\|_{L^p(U)}. \quad (3.6)$$

Remark 3.3 (Geometric regularity (3.5) for mesh elements and faces). *Elements $T \in \mathcal{T}_h$ and faces $F \in \mathcal{F}_h$ of an admissible mesh sequence satisfy the geometric regularity assumption (3.5) with $\delta = \varrho^2$ and $\delta = \varrho$ respectively.*

In the case where $W^{s,p}(U)$ is continuously embedded in $C(\overline{U})$, the following result can be found in [16, Theorem 4.4.4]. This restriction on the space $W^{s,p}$, which would prevent us from analyzing interesting cases for (1.1), is due to the very general setting chosen for analyzing the interpolation error in [16]. Because we focus here on local polynomial spaces and L^2 -projectors, we can improve this result and obtain optimal interpolation errors for any s, p . If U is an open set of \mathbb{R}^N , $s \in \mathbb{N}$ and $p \in [1, +\infty]$, we recall that $|\cdot|_{W^{s,p}(U)}$ is defined by

$$\forall v \in W^{s,p}(U), \quad |v|_{W^{s,p}(U)} := \sum_{\alpha \in \mathbb{N}^N, |\alpha|_{\ell^1} = s} \|\partial^\alpha v\|_{L^p(U)}$$

where $|\alpha|_{\ell^1} = \alpha_1 + \dots + \alpha_N$ and $\partial^\alpha = \partial_1^{\alpha_1} \dots \partial_N^{\alpha_N}$.

Lemma 3.4 ($W^{s,p}$ -approximation properties of L^2 -projectors on polynomial spaces). *Let U be an open subset of \mathbb{R}^N with diameter h_U , such that U is star-shaped with respect to a ball of radius ρh_U for some $\rho > 0$. Let $k \in \mathbb{N}$, $s \in \{1, \dots, k+1\}$ and $p \in [1, +\infty]$. Then, there exists C only depending on N , ρ , k , s and p such that*

$$\forall m \in \{0, \dots, s\}, \forall v \in W^{s,p}(U) : |v - \pi_U^k v|_{W^{m,p}(U)} \leq C h_U^{s-m} |v|_{W^{s,p}(U)}. \quad (3.7)$$

Remark 3.5. *Using [38, Section 7], the result still holds if U is a finite union of domains that are star-shaped with respect to balls of radius comparable to h_U . This enables us to use Lemma 3.4 on elements of admissible mesh sequences, which are the union of a finite number of simplices; cf. (3.3).*

The next result estimates the trace of the error, and therefore requires more geometric assumptions on the domain (which, in the following sections, will be invariably a mesh element T).




Figure 1: Degrees of freedom for $k \in \{0, 1, 2\}$. Shaded DOFs can be locally eliminated by static condensation.

Lemma 3.6 (Approximation properties of traces of L^2 -projectors on polynomial spaces). *Let T be a polyhedral subset of \mathbb{R}^N with diameter h_T , such that T is the union of disjoint simplices S of diameter h_S and inradius r_S such that $\varrho^2 h_T \leq \varrho h_S \leq r_S$ for some $\varrho > 0$. Let $k \in \mathbb{N}$, $s \in \{1, \dots, k+1\}$ and $p \in [1, +\infty]$. Then, there exists C only depending on N , ϱ , k , s and p such that*

$$\forall m \in \{0, \dots, s-1\}, \forall v \in W^{s,p}(T) : h_T^{\frac{1}{p}} |v - \pi_T^k v|_{W^{m,p}(\mathcal{F}_T)} \leq C h_T^{s-m} |v|_{W^{s,p}(T)}. \quad (3.8)$$

Here, $W^{m,p}(\mathcal{F}_T)$ is the set of functions that belong to $W^{m,p}(F)$ for any hyperplanar face F of T , with corresponding broken norm.

Finally, the triangle inequality applied to (3.7) (with $m = s$) and to (3.8) (with $m = s-1$) immediately gives the following extension of Lemma 3.2.

Corollary 3.7 ($W^{s,p}$ -stability of L^2 -projectors on polynomial spaces). *The following holds:*

(i) *Under the assumptions of Lemma 3.4, we have, with C only depending on N , ρ , k , s and p ,*

$$\forall v \in W^{s,p}(U) : |\pi_U^k v|_{W^{s,p}(U)} \leq C |v|_{W^{s,p}(U)};$$

(ii) *Under the assumptions of Lemma 3.6, we have with C only depending on N , ϱ , k , s and p ,*

$$\forall v \in W^{s,p}(T) : |\pi_T^k v|_{W^{s-1,p}(\mathcal{F}_T)} \leq C h_T^{\frac{1}{p}} |v|_{W^{s,p}(T)} + |v|_{W^{s-1,p}(\mathcal{F}_T)}.$$

4 The Hybrid High-Order method

In this section we introduce the space of degrees of freedom, define the gradient and potential reconstructions at the heart of the HHO method, state the discrete problem along with the main stability and convergence results, and provide some numerical examples.

4.1 Local degrees of freedom, interpolation and reconstructions

Let a polynomial degree $k \geq 0$ and an element $T \in \mathcal{T}_h$ be fixed. We define the local space of DOFs

$$\underline{U}_T^k := \mathbb{P}^k(T) \times \left(\prod_{F \in \mathcal{F}_T} \mathbb{P}^k(F) \right), \quad (4.1)$$

cf. Figure 1, and we use the underline notation $\underline{v}_T = (v_T, (v_F)_{F \in \mathcal{F}_T})$ for a generic element $\underline{v}_T \in \underline{U}_T^k$. We define the local interpolation operator $\underline{I}_T^k : W^{1,1}(T) \rightarrow \underline{U}_T^k$ such that, for all $v \in W^{1,1}(T)$,

$$\underline{I}_T^k v := (\pi_T^k v, (\pi_F^k v)_{F \in \mathcal{F}_T}). \quad (4.2)$$

Remark 4.1 (Domain for the interpolation operator). *The local interpolation operator is well-defined for functions $v \in W^{1,1}(T)$ since v is clearly in $L^1(T)$, the domain of π_T^k , and its trace on every face $F \in \mathcal{F}_T$ is in $L^1(F)$, the domain of π_F^k . In passing, in our convergence proofs we only need apply the interpolation operator to classically regular functions; cf., in particular, the proof of Theorem 4.5 given in Section 6.*

Based on the local DOFs, we introduce a gradient and a potential reconstructions that will be instrumental in the formulation of the method. In what follows, $(\cdot, \cdot)_T$ and $(\cdot, \cdot)_F$ denote the L^2 -inner product on T and F , respectively. The same notation is used in the vector case $(\underline{L}^2)^d$. We define the local discrete gradient operator $G_T^k : \underline{U}_T^k \mapsto \mathbb{P}^k(T)^d$ such that, if $\underline{v}_T := (v_T, (v_F)_{F \in \mathcal{F}_T}) \in \underline{U}_T^k$ then for all $\phi \in \mathbb{P}^k(T)^d$,

$$(G_T^k \underline{v}_T, \phi)_T = (\nabla v_T, \phi)_T + \sum_{F \in \mathcal{F}_T} (v_F - v_T, \phi \cdot \mathbf{n}_{TF})_F \quad (4.3a)$$

$$= -(v_T, \nabla \cdot \phi)_T + \sum_{F \in \mathcal{F}_T} (v_F, \phi \cdot \mathbf{n}_{TF})_F. \quad (4.3b)$$

Recalling the definition (4.2) of \underline{l}_T^k , and using (4.3b) together with the definition (3.4) of the L^2 -projector, one can prove that the following commuting property holds: For all $v \in W^{1,1}(T)$,

$$G_T^k \underline{l}_T^k v = \pi_T^k(\nabla v), \quad (4.4)$$

where π_T^k acts component-wise. As a result, by (3.7) and (3.8), $G_T^k \underline{l}_T^k$ has optimal approximation properties in $\mathbb{P}^k(T)^d$.

The local potential reconstruction operator $p_T^{k+1} : \underline{U}_T^k \rightarrow \mathbb{P}^{k+1}(T)$ is such that, for all $\underline{v}_T \in \underline{U}_T^k$, $p_T^{k+1} \underline{v}_T \in \mathbb{P}^{k+1}(T)$ is the unique polynomial whose gradient is the orthogonal projection of $G_T^k \underline{v}_T$ on $\nabla \mathbb{P}^{k+1}(T) \subset \mathbb{P}^k(T)^d$, and whose mean value on T is equal to that of v_T . This amounts to defining $p_T^{k+1} \underline{v}_T$ such that

$$\begin{aligned} (\nabla p_T^{k+1} \underline{v}_T, \nabla w)_T &= (\nabla v_T, \nabla w)_T + \sum_{F \in \mathcal{F}_T} (v_F - v_T, \nabla w \cdot \mathbf{n}_{TF})_F \quad \forall w \in \mathbb{P}^{k+1}(T), \\ \int_T p_T^{k+1} \underline{v}_T(\mathbf{x}) d\mathbf{x} &= \int_T v_T(\mathbf{x}) d\mathbf{x}. \end{aligned} \quad (4.5)$$

It was proved in [29, Lemma 3] that $p_T^{k+1} \underline{l}_T^k$ has optimal H^s -approximation properties in $\mathbb{P}^{k+1}(T)$. The proof hinges on the fact that, for all $v \in H^1(T)$, $p_T^{k+1} \underline{l}_T^k v$ satisfies the following Euler equation:

$$(\nabla(p_T^{k+1} \underline{l}_T^k v - v), \nabla w)_T = 0 \quad \forall w \in \mathbb{P}^{k+1}(T), \quad (4.6)$$

which shows that $p_T^{k+1} \underline{l}_T^k$ is nothing but the usual elliptic projector on $\mathbb{P}^{k+1}(T)$.

4.2 Global degrees of freedom, interpolation and reconstructions

Local DOFs are collected in the following global space obtained by patching interface values:

$$\underline{U}_h^k := \left(\times_{T \in \mathcal{T}_h} \mathbb{P}^k(T) \right) \times \left(\times_{F \in \mathcal{F}_h} \mathbb{P}^k(F) \right).$$

We use the notation $\underline{v}_h = ((v_T)_{T \in \mathcal{T}_h}, (v_F)_{F \in \mathcal{F}_h})$ for a generic element $\underline{v}_h \in \underline{U}_h^k$ and, for all $T \in \mathcal{T}_h$, it is understood that $\underline{v}_T = (v_T, (v_F)_{F \in \mathcal{F}_T})$ denotes the restriction of \underline{v}_h to \underline{U}_T^k . The global interpolation operator $\underline{l}_h^k : W^{1,1}(\Omega) \rightarrow \underline{U}_h^k$ is defined such that, for all $v \in W^{1,1}(\Omega)$,

$$(\underline{l}_h^k v)|_T = \underline{l}_T^k(v|_T). \quad (4.7)$$

Uniqueness of interface DOFs is guaranteed by the regularity of functions in $W^{1,1}(\Omega)$. With $\mathbb{P}^k(\mathcal{T}_h)$ usual broken polynomial space on \mathcal{T}_h , for all $\underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_h^k$ we denote by \mathbf{v}_h the unique function in $\mathbb{P}^k(\mathcal{T}_h)$ such that

$$\mathbf{v}_h|_T = \mathbf{v}_T \quad \forall T \in \mathcal{T}_h. \quad (4.8)$$

Finally, we introduce the global discrete gradient operator $G_h^k : \underline{\mathbf{U}}_h^k \rightarrow \mathbb{P}^k(\mathcal{T}_h)^d$ and potential reconstruction $p_h^{k+1} : \underline{\mathbf{U}}_h^k \rightarrow \mathbb{P}^{k+1}(\mathcal{T}_h)$ such that, for all $\underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_h^k$,

$$(G_h^k \underline{\mathbf{v}}_h)|_T = G_T^k \underline{\mathbf{v}}_T \quad \text{and} \quad (p_h^{k+1} \underline{\mathbf{v}}_h)|_T = p_T^{k+1} \underline{\mathbf{v}}_T \quad \forall T \in \mathcal{T}_h. \quad (4.9)$$

4.3 Discrete problem and main results

Define the following subspace of $\underline{\mathbf{U}}_h^k$ which strongly incorporates the homogeneous Dirichlet boundary condition (1.1b):

$$\underline{\mathbf{U}}_{h,0}^k := \left\{ \underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_h^k \mid \mathbf{v}_F = 0 \quad \forall F \in \mathcal{F}_h^b \right\}. \quad (4.10)$$

We consider the following approximation of (2.3):

$$\text{Find } \underline{\mathbf{u}}_h \in \underline{\mathbf{U}}_{h,0}^k \text{ such that, for any } \underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_{h,0}^k, \quad A(\underline{\mathbf{u}}_h, \underline{\mathbf{v}}_h) = \int_{\Omega} f(\mathbf{x}) \mathbf{v}_h(\mathbf{x}) d\mathbf{x}, \quad (4.11a)$$

where $A : \underline{\mathbf{U}}_h^k \times \underline{\mathbf{U}}_h^k \mapsto \mathbb{R}$ is assembled element-wise

$$A(\underline{\mathbf{u}}_h, \underline{\mathbf{v}}_h) := \sum_{T \in \mathcal{T}_h} A_T(\underline{\mathbf{u}}_T, \underline{\mathbf{v}}_T), \quad (4.11b)$$

from the local contributions $A_T : \underline{\mathbf{U}}_T^k \times \underline{\mathbf{U}}_T^k \mapsto \mathbb{R}$, $T \in \mathcal{T}_h$, defined such that

$$\begin{aligned} A_T(\underline{\mathbf{u}}_T, \underline{\mathbf{v}}_T) &:= \int_T \mathbf{a}(\mathbf{x}, \mathbf{u}_T(\mathbf{x}), G_T^k \underline{\mathbf{u}}_T(\mathbf{x})) \cdot G_T^k \underline{\mathbf{v}}_T(\mathbf{x}) d\mathbf{x} + s_T(\underline{\mathbf{u}}_T, \underline{\mathbf{v}}_T), \\ s_T(\underline{\mathbf{u}}_T, \underline{\mathbf{v}}_T) &:= \sum_{F \in \mathcal{F}_T} h_F^{1-p} \int_F |\pi_F^k(\mathbf{u}_F - P_T^k \underline{\mathbf{u}}_T)(\mathbf{x})|^{p-2} \pi_F^k(\mathbf{u}_F - P_T^k \underline{\mathbf{u}}_T)(\mathbf{x}) \pi_F^k(\mathbf{v}_F - P_T^k \underline{\mathbf{v}}_T)(\mathbf{x}) ds(\mathbf{x}), \end{aligned} \quad (4.11c)$$

with $P_T^{k+1} : \underline{\mathbf{U}}_T^k \rightarrow \mathbb{P}^{k+1}(T)$ the second potential reconstruction obtained by setting, for all $\underline{\mathbf{v}}_T \in \underline{\mathbf{U}}_T^k$,

$$P_T^{k+1} \underline{\mathbf{v}}_T := \mathbf{v}_T + (p_T^{k+1} \underline{\mathbf{v}}_T - \pi_T^k p_T^{k+1} \underline{\mathbf{v}}_T). \quad (4.11d)$$

This specific (and rather elaborate) expression for the stabilization contribution s_T aims at improving the convergence properties of the method. As a matter of fact, s_T vanishes whenever one of its arguments is a polynomial of degree $\leq (k+1)$ inside the element. More naive choices of s_T (e.g., penalizing the difference $\mathbf{u}_F - \mathbf{u}_T$ instead of $\pi_F^k(\mathbf{u}_F - P_T^{k+1} \underline{\mathbf{u}}_T)$), would only ensure that this stabilisation vanishes on polynomials of degree $\leq k$ inside the element.

Remark 4.2 (Static condensation). *Problem (4.11a) is a system of nonlinear algebraic equations, which can be solved using an iterative algorithm. When first order (Newton-like) algorithms are used, element-based DOFs can be locally eliminated at each iteration by a standard static condensation procedure.*

Remark 4.3 (Variants). *Following [22], one could replace the space $\underline{\mathbf{U}}_T^k$ of (4.1) with*

$$\underline{\mathbf{U}}_T^{l,k} := \mathbb{P}^l(T) \times \left\{ \prod_{F \in \mathcal{F}_h} \mathbb{P}^k(F) \right\},$$

for $k \geq 0$ and $l \in \{k-1, k, k+1\}$. For the sake of simplicity, we only consider here the case $l = k-1$ when $k \geq 1$. For $k = 0$ and $l = k-1$, some technical modifications (not detailed here) are required owing to the absence of element-based DOFs. The local reconstruction operators G_T^k

defined by (4.3) and p_T^{k+1} defined by (4.5) still map on $\mathbb{P}^k(T)^d$ and $\mathbb{P}^{k+1}(T)$, respectively (their domain changes, but we keep the same notation for the sake of simplicity). A close inspection shows that both key properties (4.4) and (4.6) remain valid for the proposed choices for l . The second potential reconstruction operator P_T^{k+1} defined by (4.11d), on the other hand, is replaced with $P_T^{l,k+1} : \underline{\mathbf{U}}_T^{l,k} \rightarrow \mathbb{P}^{k+1}(T)$ such that, for all $\underline{\mathbf{v}}_T \in \underline{\mathbf{U}}_T^{l,k}$, $P_T^{l,k+1} \underline{\mathbf{v}}_T := \mathbf{v}_T + (p_T^{k+1} \underline{\mathbf{v}}_T - \pi_T^l p_T^{k+1} \underline{\mathbf{v}}_T)$. The interest of the case $l = k + 1$ is that it holds, for all $\underline{\mathbf{v}}_T \in \underline{\mathbf{U}}_T^k$, $P_T^{k+1,k+1} \underline{\mathbf{v}}_T = \mathbf{v}_T$, and the stabilization contribution takes the simpler form

$$s_T(\underline{\mathbf{u}}_T, \underline{\mathbf{v}}_T) = \sum_{F \in \mathcal{F}_T} h_F^{1-p} \int_F |\pi_F^k(\mathbf{u}_F - \mathbf{u}_T)(\mathbf{x})|^{p-2} \pi_F^k(\mathbf{u}_F - \mathbf{u}_T)(\mathbf{x}) \pi_F^k(\mathbf{v}_F - \mathbf{v}_T)(\mathbf{x}) ds(\mathbf{x}).$$

This simplification, however, comes at the price of having more element-based DOFs, which leads in turn to more onerous local problems for both the computation of the operator reconstructions and the elimination of element-based unknowns by static condensation. We also notice that the choice $l = k + 1$ is close in spirit to the Hybridizable Discontinuous Galerkin methods introduced in [23] for a linear diffusion problem. The choice $l = k - 1$, on the other hand, can be related to the High-Order Mimetic method introduced in [47] in the context of linear elliptic equations.

We next state our main results for problem (4.11). The proofs are postponed to Section 6.

Theorem 4.4 (Existence of a discrete solution). *Under Assumption (2.2), if \mathcal{T}_h is a mesh, then there exists at least one solution $\underline{\mathbf{u}}_h \in \underline{\mathbf{U}}_{h,0}^k$ to (4.11).*

Theorem 4.5 (Convergence). *We assume (2.2) and we let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence. For all $h \in \mathcal{H}$, we let $\underline{\mathbf{u}}_h \in \underline{\mathbf{U}}_{h,0}^k$ be a solution to (4.11) on \mathcal{T}_h . Then up to a subsequence as $h \rightarrow 0$, recalling the definition (2.1) of the Sobolev index p^* ,*

- $\mathbf{u}_h \rightarrow u$ and $p_h^{k+1} \underline{\mathbf{u}}_h \rightarrow u$ strongly in $L^q(\Omega)$ for all $q < p^*$,
- $G_h^k \underline{\mathbf{u}}_h \rightarrow \nabla u$ weakly in $L^p(\Omega)^d$,

where $u \in W_0^{1,p}(\Omega)$ solves the weak formulation (2.3) of the PDE (1.1). If we assume, moreover, that \mathbf{a} is strictly monotone, that is the inequality in (2.2c) is strict if $\boldsymbol{\xi} \neq \boldsymbol{\eta}$, then

- $G_h^k \underline{\mathbf{u}}_h \rightarrow \nabla u$ strongly in $L^p(\Omega)^d$,

Remark 4.6 (Uniqueness). *If \mathbf{a} does not depend on s and is strictly monotone, then the solutions to both the continuous problem (2.3) and its discrete counterpart (4.11) are unique, and the whole sequence of approximate solutions converges to the weak solution of (1.1).*

4.4 Numerical examples

To close this section, we provide a few numerical examples to assess the convergence properties of the method. We consider the p -Laplacian problem, corresponding to the following choice for the function \mathbf{a} : for $1 < p < +\infty$,

$$\mathbf{a}(\mathbf{x}, u, \nabla u) = |\nabla u|^{p-2} \nabla u.$$

When $p = 2$, we recover the usual (linear) Laplace operator. We consider the two-dimensional analytical solution originally proposed in [3, Section 4], corresponding to $u(\mathbf{x}) = \exp(x_1 + \pi x_2)$ with suitable source term f inferred from (1.1a). The domain is the unit square $\Omega = (0, 1)^2$, and nonhomogeneous Dirichlet boundary conditions inferred from the expression of u are enforced on its boundary. We compute the numerical solution corresponding to polynomial degrees $k = 0, \dots, 4$. The meshes used are the triangular and Cartesian mesh families 1 and 2 from the FVCA 5 benchmark [43], and the distorted (predominantly) hexagonal mesh family of [30, Section 4.2.3]; cf. Figure 2. In Figures 3 and 4 we display the convergence of the error $\|G_h^k(\underline{\mathbf{u}}_h - \mathbb{I}_h^k u)\|_{L^p(\Omega)^d}$ for $p = 3$ and $p = 4$, respectively. In all the cases, increasing the polynomial degree k improves the convergence rate. Further numerical tests (not reported here for the sake of brevity) show that




Figure 2: Meshes used in the numerical tests of Section 4.4.




Figure 3: $\|G_h^k(\underline{u}_h - \mathbb{1}_h^k u)\|_{L^p(\Omega)^d}$ vs. h , $p = 3$.

this remains true when considering “degenerate” cases (i.e., solutions with a gradient that vanishes in part of the domain, in which case the diffusive properties of (1.1) degenerate), although the gain can sometimes be less relevant. We note, in passing, that the results obtained in [1, 3, 10] for lowest-order schemes suggest that we should not expect, in general, optimal convergence properties in $\mathbb{P}^{k+1}(\mathcal{T}_h)$. Instead, the order of convergence depends on both the regularity of the exact solution and the index p (with optimal convergence orders recovered for smooth solutions to the Poisson problem, corresponding to $p = 2$). Finally, for the sake of completeness, we report in Figure 5 the numerical results obtained for $p = 4$ with the method discussed in Remark 4.3 and corresponding to $l = k + 1$. In this case, taking the element-based DOFs in $\mathbb{P}^{k+1}(T)$ does not seem to bring any significant advantage in terms of convergence (compare with Figure 4).




Figure 4: $\|G_h^k(\underline{u}_h - \mathbb{1}_h^k u)\|_{L^p(\Omega)^d}$ vs. h , $p = 4$.




Figure 5: $\|G_h^k(\underline{u}_h - \underline{I}_h^k u)\|_{L^p(\Omega)^d}$ vs. h , $p = 4$ for the variant of the method discussed in Remark 4.3 and corresponding to $l = k + 1$.

5 Discrete functional analysis tools in hybrid polynomial spaces

This section collects discrete functional analysis results on hybrid polynomial spaces that are used in the convergence analysis of Section 6.

5.1 Discrete $W^{1,p}$ -norms

We introduce the following discrete counterpart of the $W^{1,p}$ -seminorm on \underline{U}_h^k :

$$\|\underline{v}_h\|_{1,p,h} := \left(\sum_{T \in \mathcal{T}_h} \|\underline{v}_T\|_{1,p,T}^p \right)^{\frac{1}{p}}, \quad (5.1)$$

where the local seminorm $\|\cdot\|_{1,p,T}$ on \underline{U}_T^k is defined by

$$\|\underline{v}_T\|_{1,p,T} := \left(\|\nabla \underline{v}_T\|_{L^p(T)^d}^p + \sum_{F \in \mathcal{F}_T} h_F^{1-p} \|\underline{v}_F - \underline{v}_T\|_{L^p(F)}^p \right)^{\frac{1}{p}}. \quad (5.2)$$

It is a simple matter to check that $\|\cdot\|_{1,p,h}$ defines a norm on $\underline{U}_{h,0}^k$. We next show uniform equivalence between the local seminorm defined by (5.2) and two local $W^{1,p}$ -seminorms defined using the discrete gradient and potential reconstructions (cf. (4.3a) and (4.5), respectively) and the penalty contribution s_T (cf. (4.11c)). This essentially proves stability for the discrete problem (4.11a) in terms of the $\|\cdot\|_{1,p,h}$ -norm. The argument hinges on the following direct and reverse Lebesgue embeddings, whose proof is postponed to Appendix A.1.

Lemma 5.1 (Direct and reverse Lebesgue embeddings). *Let U be a measurable subset of \mathbb{R}^N such that (3.5) holds. Let $k \in \mathbb{N}$ and $q, m \in [1, +\infty]$. Then,*

$$\forall w \in \mathbb{P}^k(U) : \|w\|_{L^q(U)} \approx |U|^{\frac{1}{q} - \frac{1}{m}} \|w\|_{L^m(U)}, \quad (5.3)$$

where $A \approx B$ means that there is a real $M > 0$ only depending on N, k, δ, q and m such that $M^{-1}A \leq B \leq MA$.

We are now ready to prove the norm equivalence.

Lemma 5.2 (Equivalence of discrete $W^{1,p}$ -seminorms). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence and $k \in \mathbb{N}$. Let $T \in \mathcal{T}_h$, $p \in [1, +\infty)$, and denote by $|\cdot|_{s,p,T}$ the local face seminorm such*

that, for all $\underline{\mathbf{v}}_T \in \underline{\mathbf{U}}_T^k$, recalling the definition (4.11c) of s_T ,

$$|\underline{\mathbf{v}}_T|_{s,p,T} := s_T(\underline{\mathbf{v}}_T, \underline{\mathbf{v}}_T)^{\frac{1}{p}} = \left(\sum_{F \in \mathcal{F}_T} \int_F h_F^{1-p} |\pi_F^k(\mathbf{v}_F - P_T^{k+1} \underline{\mathbf{v}}_T)(\mathbf{x})|^p ds(\mathbf{x}) \right)^{\frac{1}{p}}. \quad (5.4)$$

Then,

$$\|\underline{\mathbf{v}}_T\|_{1,p,T} \approx \left(\|\nabla P_T^{k+1} \underline{\mathbf{v}}_T\|_{L^p(T)^d}^p + |\underline{\mathbf{v}}_T|_{s,p,T}^p \right)^{\frac{1}{p}} \approx \left(\|G_T^k \underline{\mathbf{v}}_T\|_{L^p(T)^d}^p + |\underline{\mathbf{v}}_T|_{s,p,T}^p \right)^{\frac{1}{p}}, \quad (5.5)$$

where $A \approx B$ means that $M^{-1}A \leq B \leq MA$ for some real number $M > 0$ that may depend on Ω , ϱ , k and p , but does not otherwise depend on the mesh, T or $\underline{\mathbf{v}}_T$.

Remark 5.3 (Choice of the face seminorm). *The proof of the norm equivalence does not make use of the specific structure of s_T , and could have been proved replacing $|\cdot|_{s,p,T}$ by any other local face seminorm composed by terms scaling on each face $F \in \mathcal{F}_T$ as $h_F^{1-p} \|\cdot\|_{L^p(F)}$.*

Proof. We abridge $A \lesssim B$ the inequality $A \leq MB$ with real M only depending on Ω , ϱ , k and p .

Step 1: $p = 2$. It was proved in [29, Lemma 4] that

$$\|\underline{\mathbf{v}}_T\|_{1,2,T}^2 \approx \|\nabla P_T^{k+1} \underline{\mathbf{v}}_T\|_{L^2(T)^d}^2 + |\underline{\mathbf{v}}_T|_{s,2,T}^2, \quad (5.6)$$

which is exactly the first relation in (5.5) for $p = 2$. To prove the second, we notice that since, for all $\underline{\mathbf{v}}_T \in \underline{\mathbf{U}}_T^k$, $\nabla P_T^{k+1} \underline{\mathbf{v}}_T$ is an orthogonal projection of $G_T^k \underline{\mathbf{v}}_T$ in $L^2(T)^d$, we have $\|\nabla P_T^{k+1} \underline{\mathbf{v}}_T\|_{L^2(T)^d} \leq \|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)^d}$. Relation (5.6) therefore shows that

$$\|\underline{\mathbf{v}}_T\|_{1,2,T}^2 \lesssim \|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)^d}^2 + |\underline{\mathbf{v}}_T|_{s,2,T}^2.$$

To prove the converse estimate, we plug $\phi = G_T^k \underline{\mathbf{v}}_T$ into the definition (4.3a) of $G_T^k \underline{\mathbf{v}}_T$, and use the Cauchy–Schwarz inequality together with the discrete trace inequality [28, Lemma 1.46] and the uniform bound (3.2) on N_∂ to infer

$$\begin{aligned} \|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)^d}^2 &\lesssim \|\nabla \mathbf{v}_T\|_{L^2(T)^d} \|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)^d} + \sum_{F \in \mathcal{F}_T} h_F^{-\frac{1}{2}} \|\mathbf{v}_F - \mathbf{v}_T\|_{L^2(F)} \|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)^d} \\ &\lesssim \|\underline{\mathbf{v}}_T\|_{1,2,T} \|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)^d}. \end{aligned}$$

This estimate shows that $\|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)^d} \lesssim \|\underline{\mathbf{v}}_T\|_{1,2,T}$ and, combined with (5.6) to estimate $|\underline{\mathbf{v}}_T|_{s,2,T} \lesssim \|\underline{\mathbf{v}}_T\|_{1,2,T}$, completes the proof of the case $p = 2$.

Step 2: $p \in [1, +\infty)$. Relation (5.5) for a generic p can be deduced from the case $p = 2$ thanks to Lemma 5.1 (T and F clearly satisfy the geometric assumptions therein, cf. Remark 3.3). We only show how to do this to establish

$$\|\underline{\mathbf{v}}_T\|_{1,p,T}^p \lesssim \|G_T^k \underline{\mathbf{v}}_T\|_{L^p(T)^d}^p + |\underline{\mathbf{v}}_T|_{s,p,T}^p,$$

all the other estimates being obtained in a similar way. By admissibility of $(\mathcal{T}_h)_{h \in \mathcal{H}}$, we have $h_F |F| \approx |T|$ for any $F \in \mathcal{F}_T$. Thus, for $\underline{\mathbf{v}}_T \in \underline{\mathbf{U}}_T^k$, by Lemma 5.1,

$$\begin{aligned} \|\underline{\mathbf{v}}_T\|_{1,p,T}^p &\lesssim |T|^{1-\frac{p}{2}} \|\nabla \mathbf{v}_T\|_{L^2(T)^d}^p + \sum_{F \in \mathcal{F}_T} h_F^{1-p} |F|^{1-\frac{p}{2}} \|\mathbf{v}_F - \mathbf{v}_T\|_{L^2(F)}^p \\ &\lesssim |T|^{1-\frac{p}{2}} \left(\|\nabla \mathbf{v}_T\|_{L^2(T)^d}^2 + \sum_{F \in \mathcal{F}_T} h_F^{-1} \|\mathbf{v}_F - \mathbf{v}_T\|_{L^2(F)}^2 \right)^{\frac{p}{2}}, \end{aligned}$$

where, to pass to the second line, we used the inequality

$$\forall \theta > 0, \forall a_i \geq 0 : \sum_{i=0}^N a_i \leq N \left(\sum_{i=1}^N a_i^\theta \right)^{\frac{1}{\theta}} \quad (5.7)$$

which follows from writing $a_j = (a_j^\theta)^{\frac{1}{\theta}} \leq (\sum_{i=1}^N a_i^\theta)^{\frac{1}{\theta}}$ for all j . Apply (5.5) with $p = 2$ and use again Lemma 5.1 and the inequality (5.7) to infer

$$\begin{aligned} \|\underline{\mathbf{v}}_T\|_{1,p,T}^p &\lesssim |T|^{1-\frac{p}{2}} \left(\|G_T^k \underline{\mathbf{v}}_T\|_{L^2(T)}^2 + \sum_{F \in \mathcal{F}_T} h_F^{-1} \|\pi_F^k(\mathbf{v}_F - P_T^{k+1} \underline{\mathbf{v}}_T)\|_{L^2(F)}^2 \right)^{\frac{p}{2}} \\ &\lesssim |T|^{1-\frac{p}{2}} \left(|T|^{1-\frac{2}{p}} \|G_T^k \underline{\mathbf{v}}_T\|_{L^p(T)}^2 + \sum_{F \in \mathcal{F}_T} h_F^{-1} |F|^{1-\frac{2}{p}} \|\pi_F^k(\mathbf{v}_F - P_T^{k+1} \underline{\mathbf{v}}_T)\|_{L^p(F)}^2 \right)^{\frac{p}{2}} \\ &\lesssim |T|^{1-\frac{p}{2}} \left(|T|^{1-\frac{2}{p}} \left(\|G_T^k \underline{\mathbf{v}}_T\|_{L^p(T)}^2 + \sum_{F \in \mathcal{F}_T} h_F^{\frac{2}{p}-2} \|\pi_F^k(\mathbf{v}_F - P_T^{k+1} \underline{\mathbf{v}}_T)\|_{L^p(F)}^2 \right) \right)^{\frac{p}{2}} \\ &\lesssim \|G_T^k \underline{\mathbf{v}}_T\|_{L^p(T)}^p + \sum_{F \in \mathcal{F}_T} h_F^{1-p} \|\pi_F^k(\mathbf{v}_F - P_T^{k+1} \underline{\mathbf{v}}_T)\|_{L^p(F)}^p. \quad \square \end{aligned}$$

5.2 Discrete Sobolev embeddings

The following discrete counterpart of Sobolev embeddings will be used in Proposition 6.1 to obtain an a priori estimate of the discrete solution.

Proposition 5.4 (Discrete Sobolev embeddings). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence. Let $1 \leq q \leq p^*$ if $1 \leq p < d$ (with p^* defined by (2.1)) and $1 \leq q < +\infty$ if $p \geq d$. Then, there exists C only depending on Ω , ρ , k , q and p such that*

$$\forall \underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_{h,0}^k : \|\underline{\mathbf{v}}_h\|_{L^q(\Omega)} \leq C \|\underline{\mathbf{v}}_h\|_{1,p,h}. \quad (5.8)$$

Remark 5.5 (Discrete Poincaré). *For $q = p$ (this choice is always possible since $p \leq p^*$ for any space dimension d) this proposition states a discrete Poincaré's inequality.*

Proof. Here, $A \lesssim B$ means that $A \leq MB$ for some M only depending on Ω , ρ , k , q and p . We recall the discrete Sobolev embeddings in $\mathbb{P}^k(\mathcal{T}_h)$ from [28, Theorem 5.3] (cf. also [19, 27]):

$$\forall w \in \mathbb{P}^k(\mathcal{T}_h) : \|w\|_{L^q(\Omega)} \lesssim \|w\|_{\text{dG},p}, \quad (5.9)$$

where the discrete $W^{1,p}$ -norm on $\mathbb{P}^k(\mathcal{T}_h)$ is defined by

$$\|w\|_{\text{dG},p} := \left(\sum_{T \in \mathcal{T}_h} \|\nabla w_T\|_{L^p(T)^d}^p + \sum_{F \in \mathcal{F}_h} h_F^{1-p} \|[w]_F\|_{L^p(F)}^p \right)^{\frac{1}{p}}. \quad (5.10)$$

Here, for all $T \in \mathcal{T}_h$, $w_T := w|_T$, while $[w]_F = w_{T_1} - w_{T_2}$ is the jump of w through a face $F \in \mathcal{F}_h^i$ such that $\mathcal{T}_F = \{T_1, T_2\}$ (the sign is irrelevant). If $F \in \mathcal{F}_h^b$, then $\mathcal{T}_F = \{T\}$ and we let $[w]_F = w_T$. For $\underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_{h,0}^k$ and F a face between T_1 and T_2 , we have, using the triangle inequality,

$$\|[\underline{\mathbf{v}}_h]_F\|_{L^p(F)} \leq \|\mathbf{v}_{T_1} - \mathbf{v}_F\|_{L^p(F)} + \|\mathbf{v}_{T_2} - \mathbf{v}_F\|_{L^p(F)}.$$

Due to the strong boundary conditions, this estimate is also true if F is a boundary face and the term T_2 is removed. Hence, gathering by elements,

$$\begin{aligned} \sum_{F \in \mathcal{F}_h} h_F^{1-p} \|[w]_F\|_{L^p(F)}^p &\lesssim \sum_{F \in \mathcal{F}_h} h_F^{1-p} \sum_{T \in \mathcal{T}_F} \|\mathbf{v}_T - \mathbf{v}_F\|_{L^p(F)}^p \\ &= \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} h_F^{1-p} \|\mathbf{v}_T - \mathbf{v}_F\|_{L^p(F)}^p \leq \|\underline{\mathbf{v}}_h\|_{1,p,h}^p. \end{aligned}$$

This shows that

$$\|\mathbf{v}_h\|_{\text{dG},p} \lesssim \|\underline{\mathbf{v}}_h\|_{1,p,h}, \quad (5.11)$$

which, plugged into (5.9), concludes the proof. \square

5.3 Compactness

The second ingredient for our convergence analysis is a compactness result for sequences bounded in the $\|\cdot\|_{1,p,h}$ -norm. Before establishing compactness, we need an auxiliary result allowing us to compare, for all $\underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_h^k$, the broken polynomial function (4.8) on \mathcal{T}_h defined by element DOFs and the potential reconstruction (4.9). Instrumental to the proof is the following Poincaré–Wirtinger–Sobolev inequality on broken polynomial spaces.

Lemma 5.6 (Poincaré–Wirtinger–Sobolev inequality for broken polynomial functions with local zero average). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence, and let $p \leq q \leq p^*$ with p^* defined by (2.1). If $w \in \mathbb{P}^k(\mathcal{T}_h)$ satisfies $\int_T w = 0$ for all $T \in \mathcal{T}_h$, then there exists C only depending on Ω , ϱ , k , q and p such that (with ∇_h denoting the usual broken gradient),*

$$\|w\|_{L^q(\Omega)} \leq Ch^{1+\frac{d}{q}-\frac{d}{p}} \|\nabla_h w\|_{L^p(\Omega)^d}. \quad (5.12)$$

Remark 5.7. *If $p \leq d$, the exponent $1 + \frac{d}{q} - \frac{d}{p}$ in h is positive if $q < p^*$ and equal to 0 if $q = p^*$.*

Proof. In this proof, $A \lesssim B$ means that $A \leq MB$ for some M only depending on Ω , ϱ , k , q and p . We have $\pi_T^0 w = 0$ and therefore, by (3.7) with $k = 0$, $s = 1$ and $m = 0$, using Lemma 5.1 with $m = p$, and recalling that $|T| \lesssim h_T^d$, we write

$$\begin{aligned} \|w\|_{L^q(T)} &= \|w - \pi_T^0 w\|_{L^q(T)} \lesssim h_T \|\nabla w\|_{L^q(T)^d} \\ &\lesssim h_T |T|^{\frac{1}{q}-\frac{1}{p}} \|\nabla w\|_{L^p(T)^d} \lesssim h_T^{1+\frac{d}{q}-\frac{d}{p}} \|\nabla w\|_{L^p(T)^d}. \end{aligned} \quad (5.13)$$

If q is finite, we take the the power q of this inequality, sum over $T \in \mathcal{T}_h$, and use $\|\nabla w\|_{L^p(T)^d}^{q-p} \leq \|\nabla_h w\|_{L^p(\Omega)^d}^{q-p}$ (we have $q \geq p$) to infer

$$\begin{aligned} \|w\|_{L^q(\Omega)}^q &\lesssim h^{q+d-\frac{dq}{p}} \sum_{T \in \mathcal{T}_h} \|\nabla w\|_{L^p(T)^d}^q \leq h^{q+d-\frac{dq}{p}} \|\nabla_h w\|_{L^p(\Omega)^d}^{q-p} \sum_{T \in \mathcal{T}_h} \|\nabla w\|_{L^p(T)^d}^p \\ &= h^{q+d-\frac{dq}{p}} \|\nabla_h w\|_{L^p(\Omega)^d}^{q-p} \|\nabla_h w\|_{L^p(\Omega)^d}^p = h^{q+d-\frac{dq}{p}} \|\nabla_h w\|_{L^p(\Omega)^d}^q. \end{aligned}$$

Taking the power $1/q$ of this inequality concludes the proof. If $q = +\infty$, we apply (5.13) to $T \in \mathcal{T}_h$ such that $\|w\|_{L^\infty(T)} = \|w\|_{L^\infty(\Omega)}$ and we obtain $\|w\|_{L^\infty(\Omega)} \lesssim h^{1-\frac{d}{p}} \|\nabla w\|_{L^p(T)^d} \leq h^{1-\frac{d}{p}} \|\nabla w\|_{L^p(\Omega)^d}$. \square

Corollary 5.8 (Comparison between \mathbf{v}_h and $p_h^{k+1} \underline{\mathbf{v}}_h$). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence, and let $p \leq q \leq p^*$. Then, there exists C only depending on Ω , ϱ , k , q and p such that*

$$\forall \underline{\mathbf{v}}_h \in \underline{\mathbf{U}}_h^k : \|\mathbf{v}_h - p_h^{k+1} \underline{\mathbf{v}}_h\|_{L^q(\Omega)} \leq Ch^{1+\frac{d}{q}-\frac{d}{p}} \|\underline{\mathbf{v}}_h\|_{1,p,h}. \quad (5.14)$$

Proof. Here, $A \lesssim B$ means $A \leq MB$ for M only depending on Ω , ϱ , k , q and p . By the second equation in (4.5), the average of $\mathbf{v}_h - p_h^{k+1} \underline{\mathbf{v}}_h$ over each element of \mathcal{T}_h is zero. Hence, (5.12) gives

$$\|\mathbf{v}_h - p_h^{k+1} \underline{\mathbf{v}}_h\|_{L^q(\Omega)} \lesssim h^{1+\frac{d}{q}-\frac{d}{p}} \|\nabla_h(\mathbf{v}_h - p_h^{k+1} \underline{\mathbf{v}}_h)\|_{L^p(\Omega)^d}. \quad (5.15)$$

Recalling the definitions (4.8) of \mathbf{v}_h and (5.1) of the $\|\cdot\|_{1,p,h}$ -norm, we have

$$\|\nabla_h \mathbf{v}_h\|_{L^p(\Omega)^d}^p = \sum_{T \in \mathcal{T}_h} \|\nabla \mathbf{v}_T\|_{L^p(T)^d}^p \leq \|\underline{\mathbf{v}}_h\|_{1,p,h}^p. \quad (5.16)$$

Moreover, using the definition (4.9) of $p_h^{k+1}\underline{v}_h$ followed by the norm equivalence (5.5), and again the definition (5.1) of the $\|\cdot\|_{1,p,h}$ -norm, it is inferred that

$$\|\nabla_h p_h^{k+1}\underline{v}_h\|_{L^p(\Omega)^d}^p = \sum_{T \in \mathcal{T}_h} \|\nabla p_T^{k+1}\underline{v}_T\|_{L^p(T)^d}^p \lesssim \sum_{T \in \mathcal{T}_h} \|\underline{v}_T\|_{1,p,T}^p = \|\underline{v}_h\|_{1,p,h}^p. \quad (5.17)$$

We conclude by using the triangle inequality in the right-hand side of (5.15), and by plugging (5.16) and (5.17) into the resulting equation. \square

We are now ready to prove the compactness result.

Proposition 5.9 (Discrete compactness). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence and let $\underline{v}_h \in \underline{U}_{h,0}^k$ be such that $(\|\underline{v}_h\|_{1,p,h})_{h \in \mathcal{H}}$ is bounded. Then, there exists $v \in W_0^{1,p}(\Omega)$ such that, up to a subsequence as $h \rightarrow 0$, recalling the definition (2.1) of the Sobolev index p^* ,*

- $\underline{v}_h \rightarrow v$ and $p_h^{k+1}\underline{v}_h \rightarrow v$ strongly in $L^q(\Omega)$ for all $q < p^*$,
- $G_h^k \underline{v}_h \rightarrow \nabla v$ weakly in $L^p(\Omega)^d$.

Remark 5.10. *If $p^* < +\infty$, the discrete Sobolev embeddings of [28, Theorem 5.3] (recalled in (5.9)) and Corollary 5.8 show that both \underline{v}_h and $p_h^{k+1}\underline{v}_h$ are bounded in $L^{p^*}(\Omega)$, which proves that their convergence stated in Proposition 5.9 extends to $L^{p^*}(\Omega)$ -weak.*

Proof. By (5.11), $(\|\underline{v}_h\|_{dG,p})_{h \in \mathcal{H}}$ is bounded. The discrete Rellich–Kondrachov theorem [28, Theorem 5.6] ensures that, up to a subsequence, \underline{v}_h converges in $L^q(\Omega)$ to some v . Since $q < p^*$, Corollary 5.8 shows that $p_h^{k+1}\underline{v}_h$ also converges in this space to the same v .

It remains to establish that $v \in W_0^{1,p}(\Omega)$ and that $G_h^k \underline{v}_h$ weakly converges to ∇v . To this end, we first notice that $G_h^k \underline{v}_h$ is bounded in $L^p(\Omega)^d$ thanks to the norm equivalence (5.5). Hence, up to a subsequence, it weakly converges in $L^p(\Omega)^d$ to some \mathcal{G} . We take $\phi \in C^\infty(\mathbb{R}^d)^d$ and observe that, by definition (4.3a) of $G_T^k \underline{v}_T$,

$$\begin{aligned} \int_{\Omega} G_h^k \underline{v}_h(\mathbf{x}) \cdot \phi(\mathbf{x}) d\mathbf{x} &= \sum_{T \in \mathcal{T}_h} (G_T^k \underline{v}_T, \phi)_T \\ &= \sum_{T \in \mathcal{T}_h} (G_T^k \underline{v}_T - \nabla \underline{v}_T, \phi - \pi_T^k \phi)_T + \sum_{T \in \mathcal{T}_h} (G_T^k \underline{v}_T - \nabla \underline{v}_T, \pi_T^k \phi)_T \\ &\quad + \sum_{T \in \mathcal{T}_h} (\nabla \underline{v}_T, \phi)_T \\ &= \mathfrak{I}_1 + \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} (\underline{v}_F - \underline{v}_T, \pi_T^k \phi \cdot \mathbf{n}_{TF})_F + \sum_{T \in \mathcal{T}_h} (\nabla \underline{v}_T, \phi)_T \quad (\text{cf. (4.3a)}) \\ &= \mathfrak{I}_1 + \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} (\underline{v}_F - \underline{v}_T, (\pi_T^k \phi - \phi) \cdot \mathbf{n}_{TF})_F - \sum_{T \in \mathcal{T}_h} (\underline{v}_T, \text{div } \phi)_T \quad (\text{cf. (5.18)}) \\ &= \mathfrak{I}_1 + \mathfrak{I}_2 - \int_{\Omega} \underline{v}_h(\mathbf{x}) \text{div } \phi(\mathbf{x}) d\mathbf{x}. \end{aligned}$$

In the fourth line, we used a element-wise integration by parts, and the relation

$$\sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} (\underline{v}_F, \phi \cdot \mathbf{n}_{TF})_F = 0, \quad (5.18)$$

which follows from the homogeneous Dirichlet boundary condition incorporated in $\underline{U}_{h,0}^k$ (cf. (4.10)) and from $\mathbf{n}_{T_1 F} + \mathbf{n}_{T_2 F} = 0$ whenever $F \in \mathcal{F}_h^i$ is an interface between the two elements T_1 and T_2 . If we prove that, as $h \rightarrow 0$, $\mathfrak{I}_1 + \mathfrak{I}_2 \rightarrow 0$, then we can pass to the limit and we obtain

$$\int_{\Omega} \mathcal{G}(\mathbf{x}) \cdot \phi(\mathbf{x}) d\mathbf{x} = - \int_{\Omega} v(\mathbf{x}) \text{div } \phi(\mathbf{x}) d\mathbf{x}. \quad (5.19)$$

Taking ϕ compactly supported in Ω shows that $\mathcal{G} = \nabla v$, and hence that $v \in W^{1,p}(\Omega)$ and that $G_h^k \underline{v}_h \rightarrow \nabla v$ weakly in $L^p(\Omega)^d$. Taking then any $\phi \in C^\infty(\mathbb{R}^d)^d$ in (5.19) and using an integration by parts shows that the trace of v on $\partial\Omega$ vanishes, which establishes that $v \in W_0^{1,p}(\Omega)$.

It therefore only remains to prove that $\mathfrak{T}_1 + \mathfrak{T}_2 \rightarrow 0$. In what follows, $A \lesssim B$ means that $A \leq MB$ for some M not depending on h , ϕ or \underline{v}_T . By Lemma 3.4 (with $m = 0$, $s = 1$ and p' instead of p) we have $\|\phi - \pi_T^k \phi\|_{L^{p'}(T)^d} \lesssim h \|\phi\|_{W^{1,p'}(T)^d}$ and thus

$$|\mathfrak{T}_1| \lesssim h \left(\sum_{T \in \mathcal{T}_h} \|G_T^k \underline{v}_T - \nabla v_T\|_{L^p(T)^d}^p \right)^{1/p} \|\phi\|_{W^{1,p'}(\Omega)^d} \lesssim h (\|G_h^k \underline{v}_h\|_{L^p(\Omega)^d} + \|\nabla_h v_h\|_{L^p(\Omega)^d}).$$

Since $\|\underline{v}_T\|_{1,p,h}$ is bounded, the norm equivalence (5.5) together with the definition (5.1) of the $\|\cdot\|_{1,p,h}$ -norm show that both $\|G_h^k \underline{v}_h\|_{L^p(\Omega)^d}$ and $\|\nabla_h v_h\|_{L^p(\Omega)^d}$ remain bounded. Hence, $\mathfrak{T}_1 \rightarrow 0$ as $h \rightarrow 0$. The convergence analysis of \mathfrak{T}_2 is performed in a similar way. Using Lemma 3.6 (with p' instead of p) we have $\|\phi - \pi_T^k \phi\|_{L^{p'}(F)} \lesssim h_T^{\frac{1}{p}} \|\phi\|_{W^{1,p'}(T)^d}$ and thus, since $h_T \lesssim h_F$ whenever $F \in \mathcal{F}_T$,

$$\begin{aligned} |\mathfrak{T}_2| &\lesssim \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} h_F^{\frac{1}{p}} \|\mathbf{v}_F - \mathbf{v}_T\|_{L^p(F)} \|\phi\|_{W^{1,p'}(T)^d} \lesssim \left(\sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} h_F \|\mathbf{v}_F - \mathbf{v}_T\|_{L^p(F)}^p \right)^{\frac{1}{p}} \|\phi\|_{W^{1,p'}(\Omega)^d} \\ &\lesssim h \|\underline{v}_T\|_{1,p,h} \|\phi\|_{W^{1,p'}(\Omega)^d}. \end{aligned}$$

The convergence of \mathfrak{T}_2 to 0 follows. \square

5.4 Strong convergence of the interpolants

The proof of Theorem 4.5 relies on a weak-strong convergence argument. The following result shows strong convergence for both the discrete gradient and the stabilization contribution when their argument is the interpolate of a smooth function.

Proposition 5.11 (Strong convergence of interpolants). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence, let $p \in [1, +\infty]$ and let \mathbb{I}_T^k be defined by (4.2). Then, as $h \rightarrow 0$,*

$$\forall \varphi \in W^{1,p}(\Omega) : G_h^k \mathbb{I}_h^k \varphi \rightarrow \nabla \varphi \text{ strongly in } L^p(\Omega)^d \quad (5.20)$$

and, denoting by $W^{r,\infty}(\mathcal{T}_h)$ the space of $\varphi : \Omega \mapsto \mathbb{R}$ such that, for any $T \in \mathcal{T}_h$, $\varphi|_T \in W^{r,\infty}(T)$,

$$\forall \varphi \in W^{1,1}(\Omega) \cap W^{k+2,\infty}(\mathcal{T}_h) : \sum_{T \in \mathcal{T}_h} s_T (\mathbb{I}_T^k \varphi, \mathbb{I}_T^k \varphi) \rightarrow 0. \quad (5.21)$$

Proof. Step 1: Proof of (5.20). We write $A \lesssim B$ for $A \leq MB$ where M does not depend on h or φ . Let us first assume that φ is smooth, say $W^{2,p}(\Omega)$. By the commuting property (4.4) and the approximation property (3.7) applied to $v = \partial_i \varphi$ we have $\|G_T^k \mathbb{I}_T^k \varphi - \nabla \varphi\|_{L^p(T)^d} \lesssim h_T \|\varphi\|_{W^{2,p}(T)}$. Raising this inequality to the power p and summing over $T \in \mathcal{T}_h$ (if p is finite, otherwise taking the maximum over $T \in \mathcal{T}_h$) gives

$$\|G_T^k \mathbb{I}_T^k \varphi - \nabla \varphi\|_{L^p(\Omega)^d} \lesssim h \|\varphi\|_{W^{2,p}(\Omega)}, \quad (5.22)$$

and thus the conclusion follows. If φ only belongs to $W_0^{1,p}(\Omega)$, then we reason by density. We take $(\varphi_\epsilon)_{\epsilon>0} \subset W^{2,p}(\Omega)$ that converges to φ in $W^{1,p}(\Omega)$ as $\epsilon \rightarrow 0$ and we write, using the commuting property (4.4) of $G_T^k \mathbb{I}_T^k$ and the L^p -stability of the L^2 -projector stated in Lemma 3.2,

$$\begin{aligned} \|G_T^k \mathbb{I}_T^k \varphi - \nabla \varphi\|_{L^p(\Omega)^d} &\leq \|G_T^k \mathbb{I}_T^k (\varphi - \varphi_\epsilon)\|_{L^p(\Omega)^d} + \|G_T^k \mathbb{I}_T^k \varphi_\epsilon - \varphi_\epsilon\|_{L^p(\Omega)^d} + \|\nabla (\varphi_\epsilon - \varphi)\|_{L^p(\Omega)^d} \\ &\lesssim \|\nabla (\varphi - \varphi_\epsilon)\|_{L^p(\Omega)^d} + \|G_T^k \mathbb{I}_T^k \varphi_\epsilon - \varphi_\epsilon\|_{L^p(\Omega)^d}. \end{aligned}$$

By (5.22), the second term in this right-hand side tends to 0 as $h \rightarrow 0$. Taking (in that order) the supremum limit as $h \rightarrow 0$ and then the supremum limit as $\epsilon \rightarrow 0$ concludes the proof that $G_T^k \mathbb{1}_T^k \varphi \rightarrow \nabla \varphi$ in $L^p(\Omega)^d$.

Step 2: Proof of (5.21). It is proved in [29, Eq. (46)] that

$$h_F^{-\frac{1}{2}} \|\pi_F^k((\mathbb{1}_T^k \varphi)_F - P_T^{k+1} \mathbb{1}_T^k \varphi)\|_{L^2(F)} \lesssim h_T^{k+1} \|\varphi\|_{H^{k+2}(T)}.$$

Using Lemma 5.1, the admissibility of the mesh (which gives $h_F|F| \approx |T|$ if $F \in \mathcal{F}_T$), and the regularity assumption on φ , we infer

$$\begin{aligned} h_F^{1-p} \|\pi_F^k((\mathbb{1}_T^k \varphi)_F - P_T^{k+1} \mathbb{1}_T^k \varphi)\|_{L^p(F)}^p &\lesssim h_F^{1-\frac{p}{2}} |F|^{1-\frac{p}{2}} \left(h_F^{-\frac{1}{2}} \|\pi_F^k((\mathbb{1}_T^k \varphi)_F - P_T^{k+1} \mathbb{1}_T^k \varphi)\|_{L^2(F)} \right)^p \\ &\lesssim (h_F|F|)^{1-\frac{p}{2}} h_T^{(k+1)p} \|\varphi\|_{H^{k+2}(T)}^p \\ &\lesssim |T|^{1-\frac{p}{2}} h_T^{(k+1)p} |T|^{\frac{p}{2}} \|\varphi\|_{W^{k+2,\infty}(T)}^p \\ &\lesssim |T| h^{(k+1)p} \|\varphi\|_{W^{k+2,\infty}(\mathcal{T}_h)}^p. \end{aligned}$$

Summing this over $F \in \mathcal{F}_T$ and $T \in \mathcal{T}_h$, and recalling the uniform bound (3.2) over $\text{card}(\mathcal{F}_T)$, we get

$$\sum_{T \in \mathcal{T}_h} s_T(\mathbb{1}_T^k \varphi, \mathbb{1}_T^k \varphi) \lesssim |\Omega| h^{(k+1)p} \|\varphi\|_{W^{k+2,\infty}(\mathcal{T}_h)}^p,$$

and the proof is complete. \square

6 Convergence analysis

The following proposition contains an *a priori* estimate, uniform in h , on the solution to the discrete problem (4.11).

Proposition 6.1 (A priori estimates). *Under Assumption 3.1, if $\underline{\mathbf{u}}_h \in \underline{\mathbf{U}}_{h,0}^k$ solves (4.11) then there exists C only depending on Ω , $\lambda_{\mathbf{a}}$, ϱ , k and p such that*

$$\|\underline{\mathbf{u}}_h\|_{1,p,h} \leq C \|f\|_{L^{p'}(\Omega)}^{\frac{1}{p-1}}. \quad (6.1)$$

Proof. We write $A \lesssim B$ for $A \leq MB$ with M having the same dependencies as C in the proposition. Plugging $\underline{\mathbf{v}}_h = \underline{\mathbf{u}}_h$ into (4.11a) and using the coercivity (2.2d) of \mathbf{a} leads to

$$\lambda_{\mathbf{a}} \sum_{T \in \mathcal{T}_h} \|G_T^k \underline{\mathbf{u}}_T\|_{L^p(T)^d}^p + \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} h_F^{1-p} \|\pi_F^k(\mathbf{u}_F - P_T^{k+1} \underline{\mathbf{u}}_T)\|_{L^p(F)}^p \leq \|f\|_{L^{p'}(\Omega)} \|\underline{\mathbf{u}}_h\|_{L^p(\Omega)}.$$

Owing to the norm equivalence (5.5), and using the discrete Sobolev embeddings (5.8) with $q = p$ to estimate the second factor, this gives

$$\|\underline{\mathbf{u}}_h\|_{1,p,h}^p \lesssim \|f\|_{L^{p'}(\Omega)} \|\underline{\mathbf{u}}_h\|_{L^p(\Omega)} \lesssim \|f\|_{L^{p'}(\Omega)} \|\underline{\mathbf{u}}_h\|_{1,p,h},$$

which concludes the proof since, by assumption, $p > 1$. \square

We can now prove that the discrete problem (4.11) has at least one solution.

Proof of Theorem 4.4. We use [25, Theorem 3.3] (see also [46]): If $(E, \langle \cdot, \cdot \rangle_E, \|\cdot\|_E)$ is an Euclidean space, and $\Phi : E \mapsto E$ is continuous and satisfies $\frac{\langle \Phi(x), x \rangle_E}{\|x\|_E} \rightarrow +\infty$ as $\|x\|_E \rightarrow +\infty$, then Φ is onto.

We take $E = \underline{\mathbf{U}}_{h,0}^k$, endowed with an arbitrary inner product, and define $\Phi : \underline{\mathbf{U}}_{h,0}^k \rightarrow \underline{\mathbf{U}}_{h,0}^k$ by

$$\forall \underline{\mathbf{v}}_h, \underline{\mathbf{w}}_h \in \underline{\mathbf{U}}_{h,0}^k, \quad \langle \Phi(\underline{\mathbf{v}}_h), \underline{\mathbf{w}}_h \rangle_E = A(\underline{\mathbf{v}}_h, \underline{\mathbf{w}}_h).$$

Assumptions (2.2a) and (2.2b) show that Φ is continuous, and the coercivity (2.2d) of \mathbf{a} together with the norm equivalence (5.5) show that

$$\langle \Phi(\underline{\mathbf{v}}_h), \underline{\mathbf{v}}_h \rangle_E \geq C \|\underline{\mathbf{v}}_h\|_{1,p,h}^p \geq C_{\mathcal{T}_h} \|\underline{\mathbf{v}}_h\|_E^p,$$

where $C_{\mathcal{T}_h} > 0$ may depend on \mathcal{T}_h but does not depend on $\underline{\mathbf{v}}_h$ (we use the equivalence of all norms on $\underline{\mathbf{U}}_{h,0}^k$). Hence, Φ is onto. Let now $\underline{\mathbf{y}}_h \in \underline{\mathbf{U}}_{h,0}^k$ be such that

$$\langle \underline{\mathbf{y}}_h, \underline{\mathbf{w}}_h \rangle_E = \int_{\Omega} f(\mathbf{x}) \mathbf{w}_h(\mathbf{x}) d\mathbf{x} \quad \forall \underline{\mathbf{w}}_h \in \underline{\mathbf{U}}_{h,0}^k,$$

and take $\underline{\mathbf{u}}_h \in \underline{\mathbf{U}}_{h,0}^k$ such that $\Phi(\underline{\mathbf{u}}_h) = \underline{\mathbf{y}}_h$. By definition of Φ and $\underline{\mathbf{y}}_h$, $\underline{\mathbf{u}}_h$ is a solution to the discrete problem (4.11). \square

Let us now turn to the proof of convergence. To improve the legibility of certain formula, we drop the variable \mathbf{x} in the integral signs.

Proof of Theorem 4.5. Step 1: Existence of a limit. By Propositions 6.1 and 5.9, there exists $u \in W_0^{1,p}(\Omega)$ such that up to a subsequence as $h \rightarrow 0$, $\mathbf{u}_h \rightarrow u$ and $p_h^{k+1} \underline{\mathbf{u}}_h \rightarrow u$ in $L^q(\Omega)$ for all $q < p^*$, and $G_h^k \underline{\mathbf{u}}_h \rightarrow \nabla u$ weakly in $L^p(\Omega)^d$. Let us prove that u solves (2.3). To this end, we adapt Minty's technique [46, 49] to the discrete setting, as previously done in [32, 37].

Step 2: Identification of the limit. The growth assumption (2.2b) on \mathbf{a} ensures that $\mathbf{a}(\cdot, \mathbf{u}_h, G_h^k \underline{\mathbf{u}}_h)$ is bounded in $L^p(\Omega)^d$, and converges therefore (upon extracting another subsequence) to some χ weakly in this space. Let $\varphi \in C_c^\infty(\Omega)$. Plugging $\underline{\mathbf{v}}_h = \mathbb{1}_h^k \varphi$ into (4.11) gives

$$\int_{\Omega} \mathbf{a}(\mathbf{x}, \mathbf{u}_h, G_h^k \underline{\mathbf{u}}_h) \cdot G_h^k \mathbb{1}_h^k \varphi = \int_{\Omega} f \pi_h^k \varphi - \sum_{T \in \mathcal{T}_h} s_T(\underline{\mathbf{u}}_T, \mathbb{1}_T^k \varphi), \quad (6.2)$$

with π_h^k denoting the L^2 -projector on the broken polynomial space $\mathbb{P}^k(\mathcal{T}_h)$. Using Hölder's inequality followed by the norm equivalence (5.5) to bound the first factor, we infer

$$\begin{aligned} \left| \sum_{T \in \mathcal{T}_h} s_T(\underline{\mathbf{u}}_T, \mathbb{1}_T^k \varphi) \right| &\leq \left(\sum_{T \in \mathcal{T}_h} s_T(\underline{\mathbf{u}}_T, \underline{\mathbf{u}}_T) \right)^{\frac{1}{p'}} \left(\sum_{T \in \mathcal{T}_h} s_T(\mathbb{1}_T^k \varphi, \mathbb{1}_T^k \varphi) \right)^{\frac{1}{p}} \\ &\leq \|\underline{\mathbf{u}}_h\|_{1,p,h}^{\frac{1}{p'}} \left(\sum_{T \in \mathcal{T}_h} s_T(\mathbb{1}_T^k \varphi, \mathbb{1}_T^k \varphi) \right)^{\frac{1}{p}}. \end{aligned}$$

Recalling the a priori bound (6.1) on the exact solution and the strong convergence property (5.21), we see that this quantity tends to 0 as $h \rightarrow 0$. Additionally, by the approximation properties of the L^2 -projector stated in Lemma 3.4 together with the strong convergence property (5.20), we have $\pi_h^k \varphi \rightarrow \varphi$ in $L^p(\Omega)$ and $G_h^k \mathbb{1}_h^k \varphi \rightarrow \nabla \varphi$ in $L^p(\Omega)^d$. We can therefore pass to the limit $h \rightarrow 0$ in (6.2) and we find

$$\int_{\Omega} \chi \cdot \nabla \varphi = \int_{\Omega} f \varphi. \quad (6.3)$$

By density of $C_c^\infty(\Omega)$ in $W_0^{1,p}(\Omega)$, this relation still holds if $\varphi \in W_0^{1,p}(\Omega)$.

Let us now take $\Lambda \in L^p(\Omega)^d$ and write, using the monotonicity (2.2c) of \mathbf{a} ,

$$\int_{\Omega} [\mathbf{a}(\mathbf{x}, \mathbf{u}_h, G_h^k \underline{\mathbf{u}}_h) - \mathbf{a}(\mathbf{x}, \mathbf{u}_h, \Lambda)] \cdot [G_h^k \underline{\mathbf{u}}_h - \Lambda] \geq 0. \quad (6.4)$$

Use (4.11) and $s_T(\underline{\mathbf{u}}_T, \underline{\mathbf{u}}_T) \geq 0$ to write

$$\int_{\Omega} \mathbf{a}(\mathbf{x}, \mathbf{u}_h, G_h^k \underline{\mathbf{u}}_h) \cdot G_h^k \underline{\mathbf{u}}_h = \int_{\Omega} f \mathbf{u}_h - \sum_{T \in \mathcal{T}_h} s_T(\underline{\mathbf{u}}_T, \underline{\mathbf{u}}_T) \leq \int_{\Omega} f \mathbf{u}_h. \quad (6.5)$$

Develop (6.4) and plug this relation:

$$\int_{\Omega} f u_h - \int_{\Omega} \mathbf{a}(\mathbf{x}, \mathbf{u}_h, G_h^k \underline{\mathbf{u}}_h) \cdot \boldsymbol{\Lambda} \geq \int_{\Omega} \mathbf{a}(\mathbf{x}, \mathbf{u}_h, \boldsymbol{\Lambda}) \cdot [G_h^k \underline{\mathbf{u}}_h - \boldsymbol{\Lambda}]. \quad (6.6)$$

Since $\mathbf{u}_h \rightarrow u$ in $L^q(\Omega)$ for all $q < p^*$, the Caratheodory and growth properties (2.2a) and (2.2b) of \mathbf{a} show that $\mathbf{a}(\mathbf{x}, \mathbf{u}_h, \boldsymbol{\Lambda}) \rightarrow \mathbf{a}(\mathbf{x}, u, \boldsymbol{\Lambda})$ strongly in $L^{p'}(\Omega)^d$. We can therefore pass to the limit in (6.6):

$$\int_{\Omega} f u - \int_{\Omega} \boldsymbol{\chi} \cdot \boldsymbol{\Lambda} \geq \int_{\Omega} \mathbf{a}(\mathbf{x}, u, \boldsymbol{\Lambda}) \cdot [\boldsymbol{\nabla} u - \boldsymbol{\Lambda}]. \quad (6.7)$$

The conclusion then follows classically [46, 49]: take $v \in W_0^{1,p}(\Omega)$, apply this relation to $\boldsymbol{\Lambda} = \boldsymbol{\nabla} u \pm t \boldsymbol{\nabla} v$ for some $t > 0$, use (6.3) with $\varphi = u \pm tv$, divide by t and let $t \rightarrow 0$ using the Caratheodory and growth properties of \mathbf{a} . This leads to

$$\int_{\Omega} f v = \int_{\Omega} \mathbf{a}(\mathbf{x}, u, \boldsymbol{\nabla} u) \cdot \boldsymbol{\nabla} v,$$

and the proof that u solves (2.3) is complete.

Step 3: Convergence of the gradient. It remains to show that if \mathbf{a} is strictly monotone, then $G_h^k \underline{\mathbf{u}}_h \rightarrow \boldsymbol{\nabla} u$ strongly in $L^p(\Omega)^d$. Let

$$F_h = [\mathbf{a}(\mathbf{x}, \mathbf{u}_h, G_h^k \underline{\mathbf{u}}_h) - \mathbf{a}(\mathbf{x}, \mathbf{u}_h, \boldsymbol{\nabla} u)] \cdot [G_h^k \underline{\mathbf{u}}_h - \boldsymbol{\nabla} u] \geq 0 \quad (6.8)$$

Developing this expression and using (6.5), we can pass to the limit and use (6.3) to see that

$$\limsup_{h \rightarrow 0} \int_{\Omega} F_h \leq \int_{\Omega} f u - \int_{\Omega} \boldsymbol{\chi} \cdot \boldsymbol{\nabla} u = 0.$$

Hence, $F_h \rightarrow 0$ in $L^1(\Omega)$. Up to a subsequence, it therefore converges almost everywhere. Using the coercivity and growth assumptions (2.2d) and (2.2b) of \mathbf{a} , Young's inequality gives

$$\begin{aligned} F_h &\geq \lambda_{\mathbf{a}} |G_h^k \underline{\mathbf{u}}_h|^p - (\bar{a}(\mathbf{x}) + \beta_{\mathbf{a}} |\mathbf{u}_h|^r + \beta_{\mathbf{a}} |G_h^k \underline{\mathbf{u}}_h|^{p-1}) |\boldsymbol{\nabla} u| - (\bar{a}(\mathbf{x}) + \beta_{\mathbf{a}} |\mathbf{u}_h|^r + \beta_{\mathbf{a}} |\boldsymbol{\nabla} u|^{p-1}) |\boldsymbol{\nabla} u| \\ &\geq \frac{\lambda_{\mathbf{a}}}{2} |G_h^k \underline{\mathbf{u}}_h|^p - 2(\bar{a}(\mathbf{x}) + \beta_{\mathbf{a}} |\mathbf{u}_h|^r) |\boldsymbol{\nabla} u| - \beta_{\mathbf{a}} |\boldsymbol{\nabla} u|^p - \frac{\beta_{\mathbf{a}}^p}{p} \left(\frac{2}{p' \lambda_{\mathbf{a}}} \right)^{p-1} |\boldsymbol{\nabla} u|^p. \end{aligned} \quad (6.9)$$

Since, up to a subsequence, \mathbf{u}_h converges a.e., this relation shows that for a.e. \mathbf{x} , the sequence $(G_h^k \underline{\mathbf{u}}_h(\mathbf{x}))_{h \in \mathcal{H}}$ remains bounded. Let us show that it can only have $\boldsymbol{\nabla} u(\mathbf{x})$ as adherence value. If ζ is an adherence value of $(G_h^k \underline{\mathbf{u}}_h(\mathbf{x}))_{h \in \mathcal{H}}$, then, passing to the limit in (6.8) gives, since $F_h \rightarrow 0$ and $\mathbf{u}_h \rightarrow u$ a.e.,

$$[\mathbf{a}(\mathbf{x}, u(\mathbf{x}), \zeta) - \mathbf{a}(\mathbf{x}, u(\mathbf{x}), \boldsymbol{\nabla} u(\mathbf{x}))] \cdot [\zeta - \boldsymbol{\nabla} u(\mathbf{x})] = 0.$$

The strict monotony of \mathbf{a} then shows that $\zeta = \boldsymbol{\nabla} u(\mathbf{x})$. Hence, for a.e. \mathbf{x} , the bounded sequence $(G_h^k \underline{\mathbf{u}}_h(\mathbf{x}))_{h \in \mathcal{H}}$ has only $\boldsymbol{\nabla} u(\mathbf{x})$ as adherence value, and thus $G_h^k \underline{\mathbf{u}}_h \rightarrow \boldsymbol{\nabla} u$ a.e. on Ω .

Since $(F_h)_{h \in \mathcal{H}}$ is 1-equi-integrable (it converges in $L^1(\Omega)$) and $(|\mathbf{u}_h|^r)_{h \in \mathcal{H}}$ is p' -equi-integrable ($p'r < p^*$ and $(\mathbf{u}_h)_{h \in \mathcal{H}}$ therefore converges in $L^{p'r}(\Omega)$), (6.9) shows that $(G_h^k \underline{\mathbf{u}}_h)_{h \in \mathcal{H}}$ is p -equi-integrable. Vitali's theorem then gives the strong convergence of this sequence to $\boldsymbol{\nabla} u$ in $L^p(\Omega)^d$. \square

7 Other boundary conditions

We briefly discuss here how the HHO scheme is written for non-homogeneous Dirichlet and homogeneous Neumann boundary conditions.

7.1 Non-homogeneous Dirichlet boundary conditions

Non-homogeneous Dirichlet boundary conditions consist in replacing (1.1b) with

$$u = g \text{ on } \partial\Omega \quad (7.1)$$

with $g \in W^{1-\frac{1}{p},p}(\partial\Omega)$. Denoting by $\gamma : W^{1,p}(\Omega) \mapsto W^{1-\frac{1}{p},p}(\partial\Omega)$ the trace operator, the weak formulation becomes:

$$\begin{aligned} &\text{Find } u \in W^{1,p}(\Omega) \text{ such that } \gamma(u) = g \text{ and, for all } v \in W_0^{1,p}(\Omega), \\ &\int_{\Omega} \mathbf{a}(\mathbf{x}, u(\mathbf{x}), \nabla u(\mathbf{x})) \cdot \nabla v(\mathbf{x}) \, d\mathbf{x} = \int_{\Omega} f(\mathbf{x})v(\mathbf{x}) \, d\mathbf{x}. \end{aligned} \quad (7.2)$$

As in Remark 4.1 we notice that $\pi_F^k g$ is well defined for any $F \in \mathcal{F}_h^b$. Hence, we can define the vector $\underline{u}_{g,h} \in \underline{U}_h^k$ such that

$$\underline{u}_{g,T} = 0 \quad \forall T \in \mathcal{T}_h, \quad \underline{u}_{g,F} = 0 \quad \forall F \in \mathcal{F}_h^i, \quad \underline{u}_{g,F} = \pi_F^k g \quad \forall F \in \mathcal{F}_h^b.$$

We then set

$$\underline{U}_{h,g}^k := \underline{U}_{h,0}^k + \underline{u}_{g,h},$$

and write the discrete problem corresponding to (7.2) as

$$\text{Find } \underline{u}_h \in \underline{U}_{h,g}^k \text{ such that, for any } \underline{v}_h \in \underline{U}_{h,0}^k, A(\underline{u}_h, \underline{v}_h) = \int_{\Omega} f \underline{v}_h, \quad (7.3)$$

with A defined by (4.11b)–(4.11c). The convergence analysis for non-homogeneous Dirichlet boundary conditions is performed as usual by utilizing a lifting of the boundary conditions. We take $\tilde{g} \in W^{1,p}(\Omega)$ and let $\underline{g}_h = \mathbb{I}_h^k \tilde{g}$. Making $\underline{v}_h = \underline{u}_h - \underline{g}_h \in \underline{U}_h^k$ in (7.6) and using $\|\underline{g}_h\|_{1,p,h} \lesssim \|g\|_{W^{1,p}(\Omega)}$ (see Proposition 7.1 below) enables us to prove *a priori* estimates on $\|\underline{u}_h - \underline{g}_h\|_{1,p,h}$.

Proposition 5.11 does not rely on the homogeneous boundary conditions and therefore shows that $G_T^k \underline{g}_h \rightarrow \nabla \tilde{g}$ in $L^p(\Omega)^d$ as $h \rightarrow 0$. Since $\pi_h^k \tilde{g} \rightarrow \tilde{g}$ in $L^p(\Omega)$ (see Lemma 3.4), applying Proposition 5.9 to $\underline{v}_h = \underline{u}_h - \underline{g}_h$ shows that, for some $u \in W^{1,p}(\Omega)$ such that $u - \tilde{g} \in W_0^{1,p}(\Omega)$ (i.e. $\gamma(u) = g$), up to a subsequence $\underline{u}_h \rightarrow u$ in $L^p(\Omega)$ and $G_T^k \underline{u}_h \rightarrow \nabla u$ in $L^p(\Omega)^d$ as $h \rightarrow 0$. The proof that u is a solution to (7.2) is then done in a similar way as for homogeneous boundary conditions.

Proposition 7.1 (Discrete norm estimate for interpolate of $W^{1,p}$ functions). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence, and let $k \in \mathbb{N}$. Let $v \in W^{1,p}(\Omega)$ and let $\mathbb{I}_h^k v \in \underline{U}_h^k$ be the interpolant defined by (4.7) and (4.2). Then $\|\mathbb{I}_T^k v\|_{1,p,T} \lesssim \|v\|_{W^{1,p}(T)}$ and thus $\|\mathbb{I}_h^k v\|_{1,p,h} \lesssim \|v\|_{W^{1,p}(\Omega)}$.*

Proof. Let $\underline{v}_h = \mathbb{I}_h^k v$. Since $\mathbf{v}_T = \pi_T^k v$, Corollary 3.7 with $s = 1$ shows that $\|\nabla \mathbf{v}_T\|_{L^p(T)} \lesssim \|v\|_{W^{1,p}(T)}$. This takes care of the first term in $\|\underline{v}_h\|_{1,p,T}$. To deal with the second term, we use Lemma 3.2 with $U = F$ and then Lemma 3.6 with $m = 0$ and $s = 1$ to write

$$\|\mathbf{v}_F - \mathbf{v}_T\|_{L^p(F)} = \|\pi_F^k v - \pi_T^k v\|_{L^p(F)} = \|\pi_F^k (v - \pi_T^k v)\|_{L^p(F)} \lesssim \|v - \pi_T^k v\|_{L^p(F)} \lesssim h_T^{1-\frac{1}{p}} \|v\|_{W^{1,p}(T)}.$$

Raising this to the power p and using $h_T \lesssim h_F$ gives $h_F^{1-p} \|\mathbf{v}_F - \mathbf{v}_T\|_{L^p(F)}^p \lesssim \|v\|_{W^{1,p}(T)}^p$. \square

7.2 Homogeneous Neumann boundary conditions

We assume that

$$\int_{\Omega} f(\mathbf{x}) \, d\mathbf{x} = 0.$$

Homogeneous Neumann boundary conditions for elliptic Leray–Lions problems consist in replacing (1.1b) with

$$\mathbf{a}(\cdot, u, \nabla u) \cdot \mathbf{n} = 0 \text{ on } \partial\Omega, \quad (7.4)$$

where \mathbf{n} is the outer normal to $\partial\Omega$. The weak formulation of (1.1a)–(7.4) is

$$\begin{aligned} &\text{Find } u \in W^{1,p}(\Omega) \text{ such that } \int_{\Omega} u(\mathbf{x}) \, d\mathbf{x} = 0 \text{ and, for all } v \in W^{1,p}(\Omega), \\ &\int_{\Omega} \mathbf{a}(\mathbf{x}, u(\mathbf{x}), \nabla u(\mathbf{x})) \cdot \nabla v(\mathbf{x}) \, d\mathbf{x} = \int_{\Omega} f(\mathbf{x})v(\mathbf{x}) \, d\mathbf{x}. \end{aligned} \quad (7.5)$$

The HHO scheme for (7.5) reads

$$\text{Find } \underline{u}_h \in \underline{U}_h^k \text{ such that } \int_{\Omega} \underline{u}_h(\mathbf{x}) \, d\mathbf{x} = 0 \text{ and, for any } \underline{v}_h \in \underline{U}_h^k, A(\underline{u}_h, \underline{v}_h) = \int_{\Omega} f \underline{v}_h \quad (7.6)$$

with A still defined by (4.11b)–(4.11c).

To carry out the convergence analysis from Section 6, we need a few results. The first one is a discrete Poincaré–Wirtinger–Sobolev inequality, which bounds to the L^{p^*} -norm of discrete functions by their discrete norm. This immediately gives *a priori* estimates on the solution to the scheme (Proposition 6.1). The second result is a discrete Rellich theorem for functions with zero average and bounded discrete norm (this is the equivalent of Proposition 5.9). The proofs of both results are based on Lemma 5.6 and on a decomposition of functions in \underline{U}_h^k into low-order (piecewise-constant) vectors in \underline{U}_h^0 , and their higher order variation.

Lemma 7.2 (Discrete Poincaré–Wirtinger–Sobolev inequality for broken polynomial functions with zero global average). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence. Then, there exists C only depending on Ω , ϱ , k and p such that, for all $\underline{v}_h \in \underline{U}_h^k$ satisfying $\int_{\Omega} \underline{v}_h(\mathbf{x}) \, d\mathbf{x} = 0$, we have, recalling the definition (2.1) of the Sobolev index p^* ,*

$$\|\underline{v}_h\|_{L^{p^*}(\Omega)} \leq C \|\underline{v}_h\|_{1,h,p}. \quad (7.7)$$

Proof. Here, $A \lesssim B$ means that $A \leq MB$ with M only depending on Ω , ϱ , k and p . We define $\underline{v}_h^0 \in \underline{U}_h^0$ and $\underline{v}_h^1 \in \mathbb{P}^k(\mathcal{T}_h)$ by:

$$\begin{aligned} \underline{v}_T^0 &= \pi_T^0 \underline{v}_T \quad \forall T \in \mathcal{T}_h, & \underline{v}_F^0 &= \pi_F^0 \underline{v}_F \quad \forall F \in \mathcal{F}_h, \\ \underline{v}_T^1 &= \underline{v}_T - \pi_T^0 \underline{v}_T = \underline{v}_T - \underline{v}_T^0 \quad \forall T \in \mathcal{T}_h. \end{aligned}$$

By Lemma 5.6 we have

$$\|\underline{v}_h^1\|_{L^{p^*}(\Omega)} \lesssim \left(\sum_{T \in \mathcal{T}_h} \|\nabla \underline{v}_T\|_{L^p(T)}^p \right)^{1/p}. \quad (7.8)$$

We recall the definition of the discrete $W^{1,p}$ -norm on \underline{U}_h^0 from [35]:

$$\|\underline{v}_h^0\|_{W^{1,p}, \mathcal{T}_h} = \left(\sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} |T| \left| \frac{\underline{v}_T^0 - \underline{v}_F^0}{h_T} \right|^p \right)^{\frac{1}{p}}$$

(the genuine discrete $W^{1,p}$ -norm in [35] involves a different coefficient than $|T|$ in this sum, but under Assumption 3.1 this coefficient is $\approx |T|$). Since $\sum_{T \in \mathcal{T}_h} |T| \underline{v}_T^0 = \int_{\Omega} \underline{v}_h(\mathbf{x}) \, d\mathbf{x} = 0$, [35] gives

$$\|\underline{v}_h^0\|_{L^{p^*}(\Omega)} \lesssim \|\underline{v}_h^0\|_{W^{1,p}, \mathcal{T}_h}. \quad (7.9)$$

By noticing that $\underline{v}_h = \underline{v}_h^0 + \underline{v}_h^1$, the result follows from (7.8) and (7.9) provided that

$$\|\underline{v}_h^0\|_{W^{1,p}, \mathcal{T}_h} \lesssim \|\underline{v}_h\|_{1,h,p}. \quad (7.10)$$

An easy generalisation of [33, Lemma 6.3] and [34, Lemma 6.6] (see [35] for details) shows that

$$\left| \pi_F^0 \underline{v}_T - \pi_T^0 \underline{v}_T \right|^p = \left| \frac{1}{|F|} \int_F \underline{v}_T(\mathbf{x}) \, ds(\mathbf{x}) - \frac{1}{|T|} \int_T \underline{v}_T(\mathbf{x}) \, d\mathbf{x} \right|^p \lesssim \frac{h_T^p}{|T|} \int_T |\nabla \underline{v}_T(\mathbf{x})|^p \, d\mathbf{x}.$$

Using the triangular and Jensen's inequalities, and the relations $|T| \lesssim |F|h_F$ and $h_F \leq h_T$, we infer

$$\begin{aligned} |\mathbf{v}_F^0 - \mathbf{v}_T^0|^p &\lesssim |\pi_F^0 \mathbf{v}_F - \pi_F^0 \mathbf{v}_T|^p + \frac{h_T^p}{|T|} \int_T |\nabla \mathbf{v}_T(\mathbf{x})|^p d\mathbf{x} \\ &\lesssim \frac{1}{|F|} \int_F |\mathbf{v}_F(\mathbf{x}) - \mathbf{v}_T(\mathbf{x})|^p ds(\mathbf{x}) + \frac{h_T^p}{|T|} \|\nabla \mathbf{v}_T\|_{L^p(T)^d}^p \\ &\lesssim \frac{h_T^p}{|T|} h_F^{1-p} \|\mathbf{v}_F - \mathbf{v}_T\|_{L^p(F)}^p + \frac{h_T^p}{|T|} \|\nabla \mathbf{v}_T\|_{L^p(T)^d}^p. \end{aligned}$$

Multiplying by $\frac{|T|}{h_F^p}$ and summing over $F \in \mathcal{F}_T$ and $T \in \mathcal{T}_h$ gives (7.10). \square

Proposition 7.3 (Compactness result for broken polynomial function with zero global average). *Let $(\mathcal{T}_h)_{h \in \mathcal{H}}$ be an admissible mesh sequence and let $\mathbf{v}_T \in \mathcal{U}_h^k$ be such that $(\|\mathbf{v}_h\|_{1,h,p})_{h \in \mathcal{H}}$ is bounded and, for all $h \in \mathcal{H}$, $\int_\Omega \mathbf{v}_h(\mathbf{x}) d\mathbf{x} = 0$. Then, there exists $v \in W^{1,p}(\Omega)$ such that $\int_\Omega v(\mathbf{x}) d\mathbf{x} = 0$ and, up to a subsequence as $h \rightarrow 0$, recalling the definition (2.1) of the Sobolev index p^* ,*

- $\mathbf{v}_h \rightarrow v$ and $p_h^{k+1} \mathbf{v}_h \rightarrow v$ strongly in $L^q(\Omega)$ for all $q < p^*$,
- $G_h^k \mathbf{v}_h \rightarrow \nabla v$ weakly in $L^p(\Omega)^d$.

Proof. We use the same decomposition $\mathbf{v}_h = \mathbf{v}_h^0 + \mathbf{v}_h^1$ as in the proof of Lemma 7.2. By Lemma 5.6 we have $\|\mathbf{v}_h^1\|_{L^q(\Omega)} \leq Ch^\theta \|\mathbf{v}_h\|_{1,h,p}$ where C does not depend on h and $\theta = 1 + \frac{d}{q} - \frac{d}{p} > 0$. Hence, $\mathbf{v}_h^1 \rightarrow 0$ in $L^q(\Omega)$ as $h \rightarrow 0$. By (7.10), $(\|\mathbf{v}_h^0\|_{W^{1,p}(\mathcal{T}_h)})_{h \in \mathcal{H}}$ remains bounded. Since $\sum_{T \in \mathcal{T}_h} |T| \mathbf{v}_h^0 = 0$ for all $h \in \mathcal{H}$, the discrete compactness result for Neumann boundary conditions of [35] shows that there exists a $v \in W^{1,p}(\Omega)$ with zero average such that $\mathbf{v}_h^0 \rightarrow v$ strongly in $L^q(\Omega)$ up to a subsequence. Hence, $\mathbf{v}_h \rightarrow v$ in $L^q(\Omega)$ along the same subsequence. We then apply Corollary 5.8, which is independent of the boundary conditions, to deduce that $p_h^{k+1} \mathbf{v}_h \rightarrow v$ in $L^q(\Omega)$.

To prove that $G_T^h \mathbf{v}_h \rightarrow \nabla v$ weakly in $L^p(\Omega)^d$, we notice that by Lemma 5.2 the functions $G_T^h \mathbf{v}_h$ remain bounded in $L^p(\Omega)^d$ and therefore converge weakly to some \mathcal{G} in this space. We prove that $\mathcal{G} = \nabla v$ as in the proof of Proposition 5.9, using test functions $\phi \in C_c^\infty(\Omega)^d$ instead of $\phi \in C^\infty(\mathbb{R}^d)^d$. \square

8 Conclusion

We extended the HHO method of [29] to fully non-linear Leray–Lions equations, which include the p -Laplace model. This method can be interpreted as a higher-order extension of the classical Mimetic Finite Differences, Hybrid Finite Volumes and Mixed Finite Volumes schemes. We proved the convergence of the HHO method without assuming unrealistic regularity properties on the solution, or restrictive assumptions on the non-linear operator. To establish this convergence, we developed discrete functional analysis results that include the analysis of L^p - and $W^{s,p}$ -stability and approximation properties of L^2 -projectors on broken polynomial spaces. We provided numerical results which demonstrate the good approximation properties of the method on a variety of meshes, and for various orders (low as well as high).

A Discrete functional analysis in local polynomial spaces

This appendix collects discrete functional analysis results in local polynomial spaces that are of general interest for the analysis of polynomial-based methods for linear and nonlinear problems. Most of these results have already been stated without proof in the paper, but we restate them for the sake of easy consultation.

A.1 Estimates in local polynomial spaces

This section collects L^p - and $W^{s,p}$ -estimates in local polynomial spaces including direct and reverse Sobolev and Lebesgue embeddings.

Lemma 5.1 (Direct and reverse Lebesgue embeddings). *Let U be a measurable subset of \mathbb{R}^N such that (3.5) holds. Let $k \in \mathbb{N}$ and $q, m \in [1, +\infty]$. Then,*

$$\forall w \in \mathbb{P}^k(U) : \|w\|_{L^q(U)} \approx |U|^{\frac{1}{q} - \frac{1}{m}} \|w\|_{L^m(U)}, \quad (5.3)$$

where $A \approx B$ means that there is a real $M > 0$ only depending on N, k, δ, q and m such that $M^{-1}A \leq B \leq MA$.

Remark A.1 (Reverse embeddings). *If $q \leq m$ then this result is a classical (direct) Lebesgue embedding due to Hölder's inequality. It holds for $m < q$ solely because we consider polynomials (and we notice that the scaling $|U|^{\frac{1}{q} - \frac{1}{m}}$ explodes as $h_U \rightarrow 0$).*

Remark A.2 (Sobolev reverse embeddings). *Let U be a polyhedral set that admits a simplicial decomposition such that for any simplex S , if h_S is the diameter of S and r_S its inradius then $h_S \leq \varrho r_S$, and $h_U \leq \varrho h_S$. The following inverse inequality holds with C_{inv} depending on ϱ, k and p , but independent of h (cf. [28, Lemma 1.44] for the case $p = 2$ and use [28, Lemma 1.50] or Lemma 5.1 to deduce the general case),*

$$\forall v \in \mathbb{P}^k(U) : \|\nabla v\|_{L^p(U)} \leq C_{\text{inv}} h_U^{-1} \|v\|_{L^p(U)}. \quad (A.1)$$

Using this inequality we can easily deduce from Lemma 5.1 the following reverse Sobolev embeddings: Under the assumptions of Lemma 5.1, if U is open and $m \geq r$, then for all $w \in \mathbb{P}^k(U)$ we have

$$|w|_{W^{m,p}(U)} \lesssim h_U^{r-m} |U|^{\frac{1}{p} - \frac{1}{q}} |w|_{W^{r,q}(U)}.$$

Here \lesssim is up to a multiplicative constant only depending on k, δ, p, q and r . Note that the result obviously cannot hold if $m < r$ and $m \leq k$ (consider w polynomial of degree exactly m : the left-hand side does not vanish, but the right-hand side vanishes).

Proof of Lemma 5.1. We obviously only have to prove \lesssim since m and q play symmetrical roles in (5.3). By (3.5), there is $\mathbf{x}_U \in U$ such that $B(\mathbf{x}_U, \delta h_U) \subset U \subset B(\mathbf{x}_U, h_U)$. Let $U_0 = (U - \mathbf{x}_U)/h_U$. Using the change of variable $\mathbf{x} \in U \mapsto (\mathbf{x} - \mathbf{x}_U)/h_U \in U_0$, we see that, for $\ell \in [1, +\infty]$,

$$\|w\|_{L^\ell(U)} = h_U^{\frac{N}{\ell}} \|w_0\|_{L^\ell(U_0)} \approx |U|^{\frac{1}{\ell}} \|w_0\|_{L^\ell(U_0)}, \quad (A.2)$$

where we used $h_U^N \approx |U|$ (since $h_U \approx r_U$) and we set $w_0(\mathbf{y}) = w(\mathbf{x}_U + h_U \mathbf{y})$. Assume that there exists C_0 not depending on the geometry of U_0 but solely on δ such that

$$\forall v \in \mathbb{P}^k(U_0) : \|v\|_{L^q(U_0)} \leq C_0 \|v\|_{L^m(U_0)}. \quad (A.3)$$

Then combining this with (A.2), since $w_0 \in \mathbb{P}^k(U_0)$,

$$\|w\|_{L^q(U)} \lesssim |U|^{\frac{1}{q}} \|w_0\|_{L^q(U_0)} \lesssim |U|^{\frac{1}{q}} \|w_0\|_{L^m(U_0)} \lesssim |U|^{\frac{1}{q} - \frac{1}{m}} \|w\|_{L^m(U)},$$

and the lemma is proved.

It remains to establish (A.3). To this end, we notice that, by choice of \mathbf{x}_U , we have $B(0, \delta) \subset U_0 \subset B(0, 1)$. Since $\|\cdot\|_{L^q(B(0,1))}$ and $\|\cdot\|_{L^m(B(0,\delta))}$ are both norms on $\mathbb{P}^k(U_0)$ (any polynomial that vanishes on a ball vanishes everywhere), and since $\mathbb{P}^k(U_0)$ is a finite-dimensional vector space, we have

$$\forall v \in \mathbb{P}^k(U_0) \quad \|v\|_{L^q(B(0,1))} \lesssim \|v\|_{L^m(B(0,\delta))}, \quad (A.4)$$

with constant in \lesssim depending on δ but not on the geometry of U_0 . To prove (A.3), write

$$\|v\|_{L^q(U_0)} \leq \|v\|_{L^q(B(0,1))} \lesssim \|v\|_{L^m(B(0,\delta))} \leq \|v\|_{L^m(U_0)}. \quad \square$$

A.2 L^p -stability and $W^{s,p}$ -approximation properties of L^2 -projectors

This section collects the proofs of L^p - and $W^{s,p}$ -stability and approximation estimates for L^2 -projectors on local polynomial spaces stated in Section 3.2.

Lemma 3.2 (*L^p -stability of L^2 -projectors on polynomial spaces*). *Let U be a measurable subset of \mathbb{R}^N , with inradius r_U and diameter h_U , such that*

$$\frac{r_U}{h_U} \geq \delta > 0. \quad (3.5)$$

Let $k \in \mathbb{N}$ and $p \in [1, +\infty]$. Then, there exists C only depending on N , δ , k and p such that

$$\forall g \in L^p(U) : \|\pi_U^k g\|_{L^p(U)} \leq C \|g\|_{L^p(U)}. \quad (3.6)$$

Proof. In this proof, $A \lesssim B$ means that $A \leq MB$ for some M only depending on N , δ , k and p .

Step 0: $p = 2$. This case is trivial since π_U^k is an orthogonal projector in $L^2(U)$ and therefore satisfies (3.6) with $C = 1$.

Step 1: $p > 2$. We use Lemma 5.1 to write $\|\pi_U^k g\|_{L^p(T)} \lesssim |T|^{\frac{1}{p}-\frac{1}{2}} \|\pi_U^k g\|_{L^2(T)}$. Since $g \in L^p(T) \subset L^2(T)$, we can use (3.6) for $p = 2$ and we deduce $\|\pi_U^k g\|_{L^p(T)} \lesssim |T|^{\frac{1}{p}-\frac{1}{2}} \|g\|_{L^2(T)}$. We then conclude thanks to Hölder's inequality, valid since $p > 2$,

$$\|\pi_U^k g\|_{L^p(T)} \lesssim |T|^{\frac{1}{p}-\frac{1}{2}} |T|^{\frac{1}{2}-\frac{1}{p}} \|g\|_{L^p(T)} = \|g\|_{L^p(T)}.$$

Step 2: $p < 2$. We use a standard duality technique. Let $g \in L^p(U)$ and $w \in L^{p'}(U)$. Then by definition of π_U^k and using (3.6) with $p' > 2$ instead of p ,

$$\int_U \pi_U^k g(\mathbf{x}) w(\mathbf{x}) d\mathbf{x} = \int_U g(\mathbf{x}) \pi_U^k w(\mathbf{x}) d\mathbf{x} \leq \|g\|_{L^p(U)} \|\pi_U^k w\|_{L^{p'}(U)} \lesssim \|g\|_{L^p(U)} \|w\|_{L^{p'}(U)}.$$

Taking the supremum of this inequality over all $w \in L^{p'}(U)$ such that $\|w\|_{L^{p'}(U)} = 1$ shows that (3.6) holds. \square

Lemma 3.4 (*$W^{s,p}$ -approximation properties of L^2 -projectors on polynomial spaces*). *Let U be an open subset of \mathbb{R}^N with diameter h_U , such that U is star-shaped with respect to a ball of radius ρh_U for some $\rho > 0$. Let $k \in \mathbb{N}$, $s \in \{1, \dots, k+1\}$ and $p \in [1, +\infty]$. Then, there exists C only depending on N , ρ , k , s and p such that*

$$\forall m \in \{0, \dots, s\}, \forall v \in W^{s,p}(U) : |v - \pi_U^k v|_{W^{m,p}(U)} \leq C h_U^{s-m} |v|_{W^{s,p}(U)}. \quad (3.7)$$

Proof. Here, $A \lesssim B$ means that $A \leq MB$ with M only depending on N , ρ , k , s and p .

The proof combines averaged Taylor polynomials [16, 38] with the L^p -stability of the L^2 -projector (Lemma 3.2). Since smooth functions are dense in $W^{s,p}(U)$, we only need to prove the result for $v \in C^\infty(U) \cap W^{s,p}(U)$. The Sobolev representation of v reads [16]

$$v = Q^s v + R^s v \quad (A.5)$$

where $Q^s v$ is a polynomial of degree less than or equal to k and the remainder $R^s v$ satisfies [16, Lemma 4.3.8]

$$\forall r \in \{0, \dots, s\} : |R^s v|_{W^{r,p}(U)} \lesssim h_U^{s-r} |v|_{W^{s,p}(U)}. \quad (A.6)$$

Since $Q^s v$ is a polynomial of degree $\leq k$, $\pi_U^k(Q^s v) = Q^s v$ and therefore, from (A.5), $\pi_U^k v = Q^s v + \pi_U^k(R^s v)$. Subtracting this from (A.5), we infer $v - \pi_U^k v = R^s v - \pi_U^k(R^s v)$. Hence,

$$|v - \pi_U^k v|_{W^{m,p}(U)} \leq |R^s v|_{W^{m,p}(U)} + |\pi_U^k(R^s v)|_{W^{m,p}(U)}. \quad (A.7)$$

Iterating the inverse inequality (A.1) and using Lemma 3.2 we see that

$$|\pi_U^k(R^s v)|_{W^{m,p}(U)} \lesssim h_U^{-m} \|\pi_U^k(R^s v)\|_{L^p(U)} \lesssim h_U^{-m} \|R^s v\|_{L^p(U)}. \quad (A.8)$$

Estimate (A.6) applied to $r = m$ and $r = 0$ shows that

$$|R^s v|_{W^{m,p}(U)} + h_U^{-m} \|R^s v\|_{L^p(U)} \lesssim h_U^{s-m} |v|_{W^{s,p}(U)}. \quad (\text{A.9})$$

The result follows from (A.7), (A.8) and (A.9). \square

Lemma 3.6 (Approximation properties of traces of L^2 -projectors on polynomial spaces). *Let T be a polyhedral subset of \mathbb{R}^N with diameter h_T , such that T is the union of disjoint simplices S of diameter h_S and inradius r_S such that $\varrho^2 h_T \leq \varrho h_S \leq r_S$ for some $\varrho > 0$. Let $k \in \mathbb{N}$, $s \in \{1, \dots, k+1\}$ and $p \in [1, +\infty]$. Then, there exists C only depending on N , ϱ , k , s and p such that*

$$\forall m \in \{0, \dots, s-1\}, \forall v \in W^{s,p}(T) : h_T^{\frac{1}{p}} |v - \pi_T^k v|_{W^{m,p}(\mathcal{F}_T)} \leq C h_T^{s-m} |v|_{W^{s,p}(T)}. \quad (\text{3.8})$$

Here, $W^{m,p}(\mathcal{F}_T)$ is the set of functions that belong to $W^{m,p}(F)$ for any hyperplanar face F of T , with corresponding broken norm.

Proof. As expected $A \lesssim B$ is understood here up to a multiplicative constant that only depends on N , ϱ , k , s and p . We first recall a classical continuous trace inequality:

$$\forall w \in W^{1,p}(T) : h_T^{\frac{1}{p}} \|w\|_{L^p(\partial T)} \lesssim \|w\|_{L^p(T)} + h_T \|\nabla w\|_{L^p(T)}. \quad (\text{A.10})$$

For $p = 2$ this inequality can be deduced from [28, Lemma 1.49] and many other references. The case of a general p is less easy to find in the literature, but actually very simple to prove. Since T is the union of disjoint simplices of inradius and diameter comparable to h_T , it is sufficient to prove the result when T is one of these simplices S . For such a simplex, there exists an affine mapping $A : T \mapsto T_0$, where $T_0 = \{\mathbf{x} \in \mathbb{R}^d : x_i > 0, \sum_{i=1}^d x_i < 1\}$ is the reference simplex, such that the norms of the linear parts of A and A^{-1} are respectively of order h_T^{-1} and h_T . Consider then $w_0 \in W^{1,p}(T_0)$ defined by $w_0(\mathbf{x}) = w(A^{-1}\mathbf{x})$. On T_0 we have a trace inequality

$$\|w_0\|_{L^p(\partial T_0)} \leq C_{d,p} (\|w_0\|_{L^p(T_0)} + \|\nabla w_0\|_{L^p(T_0)}). \quad (\text{A.11})$$

By noticing that $|\nabla w_0(\mathbf{x})| \lesssim h_T |(\nabla w)(A^{-1}\mathbf{x})|$ and using changes of variables $\mathbf{x} \mapsto \mathbf{y} = A\mathbf{x}$, (A.11) gives (A.10).

Estimate (3.8) is an immediate consequence of (A.10) and of (3.7). For $m \leq s-1$, by applying (A.10) to $w = \partial^\alpha (v - \pi_T^k v) \in W^{1,p}(T)$ for all $\alpha \in \mathbb{N}^N$ of total length m we find

$$h_T^{\frac{1}{p}} |v - \pi_T^k v|_{W^{m,p}(\mathcal{F}_T)} \lesssim |v - \pi_T^k v|_{W^{m,p}(T)} + h_T |v - \pi_T^k v|_{W^{m+1,p}(T)}.$$

We then use (3.7) for m and $m+1$ on the two terms in the right-hand side to conclude. \square

References

- [1] B. Andreianov, F. Boyer, and F. Hubert. Finite volume schemes for the p -Laplacian on Cartesian meshes. *ESAIM: Math. Model Numer. Anal. (M2AN)*, 38:931–954, 2004.
- [2] B. Andreianov, F. Boyer, and F. Hubert. Besov regularity and new error estimates for finite volume approximations of the p -Laplacian. *Numer. Math.*, 100:565–592, 2005.
- [3] B. Andreianov, F. Boyer, and F. Hubert. On the finite-volume approximation of regular solutions of the p -Laplacian. *IMA J. Numer. Anal.*, 26:472–502, 2006.
- [4] B. Andreianov, F. Boyer, and F. Hubert. Discrete Duality Finite Volume schemes for Leray–Lions-type elliptic problems on general 2D meshes. *Num. Meth. PDEs*, 23:145–195, 2007.
- [5] P. F. Antonietti, N. Bigoni, and M. Verani. Mimetic finite difference approximation of quasilinear elliptic problems. *Calcolo*, 52:45–67, 2014.
- [6] P. F. Antonietti, S. Giani, and P. Houston. hp -version composite discontinuous Galerkin methods for elliptic problems on complicated domains. *SIAM J. Sci. Comput.*, 35(3):A1417–A1439, 2013.
- [7] R. Araya, C. Harder, D. Paredes, and F. Valentin. Multiscale hybrid-mixed method. *SIAM J. Numer. Anal.*, 51(6):3505–3531, 2013.

- [8] D. N. Arnold. An interior penalty finite element method with discontinuous elements. *SIAM J. Numer. Anal.*, 19:742–760, 1982.
- [9] R. E. Bank and H. Yserentant. On the H^1 -stability of the L_2 -projection onto finite element spaces. *Numer. Math.*, 126(2):361–381, 2014.
- [10] J.W. Barrett and W. Liu. Finite element approximation of degenerate quasi-linear elliptic and parabolic problems. *Pitman Res. Notes Math. Ser.*, 303:1–16, 1994.
- [11] F. Bassi, L. Botti, A. Colombo, D. A. Di Pietro, and P. Tesini. On the flexibility of agglomeration based physical space discontinuous Galerkin discretizations. *J. Comput. Phys.*, 231(1):45–65, 2012.
- [12] L. Beirão da Veiga, F. Brezzi, A. Cangiani, G. Manzini, L. D. Marini, and A. Russo. Basic principles of virtual element methods. *Math. Models Methods Appl. Sci. (M3AS)*, 199(23):199–214, 2013.
- [13] L. Beirão da Veiga, F. Brezzi, and L. D. Marini. Virtual elements for linear elasticity problems. *SIAM J. Numer. Anal.*, 2(51):794–812, 2013.
- [14] J. H. Bramble, J. E. Pasciak, and O. Steinbach. On the stability of the L^2 projection in $H^1(\Omega)$. *Math. Comp.*, 71(237):147–156 (electronic), 2002.
- [15] S. C. Brenner. Poincaré-Friedrichs inequalities for piecewise H^1 functions. *SIAM J. Numer. Anal.*, 41(1):306–324, 2003.
- [16] S. C. Brenner and L. R. Scott. *The mathematical theory of finite element methods*, volume 15 of *Texts in Applied Mathematics*. Springer, New York, third edition, 2008.
- [17] F. Brezzi, R. S. Falk, and L. D. Marini. Basic principles of mixed virtual element methods. *ESAIM: Math. Model. Numer. Anal.*, 48(4):1227–1240, 2014.
- [18] F. Brezzi, K. Lipnikov, and V. Simoncini. A family of mimetic finite difference methods on polygonal and polyhedral meshes. *Math. Models Methods Appl. Sci.*, 15(10):1533–1551, 2005.
- [19] A. Buffa and C. Ortner. Compact embeddings of broken Sobolev spaces and applications. *IMA J. Numer. Anal.*, 4(29):827–855, 2009.
- [20] E. Burman and A. Ern. Discontinuous Galerkin approximation with discrete variational principle for the nonlinear Laplacian. *C. R. Acad. Sci. Paris, Ser. I*, 346:1013–1016, 2008.
- [21] C. Carstensen. Merging the Bramble-Pasciak-Steinbach and the Crouzeix-Thomée criterion for H^1 -stability of the L^2 -projection onto finite element spaces. *Math. Comp.*, 71(237):157–163 (electronic), 2002.
- [22] B. Cockburn, D. A. Di Pietro, and A. Ern. Bridging the Hybrid High-Order and Hybridizable Discontinuous Galerkin methods. *ESAIM: Math. Model. Numer. Anal. (M2AN)*, 2015. Published online. DOI 10.1051/m2an/2015051.
- [23] B. Cockburn, J. Gopalakrishnan, and R. Lazarov. Unified hybridization of discontinuous Galerkin, mixed, and continuous Galerkin methods for second order elliptic problems. *SIAM J. Numer. Anal.*, 47(2):1319–1365, 2009.
- [24] M. Crouzeix and V. Thomée. The stability in L_p and W_p^1 of the L_2 -projection onto finite element function spaces. *Math. Comp.*, 48(178):521–532, 1987.
- [25] K. Deimling. *Nonlinear functional analysis*. Springer-Verlag, Berlin, 1985.
- [26] D. A. Di Pietro, J. Droniou, and A. Ern. A discontinuous-skeletal method for advection–diffusion–reaction on general meshes. *SIAM J. Numer. Anal.*, 2015. To appear.
- [27] D. A. Di Pietro and A. Ern. Discrete functional analysis tools for discontinuous Galerkin methods with application to the incompressible Navier–Stokes equations. *Math. Comp.*, 79:1303–1330, 2010.
- [28] D. A. Di Pietro and A. Ern. *Mathematical aspects of discontinuous Galerkin methods*, volume 69 of *Mathématiques & Applications*. Springer-Verlag, Berlin, 2012.
- [29] D. A. Di Pietro, A. Ern, and S. Lemaire. An arbitrary-order and compact-stencil discretization of diffusion on general meshes based on local reconstruction operators. *Comput. Meth. Appl. Math.*, 14(4):461–472, 2014.
- [30] D. A. Di Pietro and S. Lemaire. An extension of the Crouzeix–Raviart space to general meshes with application to quasi-incompressible linear elasticity and Stokes flow. *Math. Comp.*, 84(291):1–31, 2015.
- [31] J. I. Diaz and F. de Thelin. On a nonlinear parabolic problem arising in some models related to turbulent flows. *SIAM J. Math. Anal.*, 25(4):1085–1111, 1994.
- [32] J. Droniou. Finite volume schemes for fully non-linear elliptic equations in divergence form. *ESAIM: Math. Model. Numer. Anal. (M2AN)*, 40:1069–1100, 2006.
- [33] J. Droniou and R. Eymard. A mixed finite volume scheme for anisotropic diffusion problems on any grid. *Numer. Math.*, 105:35–71, 2006.
- [34] J. Droniou and R. Eymard. Study of the mixed finite volume method for Stokes and Navier-Stokes equations. *Numer. Methods Partial Differential Equations*, 25(1):137–171, 2009.
- [35] J. Droniou, R. Eymard, T. Gallouët, C. Guichard, and R. Herbin. Gradient schemes for elliptic and parabolic problems. 2015. In preparation.

- [36] J. Droniou, R. Eymard, T. Gallouët, and R. Herbin. A unified approach to mimetic finite difference, hybrid finite volume and mixed finite volume methods. *Math. Models Methods Appl. Sci. (M3AS)*, 20(2):1–31, 2010.
- [37] J. Droniou, R. Eymard, T. Gallouët, and R. Herbin. Gradient schemes: a generic framework for the discretization of linear, nonlinear and nonlocal elliptic and parabolic equations. *Math. Models Methods Appl. Sci. (M3AS)*, 23(13):2395–2432, 2012.
- [38] T. Dupont and R. Scott. Polynomial approximation of functions in Sobolev spaces. *Math. Comp.*, 34(150):441–463, 1980.
- [39] R. Eymard, T. Gallouët, and R. Herbin. Discretization of heterogeneous and anisotropic diffusion problems on general nonconforming meshes. SUSI: a scheme using stabilization and hybrid interfaces. *IMA J. Numer. Anal.*, 30(4):1009–1043, 2010.
- [40] V. Girault, B. Rivière, and M. F. Wheeler. A discontinuous Galerkin method with nonoverlapping domain decomposition for the Stokes and Navier-Stokes problems. *Math. Comp.*, 74(249):53–84, 2005.
- [41] R. Glowinski. *Numerical methods for nonlinear variational problems*. Springer Series in Computational Physics. Springer-Verlag, New York, 1984.
- [42] R. Glowinski and J. Rappaz. Approximation of a nonlinear elliptic problem arising in a non-Newtonian fluid flow model in glaciology. *ESAIM: Math. Model Numer. Anal. (M2AN)*, 37(1):175–186, 2003.
- [43] R. Herbin and F. Hubert. Benchmark on discretization schemes for anisotropic diffusion problems on general grids. In R. Eymard and J.-M. Hérard, editors, *Finite Volumes for Complex Applications V*, pages 659–692. John Wiley & Sons, 2008.
- [44] O. A. Karakashian and W. N. Jureidini. A nonconforming finite element method for the stationary Navier-Stokes equations. *SIAM J. Numer. Anal.*, 35(1):93–120, 1998.
- [45] A. Lasis and E. Süli. Poincaré-type inequalities for broken Sobolev spaces. Technical Report 03/10, Oxford University Computing Laboratory, Oxford, England, 2003.
- [46] J. Leray and J.-L. Lions. Quelques résultats de Višik sur les problèmes elliptiques non linéaires par les méthodes de Minty-Browder. *Bull. Soc. Math. France*, 93:97–107, 1965.
- [47] K. Lipnikov and G. Manzini. A high-order mimetic method on unstructured polyhedral meshes for the diffusion equation. *J. Comput. Phys.*, 272:360–385, 2014.
- [48] W. Liu and N. Yan. Quasi-norm a priori and a posteriori error estimates for the nonconforming approximation of p -Laplacian. *Numer. Math.*, 89:341–378, 2001.
- [49] G. J. Minty. On a “monotonicity” method for the solution of non-linear equations in Banach spaces. *Proc. Nat. Acad. Sci. U.S.A.*, 50:1038–1041, 1963.
- [50] J. Wang and X. Ye. A weak Galerkin element method for second-order elliptic problems. *J. Comput. Appl. Math.*, 241:103–115, 2013.
- [51] J. Wang and X. Ye. A weak Galerkin mixed finite element method for second order elliptic problems. *Math. Comp.*, 83(289):2101–2126, 2014.