



HAL
open science

Multiple sampling and interpolation in the classical Fock space

Alexander Borichev, Andreas Hartmann, Karim Kellay, Xavier Massaneda

► **To cite this version:**

Alexander Borichev, Andreas Hartmann, Karim Kellay, Xavier Massaneda. Multiple sampling and interpolation in the classical Fock space. *Zapiski Nauchnykh Seminarov. Sankt-Peterburgskoe Otdelenie. Matematicheskii Institut. Steklova.(POMI)*, 2015. hal-01183796v1

HAL Id: hal-01183796

<https://hal.science/hal-01183796v1>

Submitted on 11 Aug 2015 (v1), last revised 14 Dec 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MULTIPLE SAMPLING AND INTERPOLATION IN THE CLASSICAL FOCK SPACE

A. BORICHEV, A. HARTMANN, K. KELLAY, X. MASSANEDA

ABSTRACT. We study multiple sampling, interpolation and uniqueness for the classical Fock space in the case of unbounded multiplicities.

Sampling and interpolating sequences in Fock spaces were characterized by Seip and Seip–Wallstén in [3, 4] by means of a certain Beurling–type asymptotic uniform density. The case of uniformly bounded multiplicities was considered by Brekke and Seip [1] who gave a complete description in this situation. Their conditions show that it is not possible that a sequence is simultaneously sampling and interpolating.

Brekke and Seip also asked whether there exist sequences which are simultaneously sampling and interpolating when the multiplicities are unbounded.

In this research note we formulate some conditions (of geometric nature) for sampling and interpolation. They show that the answer to this question is negative when the multiplicities tend to infinity.

The detailed version of this work will be published elsewhere.

We now introduce the necessary notation. For $\alpha > 0$, define the Fock space \mathcal{F}_α^2 by

$$\mathcal{F}_\alpha^2 = \left\{ f \in \text{Hol}(\mathbb{C}) : \|f\|_2^2 = \|f\|_{\alpha,2}^2 := \frac{\alpha}{\pi} \int_{\mathbb{C}} |f(z)|^2 e^{-\alpha|z|^2} dm(z) < \infty \right\}.$$

The space \mathcal{F}_α^2 is a Hilbert space with the inner product

$$\langle f, g \rangle = \frac{\alpha}{\pi} \int_{\mathbb{C}} f(z) \overline{g(z)} e^{-\alpha|z|^2} dm(z).$$

The sequence

$$e_k(z) = \frac{\sqrt{\alpha^k}}{\sqrt{k!}} z^k, \quad k \geq 0,$$

2000 *Mathematics Subject Classification.* 30D55, 46C07, 46E22, 47B32, 47B35.

Key words and phrases. Fock space, multiple interpolation, multiple sampling.

The work was supported by Russian Science Foundation grant 14-41-00010.

defines an orthonormal basis in \mathcal{F}_α^2 .

Recall that the translations

$$T_z f(\zeta) = T_z^\alpha f(\zeta) := e^{\alpha \bar{z} \zeta - \frac{\alpha}{2} |z|^2} f(\zeta - z), \quad f \in \mathcal{F}_\alpha^2.$$

act isometrically in \mathcal{F}_α^2 .

Let us now define sampling and interpolation in the unbounded multiplicity case. Consider the divisor $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$, where Λ is a sequence of points in \mathbb{C} and $m_\lambda \in \mathbb{N}$ is the multiplicity associated with λ . The divisor X is called

- *sampling* for \mathcal{F}_α^2 if

$$\|f\|_2^2 \asymp \sum_{\lambda \in \Lambda} \sum_{k=0}^{m_\lambda-1} |\langle f, T_\lambda e_k \rangle|^2, \quad f \in \mathcal{F}_\alpha^2,$$

- *interpolating* for \mathcal{F}_α^2 if for every sequence

$$v = \{v_\lambda^{(k)}\}_{\lambda \in \Lambda, 0 \leq k < m_\lambda}$$

such that

$$\|v\|_2^2 := \sum_{\lambda \in \Lambda} \sum_{k=0}^{m_\lambda-1} |v_\lambda^{(k)}|^2 < \infty,$$

there exists a function $f \in \mathcal{F}_\alpha^2$ satisfying

$$\langle f, T_\lambda e_k \rangle = v_\lambda^{(k)}, \quad 0 \leq k < m_\lambda, \quad \lambda \in \Lambda.$$

As in the situation of classical interpolation and sampling, the separation between points in Λ plays an important role. Denote by $D(z, r)$ the disc of radius r centered at z .

- A divisor $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$ is said to satisfy the *finite overlap condition* if

$$\sup_{z \in \mathbb{C}} \sum_{\lambda \in \Lambda} \chi_{D(\lambda, \sqrt{m_\lambda/\alpha})}(z) < \infty.$$

If Λ is a finite union of Λ_j such that the discs $D(\lambda, \sqrt{m_\lambda/\alpha})$, $\lambda \in \Lambda_j$, are disjoint for every j , then X satisfies the finite overlap condition.

The following result gives geometric conditions for sampling in the case of unbounded multiplicities.

Theorem 1. (a) If $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$ is a sampling divisor for \mathcal{F}_α^2 , then X satisfies the finite overlap condition and there exists $C > 0$ such that

$$\bigcup_{\lambda \in \Lambda} D(\lambda, \sqrt{m_\lambda/\alpha} + C) = \mathbb{C}.$$

(b) Conversely, if $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$ satisfies the finite overlap condition and for every $C > 0$ there is a compact subset K of \mathbb{C} such that

$$\bigcup_{\lambda \in \Lambda, m_\lambda > \alpha C^2} D(\lambda, \sqrt{m_\lambda/\alpha} - C) = \mathbb{C} \setminus K,$$

then X is a sampling divisor for \mathcal{F}_α^2 .

Remark. Let $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$ satisfy the conditions of Theorem 1 (b). Then we can find a subset $\Lambda_1 \subset \Lambda$ such that for every $C > 0$ there is a compact subset K of \mathbb{C} satisfying

$$\bigcup_{\lambda \in \Lambda_1, m_\lambda > \alpha C^2} D(\lambda, \sqrt{m_\lambda/\alpha} - C) = \mathbb{C} \setminus K,$$

and

$$\lim_{\lambda \in \Lambda_1, |\lambda| \rightarrow \infty} m_\lambda = +\infty.$$

The following result gives geometric conditions for interpolation in the case of unbounded multiplicities.

Theorem 2. (a) If $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$ is an interpolating divisor for \mathcal{F}_α^2 , then there exists $C > 0$ such that the discs $\{D(\lambda, \sqrt{m_\lambda/\alpha} - C)\}_{\lambda \in \Lambda, m_\lambda > \alpha C^2}$ are pairwise disjoint.

(b) Conversely, if the disks $\{D(\lambda, \sqrt{m_\lambda/\alpha} + C)\}_{\lambda \in \Lambda}$ are pairwise disjoint for some $C > 0$, then X is an interpolating divisor for \mathcal{F}_α^2 .

Remark. It is easily seen that if X is a divisor such that for some $C > 0$ the discs $\{D(\lambda, \sqrt{m_\lambda/\alpha} - C)\}_{\lambda \in \Lambda, m_\lambda > \alpha C^2}$ are pairwise disjoint, and if $\lim_{|\lambda| \rightarrow \infty} m_\lambda = +\infty$, then X satisfies the finite overlap condition.

Though our geometric conditions do not characterize interpolation and sampling they allow us to deduce the following result, which gives a partial answer to the question raised by Brekke and Seip.

Corollary 3. *Let the divisor $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$ satisfy the condition $\lim_{|\lambda| \rightarrow \infty} m_\lambda = +\infty$. Then X cannot be simultaneously interpolating and sampling for \mathcal{F}_α^2 .*

The problems of sampling and interpolation are linked to that of uniqueness, and thus to zero divisors, for which some conditions are discussed in [2, 5].

We will formulate here a necessary condition for zero divisors which, apparently, does not follow from those known so far.

Proposition 4. *Let $X = \{(\lambda, m_\lambda)\}_{\lambda \in \Lambda}$. If there exists a compact subset K of \mathbb{C} such that*

$$\bigcup_{\lambda \in \Lambda} D(\lambda, \sqrt{m_\lambda/\alpha}) = \mathbb{C} \setminus K,$$

then X is not a zero divisor for \mathcal{F}_α^2 .

As a matter of fact, this result holds more generally in the weighted Fock space with uniform norm $\mathcal{F}_\alpha^\infty$ defined by

$$\mathcal{F}_\alpha^\infty = \left\{ f \in \text{Hol}(\mathbb{C}) : \|f\|_\infty = \|f\|_{\alpha, \infty} := \sup_{z \in \mathbb{C}} |f(z)| e^{-\frac{\alpha}{2}|z|^2} < \infty \right\}.$$

REFERENCES

- [1] S. Brekke, K. Seip, Density theorems for sampling and interpolation in the Bargmann–Fock space III, *Math. Scand.* **73** (1993), 112–126.
- [2] B. Ya. Levin, *Lectures on entire functions*, Translations of Mathematical Monographs, 150, American Mathematical Society, Providence, RI, 1996.
- [3] K. Seip, Density theorems for sampling and interpolation in the Bargmann–Fock space I, *J. Reine Angew. Math.* **429** (1992), 91–106.
- [4] K. Seip, R. Wallstén, Density theorems for sampling and interpolation in the Bargmann–Fock space II, *J. Reine Angew. Math.* **429** (1992), 107–113.
- [5] K. Zhu, *Analysis on Fock spaces*, Graduate Texts in Mathematics, 263, Springer, New York, 2012.

A.BORICHEV: AIX MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, 13453 MARSEILLE, FRANCE

E-mail address: alexander.borichev@math.cnrs.fr

A.HARTMANN: UNIVERSITÉ DE BORDEAUX, IMB, 351 COURS DE LA LIBÉRATION, 33405 TALENCE, FRANCE

E-mail address: Andreas.Hartmann@math.u-bordeaux1.fr

K.KELLAY: UNIVERSITÉ DE BORDEAUX, IMB, 351 COURS DE LA LIBÉRATION, 33405 TALENCE, FRANCE

E-mail address: karim.kellay@math.u-bordeaux1.fr

X.MASSANEDA: UNIVERSITAT DE BARCELONA, DEPARTAMENT DE MATEMÀ-
TICA APLICADA I ANÀLISI, GRAN VIA 585, 08007-BARCELONA, SPAIN
E-mail address: `xavier.massaneda@ub.edu`