



HAL
open science

Arrow Type Impossibility Theorems over Median Algebras

Miguel Couceiro, Stephan Foldes, Gerasimos Meletiou

► **To cite this version:**

Miguel Couceiro, Stephan Foldes, Gerasimos Meletiou. Arrow Type Impossibility Theorems over Median Algebras. 2015. hal-01185977

HAL Id: hal-01185977

<https://hal.science/hal-01185977v1>

Preprint submitted on 22 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

ARROW TYPE IMPOSSIBILITY THEOREMS OVER MEDIAN ALGEBRAS

MIGUEL COUCEIRO, STEPHAN FOLDES, AND GERASIMOS C. MELETIOU

ABSTRACT. We characterize trees as median algebras and semilattices by relaxing conservativeness. Moreover, we describe median homomorphisms between products of median algebras and show that Arrow type impossibility theorems for mappings from a product $\mathbf{A}_1 \times \cdots \times \mathbf{A}_n$ of median algebras to a median algebra \mathbf{B} are possible if and only if \mathbf{B} is a tree, when thought of as an ordered structure.

1. INTRODUCTION

Median algebras have been investigated by several authors in the context of distributive algebras, semilattices, graphs and hypergraphs and in the framework of convex and metric spaces (see, e.g., [2, 3, 5, 6, 10, 14, 15, 16, 18, 19, 20]). A *median algebra* is a structure $\mathbf{A} = \langle A, \mathbf{m} \rangle$ for a set A and a ternary symmetric operation $\mathbf{m}: A^3 \rightarrow A$, called *median*, such that

$$\begin{aligned} \mathbf{m}(x, x, y) &= x, \\ \mathbf{m}(\mathbf{m}(x, y, z), t, u) &= \mathbf{m}(x, \mathbf{m}(y, t, u), \mathbf{m}(z, t, u)). \end{aligned}$$

In fact, axiom systems using only 4 variables are also known [3, 12, 13]. For instance, in [13] it was shown that

$$\mathbf{m}(x, x, y) = x \quad \text{and} \quad \mathbf{m}(\mathbf{m}(x, w, z), y, z) = \mathbf{m}(\mathbf{m}(y, z, w), x, z)$$

suffice to axiomatize median algebras. In particular, it immediately follows that

$$\mathbf{m}(\mathbf{m}(x, y, z), y, z) = \mathbf{m}(x, y, z).$$

It is well known [18] that each element a of a median algebra \mathbf{A} gives rise to a *median semilattice* $\langle A, \leq_a \rangle$ where \leq_a is given by

$$x \leq_a y \iff \mathbf{m}(a, x, y) = x.$$

In this semilattice a is the bottom element and the associated binary operation, denoted \wedge_a , is defined by $x \wedge_a y = \mathbf{m}(a, x, y)$. As shown in [18], median semilattices coincide exactly with those \wedge -semilattices whose principal ideals

$$\downarrow x := \{y \in A : y \leq_a x\}$$

are distributive lattices, and such that for any $a, b, c \in A$, $a \wedge b, b \wedge c, c \wedge a$ have a supremum whenever each pair of these meets is bounded above. In such a case, we can define a median operation by

$$(1.1) \quad \mathbf{m}_{\leq_a}(x, y, z) = (x \wedge_a y) \vee (x \wedge_a z) \vee (z \wedge_a y),$$

for every $x, y, z \in A$. Moreover, $\mathbf{m} = \mathbf{m}_{\leq_a}$ for every median algebra $\mathbf{A} = \langle A, \mathbf{m} \rangle$, and every $a \in A$ [2]. Similarly, every distributive lattice gives rise to a median algebra using (1.1) and the converse also holds whenever there are $a, b \in A$ such that $\mathbf{m}(a, x, b) = x$ for every $x \in A$.

Another noteworthy connection is to *median graphs*, i.e., connected graphs having the property that for any three vertices a, b, c , there is exactly one vertex x in the intersection of the shortest paths between any two vertices in $\{a, b, c\}$. As shown in [2], the *covering graph* (i.e., undirected Hasse diagram) of every median semilattice whose intervals are finite is a median graph. Conversely, every median graph is the covering graph of a median semilattice. For further background see, e.g., [3].

In [8] the authors studied “conservative” median algebras, i.e., median algebras that satisfy $\mathbf{m}(x, y, z) \in \{x, y, z\}$, for all $x, y, z \in A$. Apart from the 4-element Boolean algebra, it was shown that conservative median algebras can be represented by two lower bounded chains whose least elements are identified (thus, they can always be regarded as chains). It is noteworthy to observe that they are exactly those median algebras whose subsets are themselves median (sub)algebras (in the terminology of [4], those with median stabilization degree equal to 0). Equivalently, they were shown to have the 4-element star as a forbidden subgraph. The authors in [8] also provided complete descriptions of median-homomorphisms between conservative median algebras (with at least 5 elements).

Theorem 1.1 ([8]). *Let \mathbf{A} and \mathbf{B} be two conservative median algebras (thought of as chains) with at least five elements. A map $f : \mathbf{A} \rightarrow \mathbf{B}$ is a median homomorphism if and only if f is monotone (i.e., order-preserving or order-reversing).*

This result was then lifted, by making use of dual topological spaces of median algebras, to median-homomorphisms between products of conservative median algebras. For each positive integer n , we set $[n] = \{1, \dots, n\}$.

Theorem 1.2 ([8]). *Let $\mathbf{A} = \mathbf{C}_1 \times \dots \times \mathbf{C}_n$ and $\mathbf{B} = \mathbf{D}_1 \times \dots \times \mathbf{D}_k$ be two finite products of conservative median algebras (thought of as chains). Then $f : \mathbf{A} \rightarrow \mathbf{B}$ is a median homomorphism if and only if there exist $\sigma : [k] \rightarrow [n]$ and monotone maps $f_i : \mathbf{C}_{\sigma(i)} \rightarrow \mathbf{D}_{(i)}$ for $i \in [k]$, such that $f = (f_{\sigma(1)}, \dots, f_{\sigma(k)})$.*

In the particular case when $k = 1$,

$$f : \mathbf{C}_1 \times \dots \times \mathbf{C}_n \rightarrow \mathbf{D}$$

is a median homomorphism if and only if there is $j \in [n]$ and a monotone map $g : \mathbf{C}_j \rightarrow \mathbf{D}$ such that $f = g \circ \pi_j$.

In this paper we are interested in median algebras \mathbf{A} that are trees (i.e., acyclic undirected graphs) when thought of as semilattices: a \wedge -semilattice (resp. \vee -semilattice) is said to be a *tree* if no pair of incomparable elements have an upper (resp. lower) bound. As we will see, such median algebras are obtained by relaxing the condition of being conservative: we say that a median algebra \mathbf{A} is a *(2 : 3)-median semilattice* if $\mathbf{m}(x, y, z) \in \{x \wedge y, y \wedge z, z \wedge x\}$ for all $x, y, z \in A$. Following the same structure as in [8], we will then proceed to describing the median homomorphisms between (products of) such median algebras. As conservative median algebras with more than 4 elements are particular cases of (2 : 3)-median semilattices, these descriptions properly extend results in [8]. Unlike in the latter paper, where an extensive use of dual spaces of median algebras took place, in the current paper we take a purely algebraic approach.

The paper is organized as follows. After recalling some terminology and basic results on median algebras in Section 2, we introduce the notion of (2 : 3)-median semilattice and show, in Section 3, that it captures those median semilattices that are trees. The problem of describing homomorphisms between products of median algebras is then tackled in Section 4. We show that such median preserving mappings can be decomposed into homomorphisms of the form

$$(1.2) \quad f : \mathbf{A}_1 \times \dots \times \mathbf{A}_n \rightarrow \mathbf{B},$$

and explicitly describe them in the case when \mathbf{B} is a tree. Our results are illustrated by several examples and their limitations by counter-examples. In Section 5 we comment on natural interpretations of our results, namely, when looked at as impossibility theorems. In particular, we show that Arrow type impossibility results for median-preserving procedures (1.2) (that basically state that they depend on at most one argument) can hold if and only if \mathbf{B} is a tree.

2. PRELIMINARY RESULTS

In this section we introduce basic notions, terminology and notation, as well as recall some basic facts and well-known results about median algebras as ordered sets, that will be used throughout the paper. To avoid a lengthy preliminary section, we will restrict ourselves to strictly necessary background, and we refer the reader to [7, 9, 11] for further background.

As discussed in Section 1, when there is no danger of ambiguity, we will not distinguish between median algebras and median semilattices. They will be denoted by bold roman capital letters $\mathbf{A}, \mathbf{B}, \dots$, while their universes will be denoted by italic roman capital letters A, B, \dots . We will assume that universes of structures are nonempty, possibly infinite, sets.

Let $\mathbf{A} = \langle A, \mathbf{m} \rangle$ be a median algebra and let $a, b \in A$. The *convex hull* of $\{a, b\}$ or the *interval* from a to b , denoted by $[a, b]$, is defined by

$$[a, b] := \{t \in A : t = \mathbf{m}(a, t, b)\}.$$

Intuitively, it is the set of all elements of A in the “shortest” paths from a to b , thus explaining our choice of notation.

Proposition 2.1. *Let $\mathbf{A} = \langle A, \mathbf{m} \rangle$ be a median algebra and let $a, b \in A$. Then*

$$[a, b] = \{\mathbf{m}(a, t, b) : t \in A\}.$$

Furthermore, we also have the following useful result.

Proposition 2.2 ([6]). *Let $\mathbf{A} = \langle A, \mathbf{m} \rangle$ be a median algebra, and consider $a, b, c \in A$. Then*

$$[a, b] \cap [b, c] \cap [a, c] = \{\mathbf{m}(a, b, c)\}.$$

Let $\mathbf{A} = \langle A, \mathbf{m} \rangle$ be a median algebra and let $a, b \in A$. Each convex hull of a 2-element set $\{a, b\}$ can be endowed with a distributive lattice structure. To this purpose, for $a \in A$ recall the binary operation $\wedge_a : A^2 \rightarrow A$ defined by:

$$x \wedge_a y := \mathbf{m}(a, x, y), \quad \text{for } (x, y) \in A^2.$$

Note that such an operation is idempotent, commutative and associative. Furthermore, it is not difficult to see that on $[a, b]$, \wedge_a and \wedge_b verify the absorption laws, and that for every $s \in [a, b]$, we have

$$a \leq_a s \leq_a b \quad \text{and} \quad b \leq_b s \leq_b a.$$

In fact, we have the following well-known result; see, e.g., [3].

Proposition 2.3. *Let $\mathbf{A} = \langle A, \mathbf{m} \rangle$ be a median algebra. For every $a, b \in A$, $\langle [a, b], \wedge_a, \wedge_b \rangle$ is a distributive lattice with a and b as the least and greatest elements, respectively.*

Moreover, we also have a general description of median-preserving mappings between two median algebras.

Theorem 2.4. *Let \mathbf{A} and \mathbf{B} be median algebras, and consider a mapping $f : A \rightarrow B$. Then the following assertions are equivalent.*

- (i) *f is a median-homomorphism.*

- (ii) For all $p \in A$, $f: \langle A, \wedge_p \rangle \rightarrow \langle B, \wedge_{f(p)} \rangle$ is a semilattice-homomorphism.
- (iii) For all $p \in A$, $f: \langle A, \leq_p \rangle \rightarrow \langle B, \leq_{f(p)} \rangle$ is an order-homomorphism.
- (iv) For all $a, b \in A$, $f([a, b]) \subseteq [f(a), f(b)]$.

Proof. We prove the following sequence of implications

$$(i) \implies (ii) \implies (iii) \implies (iv) \implies (i)$$

thus showing that they are all equivalent.

(i) \implies (ii): Suppose that f is a median-homomorphism and $p \in A$. Then

$$f(a \wedge_p b) = f(\mathbf{m}(p, a, b)) = \mathbf{m}(f(p), f(a), f(b)) = f(a) \wedge_{f(p)} f(b).$$

(ii) \implies (iii): Straightforward.

(iii) \implies (iv): Let $t \in [a, b]$. Then $a \leq_a t \leq_a b$, and thus $f(a) \leq_{f(a)} f(t) \leq_{f(a)} f(b)$ from which it follows that $f(t) \in [f(a), f(b)]$. In other words, $f([a, b]) \subseteq [f(a), f(b)]$.

(iv) \implies (i): Let $a, b, c \in A$. By Proposition 2.2, we have

$$\{\mathbf{m}(a, b, c)\} = [a, b] \cap [b, c] \cap [c, a].$$

By (iv), we then conclude that

$$f(\mathbf{m}(a, b, c)) \in [f(a), f(b)] \cap [f(b), f(c)] \cap [f(c), f(a)] = \{\mathbf{m}(f(a), f(b), f(c))\}.$$

Therefore, $f(\mathbf{m}(a, b, c)) = \mathbf{m}(f(a), f(b), f(c))$. \square

Corollary 2.5. *Let $f: A \rightarrow B$ be a median homomorphism between two median algebras \mathbf{A} and \mathbf{B} . Then, for every $a, b \in A$, f is also a lattice homomorphism from $\langle [a, b], \wedge_a, \wedge_b \rangle$ to $\langle [f(a), f(b)], \wedge_{f(a)}, \wedge_{f(b)} \rangle$.*

3. TREES AS MEDIAN ALGEBRAS

In this section we focus on median algebras that are trees. In the particular case of chains, it was shown that these median algebras are exactly those that are conservative, i.e.,

$$\mathbf{m}(x, y, z) \in \{x, y, z\}, \quad x, y, z \in A.$$

To identify median semilattices that are trees, we propose the following relaxation of conservativeness. We say that a median algebra, thought of as a median semilattice $\mathbf{A} = \langle A, \wedge \rangle$, is a $(2 : 3)$ -median semilattice if for every $x, y, z \in A$, we have

$$\mathbf{m}(x, y, z) := (x \wedge y) \vee (x \wedge z) \vee (z \wedge y) \in \{x \wedge y, y \wedge z, z \wedge x\}.$$

We start with a simple yet useful observation.

Remark 3.1. Let $\mathbf{A} = \langle A, \mathbf{m} \rangle$ be a median algebra such that $\langle A, \leq_c \rangle$ is a tree for some (or, equivalently, all) $c \in A$. Then, for all $a, b \in A$, the convex hull $[a, b]$ is a chain from a to b .

Theorem 3.2. *Let $\mathbf{A} = \langle A, \mathbf{m} \rangle$ be a median algebra. Then the following assertions are equivalent.*

- (i) There is $p \in A$ such that $\mathbf{A} = \langle A, \leq_p \rangle$ is a $(2 : 3)$ -median semilattice.
- (ii) For all $p \in A$, $\mathbf{A} = \langle A, \leq_p \rangle$ is a $(2 : 3)$ -median semilattice.
- (iii) There is $p \in A$ such that $\mathbf{A} = \langle A, \leq_p \rangle$ is a tree.
- (iv) For all $p \in A$, $\mathbf{A} = \langle A, \leq_p \rangle$ is a tree.
- (v) For all $a, b \in A$, the bounded distributive lattice $\langle [a, b], \wedge_a, \wedge_b \rangle$ is a chain.

Proof. We prove the following sequence of implications

$$(ii) \implies (i) \implies (iii) \implies (iv) \implies (v) \implies (iv) \implies (ii),$$

thus showing that they are all equivalent.

(ii) \implies (i): Straightforward.

(i) \implies (iii): Suppose that $\mathbf{A} = \langle A, \leq_c \rangle$ is a $(2 : 3)$ -median semilattice. To show that it is a tree it suffices to show that for every pair $a, b \in A$ with a common upper bound, we have $a \leq_p b$ or $b \leq_p a$.

So suppose that $c \in A$ is a common upper bound of a and b , that is, $\mathbf{m}(p, a, c) = a$ and $\mathbf{m}(p, b, c) = b$. Since $\mathbf{A} = \langle A, \leq_c \rangle$ is a $(2 : 3)$ -median semilattice and c is an upper bound of a and b ,

$$(3.1) \quad \mathbf{m}(c, a, b) \in \{\mathbf{m}(p, a, b), a, b\}.$$

Note that since \mathbf{m} is a median,

$$\mathbf{m}(p, \mathbf{m}(a, c, b), a) = \mathbf{m}(b, \mathbf{m}(a, a, p), \mathbf{m}(c, a, p)) = \mathbf{m}(b, a, a) = a,$$

and thus $a \leq_p \mathbf{m}(c, a, b)$. Similarly, we have $b \leq_p \mathbf{m}(c, a, b)$.

Now, if $\mathbf{m}(c, a, b) = \mathbf{m}(p, a, b)$ in (3.1), then $\mathbf{m}(c, a, b) \leq_p a$ since $\mathbf{m}(p, a, b) \leq_p a$. Hence, $a = \mathbf{m}(c, a, b)$. Similarly, we also have $b = \mathbf{m}(c, a, b)$, and thus $a = b$.

If $\mathbf{m}(c, a, b) = a$ in (3.1), then

$$\mathbf{m}(p, a, b) = \mathbf{m}(\mathbf{m}(c, a, b), p, b) = \mathbf{m}(a, \mathbf{m}(c, p, b), \mathbf{m}(b, p, b)) = \mathbf{m}(a, b, b) = b,$$

and thus $b \leq_p a$. Similarly, if $\mathbf{m}(c, a, b) = b$ in (3.1), we conclude that $a \leq_p b$.

Since in all possible cases we have that $a \leq_p b$ or $b \leq_p a$, $\mathbf{A} = \langle A, \leq_p \rangle$ is a tree.

(iii) \implies (iv): Straightforward.

(iv) \implies (v): Let $s, t \in [a, b]$. Thus $a \leq_a s, t \leq_a b$, and s, t have a common upper bound in $\langle A, \leq_a \rangle$. As $\langle A, \leq_a \rangle$ is a tree, s and t cannot be incomparable, and hence the interval $[a, b]$ is a chain.

(v) \implies (iv): Let $a, b, p \in A$ and suppose that a and b are incomparable in $\mathbf{A} = \langle A, \leq_p \rangle$. Note that $p \leq_p a, b$.

For the sake of a contradiction, suppose that a and b have a common upper bound $d \in A$, that is, $p \leq_p a, b \leq_p d$. However, by (v) the interval $[p, d]$ is a chain and thus we must have $a \leq_p b$ or $b \leq_p a$, which constitutes the desired contradiction.

(iv) \implies (ii): Let $p \in A$. As $\mathbf{A} = \langle A, \leq_p \rangle$ is a tree, for every $a, b, c \in A$, we have that $a \wedge b, b \wedge c$ and $c \wedge a$ are pairwise comparable, and thus

$$\mathbf{m}(a, b, c) \in \{a \wedge b, b \wedge c, c \wedge a\}.$$

Hence, $\mathbf{A} = \langle A, \leq_p \rangle$ is a $(2 : 3)$ -median semilattice. \square

Remark 3.3. Other equivalent descriptions of trees as median semilattices are given in [3, 18].

Remark 3.4. As mentioned, conservative median semilattices \mathbf{A} are $(2 : 3)$ -median semilattices, whenever $|A| \geq 5$. Hence, Lemma 3 and Theorem 3 in [8] follow as corollaries of Theorem 3.2. In fact, the conservative median semilattice $\{0, 1\}^2$ is the only ‘‘pathological’’ case.

4. MEDIAN-HOMOMORPHISMS OVER TREES

Let $\mathbf{A}_1, \dots, \mathbf{A}_n$ and $\mathbf{B}_1, \dots, \mathbf{B}_k$ be median algebras. We seek to describe median-homomorphisms of the form

$$(4.1) \quad f: \mathbf{A}_1 \times \dots \times \mathbf{A}_n \rightarrow \mathbf{B}_1 \times \dots \times \mathbf{B}_k.$$

For each $j \in [k]$, let π_j denote the j -th projection on $\mathbf{B}_1 \times \dots \times \mathbf{B}_k$, i.e., the mapping

$$\pi_j: \mathbf{B}_1 \times \dots \times \mathbf{B}_k \rightarrow \mathbf{B}_j.$$

It is easy to see that if f is a median-homomorphism, then the composition

$$g_j = \pi_j \circ f: \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}_j$$

is a median-homomorphism. As the converse holds trivially, possibly with repeated g_j 's (as they are not necessarily pairwise distinct), we get the following result.

Lemma 4.1. *Let $\mathbf{A}_1, \dots, \mathbf{A}_n$ and $\mathbf{B}_1, \dots, \mathbf{B}_k$ be median algebras. A mapping $f: \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}_1 \times \cdots \times \mathbf{B}_k$ is a median-homomorphism if and only if there are median-homomorphisms*

$$g_j: \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}_j, \quad j \in [k],$$

such that $f = (g_1, \dots, g_k)$. Moreover, we have $g_j = \pi_j \circ f$ for the projection

$$\pi_j: \mathbf{B}_1 \times \cdots \times \mathbf{B}_k \rightarrow \mathbf{B}_j, \quad j \in [k].$$

Hence the description of median-homomorphisms (4.1) boils down to describing median-homomorphisms of the form

$$f: \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$$

from a finite product of median algebras $\mathbf{A}_1, \dots, \mathbf{A}_n$ (thought of as median semilattices) to a median algebra \mathbf{B} . The general answer to this question still eludes us, but we can provide explicit descriptions of such mappings when \mathbf{B} is of a special type. We start by making some useful observations.

A function $f: \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$ is called an *n*-median-homomorphism if its “unary sections” $f_i: \mathbf{A}_i \rightarrow \mathbf{B}$ (obtained from f by fixing all but the i -th argument) are median-homomorphisms. It is easy to verify that every median-homomorphism is an n -median-homomorphism. However, the converse is not true. To see this, let $\mathbf{A}_1 = \mathbf{A}_2 = \mathbf{B}$ be the two element median algebra, and consider $f: \mathbf{A}_1 \times \mathbf{A}_2 \rightarrow \mathbf{B}$ given by $f(1,1) = f(0,1) = f(1,0) = 1$ and $f(0,0) = 0$. Clearly, f is a 2-median homomorphism but it is not a median-homomorphism: for $\mathbf{a} = (0,0)$, $\mathbf{b} = (0,1)$ and $\mathbf{c} = (1,0)$ we have

$$f(\mathbf{m}(\mathbf{a}, \mathbf{b}, \mathbf{c})) = f(0,0) \neq \mathbf{m}(1,1,0) = \mathbf{m}(f(0,1), f(1,0), f(0,0)).$$

Now we consider some particular cases, namely, when \mathbf{B} is a chain (or, equivalently, a conservative median algebra) and when it is a (2 : 3)-semilattice (or, equivalently, when \mathbf{B} is a tree).

We start with the case when \mathbf{B} (thought of as median semilattice) is a chain.

Proposition 4.2. *Let $\mathbf{A}_1, \dots, \mathbf{A}_n$ be median algebras and let \mathbf{B} be a chain with $|\mathbf{B}| \geq 2$. A mapping $f: \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$ is a median-homomorphism if and only if there is an $i \in [n]$ and a median-homomorphism $g: \mathbf{A}_i \rightarrow \mathbf{B}$ such that*

$$f(x_1, \dots, x_n) = g(x_i), \quad \text{for all } (x_1, \dots, x_n) \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n.$$

In other words, f is an essentially unary median-homomorphism.

Proof. Clearly, sufficiency holds. To prove necessity we show that f cannot depend on two different arguments.

For the sake of contradiction, suppose that f depends on at least 2 of its arguments. Without loss of generality, we may assume that they are the first two. Hence, there are $a, b \in \mathbf{A}_1$ and $\mathbf{c} \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$ such that $f(\mathbf{c}_1^a) > f(\mathbf{c}_1^b)$, where \mathbf{c}_i^x denotes the tuple obtained from \mathbf{c} by setting its i -th component to x .

Claim 4.3. For every $\mathbf{d} \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$, $f(\mathbf{d}_1^a) > f(\mathbf{d}_1^b)$.

Proof. Just consider the tuples $\mathbf{c}_1^a, \mathbf{c}_1^b, \mathbf{d}_1^a$ and $\mathbf{c}_1^a, \mathbf{c}_1^b, \mathbf{d}_1^b$, and use the fact that f is a median homomorphism. \square




FIGURE 1. A square as in proof of Proposition 4.5.

Similarly, there are $a', b' \in \mathbf{A}_2$ and $\mathbf{c}' \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$ such that $f(\mathbf{c}'_2^{a'}) > f(\mathbf{c}'_2^{b'})$. Again, we conclude that for every $\mathbf{d} \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$, we have

$$f(\mathbf{d}_2^{a'}) > f(\mathbf{d}_2^{b'}).$$

Observe that, in particular, $f(\mathbf{d}_{1,2}^{a,a'}) \geq f(\mathbf{d}_{1,2}^{b,a'})$, $f(\mathbf{d}_{1,2}^{a,b'}) \geq f(\mathbf{d}_{1,2}^{b,b'})$. As \mathbf{B} is a chain, we must have

$$f(\mathbf{d}_{1,2}^{b,a'}) \geq f(\mathbf{d}_{1,2}^{a,b'}) \quad \text{or} \quad f(\mathbf{d}_{1,2}^{b,a'}) \leq f(\mathbf{d}_{1,2}^{a,b'}).$$

Without loss of generality, suppose that \geq holds.

Claim 4.4. If f is a median homomorphism, then $f(\mathbf{d}_{1,2}^{a,a'}) = f(\mathbf{d}_{1,2}^{b,a'})$. Similarly, If f is a median homomorphism, then $f(\mathbf{d}_{1,2}^{a,b'}) = f(\mathbf{d}_{1,2}^{b,b'})$.

Proof. We prove the first claim; the second follows analogously. To see that we cannot have $>$, consider the tuples

$$f(\mathbf{d}_{1,2}^{a,a'}) > f(\mathbf{d}_{1,2}^{b,a'}) \geq f(\mathbf{d}_{1,2}^{a,b'})$$

to conclude that f cannot then be a median-homomorphism. \square

As Claim 4.3 and Claim 4.4 contradict one another, the proof of Proposition 4.2 is now complete. \square

The case when \mathbf{B} is a $(2 : 3)$ -semilattice (or, equivalently, when \mathbf{B} is a tree), follows from Proposition 4.2 by observing that if a and b are distinct elements of \mathbf{B} , then $[a, b]$ can be thought of as a chain (see Remark 3.1). In this case, we can reason as in the proof of Proposition 4.2 to obtain a more general result, namely, when \mathbf{B} is not necessarily a chain but a median algebra whose median graph is a tree.

Proposition 4.5. *Let $\mathbf{A}_1, \dots, \mathbf{A}_n$ be median algebras and let \mathbf{B} be a tree. A mapping $f : \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$ is a median-homomorphism if and only if there is an $i \in [n]$ and a median-homomorphism $g : \mathbf{A}_i \rightarrow \mathbf{B}$ such that*


$$f(x_1, \dots, x_n) = g(x_i), \quad \text{for all } (x_1, \dots, x_n) \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n.$$

In other words, f is an essentially unary median-homomorphism.

Proof. We provide an additional proof, alternative to that indicated in the paragraph preceding Proposition 4.5.

As in the proof of Proposition 4.2, suppose that f depends on at least 2 of its arguments. Without loss of generality, we may assume that they are the first two. Hence, there are $a, b \in \mathbf{A}_1$ and $\mathbf{c} \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$ such that $f(\mathbf{c}_1^a) \neq f(\mathbf{c}_1^b)$. As in Claim 4.3, it is easy to see that the same holds for all $\mathbf{d} \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$.

Similarly, if the second argument of f is essential, then are $a', b' \in \mathbf{A}_2$ and $\mathbf{c}' \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$ such that $f(\mathbf{c}'_2^{a'}) \neq f(\mathbf{c}'_2^{b'})$. Once again, we have in fact that $f(\mathbf{d}'_2^{a'}) \neq f(\mathbf{d}'_2^{b'})$, for all $\mathbf{d}' \in \mathbf{A}_1 \times \cdots \times \mathbf{A}_n$.

FIGURE 2. Examples of \wedge -semilattices.

Hence, we have that $f(\mathbf{c}_{1,2}^{x,y})$ for $(x, y) \in \{a, b\} \times \{a', b'\}$, are pairwise distinct. By picking p of the form $f(\mathbf{c}_{1,2}^{x,y})$, it then follows that the four points $f(\mathbf{c}_{1,2}^{x,y}) \in \mathbf{B}$ form a square as in Figure 4. This contradicts the fact that \mathbf{B} is a tree. \square

By Proposition 4.5 to describe median-homomorphisms of the form

$$f: \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$$

in the case when $\mathbf{A}_1, \dots, \mathbf{A}_n$ are median algebras and \mathbf{B} is a tree, it suffices to describe median homomorphisms $f: \mathbf{A} \rightarrow \mathbf{B}$ for a median algebra \mathbf{A} and a tree \mathbf{B} . Such descriptions follow from Theorem 2.4.

In the case when both \mathbf{A} and \mathbf{B} are trees, Theorem 2.4 together with Corollary 2.5 and Theorem 3.2, imply the following proposition.

Proposition 4.6. *Suppose that the median algebras \mathbf{A} and \mathbf{B} are trees, and consider a mapping $f: A \rightarrow B$. Then the following assertions are equivalent.*

- (i) f is a median-homomorphism.
- (ii) For all $a, b \in A$, the induced mapping $f: \langle [a, b], \leq_a \rangle \rightarrow \langle [f(a), f(b)], \leq_{f(a)} \rangle$ is an isotone function between chains.

In the case when \mathbf{A} is a tree and \mathbf{B} is an arbitrary median algebra, Theorem 2.4 can be strengthened by an additional and equivalent assertion, namely, the existential variant of its assertion (ii).

Theorem 4.7. *Let \mathbf{A} be a tree and \mathbf{B} an arbitrary median algebra. Then we have that one (or, equivalently, all) assertions in Theorem 2.4 are equivalent to*

- (ii') *There is a $p \in A$ such that $f: \langle A, \wedge_p \rangle \rightarrow \langle B, \wedge_{f(p)} \rangle$ is a semilattice-homomorphism.*

Proof. We show that (ii') \implies (i). In this way we have

$$(ii) \implies (ii') \implies (i) \implies (ii)$$

and thus they are all equivalent. So suppose that $\langle A, \wedge_p \rangle$ is a tree and that

$$f: \langle A, \wedge_p \rangle \rightarrow \langle B, \wedge_{f(p)} \rangle$$

is a semilattice-homomorphism. In particular, for every $a, b, c \in A$, we have that

$$\mathbf{m}(a, b, c) \in \{a \wedge_p b, b \wedge_p c, c \wedge_p a\},$$

and that $f: \langle A, \leq_p \rangle \rightarrow \langle B, \leq_{f(p)} \rangle$ is an order-homomorphism.

Without loss of generality, assume that $\mathbf{m}(a, b, c) = a \wedge_p b$. Hence,

$$a \wedge_p b = \mathbf{m}(a, b, c) = (a \wedge_p b) \vee_p (b \wedge_p c) \vee_p (c \wedge_p a) \geq_p b \wedge_p c, c \wedge_p a,$$

and, from the assumption that f is a semilattice-homomorphism,

$$f(\mathbf{m}(a, b, c)) = f(a \wedge_p b) = f(a) \wedge_{f(p)} f(b) \geq_{f(p)} f(b) \wedge_{f(p)} f(c), f(c) \wedge_{f(p)} f(a).$$

Therefore,

$$\begin{aligned} & \mathbf{m} \quad (f(a), f(b), f(c)) \\ & = \quad (f(a) \wedge_{f(p)} f(b)) \vee_{f(p)} (f(b) \wedge_{f(p)} f(c)) \vee_{f(p)} (f(c) \wedge_{f(p)} f(a)) \\ & = \quad f(a) \wedge_{f(p)} f(b) = f(\mathbf{m}(a, b, c)) \end{aligned}$$

and the proof of (ii') \implies (i), and thus of the theorem, is now complete. \square

Remark 4.8. Note that the existential variant of (iii) in Theorem 2.4 is not equivalent to (i) even in the case when \mathbf{A} is a tree. To see this, consider the function on \mathbf{A}_1 (see Figure 2(a)) that maps a, b to a , and leaves c and d fixed. Then it is an order-homomorphism for $p = a$, but it is not a median-homomorphism.

Remark 4.9. Note also that Theorem 4.7 does not necessarily hold in the case when \mathbf{A} is not a tree, even if \mathbf{B} is conservative. For instance, let \mathbf{A} be the median algebra given in Figure 2(b) and \mathbf{B} the 4-element chain $1, 2, 3, 4$, and consider the mapping that sends d, d' to 4, b, c to 2, and a to 1. Then, for all $p \in A$, $f: \langle A, \wedge_p \rangle \rightarrow \langle B, \wedge_{f(p)} \rangle$ is an order-homomorphism, but it is not a semilattice-homomorphism for $p = a$ (although it is a semilattice-homomorphism for $p = d'$).

Problem 1. Given an arbitrary median algebra \mathbf{B} , describe those median algebras \mathbf{A} for which Theorem 4.7 holds.

5. CONCLUDING REMARKS

In this paper we proposed a natural relaxation of conservativeness as considered in [8], which is of quite different flavour than that proposed in [3, 18], and showed that it axiomatizes the class of median algebras that are trees. Moreover, we considered mappings

$$f : \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}_1 \times \cdots \times \mathbf{B}_k$$

from products of arbitrary median algebras to products of $(2 : 3)$ -semilattices (or, equivalently, trees) and showed that those that preserve the median structure of $\mathbf{A}_1 \times \cdots \times \mathbf{A}_n$ can be decomposed into median-homomorphisms $g: \mathbf{A}_i \rightarrow \mathbf{B}_j$. The latter are described in Theorem 2.4 and, in the particular case when \mathbf{A}_i is a tree, also in Proposition 4.6 and in Theorem 4.7.

In the way, we looked into mappings

$$f : \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B},$$

where $\mathbf{A}_1, \dots, \mathbf{A}_n$ are arbitrary but where \mathbf{B} is a $(2 : 3)$ -semilattice, that is, \mathbf{B} is a tree. The description of such median-homomorphisms was then given in Proposition 4.5, from which it follows that they are essentially unary.

Looking at them as aggregation procedures that are not dictatorial (i.e., that depend on at least two arguments), this translates into an impossibility theorem variant to that of Arrow [1].

Now the natural question is to determine whether Proposition 4.5 still holds for an arbitrary median algebra \mathbf{B} . More precisely:

Problem 2. Given arbitrary median algebras $\mathbf{A}_1, \dots, \mathbf{A}_n$, describe those median algebras \mathbf{B} for which all median-homomorphisms $f : \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$ are trivial, i.e., essentially unary.

Towards a solution to this problem, consider the following example. Suppose that \mathbf{B} is an arbitrary median algebra thought as a \wedge -semilattice that is not a tree. Thus, there are $a, b \in B$ with a nontrivial upper bound c , i.e., $c > a, b > a \wedge b$. Set

$p = a \vee b$, $q = a \wedge b$. Note that, together with a and b , they form a square. Now, consider the median algebras $\mathbf{A}_1 = \mathbf{A}_2 = \{0, 1\}$. Then $f : \mathbf{A}_1 \times \mathbf{A}_2 \rightarrow \mathbf{B}$ given by

$$f(1, 1) = p, \quad f(0, 1) = a, \quad f(1, 0) = b, \quad f(0, 0) = q$$

is a median homomorphism that depends on both of its variables. In other words, this is a counter-example to Problem 2.

Now this example can be easily extended to mappings $f : \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$ and, as we have seen, the condition that \mathbf{B} is a $(2 : 3)$ -semilattice (i.e., a tree or, equivalently, does not contain a square as an order substructure) forces such median-preserving mappings to be essentially unary. From these considerations, we can thus provide an answer to Problem 2, namely:

Corollary 5.1. *All median-homomorphisms $f : \mathbf{A}_1 \times \cdots \times \mathbf{A}_n \rightarrow \mathbf{B}$ are essentially unary if and only if \mathbf{B} is a tree.*

REFERENCES

- [1] K.J. Arrow. A difficulty in the concept of social welfare. *Journal of Political Economy*, **58**(4):328–346, 1950.
- [2] S. P. Avann. Metric ternary distributive semi-lattices. *Proceedings of the American Mathematical Society*, **12**:407–414, 1961.
- [3] H. J. Bandelt and J. Hedlíková. Median algebras. *Discrete mathematics*, **45**:1–30, 1983.
- [4] H.-J Bandelt, M. Van De Vel. The median stabilization degree of a median algebra. *Journal of Algebraic Combinatorics*, **9**(2):115–127, 1999.
- [5] J.-P. Barthélemy. Caractérisation médiane des arbres. *Annals Discrete Math.*, **17**:39–46, 1983.
- [6] G. Birkhoff and S. A. Kiss. A ternary operation in distributive lattices. *Bulletin of the American Mathematical Society*, **53**:749–752, 1947.
- [7] G. Birkhoff. *Lattice Theory*, volume 25 of *American Mathematical Society Colloquium Publications*, revised edition. American Mathematical Society, New York, 1948.
- [8] M. Couceiro, J.-L. Marichal and B. Teheux, Conservative median algebras and semilattices. To appear in *Order*.
- [9] B. A. Davey and H. A. Priestley. *Introduction to lattices and order*. Cambridge University Press, New York, second edition, 2002.
- [10] E. Evans. *Median lattices and convex subalgebras*. In *Universal Algebra*, volume 29 of *Colloquia Mathematica Societatis János Bolyai*, pages 225–240. North-Holland, Amsterdam, 1982.
- [11] G. Grätzer. *General lattice theory*. Birkhäuser Verlag, Basel, second edition, 1998. New appendices by the author with B. A. Davey, R. Freese, B. Ganter, M. Greferath, P. Jipsen, H. A. Priestley, H. Rose, E. T. Schmidt, S. E. Schmidt, F. Wehrung and R. Wille.
- [12] J. R. Isbell. Median algebra. *Transactions of the American Mathematical Society*, **260**(2):319–362, 1980.
- [13] M. Kolibiar, T. Marcisová. On a question of J. Hashimoto. *Mat. Časopis*, **24**:179–185, 1974.
- [14] B. Monjardet. Caractérisations métriques des ensembles ordonnés semi-modulaires. *Math. Sci. Hum.*, **56**:77–87, 1976.
- [15] B. Monjardet. Théorie et applications de la médiane dans les treillis distributifs finis. *Annals Discrete Math.*, **9**:87–91, 1980.
- [16] H.M. Mulder, A. Schrijver. Median graphs and Helly hypergraphs. *Discrete Math.*, **25**, 41–50, 1979.
- [17] M. Sholander. Trees, lattices, order, and betweenness. *Proceedings of the American Mathematical Society*, **3**(3):369–381, 1952.
- [18] M. Sholander. Medians, lattices, and trees. *Proceedings of the American Mathematical Society*, **5**(5):808–812, 1954.
- [19] P.J. Slater Medians of arbitrary graphs *J. Graph Theory*, **4**:389–392, 1980.
- [20] E.R. Verheul. Multimedians in Metric and Normed Spaces. *CWI Tract*, Vol. 91, Centrum voor Wiskunde en Informatika, Amsterdam, The Netherlands, 1993.

LORIA (CNRS - INRIA NANCY GRAND EST - UNIVERSITÉ DE LORRAINE), ÉQUIPE OR-
PAILLEUR, BATIMENT B, CAMPUS SCIENTIFIQUE, B.P. 239, F-54506 VANDOEUVRE-LÈS-NANCY,
FRANCE

E-mail address: miguel.couceiro[at]inria.fr

TAMPERE UNIVERSITY OF TECHNOLOGY, PL 553, 33101 TAMPERE, FINLAND

E-mail address: sf[at]tut.fi

TEI OF EPIRUS, PO BOX 110, 47100 ARTA, GREECE

E-mail address: gmelet[at]teiep.gr