



HAL
open science

A compactness result for an equation with Holderian condition.

Samy Skander Bahoura

► **To cite this version:**

Samy Skander Bahoura. A compactness result for an equation with Holderian condition.. 2016.
hal-01187668v2

HAL Id: hal-01187668

<https://hal.science/hal-01187668v2>

Preprint submitted on 2 May 2016 (v2), last revised 18 Jan 2018 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A COMPACTNESS RESULT FOR AN EQUATION WITH HOLDERIAN CONDITION.

SAMY SKANDER BAHOURA

ABSTRACT. We give blow-up analysis for a Brezis and Merle's problem with Dirichlet and Hölderian condition. Also we derive a compactness criterion as in the work of Brezis and Merle.

Keywords: blow-up, boundary, Dirichlet condition, a priori estimate, Hölder condition.

1. INTRODUCTION AND MAIN RESULTS

We set $\Delta = -(\partial_{11} + \partial_{22})$ on open set Ω of \mathbb{R}^2 with a smooth boundary.

We consider the following equation:

$$(P) \begin{cases} \Delta u = V(1 + |x|^{2\beta})e^u & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

Here, we assume that:

$$0 \leq V \leq b < +\infty, \quad e^u \in L^1(\Omega) \text{ and } u \in W_0^{1,1}(\Omega),$$

and,

$$0 \in \partial\Omega, \quad \beta \in [0, 1/2).$$

We can see in [6] a nice formulation to this problem (P) in the sens of the distributions. This Problem arises in geometrical and physical problems, see for example [1, 2, 17, 18]. The above equation was studied by many authors, with or without the boundary condition, also for Riemannian surfaces, see [1-18], where one can find some existence and compactness results. In [5] we have the following important Theorem,

Theorem A(Brezis-Merle [5]). For $(u_i)_i$ and $(V_i)_i$ two sequences of functions relative to (P) with,

$$0 < a \leq V_i \leq b < +\infty$$

then it holds,

$$\sup_K u_i \leq c,$$

with c depending on a, b, β, K and Ω .

One can find in [5] an interior estimate if we assume $a = 0$, but we need an assumption on the integral of e^{u_i} , namely, we have:

Theorem B(Brezis-Merle [5]).*For $(u_i)_i$ and $(V_i)_i$ two sequences of functions relative to the problem (P) with,*

$$0 \leq V_i \leq b < +\infty \text{ and } \int_{\Omega} e^{u_i} dy \leq C,$$

then it holds;

$$\sup_K u_i \leq c,$$

with c depending on b, β, C, K and Ω .

We look to the uniform boundedness in all $\bar{\Omega}$ of the solutions of the Problem (P). When $a = 0$, the boundedness of $\int_{\Omega} e^{u_i}$ is a necessary condition in the problem (P) as showed in [5] by the following counterexample.

Theorem C(Brezis-Merle [5]).*There are two sequences $(u_i)_i$ and $(V_i)_i$ of the problem (P) with,*

$$0 \leq V_i \leq b < +\infty \text{ and } \int_{\Omega} e^{u_i} dy \leq C,$$

such that,

$$\sup_{\Omega} u_i \rightarrow +\infty.$$

When $\beta = 0$, the above equation has many properties in the constant and the Lipschitzian cases:

Note that for the problem (P) ($\beta = 0$), by using the Pohozaev identity, we can prove that $\int_{\Omega} e^{u_i}$ is uniformly bounded when $0 < a \leq V_i \leq b < +\infty$ and $\|\nabla V_i\|_{L^\infty} \leq A$ and Ω starshaped, when $a = 0$ and $\nabla \log V_i$ is uniformly bounded, we can bound uniformly $\int_{\Omega} V_i e^{u_i}$. In [14], Ma-Wei have proved that those results stay true for all open sets not necessarily starshaped.

In [8] ($\beta = 0$) Chen-Li have proved that if $a = 0$ and $\nabla \log V_i$ is uniformly bounded, then the functions are uniformly bounded near the boundary.

In [8] ($\beta = 0$) Chen-Li have proved that if $a = 0$ and $\int_{\Omega} e^{u_i}$ is uniformly bounded and $\nabla \log V_i$ is uniformly bounded, then we have the compactness result directly. Ma-Wei in [14], extend this result in the case where $a > 0$.

If we assume V more regular, we can have another type of estimates, a sup + inf type inequalities. It was proved by Shafrir see [16], that, if $(u_i)_i, (V_i)_i$ are two sequences of functions solutions of the previous equation without assumption on the boundary and, $0 < a \leq V_i \leq b < +\infty$, then we have the following interior estimate:

$$C \left(\frac{a}{b} \right) \sup_K u_i + \inf_{\Omega} u_i \leq c = c(a, b, K, \Omega).$$

One can see in [9] an explicit value of $C \left(\frac{a}{b} \right) = \sqrt{\frac{a}{b}}$. In his proof, Shafrir has used a blow-up function, the Stokes formula and an isoperimetric inequality, see [2]. For Chen-Lin, they have used the blow-up analysis combined with some geometric type inequality for the integral curvature.

Now, if we suppose $(V_i)_i$ uniformly Lipschitzian with A the Lipschitz constant, then, $C(a/b) = 1$ and $c = c(a, b, A, K, \Omega)$, see Brezis-Li-Shafrir [4]. This result was extended for Hölderian sequences $(V_i)_i$ by Chen-Lin, see [9]. Also, one can see in [13], an extension of the Brezis-Li-Shafrir result to compact Riemannian surfaces without boundary. One can see in [14] explicit form, $(8\pi m, m \in \mathbb{N}^*$ exactly), for the numbers in front of the Dirac masses when the solutions blow-up. Here, the notion of isolated blow-up point is used.

In [7] we have some a priori estimates on the 2 and 3-spheres $\mathbb{S}_2, \mathbb{S}_3$.

Here we give the behavior of the blow-up points on the boundary and a proof of Brezis-Merle Problem when $\beta \geq 0$.

The Brezis-Merle Problem (see [5]) is:

Problem. Suppose that $V_i \rightarrow V$ in $C^0(\bar{\Omega})$ with $0 \leq V_i \leq b$ for some positive constant b . Also, we consider a sequence of solutions (u_i) of (P) relative to (V_i) such that,

$$\int_{\Omega} e^{u_i} dx \leq C,$$

is it possible to have:

$$\|u_i\|_{L^\infty} \leq C = C(b, \beta, C, V, \Omega)?$$

Here we give blow-up analysis for a sequence of solutions of the Problem (P) and a proof of compactness result for the Brezis-Merle's Problem when $\beta \geq 0$. We extend the result of Chen-Li [8]. For the blow-up analysis we assume that:

$$0 \leq V_i \leq b,$$

The condition $V_i \rightarrow V$ in $C^0(\bar{\Omega})$ is not necessary, but for the proof of the compactness result we assume that:

$$\|\nabla V_i\|_{L^\infty} \leq A.$$

Our main result are:

Theorem 1.1. *Assume that $\max_\Omega u_i \rightarrow +\infty$, where (u_i) are solutions of the problem (P) with:*

$$0 \leq V_i \leq b \text{ and } \int_\Omega e^{u_i} dx \leq C, \quad \forall i,$$

then, after passing to a subsequence, there is a function u , there is a number $N \in \mathbb{N}$ and N points $x_1, \dots, x_N \in \partial\Omega$, such that,

$$\partial_\nu u_i \rightarrow \partial_\nu u + \sum_{j=1}^N \alpha_j \delta_{x_j}, \quad \alpha_j \geq 4\pi, \text{ weakly in the sense of measures.}$$

$$u_i \rightarrow u \text{ in } C_{loc}^1(\bar{\Omega} - \{x_1, \dots, x_N\}).$$

Theorem 1.2. *Assume that (u_i) are solutions of (P) relative to (V_i) with the following conditions:*

$$0 \leq V_i \leq b, \quad \|\nabla V_i\|_{L^\infty} \leq A \text{ and } \int_\Omega e^{u_i} \leq C,$$

we have,

$$\|u_i\|_{L^\infty} \leq c(b, \beta, A, C, \Omega),$$

In the last theorem we extend the result of Chen-Li ($\beta = 0$). The proof of Chen-Li and Ma-Wei [8,15], use the moving-plane method ($\beta = 0$).

2. PROOF OF THE THEOREMS

Proof of theorem 1.1:

We have:

$$u_i \in W_0^{1,1}(\Omega).$$

Since $e^{u_i} \in L^1(\Omega)$ by the corollary 1 of Brezis-Merle's paper (see [4]) we have $e^{u_i} \in L^k(\Omega)$ for all $k > 2$ and the elliptic estimates of Agmon and the Sobolev embedding (see [1]) imply that:

$$u_i \in W^{2,k}(\Omega) \cap C^{1,\epsilon}(\bar{\Omega}).$$

We denote by $\partial_\nu u_i$ the inner normal derivative. By the maximum principle we have, $\partial_\nu u_i \geq 0$.

By the Stokes formula we have,

$$\int_{\partial\Omega} \partial_\nu u_i d\sigma \leq C,$$

We use the weak convergence in the space of Radon measures to have the existence of a nonnegative Radon measure μ such that,

$$\int_{\partial\Omega} \partial_\nu u_i \varphi d\sigma \rightarrow \mu(\varphi), \quad \forall \varphi \in C^0(\partial\Omega).$$

We take an $x_0 \in \partial\Omega$ such that, $\mu(x_0) < 4\pi$. For $\epsilon > 0$ small enough set $I_\epsilon = B(x_0, \epsilon) \cap \partial\Omega$. We choose a function η_ϵ such that,

$$\begin{cases} \eta_\epsilon \equiv 1, & \text{on } I_\epsilon, \quad 0 < \epsilon < \delta/2, \\ \eta_\epsilon \equiv 0, & \text{outside } I_{2\epsilon}, \\ 0 \leq \eta_\epsilon \leq 1, \\ \|\nabla \eta_\epsilon\|_{L^\infty(I_{2\epsilon})} \leq \frac{C_0(\Omega, x_0)}{\epsilon}. \end{cases}$$

We take a $\tilde{\eta}_\epsilon$ such that,

$$\begin{cases} \Delta \tilde{\eta}_\epsilon = 0 & \text{in } \Omega \subset \mathbb{R}^2, \\ \tilde{\eta}_\epsilon = \eta_\epsilon & \text{in } \partial\Omega. \end{cases}$$

Remark: We use the following steps in the construction of $\tilde{\eta}_\epsilon$:

We take a cutoff function η_0 in $B(0, 2)$ and we set $\eta_\epsilon(x) = \eta_0(|x - x_0|/\epsilon)$. We solve the Dirichlet Problem:

$$\begin{cases} \Delta \bar{\eta}_\epsilon = \Delta \eta_\epsilon & \text{in } \Omega \subset \mathbb{R}^2, \\ \bar{\eta}_\epsilon = 0 & \text{in } \partial\Omega. \end{cases}$$

and finally we set, $\tilde{\eta}_\epsilon = -\bar{\eta}_\epsilon + \eta_\epsilon$. Also, by the maximum principle and the elliptic estimates we have :

$$\|\nabla \tilde{\eta}_\epsilon\|_{L^\infty} \leq C(\|\eta_\epsilon\|_{L^\infty} + \|\nabla \eta_\epsilon\|_{L^\infty} + \|\Delta \eta_\epsilon\|_{L^\infty}) \leq \frac{C_1}{\epsilon^2},$$

with C_1 depends on Ω .

We use the following estimate, see [3, 6, 11, 18],

$$\|\nabla u_i\|_{L^q} \leq C_q, \quad \forall i \text{ and } 1 < q < 2.$$

We deduce from the last estimate that, (u_i) converge weakly in $W_0^{1,q}(\Omega)$, almost everywhere to a function $u \geq 0$ and $\int_{\Omega} e^u < +\infty$ (by Fatou lemma). Also, V_i weakly converge to a nonnegative function V in L^∞ . The function u is in $W_0^{1,q}(\Omega)$ solution of :

$$\begin{cases} \Delta u = V(1 + |x|^{2\beta})e^u \in L^1(\Omega) & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

According to the corollary 1 of Brezis-Merle's result, see [5], we have $e^{ku} \in L^1(\Omega)$, $k > 1$. By the elliptic estimates, we have $u \in C^1(\bar{\Omega})$.

For two C^1 functions f and g we denote by $\langle \nabla f | \nabla g \rangle$ the inner product of ∇f and ∇g .

We can write:

$$\Delta((u_i - u)\tilde{\eta}_\epsilon) = (1 + |x|^{2\beta})(V_i e^{u_i} - V e^u)\tilde{\eta}_\epsilon - 2 \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle. \quad (1)$$

We use the interior estimate of Brezis-Merle, see [5],

Step 1: Estimate of the integral of the first term of the right hand side of (1).

We use the Green formula between $\tilde{\eta}_\epsilon$ and u , we obtain,

$$\int_{\Omega} (1 + |x|^{2\beta})V e^u \tilde{\eta}_\epsilon dx = \int_{\partial\Omega} \partial_\nu u \eta_\epsilon \leq 4\epsilon \|\partial_\nu u\|_{L^\infty} = C\epsilon \quad (2)$$

We have,

$$\begin{cases} \Delta u_i = (1 + |x|^{2\beta})V_i e^{u_i} & \text{in } \Omega \subset \mathbb{R}^2, \\ u_i = 0 & \text{in } \partial\Omega. \end{cases}$$

We use the Green formula between u_i and $\tilde{\eta}_\epsilon$ to have:

$$\int_{\Omega} (1 + |x|^{2\beta})V_i e^{u_i} \tilde{\eta}_\epsilon dx = \int_{\partial\Omega} \partial_\nu u_i \eta_\epsilon d\sigma \rightarrow \mu(\eta_\epsilon) \leq \mu(I_{2\epsilon}) \leq 4\pi - \epsilon_0, \quad \epsilon_0 > 0 \quad (3)$$

From (2) and (3) we have for all $\epsilon > 0$ there is $i_0 = i_0(\epsilon)$ such that, for $i \geq i_0$,

$$\int_{\Omega} |(1 + |x|^{2\beta})(V_i e^{u_i} - V e^u)\tilde{\eta}_\epsilon| dx \leq 4\pi - \epsilon_0 + C\epsilon \quad (4)$$

Step 2: Estimate of integral of the second term of the right hand side of (1).

Let $\Sigma_\epsilon = \{x \in \Omega, d(x, \partial\Omega) = \epsilon^3\}$ and $\Omega_{\epsilon^3} = \{x \in \Omega, d(x, \partial\Omega) \geq \epsilon^3\}$, $\epsilon > 0$. Then, for ϵ small enough, Σ_ϵ is hypersurface.

The measure of $\Omega - \Omega_{\epsilon^3}$ is $k_2\epsilon^3 \leq \mu_L(\Omega - \Omega_{\epsilon^3}) \leq k_1\epsilon^3$.

Remark: for the unit ball $\bar{B}(0, 1)$, our new manifold is $\bar{B}(0, 1 - \epsilon^3)$.

We write,

$$\int_{\Omega} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx = \int_{\Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx + \int_{\Omega - \Omega_{\epsilon^3}} \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx. \quad (5)$$

Step 2.1: Estimate of $\int_{\Omega - \Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx$.

First, we know from the elliptic estimates that $\|\nabla \tilde{\eta}_\epsilon\|_{L^\infty} \leq C_1/\epsilon^2$, C_1 depends on Ω

We know that $(|\nabla u_i|)_i$ is bounded in L^q , $1 < q < 2$, we can extract from this sequence a subsequence which converge weakly to $h \in L^q$. But, we know that we have locally the uniform convergence to $|\nabla u|$ (by Brezis-Merle's theorem), then, $h = |\nabla u|$ a.e. Let q' be the conjugate of q .

We have, $\forall f \in L^{q'}(\Omega)$

$$\int_{\Omega} |\nabla u_i| f dx \rightarrow \int_{\Omega} |\nabla u| f dx$$

If we take $f = 1_{\Omega - \Omega_{\epsilon^3}}$, we have:

$$\text{for } \epsilon > 0 \exists i_1 = i_1(\epsilon) \in \mathbb{N}, \quad i \geq i_1, \quad \int_{\Omega - \Omega_{\epsilon^3}} |\nabla u_i| \leq \int_{\Omega - \Omega_{\epsilon^3}} |\nabla u| + \epsilon^3.$$

Then, for $i \geq i_1(\epsilon)$,

$$\int_{\Omega - \Omega_{\epsilon^3}} |\nabla u_i| \leq \text{mes}(\Omega - \Omega_{\epsilon^3}) \|\nabla u\|_{L^\infty} + \epsilon^3 = \epsilon^3(k_1 \|\nabla u\|_{L^\infty} + 1).$$

Thus, we obtain,

$$\int_{\Omega - \Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx \leq \epsilon C_1 (2k_1 \|\nabla u\|_{L^\infty} + 1) \quad (6)$$

The constant C_1 does not depend on ϵ but on Ω .

Step 2.2: Estimate of $\int_{\Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx$.

We know that, $\Omega_\epsilon \subset\subset \Omega$, and (because of Brezis-Merle's interior estimates) $u_i \rightarrow u$ in $C^1(\Omega_{\epsilon^3})$. We have,

$$\|\nabla(u_i - u)\|_{L^\infty(\Omega_{\epsilon^3})} \leq \epsilon^3, \text{ for } i \geq i_3 = i_3(\epsilon).$$

We write,

$$\int_{\Omega_{\epsilon^3}} |\langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle| dx \leq \|\nabla(u_i - u)\|_{L^\infty(\Omega_{\epsilon^3})} \|\nabla \tilde{\eta}_\epsilon\|_{L^\infty} \leq C_1 \epsilon \text{ for } i \geq i_3,$$

For $\epsilon > 0$, we have for $i \in \mathbb{N}$, $i \geq \max\{i_1, i_2, i_3\}$,

$$\int_{\Omega} |\langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle| dx \leq \epsilon C_1 (2k_1 \|\nabla u\|_{L^\infty} + 2) \quad (7)$$

From (4) and (7), we have, for $\epsilon > 0$, there is $i_3 = i_3(\epsilon) \in \mathbb{N}$, $i_3 = \max\{i_0, i_1, i_2\}$ such that,

$$\int_{\Omega} |\Delta[(u_i - u)\tilde{\eta}_\epsilon]| dx \leq 4\pi - \epsilon_0 + \epsilon 2C_1 (2k_1 \|\nabla u\|_{L^\infty} + 2 + C) \quad (8)$$

We choose $\epsilon > 0$ small enough to have a good estimate of (1).

Indeed, we have:

$$\begin{cases} \Delta[(u_i - u)\tilde{\eta}_\epsilon] = g_{i,\epsilon} & \text{in } \Omega \subset \mathbb{R}^2, \\ (u_i - u)\tilde{\eta}_\epsilon = 0 & \text{in } \partial\Omega. \end{cases}$$

with $\|g_{i,\epsilon}\|_{L^1(\Omega)} \leq 4\pi - \frac{\epsilon_0}{2}$.

We can use Theorem 1 of [5] to conclude that there are $q \geq \tilde{q} > 1$ such that:

$$\int_{V_\epsilon(x_0)} e^{\tilde{q}(u_i - u)} dx \leq \int_{\Omega} e^{q(u_i - u)\tilde{\eta}_\epsilon} dx \leq C(\epsilon, \Omega).$$

where, $V_\epsilon(x_0)$ is a neighborhood of x_0 in $\bar{\Omega}$. Here we have used that in a neighborhood of x_0 by the elliptic estimates, $1 - C\epsilon \leq \tilde{\eta}_\epsilon \leq 1$. (We can take $(\Omega - \Omega_{\epsilon^3}) \cap B(x_0, \epsilon^3)$).

Thus, for each $x_0 \in \partial\Omega - \{\bar{x}_1, \dots, \bar{x}_m\}$ there is $\epsilon_{x_0} > 0$, $q_{x_0} > 1$ such that:

$$\int_{B(x_0, \epsilon_{x_0})} e^{q_{x_0} u_i} dx \leq C, \quad \forall i. \quad (9)$$

By the elliptic estimates (see [12]) $(u_i)_i$ is uniformly bounded in $W^{2,q_1}(V_\epsilon(x_0))$ and also, in $C^1(V_\epsilon(x_0))$. Finally, we have, for some $\epsilon > 0$ small enough,

$$\|u_i\|_{C^{1,\theta}[B(x_0,\epsilon)]} \leq c_3 \quad \forall i.$$

We have proved that, there is a finite number of points $\bar{x}_1, \dots, \bar{x}_m$ such that the sequence $(u_i)_i$ is locally uniformly bounded (in $C^{1,\theta}$, $\theta > 0$) in $\bar{\Omega} - \{\bar{x}_1, \dots, \bar{x}_m\}$.

Proof of theorem 1.2:

Without loss of generality, we can assume that 0 is a blow-up point. Also, by a conformal transformation, we can assume that $\Omega = B_1^+$, the half ball, and $\partial^+ B_1^+$ is the exterior part, a part which not contain 0 and on which u_i converge in the C^1 norm to u . Let us consider B_ϵ^+ , the half ball with radius $\epsilon > 0$.

We know that:

$$u_i \in W^{2,k}(\Omega) \cap C^{1,\epsilon}(\bar{\Omega}).$$

Thus we can use integrations by parts. The second Pohozaev identity applied around the blow-up 0 see for example [15] gives :

$$\int_{B_\epsilon^+} \Delta u_i \langle x | \nabla u_i \rangle dx = - \int_{\partial^+ B_\epsilon^+} g(\nabla u_i) d\sigma, \quad (10)$$

with,

$$g(\nabla u_i) = \langle \nu | \nabla u_i \rangle \langle x | \nabla u_i \rangle - \langle x | \nu \rangle \frac{|\nabla u_i|^2}{2}.$$

Thus,

$$\int_{B_\epsilon^+} V_i(1 + |x|^{2\beta})e^{u_i} \langle x | \nabla u_i \rangle dx = - \int_{\partial^+ B_\epsilon^+} g(\nabla u_i) d\sigma, \quad (11)$$

After integration by parts, we obtain:

$$\begin{aligned} \int_{B_\epsilon^+} 2V_i(1 + (1+\beta)|x|^{2\beta})e^{u_i} dx + \int_{B_\epsilon^+} \langle x | \nabla V_i \rangle (1 + |x|^{2\beta})e^{u_i} dx + \int_{\partial^+ B_\epsilon^+} \langle \nu | x \rangle (1 + |x|^{2\beta})V_i e^{u_i} d\sigma = \\ = \int_{\partial^+ B_\epsilon^+} g(\nabla u_i) d\sigma, \end{aligned} \quad (12)$$

Also, for u we have:

$$\begin{aligned} \int_{B_\epsilon^+} 2V(1 + |x|^{2\beta})e^u dx + \int_{B_\epsilon^+} \langle x|\nabla V \rangle (1 + |x|^{2\beta})e^u dx + \int_{\partial B_\epsilon^+} \langle \nu|x \rangle (1 + |x|^{2\beta})Ve^u d\sigma = \\ = \int_{\partial^+ B_\epsilon^+} g(\nabla u) d\sigma, \end{aligned} \quad (13)$$

We use the fact that $u_i = u = 0$ on $\{x_1 = 0\}$ and u_i, u are bounded in the C^1 norm outside a neighborhood of 0 to obtain:

$$(1 + o(\epsilon)) \left(\int_{B_\epsilon^+} V_i(1 + |x|^{2\beta})e^{u_i} dx \right) = \alpha_1 + o(\epsilon) + o(1) = o(1). \quad (14)$$

a contradiction.

REFERENCES

- [1] T. Aubin. Some Nonlinear Problems in Riemannian Geometry. Springer-Verlag, 1998.
- [2] C. Bandle. Isoperimetric Inequalities and Applications. Pitman, 1980.
- [3] L. Boccardo, T. Gallouet. Nonlinear elliptic and parabolic equations involving measure data. J. Funct. Anal. 87 no 1, (1989), 149-169.
- [4] H. Brezis, YY. Li and I. Shafrir. A sup+inf inequality for some nonlinear elliptic equations involving exponential nonlinearities. J.Funct.Anal.115 (1993) 344-358.
- [5] H. Brezis, F. Merle. Uniform estimates and Blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimension. Commun. in Partial Differential Equations, 16 (8 and 9), 1223-1253(1991).
- [6] H. Brezis, W. A. Strauss. Semi-linear second-order elliptic equations in L^1 . J. Math. Soc. Japan 25 (1973), 565-590.
- [7] Chang, Sun-Yung A, Gursky, Matthew J, Yang, Paul C. Scalar curvature equation on 2- and 3-spheres. Calc. Var. Partial Differential Equations 1 (1993), no. 2, 205-229.
- [8] W. Chen, C. Li. A priori estimates for solutions to nonlinear elliptic equations. Arch. Rational. Mech. Anal. 122 (1993) 145-157.
- [9] C-C. Chen, C-S. Lin. A sharp sup+inf inequality for a nonlinear elliptic equation in \mathbb{R}^2 . Commun. Anal. Geom. 6, No.1, 1-19 (1998).
- [10] D.G. De Figueiredo, P.L. Lions, R.D. Nussbaum, A priori Estimates and Existence of Positive Solutions of Semilinear Elliptic Equations, J. Math. Pures et Appl., vol 61, 1982, pp.41-63.
- [11] Ding.W, Jost. J, Li. J, Wang. G. The differential equation $\Delta u = 8\pi - 8\pi h e^u$ on a compact Riemann surface. Asian J. Math. 1 (1997), no. 2, 230-248.
- [12] D. Gilbarg, N. S. Trudinger. Elliptic Partial Differential Equations of Second order, Berlin Springer-Verlag.
- [13] YY. Li. Harnack Type Inequality: the method of moving planes. Commun. Math. Phys. 200,421-444 (1999).
- [14] YY. Li, I. Shafrir. Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two. Indiana. Math. J. Vol 3, no 4. (1994). 1255-1270.
- [15] L. Ma, J-C. Wei. Convergence for a Liouville equation. Comment. Math. Helv. 76 (2001) 506-514.
- [16] I. Shafrir. A sup+inf inequality for the equation $-\Delta u = Ve^u$. C. R. Acad.Sci. Paris Sér. I Math. 315 (1992), no. 2, 159-164.
- [17] Nagasaki, K, Suzuki,T. Asymptotic analysis for two-dimensional elliptic eigenvalue problems with exponentially dominated nonlinearities. Asymptotic Anal. 3 (1990), no. 2, 173-188.
- [18] Tarantello, G. Multiple condensate solutions for the Chern-Simons-Higgs theory. J. Math. Phys. 37 (1996), no. 8, 3769-3796.

DEPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ PIERRE ET MARIE CURIE, 2 PLACE JUSSIEU, 75005, PARIS, FRANCE.

E-mail address: samybahoura@yahoo.fr