



HAL
open science

Equidistribution of the conormal cycle of random nodal sets

Nguyen Viet Dang, Gabriel Riviere

► **To cite this version:**

Nguyen Viet Dang, Gabriel Riviere. Equidistribution of the conormal cycle of random nodal sets. 2015. hal-01189071v1

HAL Id: hal-01189071

<https://hal.science/hal-01189071v1>

Preprint submitted on 1 Sep 2015 (v1), last revised 31 Mar 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EQUIDISTRIBUTION OF THE CONORMAL CYCLE OF RANDOM NODAL SETS

NGUYEN VIET DANG AND GABRIEL RIVIÈRE

ABSTRACT. We study the asymptotic properties of the conormal cycle of nodal sets associated to a random superposition of eigenfunctions of the Laplacian on a smooth compact Riemannian manifold without boundary. In the case where the dimension is odd, we show that the expectation of the corresponding current of integration equidistributes on the fibers of the cotangent bundle. In the case where the dimension is even, we obtain an upper bound of lower order on the expectation. Using recent results of Alesker, we also deduce some properties on the asymptotic expectation of any smooth valuation including the Euler characteristic of random nodal sets.

1. INTRODUCTION

We consider (M, g) a smooth (\mathcal{C}^∞) compact connected oriented Riemannian manifold without boundary, and we study the solutions of

$$(1) \quad -\Delta_g e_\lambda = \lambda^2 e_\lambda,$$

where Δ_g is the Laplace-Beltrami operator induced by the Riemannian metric g and $\lambda > 0$. Our geometric assumption implies that there exists a non decreasing sequence

$$0 = \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \leq \lambda_j \dots \longrightarrow +\infty,$$

and an orthonormal basis $(e_j)_{j \geq 1}$ of $L^2(M)$ such that

$$\forall j \geq 1, \quad -\Delta_g e_j = \lambda_j^2 e_j.$$

More generally, we are in fact interested in finite superpositions of such solutions, i.e.

$$(2) \quad f \in \mathcal{H}_\Lambda := \mathbf{1}_{[0, \Lambda^2]}(-\Delta_g)L^2(M).$$

Recall that, if we denote by $N(\Lambda)$ the dimension of \mathcal{H}_Λ , Hörmander showed [38]

$$(3) \quad N(\Lambda) = \frac{\text{Vol}_g(M)}{2^n \pi^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} \Lambda^n + \mathcal{O}(\Lambda^{n-1}),$$

where n is the dimension of M and Vol_g is the Riemannian volume induced by g . In the present article, we aim at studying elements in \mathcal{H}_Λ in the high-frequency limit $\Lambda \rightarrow +\infty$. There are several natural ways to study these objects: among others, one can for instance look at their L^p norms, their nodal sets, their nodal domains, or their quantum limits. In the present article, we will address the question of the geometry of the nodal sets,

$$\forall f \in \mathcal{H}_\Lambda, \quad \mathcal{N}_f := \{x \in M : f(x) = 0\}.$$

Again, there are many natural questions regarding the geometry of a nodal set \mathcal{N}_f or its distribution on M – see for instance the book of Han and Lin [34] or the recent survey of Zelditch [61] on nodal sets of eigenfunctions. For instance, one can try to compute their Hausdorff volume, the number of their connected components, or more general geometric quantities like their Betti numbers.

The simplest question seems to be a priori the computation of the volume. In fact, in this case, it was conjectured by Yau [59] that there exist two constants $0 < c_g \leq C_g < +\infty$ such that, for every e_λ solution of (1),

$$c_g \lambda \leq \mathcal{V}^{n-1}(\mathcal{N}_{e_\lambda}) \leq C_g \lambda,$$

where \mathcal{V}^{n-1} is the Riemannian hypersurface volume. In the case of real-analytic metrics, this conjecture was proved by Donnelly and Fefferman [25] – see also [40] for the case of arbitrary elements in \mathcal{H}_Λ . In the \mathcal{C}^∞ case, the lower bound was proved earlier in dimension 2 by Brüning [16] while in dimension $n \geq 3$, the best known lower bounds are of order $\lambda^{\frac{3-n}{2}}$ as recently proved by Colding and Minicozzi [21] and Hezari, Sogge and Zelditch [37, 56]. Concerning the upper bound in the \mathcal{C}^∞ case, Dong, Donnelly and Fefferman obtained an upper bound of order $\lambda^{\frac{3}{2}}$ in dimension 2 [26, 24], and in dimension $n \geq 3$, Hardt and Simon proved an upper bound of order λ^{c_λ} for some $c > 0$ [35]. Thus, even the a priori simpler question of estimating the volume of nodal sets remains to our knowledge far from being completely understood in a general setting.

A natural approach is to ask whether the properties expected for any nodal set are valid for a “generic” solution of (1) or at least for a “generic” element f of \mathcal{H}_Λ . In order to formulate a notion of genericity in \mathcal{H}_Λ , we introduce the following Gaussian probability measure on the space \mathcal{H}_Λ :

$$\mu_\Lambda(df) = d\mu_\Lambda(f) := e^{-\frac{N(\Lambda)\|f\|^2}{2}} \left(\frac{N(\Lambda)}{2\pi} \right)^{\frac{N(\Lambda)}{2}} dc_1 \dots dc_{N(\Lambda)}, \quad \text{with } f = \sum_{j=1}^{N(\Lambda)} c_j e_j.$$

Remark 1.1. We note that, with this convention, the expectation of the L^2 norm is

$$\int_{\mathcal{H}_\Lambda} \|f\|_{L^2(M)}^2 d\mu_\Lambda(f) = 1.$$

In this setting, Bérard gave an “average” version [3] of Yau’s conjecture, i.e. one has

$$\int_{\mathcal{H}_\Lambda} \mathcal{V}^{n-1}(\mathcal{N}_f) d\mu_\Lambda(f) = \frac{\text{Vol}(\mathbb{S}^{n-1})}{\text{Vol}(\mathbb{S}^n)} \frac{\Lambda}{\sqrt{n+2}} \text{Vol}_g(M) (1 + o(1)).$$

More recently, Rudnick and Wigman also estimated the variance in the case where (M, g) is a rational torus [51] or the canonical sphere [58], and Letendre generalized Bérard’s result to the intersection of several “random” nodal sets [45]. In [60], Zelditch also showed that something slightly stronger than Bérard’s result holds. More precisely, he proved that, given any smooth function ω on M , one has

$$(4) \quad \int_{\mathcal{H}_\Lambda} \langle f^*(\delta_0) \|df\|, \omega \rangle d\mu_\Lambda(f) = \frac{\text{Vol}(\mathbb{S}^{n-1})}{\text{Vol}(\mathbb{S}^n)} \frac{\Lambda}{\sqrt{n+2}} \int_M \omega(x) d\text{Vol}_g(M) (1 + o(1)),$$

where $f^*(\delta_0)$ is the pullback by f of the Dirac distribution, and $\langle \cdot, \cdot \rangle$ is the duality bracket. This result shows that, if we average over the Gaussian measure, the nodal sets become equidistributed in M .

Using this probabilistic approach, we can in fact say much more on the geometry and the topology of the nodal sets. In [47, 55, 48], Nazarov and Sodin computed the average value of the number of connected components of \mathcal{N}_f . More precisely, if we define by $N(f)$ the number of connected components of \mathcal{N}_f , one has

$$\forall \delta > 0, \quad \lim_{\Lambda \rightarrow +\infty} \mu_\Lambda \left(\left\{ f \in \mathcal{H}_\Lambda : \left| \frac{N(f)}{\Lambda^n} - a_g \right| > \delta \right\} \right) = 0,$$

for some *non explicit constant* $a_g > 0$ depending only on (M, g) . This is a very strong result in the sense that it does not only give an asymptotic value for the expectation but also a large deviation estimate. It is also slightly more precise in the sense that the result remains true if we count the number of connected components in some rescaled geodesic ball $B_g(x, R\Lambda^{-1})$ for some large $R > 0$ – see Theorem 5 in [55]. In [49], Nicolaescu also estimated the number of critical points of a random function f in \mathcal{H}_Λ : it gives in particular an upper bound on the value of a_g . Gayet and Welschinger improved some aspects of Nazarov and Sodin results as follows. They proved that, on any smooth compact connected Riemannian manifold (M, g) without boundary, one can find constants $0 < b(M, g) \leq B(M, g) < +\infty$ such that

$$\forall 0 \leq i \leq n-1, \quad b(M, g)\Lambda^n \leq \int_{\mathcal{H}_\Lambda} b_i(\mathcal{N}_f) d\mu_\Lambda(f) \leq B(M, g)\Lambda^n,$$

where b_i is the i -th Betti number [30, 31]. Recall that $b_0(\mathcal{N}_f) = N(f)$. Their result is in fact more general than what we claim in the sense that it is valid for any elliptic pseudo-differential operator of positive order¹ m , and that they obtain some explicit bounds for the values of $b(M, g)$ and $B(M, g)$. In the case where n is odd, Letendre [45] showed the following asymptotic property of the Euler characteristic $\chi(\mathcal{N}_f)$ of a random nodal set:

$$(5) \quad \int_{\mathcal{H}_\Lambda} \chi(\mathcal{N}_f) d\mu_\Lambda(f) = \frac{2(-1)^{\frac{n-1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})} \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \text{Vol}_g(M) + \mathcal{O}(\Lambda^{n-1}).$$

We recall that the Euler characteristic is given by the alternate sum of Betti numbers. Finally, Canzani, Sarnak and Wigman recently studied the universal laws satisfied by the topologies of nodal sets [52, 17].

These different results show that much more can be said on the geometry of nodal sets if one only aims at probabilistic results. Even if we do not address this kind of questions in the present article, we note that probabilistic approaches have been considered for a long time in algebraic geometry – see for instance [41] where Kac estimated the number of real zeros of a random polynomial as its degree goes to infinity. In “random” algebraic geometry, one aims at studying the zero set of random polynomials of several variables instead of random superposition of eigenfunctions, and the spectral parameter Λ is replaced by the degree of the polynomials [44, 54, 9, 10, 29, 45].

¹In this case, the asymptotics are of order $\Lambda^{\frac{2n}{m}}$.

So far, we have only discussed the mathematical point of view of random Gaussian superposition of high-frequency eigenfunctions but we emphasize that they play an important role in the physics literature. For instance, in quantum chaos, one is interested in understanding the semiclassical properties of a quantum system whose underlying classical system enjoys chaotic features such as ergodicity or mixing. In [6], Berry predicted that semiclassical eigenmodes of chaotic systems should exhibit the same behaviour as a random superposition of waves. This is known as the *Berry random wave conjecture*. In particular, any result on random superposition of waves should provide an intuition on what one could expect for a chaotic system. In our framework, the main example of chaotic system is given by manifolds with negative curvature whose geodesic flows are exponentially mixing. Motivated by this conjecture of Berry, random nodal sets (and nodal domains²) were also extensively studied in the physics literature especially in the last fifteen years starting from the works of Blum, Gnutzmann and Smilansky [11] on nodal domains. The case of nodal sets was studied by Berry in [7] where he considered (on 2-dimensional domains) similar questions as the ones mentioned above on the volume and the curvature of random nodal sets. In the 2-dimensional case, Bogomolny, Dubertrand and Schmit conjectured that random nodal sets are in fact well described in terms of percolation processes and Schramm-Loewner Evolution models [12].

2. STATEMENT OF THE MAIN RESULTS

In the sequel, we denote by T^*M the cotangent bundle of M and an element of T^*M is denoted by $(x; \xi)$ for $x \in M, \xi \in T_x^*M$. From the point of view of “microlocal geometry”, it is often more natural to consider the conormal cycle of $\mathcal{N}_f = \{f = 0\}$ rather than the set \mathcal{N}_f itself [43, Chapter 9]. Recall that the conormal cycle is defined as follows, for every f in \mathcal{H}_Λ ,

$$(6) \quad N^*(\{f = 0\}) = \{(x; \xi) \in T^*M \text{ s.t. } f(x) = 0, \xi = td_x f \text{ for some } t \neq 0\} \subset T^*M.$$

Note that the above defines the conormal only in the set theoretical sense and we will later discuss the central issue of orientation. Outside of its singular points, this subset of T^*M is a *Lagrangian conical submanifold* and it contains in some sense more informations on the geometry of the nodal sets. For that reason, it sounds natural to us to focus on the properties of this set. Before giving more explanations on the relevance of the conormal cycle in “microlocal geometry”, we mention, as a first motivation, a celebrated index Theorem of Kashiwara [43, Thm 9.5.3 p. 385]. In the real analytic case, this result expresses the Euler characteristic of \mathcal{N}_f as a Lagrangian intersection in T^*M between the conormal cycle of \mathcal{N}_f and the graph of dg for a generic function g on M :

$$(7) \quad [\text{Graph } dg] \cap [N^*(\{f = 0\})] = \chi(\mathcal{N}_f)$$

where the intersection is in the oriented sense.

To our knowledge, the properties of this set have not been studied neither from the deterministic point of view, nor from the probabilistic one. In this article, we aim at

²This means the subsets of M where f has a fixed sign.

studying the probabilistic properties of the set $N^*(\{f = 0\})$ which are somewhat easier to consider. For that purpose, we need to introduce the following subset of \mathcal{H}_Λ :

$$D_\Lambda := \{f \in \mathcal{H}_\Lambda : \exists x \in \mathcal{N}_f \text{ such that } d_x f = 0\}.$$

By a Sard type argument – see for instance paragraph 2.3 in [45], one can verify that $\mu_\Lambda(D_\Lambda) = 0$ for Λ large enough. Thus, for μ_Λ -a.e. f in \mathcal{H}_Λ , $N^*(\{f = 0\})$ is a smooth n -dimensional submanifold of

$$T^\bullet M := \{(x, \xi) \in T^*M : \xi \neq 0\}.$$

In particular, once we choose an orientation, $N^*(\{f = 0\})$ can be viewed as a n -current $[N^*(\{f = 0\})]$ in the sense of de Rham [22, 53]. It means that, for every smooth compactly supported n -form ω on $T^\bullet M$, we define

$$\langle [N^*(\{f = 0\})], \omega \rangle = \int_{N^*(\{f=0\})} \omega.$$

Remark 2.1. We note that $N^*(\{f = 0\})$ has two connected components for f in D_Λ :

$$N_\pm^*(\{f = 0\}) := \{(x, \xi) \in T^\bullet M \text{ s.t. } f(x) = 0 \text{ and } \xi = td_x f \text{ for some } t \in \mathbb{R}_\pm \setminus \{0\}\}.$$

We will discuss precisely orientability questions in paragraph 3.2. We will in fact use the classical conventions of [20, p. 682].

Our first main result shows that this defines in fact an L^1 function:

Theorem 2.2. *Let (M, g) be a smooth oriented connected compact Riemannian manifold without boundary of dimension n . Then, the map*

$$(8) \quad f \in \mathcal{H}_\Lambda \mapsto [N^*(\{f = 0\})] \in \mathcal{D}'_n(T^\bullet M)$$

is integrable with respect to the Gaussian measure μ_Λ . Equivalently, for every test form ω on $T^\bullet M$, the map

$$f \in \mathcal{H}_\Lambda \mapsto \langle [N^*(\{f = 0\})], \omega \rangle \in \mathbb{R}$$

belongs to $L^1(\mathcal{H}_\Lambda, d\mu_\Lambda)$.

In order to state our second result, let us denote by Ω_g the Riemannian volume form on (M, g) , by $\pi : T^*M \rightarrow M$ the natural projection, and by $\pi^*\Omega_g$ the pull-back of the Riemannian volume on T^*M . Then, our second theorem shows the following universal behaviour:

Theorem 2.3. *Let (M, g) be a smooth oriented connected compact Riemannian manifold without boundary of dimension n . Then, one has*

$$(9) \quad \int_{\mathcal{H}_\Lambda} [N^*(\{f = 0\})] d\mu_\Lambda(f) = C_n \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \pi^*\Omega_g + \mathcal{O}(\Lambda^{n-1}),$$

with

$$C_n = \frac{2(-1)^{\frac{n+1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})} \text{ if } n \text{ is odd, and } C_n = 0 \text{ otherwise.}$$

Equivalently, it means that, for every test form ω on T^*M , one has

$$\int_{\mathcal{H}_\Lambda} \langle [N^*(\{f=0\})], \omega \rangle d\mu_\Lambda(f) = C_n \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \int_{T^*M} \pi^* \Omega_g \wedge \omega + \mathcal{O}(\Lambda^{n-1}).$$

We note that an important feature of this result is that it shows a very different behaviour depending on whether the dimension is odd or not. In the case where n is odd, our theorem shows that *the submanifold $N^*(\{f=0\})$ becomes uniformly equidistributed on the fibers T_x^*M while, when n is even, we only obtain an upper bound of lower order.*

2.1. Applications: smooth valuations on manifolds. We will now explain the interest of the concept of conormal cycle by giving some applications of our equidistribution Theorem. First, we recall a celebrated result of Chern [20, equation (20) p. 679–683] later generalized by Fu for subanalytic sets [28, p. 832] (see also [50, Definition 4.4.1 and Proposition 4.4.2] for a detailed exposition). In some sense, the following result can be understood as a “microlocal version” of Gauss-Bonnet Theorem:

Theorem 2.4 (Chern [20]). *Let (M, g) be a smooth compact Riemannian manifold of odd dimension n . Then, there exists a smooth, compactly supported and closed n -form³ θ so that $\int_{\pi^{-1}(x)} \theta = 1, \forall x \in M$, and, for any smooth submanifold of codimension ≥ 1 , one has*

$$(10) \quad \chi(X) = \langle [N^*(X)], \theta \rangle$$

where $[N^*(X)]$ is the conormal cycle of X and where χ is its Euler characteristic.

Remark 2.5. In his seminal article [20], Chern considered in fact cycles on the unit tangent bundle $\mathbb{U}M$ called normal cycles and the geodesic curvature form is a $(n-1)$ form Π on $\mathbb{U}M$ so that $d\Pi = 0, \int_{\pi^{-1}(x)} \Pi = 1, \forall x \in M$. Since M is Riemannian, there is a fibration $p : T^*M \mapsto \mathbb{U}M$ whose fiber is the group $(\mathbb{R}_{>0}, \times)$. Choose any smooth function $\varphi \geq 0$ on $\mathbb{R}_{>0}$ s.t. $d\varphi$ is compactly supported and $\int_0^{+\infty} d\varphi = 1$, then the n -form $\theta = d\varphi \wedge p^*\Pi$ satisfies the assumptions of the Theorem.

From this result, we deduce an alternative proof of equality (5) which gives the mean Euler characteristic of random nodal sets :

Theorem 2.6 (Letendre [45]). *Let (M, g) be a smooth oriented compact Riemannian manifold of odd dimension n , for any function f set $\mathcal{N}_f = \{f=0\}$ then*

$$(11) \quad \int_{\mathcal{H}_\Lambda} \chi(\mathcal{N}_f) d\mu_\Lambda(f) = \frac{2(-1)^{\frac{n-1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})} \text{Vol}_g(M) \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n + \mathcal{O}(\Lambda^{n-1}).$$

Proof. Let θ be the n -form of Theorem 2.4. Then, one finds

$$\int_{\mathcal{H}_\Lambda} \chi(\mathcal{N}_f) d\mu_\Lambda(f) = \int_{\mathcal{H}_\Lambda} \langle [N^*(\{f=0\})], \theta \rangle d\mu_\Lambda(f)$$

³Such a form is often called a geodesic curvature form.

since the map $f \mapsto \langle [N^*(\{f = 0\})], \theta \rangle$ is integrable with respect to the measure $d\mu_\Lambda$ by Theorem 2.2. Hence by Theorem 2.3, we obtain an integral on the cotangent cone :

$$\begin{aligned} \int_{\mathcal{H}_\Lambda} \chi(\mathcal{N}_f) d\mu_\Lambda(f) &= \int_{T^\bullet M} C_n \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \pi^* \Omega_g \wedge \theta + \mathcal{O}(\Lambda^{n-1}) \\ &= \int_{x \in M} \left(\int_{T_x^\bullet M} (-1)^n \theta \right) \wedge C_n \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \pi^* \Omega_g + \mathcal{O}(\Lambda^{n-1}) \\ &= (-1)^n \int_{x \in M} C_n \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \Omega_g + \mathcal{O}(\Lambda^{n-1}) \\ &= \frac{2(-1)^n (-1)^{\frac{n+1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})} \text{Vol}(M) \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n + \mathcal{O}(\Lambda^{n-1}) \end{aligned}$$

where we used a Fubini Theorem on the cotangent cone $T^\bullet M$ and we integrated on the fibers of $T^\bullet M$ first. Note that since n is odd, $(-1)^n (-1)^{\frac{n+1}{2}} = (-1)^{\frac{3n+1}{2}} = (-1)^{\frac{n-1}{2}}$ which explains the constant found in the statement of our Theorem. \square

In [45], Letendre computed the Euler characteristic by expressing it as the integral over $\{f = 0\}$ of a certain curvature form *depending on f* as for instance in [19, Eq. (9)]. Here, instead of this approach, we use Theorem 2.4 which gives the Euler characteristic as the integral of a *fixed curvature form* over $N^*(\{f = 0\})$. In some sense, this point of view allows us to extend Letendre's Theorem into an equidistribution result in the same way as Zelditch's equidistribution result (4) generalized Bérard's result on the volume.

Chern's formula for the Euler characteristic of a smooth submanifold can in fact be understood in the more general framework of the theory of smooth valuations recently developed by Alesker [2] and that we will now briefly review. We consider X a smooth oriented manifold (not necessarily endowed with a Riemannian structure) and we denote by $\mathcal{P}(X)$ the set of all compact submanifolds of M with corners. In the terminology of [2], we say that a map

$$\phi : \mathcal{P}(X) \rightarrow \mathbb{C}$$

is a smooth valuation if it is a finitely additive functional, and if it satisfies certain continuity properties. For simplicity of exposition, we remain vague on these two notions which need to be defined carefully on $\mathcal{P}(X)$ – see Part II of [1] for details. As was already mentioned, smooth valuations on manifolds generalize classical concepts from integral geometry such as volumes, Euler characteristic and mixed volumes. One of the remarkable property of these valuations is that they can be represented as follows : for every smooth valuation ϕ , there exists a smooth differential n -form ω_ϕ on T^*X such that, for any P in $\mathcal{P}(X)$, one has

$$\phi(P) = \int_{N^*(P)} \omega_\phi,$$

where $N^*(P)$ is the conormal cycle of P [2]. We emphasize that the converse statement is also true, and that the n -form is a priori non unique. In the example of the Euler characteristic χ , Theorem 2.4 gives such a n -form as soon as we have fixed some arbitrary Riemannian metric on X .

Thanks to this interpretation of smooth valuations as integrals over conormal cycles, Theorems 2.2 and 2.3 imply the following corollary on smooth valuations:

Corollary 2.7. *Let (M, g) be a smooth connected compact Riemannian manifold without boundary of dimension n . Let ϕ be a smooth valuation in the sense of [1] such that ω_ϕ can be chosen compactly supported in $T^\bullet M$. Then, the map*

$$f \mapsto \phi(\mathcal{N}_f)$$

belongs to $L^1(\mathcal{H}_\Lambda, d\mu_\Lambda)$. Moreover, one has

$$(12) \quad \int_{\mathcal{H}_\Lambda} \phi(\mathcal{N}_f) d\mu_\Lambda(f) = C_n \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \int_{T^*M} \pi^* \Omega_g \wedge \omega_\phi + \mathcal{O}(\Lambda^{n-1}),$$

where

$$C_n = \frac{2(-1)^{\frac{n+1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})} \text{ if } n \text{ is odd, and } C_n = 0 \text{ otherwise.}$$

2.2. Further questions. Let us now mention natural questions that can be asked about the conormal cycles attached to nodal sets.

- In the spirit of [45], it would be natural to consider intersections of nodal sets and compute the expectation of their conormal cycle. This could probably be directly obtained with the methods of the article but would require some slightly more complicated combinatorial arguments. For simplicity of exposition, we decided to consider only one nodal set.
- Regarding Theorem 2.3, it would be natural to understand the asymptotics in the case where n is even. At least in the case where the set of closed geodesics is of zero Liouville measure, we believe that one should be able to compute the term of order Λ^{n-1} in the asymptotics using the fact that in this geometric framework, the remainders in Weyl's asymptotics are of order $o(\Lambda^{n-1})$. Understanding the geometric meaning of the lower order term (and also checking that it is a nonzero term) is probably more subtle than computing the leading term we obtain here.
- In this article, we only computed the expectation of the conormal cycle and it would be of course natural to look for variance or large deviations estimates in the spirit of [7, 51, 47, 58, 48].
- Finally, it would be natural to understand what happens in the deterministic case. Can one obtain at least some upper bounds on the conormal cycle of a deterministic nodal set? If yes, what are the rates, and do they also depend on the dimension? This kind of questions would require completely different techniques like the ones in [34].

2.3. Organization of the article. Our first task in the present article is to express the conormal cycle $[N^*(f = 0)]$ explicitly in terms of Dirac distributions and differential forms. This kind of representation appears in the book of Schwartz [53] and it is in some sense slightly more adapted to our problem than the geometrically appealing definitions appearing in the works of Kashiwara–Schapira, Fu, Alesker, Bernig ([5, 43, 42, 28]). We

give in section 3 a brief account on the properties of currents needed for our purpose and carefully justify the meaning of our formula using tools from microlocal analysis.

In section 4 which forms the core of our paper, we prove Theorems 2.2 and 2.3 using microlocal analysis and combinatorics.

Then, in section 5, building on tools from quantum field theory, we give a new integral formula which expresses the conormal cycle of a nodal set as an oscillatory integral in bosonic (odd) and fermionic (even) variables. Our formula makes use of the so called *Berezin integral* [3, 33, 23, 57]. This formula is inspired from a very general formula expressing integration currents as oscillatory integrals on odd and even variables used by A. Losev et al. in their works on instantonic quantum field theories [27, 46]. We give a leisurely introduction to the necessary tools to grasp the meaning of our formula. Using this very simple integral formula representing $[N^*(f = 0)]$, we give a fast derivation of the leading term of the asymptotics of $\int_{\mathcal{H}_\Lambda} [N^*(\{f = 0\})] d\mu_\Lambda(f)$ which becomes a simple exercise of Gaussian integration with respect to odd and even variables and completely avoids the heavy combinatorics of the second part. The introduction of even (fermionic) variables allows us to avoid complicated combinatorics involving sums over partitions at the expense of a little bit of abstraction. The calculations are similar to those encountered in quantum field theory on curved Riemannian manifolds. This somewhat alternative derivation of the main asymptotic formula is presented in this last part in a purely formal way and it relies on several inversions of integrals and a priori estimates on remainder terms which could in fact be justified by using the arguments of the second part of the article.

Finally, in the appendix, we briefly discuss a question of independent interest. More precisely, building on the proof of lemma 4.12, we give a functional analytic proof of Whitney embedding theorem.

Acknowledgements. We warmly thank Semyon Alesker and Stéphane Nonnenmacher for very useful discussions related to this work. We also thank Didier Robert and Jean-Yves Welschinger for giving inspiring talks in Lille on random waves from which this work started, and Patrick Popescu Pampu for explanations related to the Euler characteristic. This work was done while the first author was a post-doc in the framework of the CEMPI program (ANR-11-LABX-0007-01) from the Agence Nationale de la Recherche. The second author is also partially supported by the Agence Nationale de la Recherche through the Labex CEMPI (ANR-11-LABX-0007-01) and the ANR project GERASIC (ANR-13-BS01-0007-01).

3. A REPRESENTATION FORMULA FOR $[N^*(\{f = 0\})]$.

In this section, we give a representation formula for the conormal cycle in terms of pull-back of Dirac distributions. More generally, we give formulas for the current of integration over submanifolds defined by systems of equations of the form $g_1 = g_2 = \dots = g_d = 0$ for some sufficiently nice functions g_1, \dots, g_d . This kind of formulas can be found for instance in Schwartz's book [53] – section V.5 (example 3). The advantage of representing currents of integration with Dirac distributions is that they will then be easily integrated against

the Gaussian measure $d\mu_\Lambda$ on \mathcal{H}_Λ – see section 4. For an introduction to the theory of currents, we also refer to the books of de Rham [22] and Giaquinta–Modica–Soucek [32].

Before being more precise, let us introduce some terminology. For any smooth oriented manifold X of dimension m , we will denote by $(\mathcal{D}_d(X))_{0 \leq d \leq m}$ the smooth compactly supported d -forms on X and by $(\mathcal{D}'_d(X))_{0 \leq d \leq m}$ its dual, i.e. the set of currents of dimension d on X .

3.1. Representing submanifolds defined by equations. Before we discuss the representation of $[N^*(\{f = 0\})]$, we start mildly with the problem of representing the current associated to the zero set of a family of functions $(g_i)_{1 \leq i \leq d}$.

3.1.1. Currents on zero sets. In mathematical physics, one is often interested in studying subsets in X given by equations $\{G = 0\}$ for some $G = (g_i)_{1 \leq i \leq d}$. In our article, we will encounter two classes of currents associated to this type of submanifolds. The first class are currents *representable by integration on some smooth submanifolds* which form a subclass of the De Rham chains – see chapter 2 in [22]. Let X be a smooth oriented manifold of dimension m , let S be a smooth, oriented, compact submanifold of X of dimension $d \leq m$. This defines an integration current $[S]$ in $\mathcal{D}'_d(X)$ which acts on test forms $\omega \in \mathcal{D}_d(X)$ by

$$(13) \quad [S](\omega) = \int_S \omega.$$

The second class consists in Dirac distributions which are defined as follows. We denote by δ_0^d the usual Dirac distribution at $\{0\}$ on \mathbb{R}^d . This is defined by

$$(14) \quad \forall \varphi(x) dx^1 \wedge \cdots \wedge dx^d \in \mathcal{D}_d(\mathbb{R}^d), \langle \delta_0^d, \varphi dx^1 \wedge \cdots \wedge dx^d \rangle_{\mathbb{R}^d} = \varphi(0).$$

Now let us define the Dirac distribution $G^*(\delta_0^d) = \delta_0^d \circ G$ supported by the zeros of a function $G : X \rightarrow \mathbb{R}^d$.

Proposition 3.1. *Let $G = (g_i)_{1 \leq i \leq d} \in C^\infty(X)$ be such that, for all $x \in X$ satisfying $g_1(x) = \cdots = g_d(x) = 0$, the linear forms $d_x g_1(x), \dots, d_x g_d(x)$ are linearly independent. Then*

- (1) *the pull-back of $\delta_0^d \in \mathcal{D}'_d(\mathbb{R}^d)$ by G which is denoted by $G^* \delta_0^d$ (or also $\delta_0^d \circ G$) is well-defined,*
- (2) *the wave front set of $G^* \delta_0^d$ is contained in*

$$N^*(\{G = 0\}) = \{(x, \xi) \in T^*X : G(x) = 0, \xi = \tau \circ d_x G \text{ for some } \tau \neq 0 \in \mathbb{R}^d\}$$

- (3) *for every sequence $(T_p)_p$ which converges to δ_0^d in $\mathcal{D}'_{\{0\} \times (\mathbb{R}^d)^*}(\mathbb{R}^d)$, $G^* T_p \rightarrow G^* \delta_0^d$ in the normal topology of $\mathcal{D}'_{N_0^* \{G=0\}}(X)$ which implies convergence in the weak topology of $\mathcal{D}'(X)$.*

Remark 3.2. For the notion of wave front set and for the precise definition of the set $\mathcal{D}'_\Gamma(X)$ of distributions whose wave front set is contained in Γ , we refer the reader to section 8.2 of [39] (see also [14]) and for the definition of the normal topology on $\mathcal{D}'_\Gamma(X)$, we refer to [15].

Proof. Let $N_G = \{(y; \eta) \in T^*\mathbb{R} \text{ s.t. } G(x) = y \text{ and } \eta \circ d_x G = 0 \text{ for some } x \in X\}$ be the normal set of the map G . By assumption, N_G does not meet the set $\{0\} \times (\mathbb{R}^d)^* \subset T^*\mathbb{R}^d$ which is the wave front set of δ_0^d . Therefore by the pull-back Theorem of Hörmander – Th. 4.10 below, $G^*\delta_0^d = \delta^d \circ G$ is well-defined and its wave front set is included $G^*(\{0\} \times (\mathbb{R}^d)^*) = N_0^*(\{G = 0\})$. Finally, the last claim follows from the sequential continuity of the pull-back operation – Th. 8.2.4 in [39] or [15]. \square

Now the natural question which arises is how the integration current on $S = \{G = 0\}$ and the delta form $G^*\delta_0^d = \delta^d \circ G$ are related to each other. Note that the Dirac distribution of the submanifold $S = \{G = 0\}$ is obviously not the same as the integration current $[S]$ because we integrate **top forms** against $\delta_0^d \circ G$ but only $(m - d)$ forms on $[S]$.

3.1.2. Orientations on submanifolds. Before comparing these currents, we need to discuss briefly orientability questions. For any smooth manifold X (orientable or not), there exists a smooth principal \mathbb{Z}_2 bundle $O_X \mapsto X$ called the orientation bundle. Recall that if X is an orientable manifold, then there exists a volume form $\omega \in \Gamma(X, \Lambda^n T^*X)$ where $n = \dim X$ or equivalently there exists a global section of O_X . Indeed, every choice of volume form ω on X defines canonically a global section $[\omega]$ of the orientation bundle O_X , also $[\varphi\omega] = [\omega]$ for all $\varphi > 0$. Now, if we are given a submanifold $S \subset X$ defined by equations $g_1 = \dots = g_d = 0$ and if X is oriented by $[\omega_1] \in O_X$ and S by $[\omega_2] \in O_S$, we choose representatives $\omega_1 \in \Gamma(X, \Lambda^n T^*X)$ and $\omega_2 \in \Gamma(S, \Lambda^{n-d} T^*S)$. Then we say that the orientation of $dg_1 \wedge \dots \wedge dg_d$ is compatible with the pair of orientations $[\omega_1], [\omega_2]$ if $dg_1 \wedge \dots \wedge dg_d \wedge \omega_2|_S = \varphi\omega_1|_S$ for some smooth function φ s.t. $\varphi|_S > 0$.

3.1.3. Relation between the different types of currents. In the next proposition, we give a concrete description of the action of Dirac distributions $G^*\delta_0^d$ on differential forms.

Proposition 3.3. *Let $G = (g_i)_{1 \leq i \leq d}$ be a smooth map from X to \mathbb{R}^d such that the differentials $d_x g_i$ are linearly independent for every x satisfying $g_1(x) = \dots = g_d(x) = 0$.*

Then, for any test form $\omega \in \mathcal{D}_m(X)$ and for any form α in $\mathcal{D}_{m-d}(X)$ such that $dg_1 \wedge \dots \wedge dg_d \wedge \alpha = \omega$, one has

$$(15) \quad \langle G^*\delta_0^d, \omega \rangle = \int_{\{G=0\}} \alpha.$$

Proof. We first prove the claim in the case where $g = (g_i)_{1 \leq i \leq d}$ is a linear map from \mathbb{R}^n to \mathbb{R}^d of maximal rank. Then one can choose (g_1, \dots, g_d) to be the first d components of some system of linear coordinate functions $(g_1, \dots, g_d, \dots, g_n)$ on \mathbb{R}^n , in this case G is just the linear projection $\mathbb{R}^n \mapsto \mathbb{R}^d$ on the first d components and the zero set $\{G = 0\}$ is just $\{0\} \times \mathbb{R}^{n-d} \subset \mathbb{R}^n$. For any function ψ on \mathbb{R}^d , the pull-back of ψ by G simply reads

$$G^*\psi(g_1, \dots, g_n) = \psi(g_1, \dots, g_d) \otimes 1(g_{d+1}, \dots, g_n)$$

and therefore, by continuity of the pull-back operation [14], one has $\delta_0^d \circ G = \delta_0^d \otimes 1$. Any form ω has a unique representation in the basis $dg_1 \wedge \dots \wedge dg_n$ as

$$\omega = \omega_{1\dots n}(g_1, \dots, g_n) dg_1 \wedge \dots \wedge dg_n$$

therefore :

$$\begin{aligned} [\delta_0^d \circ G](\omega) &= [\delta_0^d \otimes 1(g_{d+1}, \dots, g_n)](\omega) \\ &= \int_{\mathbb{R}^{n-d}} \omega_{1\dots n}(0, \dots, 0, g_{d+1}, \dots, g_n) dg_{d+1} \wedge \dots \wedge dg_n. \end{aligned}$$

Note that any form α such that $dg_1 \wedge \dots \wedge dg_d \wedge \alpha = \omega$ will coincide with

$$\omega_{1\dots n}(0, \dots, 0, g_{d+1}, \dots, g_n) dg_{d+1} \wedge \dots \wedge dg_n$$

when restricted to $\{G = 0\} = \{0\} \times \mathbb{R}^{n-d}$. It follows that the pairing does not depend on the choice of α .

Let us go back to the manifold case. For any $m \in \{G = 0\}$, let U be some neighborhood of m on which the differentials dg_i are linearly independent. Then one can choose (g_1, \dots, g_d) to be the first d components of some system of local coordinate functions $(g_1, \dots, g_d, \dots, g_n)$ on U . What we just did was to define some local chart $\Phi : m \in U \subset X \mapsto (g_1(m), \dots, g_n(m)) \in \mathbb{R}^n$ on $U \subset X$ such that $G \circ \Phi^{-1} : \mathbb{R}^n \mapsto \mathbb{R}^d$ is just the linear projection on the first d components. Then the distribution $\delta_0^d \circ (G \circ \Phi^{-1})$ is well defined by our previous discussion since $G \circ \Phi^{-1}$ is linear. Then we have the identity :

$$\begin{aligned} \langle (G \circ \Phi^{-1})^* \delta_0^d, \omega \rangle &= \int_{\{G \circ \Phi^{-1} = 0\}} \alpha \\ \forall \alpha \text{ s.t. } dg_1 \circ \Phi^{-1} \wedge \dots \wedge dg_d \circ \Phi^{-1} \wedge \alpha &= \omega. \end{aligned}$$

We just constructed a delta distribution $(G \circ \Phi^{-1})^* \delta_0^d$ supported by some linear subspace and finally, to go back to the open subset U in the manifold X , we must pull-back the distribution $(G \circ \Phi^{-1})^* \delta_0^d$ by Φ . By the pull-back Theorem 4.10, this yields $\Phi^* (G \circ \Phi^{-1})^* \delta_0^d = G^* \delta_0^d$ and, for every α s.t. $dg_1 \wedge \dots \wedge dg_d \wedge \alpha = \omega$,

$$\langle G^* \delta_0^d, \omega \rangle = \int_{\{G=0\}} \alpha.$$

□

In fact, using the notations of paragraph 3.1.2 concerning the orientation of submanifolds defined by systems of equations, the following corollary holds :

Corollary 3.4. *Let X be a smooth oriented manifold with orientation $[\omega_1]$, let*

$$G := (g_i)_{1 \leq i \leq d} : X \rightarrow \mathbb{R}^d$$

be a smooth function such that the differentials $d_x g_i$ are linearly independent for every $x \in X$ satisfying $g_1(x) = \dots = g_d(x) = 0$. Set S to be the submanifold in X defined by the regular system of equations $\{g_1 = \dots = g_d = 0\}$ and oriented by $[\omega_2]$. If $dg_1 \wedge \dots \wedge dg_d$ has orientation compatible with the pair of orientations $[\omega_1], [\omega_2]$ then one has

$$(16) \quad [S] = G^*(\delta_0^d) dg_1 \wedge \dots \wedge dg_d.$$

Proof. We note that this corollary is proved in [53] (Section V.5) when $d = 1$. It is in fact a direct consequence of the property of the Dirac distribution $\delta_0^d \circ G$ established in proposition 3.3, i.e.

$$\langle G^* \delta_0^d dg_1 \wedge \cdots \wedge dg_d, \alpha \rangle = \langle G^* \delta_0^d, dg_1 \wedge \cdots \wedge dg_d \wedge \alpha \rangle = \int_S \alpha.$$

□

3.2. The conormal cycle. We will now explain how to use the above results in order to represent the conormal cycle $N^*(\{f = 0\})$ in terms of Dirac distributions. For that purpose, we first need to recall the definition of push-forward in the context of currents. Let $\lambda : X \mapsto Y$ be a proper smooth map between two smooth oriented manifolds X and Y , and let T be a current in $\mathcal{D}'_d(X)$, where $d \leq \dim Y$. Then the push-forward $\lambda_*(T)$ is defined by duality from the pull-back :

$$\forall \omega \in \mathcal{D}_d(Y), \langle \lambda_* T, \omega \rangle_Y = \langle T, \lambda^* \omega \rangle_X.$$

An important fact concerning the push-forward operation is that an orientation of T induces canonically an orientation of $\lambda_* T$.

3.2.1. Precise definition of the current of integration. Let us start by recalling the definition of the conormal cycle of $\{f = 0\}$:

Definition 3.5. Let M be a smooth connected compact oriented manifold of dimension n without boundary. Let $f \in C^\infty(M)$, if $d_x f$ never vanishes on $\{f = 0\}$ then we define the conormal cycle $[N^*(\{f = 0\})]$ as the integration current on the conical Lagrangian submanifold $\{(x; \xi) \text{ s.t. } f(x) = 0 \text{ and } \xi = td_x f \text{ for some } t \in \mathbb{R} \setminus \{0\}\}$ in $\mathcal{D}'_n(T^*M)$.

A subtle fact we must add to our definition concerns the **orientation of the conormal cycle**. First, note that the conormal cycle contains two connected components

$$N^*_\pm(\{f = 0\}) = \{(x; \xi) \text{ s.t. } f(x) = 0 \text{ and } \xi = td_x f \text{ for some } t \in \mathbb{R}_\pm \setminus \{0\}\}$$

since it does not meet the zero section of the cotangent bundle. Hence an orientation of $[N^*(\{f = 0\})]$ consists of a choice of an orientation for each connected component separately. In the sequel, we denote by \mathbb{R}^* the set $\mathbb{R} \setminus \{0\}$. Let us give another description of $[N^*(\{f = 0\})]$ in terms of Lagrange immersion which will be useful in the sequel since we will interpret the conormal cycle $[N^*(\{f = 0\})]$ as the push-forward of a cone $S \times \mathbb{R}^*$ and the orientation of $[N^*(\{f = 0\})]$ will be induced by an orientation of $S \times \mathbb{R}^*$:

Lemma 3.6. *Let $f \in C^\infty(M)$ where df never vanishes on $\{f = 0\}$. Set S to be the abstract manifold $\{f = 0\}$ and $i : S \hookrightarrow M$ the canonical immersion of S in M . Consider the cone $S \times \mathbb{R}^*$. Then, the conormal cycle $N^*(\{f = 0\})$ is the image of the immersion :*

$$(17) \quad \lambda : (x; t) \in S \times \mathbb{R}^* \mapsto (i(x); td_{i(x)} f) \in T^*M,$$

and, at the level of currents, we find that

$$(18) \quad [N^*(\{f = 0\})] = [\lambda(S \times \mathbb{R}^*)] = \lambda_*[S \times \mathbb{R}^*],$$

where we orient the current $[S \times \mathbb{R}^*]$ by any differential form $\alpha \in \Omega^n(S \times \mathbb{R}^*)$ such that $df \wedge \alpha|_{S \times \mathbb{R}_{>0}} = \Omega_g \wedge dt|_{S \times \mathbb{R}_{>0}}$ and $df \wedge \alpha|_{S \times \mathbb{R}_{<0}} = -\Omega_g \wedge dt|_{S \times \mathbb{R}_{<0}}$.

Remark 3.7. The cone $S \times \mathbb{R}^*$ contains two connected components $S \times \mathbb{R}_{>0}$ and $S \times \mathbb{R}_{<0}$ that we should orient separately.

Proof. By definition, $[N^*(\{f = 0\})]$ is just the integration current on $N^*(\{f = 0\})$ which is equal to $\lambda(S \times \mathbb{R}^*)$. Therefore, for any test form ω in $\mathcal{D}_n(T^\bullet M)$:

$$\begin{aligned} \langle [N^*(\{f = 0\})], \omega \rangle &= \int_{N^*(\{f=0\})} \omega = \int_{\lambda(S \times \mathbb{R}^*)} \omega \\ &= \int_{(S \times \mathbb{R}^*)} \lambda^* \omega = \langle [S \times \mathbb{R}^*], \lambda^* \omega \rangle = \langle \lambda_* [S \times \mathbb{R}^*], \omega \rangle \end{aligned}$$

by definition of the push-forward which yields the claim. \square

3.2.2. Expression in local coordinates. Our goal in this paragraph is to show that the conormal cycle $[N^*(\{f = 0\})]$ can be written (in local coordinates) as the current

$$(19) \quad \int_{t \in \mathbb{R}^*} \delta_0^{n+1} \circ \left(f, \xi - \frac{t}{\Lambda} df \right) df \wedge \bigwedge_{i=1}^n d \left(\frac{t}{\Lambda} \partial_{x_i} f - \xi_i \right),$$

where Λ is any positive number. We need to do this progressively and the first step is to define some integration current $[Z_{f,\Lambda}]$ on $\mathbb{R}^* \times T^\bullet M$ whose push-forward along the first factor \mathbb{R}^* yields the conormal cycle $[N^*(\{f = 0\})]$. For that purpose, we set

$$(20) \quad Z_{f,\Lambda} := \left\{ (t; x, \xi) \in \mathbb{R}^* \times T^\bullet M : f(x) = 0, \text{ and } \xi = \frac{t}{\Lambda} d_x f \right\}.$$

The following lemma holds:

Lemma 3.8. *Let $f \in C^\infty(M)$ where df never vanishes on $\{f = 0\}$. Let $[Z_{f,\Lambda}]$ be the integration current on the set $Z_{f,\Lambda}$ defined by equation (20) and π the projection $(t; x, \xi) \in \mathbb{R}^* \times T^\bullet M \mapsto (x, \xi) \in T^\bullet M$. Then*

$$(21) \quad [N^*(\{f = 0\})] = \pi_* [Z_{f,\Lambda}].$$

Proof. Let $\omega(x, \xi; dx, d\xi)$ be an element in $\mathcal{D}^n(T^*M)$. One has :

$$\begin{aligned}
 \langle \pi_*[Z_{f,\Lambda}], \omega \rangle &= \langle [Z_{f,\Lambda}], \pi^*\omega \rangle \\
 &= \int_{\{f(x)=0, \xi=\frac{t}{\Lambda}d_x f, t \in \mathbb{R}^*\}} \omega(x, \xi; dx, d\xi) \\
 &= \int_{\{f(x)=0, \xi=td_x f, t \in \mathbb{R}^*\}} \omega(x, \xi; dx, d\xi) \\
 &\quad \text{by variable change} \\
 &= \int_{\{f(x)=0, t \in \mathbb{R}^*\}} \omega(x, td_x f; dx, d(td_x f)) \\
 &= \int_{S \times \mathbb{R}^*} (\lambda^*\omega)(t, x; dt, dx) \\
 &\quad \text{since } (\lambda^*\omega)(t, x; dt, dx) = \omega(x, td_x f; dx, d(td_x f)) \\
 &= \lambda_*[S \times \mathbb{R}^*](\omega) = [N^*(\{f = 0\})](\omega) \\
 &\quad \text{by lemma 3.6.}
 \end{aligned}$$

□

The next lemma aims to represent the current $[Z_{f,\Lambda}]$ by Dirac distributions in a local coordinate chart. Let (U, ϕ) be a sufficiently small coordinate chart centered at a point x_0 in M . It induces a coordinate chart on T^*M as follows:

$$\Phi := T^*\phi : T^*U \rightarrow \phi(U) \times \mathbb{R}^n, \quad (x, \xi) \mapsto (y, \eta) = (\phi(x), (d_x \phi^*)^{-1}\xi).$$

We also set

$$\mathring{T}^*U := \{(y, \eta) \in \phi(U) \times \mathbb{R}^n : \eta \neq 0\}.$$

In the following of the article, we will use these conventions. In particular, (x, ξ) will always denote a point in T^*M and (y, η) its image in local coordinates. Using these conventions, one has

Lemma 3.9. *Let $f \in C^\infty(M)$ where df never vanishes on $\{f = 0\}$. Then, for any system of local coordinates $(y; \eta)$ on \mathring{T}^*U where $U \subset M$ is some bounded open subset, one finds that, in these local coordinates,*

$$(22) \quad [Z_{f,\Lambda}]|_{\mathbb{R}^* \times \mathring{T}^*U} = \delta_0^{n+1} \circ \left(f(y), \frac{t}{\Lambda} d_y f - \eta \right) df \wedge \bigwedge_{i=1}^n d \left(\frac{t}{\Lambda} \partial_{y_i} f - \eta_i \right)$$

Proof. The submanifold $Z_{f,\Lambda}$ in $\mathbb{R}^* \times T^*U$ is defined in the local coordinates (y, η) by the system of $(n+1)$ equations :

$$\begin{aligned}
 f(y) &= 0 \\
 \frac{t}{\Lambda} \frac{\partial f}{\partial y^i} - \eta_i &= 0, 1 \leq i \leq n.
 \end{aligned}$$

The collection of one forms $\left(d_y f, d\eta_i - d\left(\frac{t}{\Lambda} \frac{\partial f}{\partial y^i}\right)\right)_{1 \leq i \leq n}$ is linearly independent along $Z_{f,\Lambda}$ as

$$d_y f \wedge \bigwedge_{i=1}^n \left(d\eta_i - d\left(\frac{t}{\Lambda} \frac{\partial f}{\partial y^i}\right)\right) = d_y f \wedge d\eta_1 \wedge \dots \wedge d\eta_n + \text{lower order terms in } d\eta,$$

where $d_y f \wedge d\eta_1 \wedge \dots \wedge d\eta_n$ does not vanish on $Z_{f,\Lambda}$ since df does not vanish on $\{f = 0\}$. Then by Corollary 3.4, the current $[Z_{f,\Lambda}]$ is represented in local coordinates (y, η) by the formula :

$$(23) \quad [Z_{f,\Lambda}]|_{\mathbb{R}^* \times \dot{T}^*U} = \delta_0^{n+1} \circ \left(f(y), \frac{t}{\Lambda} d_y f - \eta\right) d_y f \wedge \bigwedge_{i=1}^n d\left(\frac{t}{\Lambda} \partial_{y^i} f - \eta_i\right).$$

□

Then combining the above Lemma and Lemma 3.8 which claimed the identity $[N^*(\{f = 0\})] = \pi_*[Z_{f,\Lambda}]$, we conclude this first part by the following :

Corollary 3.10. *Let $f \in C^\infty(M)$ where df never vanishes on $\{f = 0\}$. Then, for any system of local coordinates (y, η) on \dot{T}^*U where $U \subset M$ is some bounded open subset, the restriction of the conormal cycle $[N^*(\{f = 0\})]$ on $T^\bullet U$ can be described in the local coordinates (y, η) by the integral formula :*

$$(24) \quad [N^*(\{f = 0\})]|_{T^\bullet U} = \int_{t \in \mathbb{R}^*} \delta_0^{n+1} \circ \left(f(y), \frac{t}{\Lambda} d_y f - \eta\right) d_y f \wedge \bigwedge_{i=1}^n d\left(\frac{t}{\Lambda} \partial_{y^i} f - \eta_i\right)$$

where we use the convention $\int_{t \in \mathbb{R}^*} = \int_0^{+\infty} + \int_0^{-\infty}$.

Our convention for the integral $\int_{t \in \mathbb{R}^*} = \int_0^{+\infty} + \int_0^{-\infty}$ reflects our choice of orientation for the conormal cycle. We note that we use in fact the same conventions as in [20, p. 682].

3.2.3. Discussion on the orientation of the conormal cycle. We should finally give convincing arguments that our choice of orientation of \mathbb{R}^* is the right one. There are four orientation choices that we should list :

$$(25) \quad \int_{\mathbb{R}^*} = \int_0^{-\infty} + \int_0^{+\infty}$$

$$(26) \quad \int_{\mathbb{R}^*} = \int_{-\infty}^0 + \int_{+\infty}^0$$

$$(27) \quad \int_{\mathbb{R}^*} = \int_{-\infty}^0 + \int_0^{+\infty}$$

$$(28) \quad \int_{\mathbb{R}^*} = \int_0^{-\infty} + \int_{+\infty}^0.$$

First note that equations (25),(26) and equations (27),(28) define opposite pairs of orientations. Let us explain why we should exclude the pair (27),(28). Assume that $[N^*\{f =$

$0\}$] = $\int_{-\infty}^0 [Z_{f,\Lambda}] + \int_0^{+\infty} [Z_{f,\Lambda}]$, then a quick calculation shows that :

$$\begin{aligned}
[N^*\{-f = 0\}] &= \int_{-\infty}^0 [Z_{-f,\Lambda}] + \int_0^{+\infty} [Z_{-f,\Lambda}] \\
&= \int_{-\infty}^{+\infty} \delta_0^{n+1} \circ \left(-f(y), \frac{t}{\Lambda}(-d_y f) - \eta \right) (-d_y f) \wedge \bigwedge_{i=1}^n d \left(\frac{t}{\Lambda}(-\partial_{y^i} f) - \eta_i \right) \\
&= \int_{-\infty}^{+\infty} \delta_0^{n+1} \circ \left(f(y), \frac{-t}{\Lambda}d_y f - \eta \right) (d_y f) \wedge \bigwedge_{i=1}^n d \left(\frac{-t}{\Lambda}\partial_{y^i} f - \eta_i \right) \\
&= \int_{+\infty}^{-\infty} \delta_0^{n+1} \circ \left(f(y), \frac{t}{\Lambda}d_y f - \eta \right) d_y f \wedge \bigwedge_{i=1}^n d \left(\frac{t}{\Lambda}\partial_{y^i} f - \eta_i \right) \\
&= -[N^*\{f = 0\}]
\end{aligned}$$

The same relation $[N^*\{-f = 0\}] = -[N^*\{f = 0\}]$ would hold true if we chose the orientation defined by equation (28). But calculating the expectation of $N^*\{f = 0\}$ with respect to the Gaussian measure, we would then obtain :

$$\int_{\mathcal{H}_\Lambda} [N^*\{f = 0\}]d\mu(f) = \int_{\mathcal{H}_\Lambda} [N^*\{-f = 0\}]d\mu(f) = - \int_{\mathcal{H}_\Lambda} [N^*\{f = 0\}]d\mu(f) = 0$$

which is not interesting for us. Finally, the two relevant orientations in our case are given by the pair (25),(26) and we made the arbitrary choice of the orientation (25).

4. PROOF OF THE MAIN RESULTS

We will now give the complete proof of our two main results, namely Theorems 2.2 and 2.3. By a partition of unity argument, it is sufficient to prove these results in a local chart. Let (U, ϕ) be a sufficiently small geodesic normal coordinate chart centered at x_0 in M – see remark 4.5 below. We refer to paragraph 3.2 for the choice of conventions for local charts. For every f in \mathcal{H}_Λ , we define the following map on $\mathcal{H}_\Lambda \times \phi(U) \times \mathbb{R}^n \times \mathbb{R}^*$:

$$G_\Lambda : (f, y, \eta, t) \in \mathcal{H}_\Lambda \times \phi(U) \times \mathbb{R}^n \times \mathbb{R}^* \longmapsto \left(f(y); \frac{t}{\Lambda}\partial_{y_1} f - \eta_1, \dots, \frac{t}{\Lambda}\partial_{y_n} f - \eta_n \right) \in \mathbb{R} \times \mathbb{R}^n,$$

and we introduce the following subset of \mathcal{H}_Λ :

$$D_\Lambda := \{f \in \mathcal{H}_\Lambda : \exists y \in f^{-1}(0) \text{ such that } d_y f = 0\}.$$

By a Sard type argument – see for instance paragraph 2.3 in [45], one can verify that $\mu_\Lambda(D_\Lambda) = 0$ for Λ large enough. One can also verify that $\Omega_\Lambda = \mathcal{H}_\Lambda \setminus D_\Lambda$ is an open subset of \mathcal{H}_Λ . We now define the following subset of \mathring{T}^*U :

$$\forall f \in \mathcal{H}_\Lambda, N^*(\{f = 0\})_U := \left\{ (y, \eta) \in \mathring{T}^*U : G_\Lambda(f, y, \eta, t) = 0, \text{ for some } t \neq 0 \right\},$$

which is just the conormal cycle of $f^{-1}(0)$ restricted over U and which was defined as the push-forward of a certain set $Z_{f,\Lambda}$ defined in paragraph 3.2. For simplicity, we will make in the following a small abuse of notations and omit the subscript U .

Recall that we aim at computing the expectation of the current of integration on the submanifold $N^*(\{f=0\})$. According to corollary 3.10, one knows that, for every f in Ω_Λ , and for every ω in $\mathcal{D}_n(\dot{T}^*U)$, one has

$$\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) = \left\langle G_\Lambda(f)^*(\delta_0^{n+1}), df \wedge \bigwedge_{j=1}^n d\left(\frac{t}{\Lambda} \partial_{y_j} f - \eta_j\right) \wedge \omega(y, \eta, dy, d\eta) \right\rangle_{\dot{T}^*U \times \mathbb{R}^*},$$

where $\langle \cdot, \cdot \rangle_{\dot{T}^*U \times \mathbb{R}^*}$ is the duality bracket in $\mathcal{D}'(\dot{T}^*U \times \mathbb{R}^*) \times \mathcal{D}(\dot{T}^*U \times \mathbb{R}^*)$. Recall that we chose a particular orientation for \mathbb{R}^* in paragraph 3.2.

The main result of this section is the following Theorem from which Theorems 2.2 and 2.3 follow by partition of unity:

Theorem 4.1. *We use the above notations. For any element*

$$\omega := \sum_{|\alpha|+|\beta|=n} \omega_{\alpha,\beta}(y, \eta) dy^\alpha \wedge d\eta_\beta$$

in $\mathcal{D}_n(\dot{T}^*U)$, let

$$\|\omega\| = \sup_{\alpha,\beta,(y,\eta) \in \dot{T}^*U} |\omega_{\alpha,\beta}(y, \eta)|.$$

For every compact set $K \subset \dot{T}^*U$, there exists a constant C_K such that for all test form ω supported in K

$$\int_{\mathcal{H}_\Lambda} \left| \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) \right| d\mu_\Lambda \leq C_K \|\omega\|$$

and

$$\begin{aligned} & \int_{\mathcal{H}_\Lambda} \left(\int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) \right) d\mu_\Lambda(f) \\ &= C_n \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \int_{\dot{T}^*U} dy^1 \wedge dy^2 \dots \wedge dy^n \wedge \omega(y, dy, \eta, d\eta) + \mathcal{O}(\Lambda^{n-1}), \end{aligned}$$

with $C_n = 0$ if $n \equiv 0 \pmod{2}$ and

$$C_n := \frac{2(-1)^{\frac{n+1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})}$$

if $n \equiv 1 \pmod{2}$.

Remark 4.2. It follows from the first statement of Theorem 4.1 that for every test form ω , the map

$$f \mapsto \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta)$$

belongs to $L^1(\mathcal{H}_\Lambda, d\mu_\Lambda)$. For any integral current T , let $\mathbb{M}(T)$ denote the *mass* of the current T [32]. Our result is in fact slightly stronger than what we claimed in Theorem 2.2

as it also means that, for any smooth compactly supported cut-off function $\varphi \in \mathcal{D}^0(T^*M)$,

$$\int_{\mathcal{H}_\Lambda} \mathbb{M}(\varphi[N^*(\{f=0\})]) < +\infty.$$

In other words, the mass of the cut-off conormal cycle $\varphi[N^*(\{f=0\})]$ is a L^1 function of $f \in \mathcal{H}_\Lambda$.

The purpose of this section is to prove this Theorem. We emphasize that *there are two parts in this statement*. On the one hand, we have to show the integrability property and on the other hand, we have to compute the precise value of the expectation. The first part of the statement requires a delicate analysis which is carried out in the first part of this section. After that step, we can combine this first part to some combinatorial arguments in order to obtain the second part of the Theorem.

More precisely, the proof is organized as follows. First, we recall in paragraph 4.1 some results on the kernel of the spectral projector associated to the space \mathcal{H}_Λ . Then, in paragraph 4.2, we write a formal asymptotic expansion in powers of Λ^{-1} of

$$\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta).$$

Paragraph 4.4 is the more delicate part where we prove the first part of Theorem 4.1 concerning the integrability of the conormal cycle as a function of $f \in \mathcal{H}_\Lambda$ with respect to the measure $d\mu_\Lambda$. After that, in paragraph 4.5, we compute explicitly the expression of the leading term and in 4.6, we compute the combinatorial constant appearing in the leading term. Finally, in paragraph 4.8, we conclude the proof of Theorem 4.1.

4.1. Spectral asymptotics. Recall that $(e_j)_{1 \leq j \leq N(\Lambda)}$ is an orthonormal basis of \mathcal{H}_Λ made of eigenfunctions of Δ . We define the normalized projection kernel:

$$C_\Lambda(y, z) := \frac{1}{N(\Lambda)} \sum_{j=1}^{N(\Lambda)} e_j(y)e_j(z),$$

where $N(\Lambda) = \dim \mathcal{H}_\Lambda$. The reason why we can perform some computations with respect to the Gaussian measure μ_Λ is that we have very precise asymptotic informations on the kernel $C_\Lambda(y, z)$, at least on the diagonal. Precisely, Bin proved the following result [8] building on earlier arguments of Hörmander [38]:

Theorem 4.3. *Let (U, ϕ) be a sufficiently small geodesic normal coordinate chart. For every multiindices α and β in \mathbb{Z}_+^n , the following estimates hold uniformly for y in $\phi(U)$, as $\Lambda \rightarrow +\infty$,*

$$\partial_y^\alpha \partial_z^\beta C_\Lambda(y, z)|_{y=z} = C_{n,\alpha,\beta} \Lambda^{|\alpha+\beta|} + \mathcal{O}(\Lambda^{|\alpha+\beta|-1}),$$

where, for $\alpha = \beta = 0$,

$$C_{n,0,0} := \frac{1}{\text{Vol}_g(M)},$$

for $\alpha \equiv \beta \pmod{2}$ and $(\alpha, \beta) \neq (0, 0)$,

$$C_{n,\alpha,\beta} := (-1)^{\frac{|\alpha|+|\beta|}{2}} \frac{\prod_{j=1}^n (\alpha_j + \beta_j - 1)!!}{\text{Vol}_g(M) (|\alpha + \beta| + n) \dots (n + 2)},$$

and

$$C_{n,\alpha,\beta} := 0 \quad \text{otherwise.}$$

Remark 4.4. We used the following conventions:

$$(-1)!! := 1 \text{ and } (2m - 1)!! := (2m - 1)(2m - 3) \dots 3 \times 1.$$

Remark 4.5. By geodesic normal coordinate chart, we mean the following. Consider the exponential map Exp induced by the Riemannian metric on M . There exists a neighborhood \mathcal{U} of $M \times \{0\}$ in TM such that Exp induces a smooth diffeomorphism $(x, v) \mapsto (x, \exp_x(v))$ from \mathcal{U} onto a small neighborhood of the diagonal in $M \times M$. Then, we fix a point x_0 in M , and we consider the preimage of $T_{x_0}M$ under this diffeomorphism. This defines a local chart around x_0 into $T_{x_0}M$ which can be identified with \mathbb{R}^n via a measure preserving linear isomorphism.

Remark 4.6. It follows from the previous theorem that one has the following asymptotic formulas that will be useful in the following:

- for derivatives of order 0, one has

$$C_\Lambda(y, z)|_{y=z} = \frac{1}{\text{Vol}_g(M)} + \mathcal{O}(\Lambda^{-1}),$$

- for every $1 \leq j \leq n$, one has

$$\frac{1}{\Lambda^2} \partial_{y^j}^2 C_\Lambda(y, z)|_{y=z} = -\frac{1}{(n+2) \text{Vol}_g(M)} + \mathcal{O}(\Lambda^{-1}),$$

- for every $1 \leq j \leq n$, one has

$$\frac{1}{\Lambda^2} \partial_{y^j} \partial_{z^j} C_\Lambda(y, z)|_{y=z} = \frac{1}{(n+2) \text{Vol}_g(M)} + \mathcal{O}(\Lambda^{-1}),$$

- for every $1 \leq j \neq k \leq n$, one has

$$\frac{1}{\Lambda^4} \partial_{y^j}^2 \partial_{z^k}^2 C_\Lambda(y, z)|_{y=z} = \frac{1}{\Lambda^4} \partial_{y^j y^k}^2 \partial_{z^j z^k}^2 C_\Lambda(y, z)|_{y=z} = \frac{1}{(n+2)(n+4) \text{Vol}_g(M)} + \mathcal{O}(\Lambda^{-1}),$$

- for every $1 \leq j \leq n$, one has

$$\frac{1}{\Lambda^4} \partial_{y^j}^2 \partial_{z^j}^2 C_\Lambda(y, z)|_{y=z} = \frac{3}{(n+2)(n+4) \text{Vol}_g(M)} + \mathcal{O}(\Lambda^{-1}),$$

- for every $\alpha \neq \beta \pmod{2}$, one has

$$\frac{1}{\Lambda^{|\alpha+\beta|}} \partial_y^\alpha \partial_z^\beta C_\Lambda(y, z)|_{y=z} = \mathcal{O}(\Lambda^{-1}).$$

4.2. Preliminary simplification. Before getting into the details of the proof of Theorem 4.1, we start by making a few reduction that will make the calculation slightly simpler:

Lemma 4.7. *Let*

$$\omega := \sum_{|\alpha|+|\beta|=n} \omega_{\alpha,\beta}(y, \eta) dy^\alpha \wedge d\eta_\beta$$

be an element in $\mathcal{D}_n(\mathring{T}^*U)$. With the above conventions, one has, for every Λ and for every f in Ω_Λ ,

$$\begin{aligned} & \frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) \\ &= - \sum_{k=0}^{n-1} \Lambda^{-k} \langle G_\Lambda(f)^*(\delta_0^{n+1}), t^{n-3-k} P_k((\eta_j)_j, (\partial_{y^j}^2 f / \Lambda^2)_{j,k}, (\omega_{\alpha,\beta})_{\alpha,\beta}) \rangle_{\mathring{T}^*U \times \mathbb{R}^*}, \end{aligned}$$

where

- for every $0 \leq k \leq n-1$, $P_k(R, S, T)$ is a polynomial which does not depend on Λ , t and f ,
- for every $0 \leq k \leq n-1$, $P_k(R, S, T)$ is homogeneous of degree 2 in the variables R , homogeneous of degree $n-1-k$ in the variables S , homogeneous of degree 1 in the T variables,
- in the case $k=0$, one has

$$P_0((R_j)_j, (S_{j,k})_{j,k}, (T_{\alpha,\beta})_{\alpha,\beta}) = \left(\sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) R_p R_{\sigma(p)} \prod_{j \neq p} S_{j,\sigma(j)} \right) T_{(0,\dots,0),(1,\dots,1)}.$$

At first sight, this lemma seems rather technical; yet, writing the integrals this way will simplify the presentation afterwards. This lemma gives at least a formal expansion in powers of Λ^{-1} , and we will verify in the following paragraphs that each term in the sum is in fact integrable and has uniformly bounded L^1 norm in \mathcal{H}_Λ .

Proof. Write first that

$$\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) = \sum_{p=1}^n (-1)^p \frac{1}{\Lambda^n} \int_{\mathbb{R}_+} \tilde{I}_\Lambda^{(p)}(f, t) dt,$$

where

$$\tilde{I}_\Lambda^{(p)}(f, t) = \left\langle G_\Lambda(f)^*(\delta_0^{n+1}), \frac{\partial_{y_p} f}{\Lambda} df \wedge \bigwedge_{j \neq p} d \left(\frac{t}{\Lambda} \partial_{y_j} f - \eta_j \right) \wedge \omega(y, \eta, dy, d\eta) \right\rangle_{\mathring{T}^*U}.$$

Expanding the df term in the wedge product, one can also write

$$\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) = \sum_{p,q=1}^n (-1)^{n+p-1} \int_{\mathbb{R}_+} \tilde{I}_\Lambda^{(p,q)}(f, t) dt,$$

where

$$\tilde{I}_\Lambda^{(p,q)}(f, t) := \left\langle G_\Lambda(f)^*(\delta_0^{n+1}), \frac{\partial_{y_p} f}{\Lambda} \frac{\partial_{y_q} f}{\Lambda} \bigwedge_{j \neq p} d \left(\frac{t}{\Lambda^2} \partial_{y_j} f - \frac{\eta_j}{\Lambda} \right) \wedge dy_q \wedge \omega(y, \eta, dy, d\eta) \right\rangle_{\hat{T}^*U}.$$

Thanks to the definition of $G_\Lambda(f)^*(\delta_0^{n+1})$, this can be rewritten as

$$\tilde{I}_\Lambda^{(p,q)}(f, t) = \frac{1}{t^2} \left\langle G_\Lambda(f)^*(\delta_0^{n+1}), \eta_p \eta_q \bigwedge_{j \neq p} d \left(\frac{t}{\Lambda^2} \partial_{y_j} f - \frac{\eta_j}{\Lambda} \right) \wedge dy_q \wedge \omega(y, \eta, dy, d\eta) \right\rangle_{\hat{T}^*U}.$$

Write now

$$\omega := \sum_{|\alpha|+|\beta|=n} \omega_{\alpha,\beta}(y, \eta) dy^\alpha \wedge d\eta_\beta,$$

where $\omega_{\alpha,\beta}$ is compactly supported in \hat{T}^*U . By expanding the wedge product involved in the integrals, we can rearrange the sum in the following way:

$$\begin{aligned} & \frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) \\ &= - \sum_{k=0}^{n-1} \Lambda^{-k} \langle G_\Lambda(f)^*(\delta_0^{n+1}), t^{n-3-k} P_k((\eta_j)_j, (\partial_{y^j y^k}^2 f / \Lambda^2)_{j,k}, (\omega_{\alpha,\beta})_{\alpha,\beta}) \rangle_{\hat{T}^*U \times \mathbb{R}^*}, \end{aligned}$$

where

- for every $0 \leq k \leq n-1$, $P_k(R, S, T)$ is a polynomial which does not depend on Λ , t and f ,
- for every $0 \leq k \leq n-1$, $P_k(R, S, T)$ is homogeneous of degree 2 in the variables R , homogeneous of degree $n-1-k$ in the variables S , and homogeneous of degree 1 in the T variables.
- in the case $k=0$, one has

$$P_0((R_j)_j, (S_{j,k})_{j,k}, (T_{\alpha,\beta})_{\alpha,\beta}) = \left(\sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) R_p R_{\sigma(p)} \prod_{j \neq p} S_{j,\sigma(j)} \right) T_{(0,\dots,0),(1,\dots,1)}.$$

This concludes the proof of the lemma. \square

4.3. Notations and conventions. In the following, we fix the following convention, for every $0 \leq k \leq n-1$,

$$I_\Lambda^{(k)}(f, \omega, U) = \langle G_\Lambda(f)^*(\delta_0^{n+1}), t^{n-3-k} P_k((\eta_j)_j, (\partial_{y^j y^k}^2 f / \Lambda^2)_{j,k}, (\omega_{\alpha,\beta})_{\alpha,\beta}) \rangle_{\hat{T}^*U \times \mathbb{R}^*}.$$

In particular, one has

$$(29) \quad \frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) = - \sum_{k=0}^{n-1} \Lambda^{-k} I_\Lambda^{(k)}(f, \omega, U).$$

4.4. **Integrability.** For any element $\omega := \sum_{|\alpha|+|\beta|=n} \omega_{\alpha,\beta}(y, \eta) dy^\alpha \wedge d\eta_\beta$ in $\mathcal{D}_n(\mathring{T}^*U)$, let

$$\|\omega\| = \sup_{\alpha,\beta,(y,\eta) \in \mathring{T}^*U} |\omega_{\alpha,\beta}(y, \eta)|.$$

Using the conventions of paragraph 4.3, we will now prove the first part of Theorem 4.1 which follows immediately from the following proposition

Proposition 4.8. *Let ω be an element in $\mathcal{D}_n(\mathring{T}^*U)$. Then, for every $0 \leq k \leq n-1$, one has*

$$f \mapsto I_\Lambda^{(k)}(f, \omega, U)$$

belongs to $L^1(\mathcal{H}_\Lambda, d\mu_\Lambda)$ and for every compact set K , there exists a constant $C_{K,k}$ such that for any test form ω in $\mathcal{D}_n(\mathring{T}^*U)$ supported in K ,

$$\int_{\Omega_\Lambda} |I_\Lambda^{(k)}(f, \omega, U)| d\mu_\Lambda(f) \leq C_{K,k} \|\omega\|.$$

Showing this proposition is our more delicate task from the analytical point of view as we have to justify carefully several inversions and convergences of integrals.

Remark 4.9. As was already mentioned, a direct consequence of this lemma is that, for every ω in $\mathcal{D}_n(\mathring{T}^*U)$,

$$f \mapsto \frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta)$$

belongs to $L^1(\mathcal{H}_\Lambda, d\mu_\Lambda)$ which is exactly the content of the first part of Theorem 4.1. In fact, we have something slightly more precise in the sense that

$$\int_{\Omega_\Lambda} \left(\int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) \right) d\mu_\Lambda(f) = -\Lambda^n \int_{\Omega_\Lambda} I_\Lambda^{(0)}(f, \omega, U) d\mu_\Lambda(f) + \mathcal{O}(\Lambda^{n-1}).$$

Recall from paragraph 4.3 that

$$I_\Lambda^{(0)}(f, \omega, U) := \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) \left\langle G_\Lambda(f)^*(\delta_0^{n+1}), t^{n-3} \eta_p \eta_{\sigma(p)} \left(\prod_{j \neq p} \frac{\partial_{y^j, y^{\sigma(j)}}^2 f}{\Lambda^2} \right) \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) \right\rangle_{\mathring{T}^*U \times \mathbb{R}^*}.$$

In particular, the second part of Theorem 4.1 is reduced to the computation of the expectation of $I_\Lambda^{(0)}(f, \omega, U)$ with respect to the Gaussian measure $d\mu_\Lambda(f)$. This computation will be performed in paragraphs 4.5 and 4.6.

We will now prove proposition 4.8 in several steps. We will start with general considerations on pull-back of Dirac distributions that we will apply in a second part to derive the proof of the proposition.

4.4.1. *Pullback of distributions and Fubini results.* We fix $0 \leq k \leq n-1$ and f in $\Omega_\Lambda = \mathcal{H}_\Lambda \setminus D_\Lambda$. Recall that D_Λ is a zero measure subset of \mathcal{H}_Λ and that it is open. In the definition of $I_\Lambda^{(k)}(f, \omega, U)$, we have to consider the pulled-back distribution $G_\Lambda(f)^* \delta_0^{n+1}$. Given any test function $\Psi(f, y, \eta, t)$ in $\mathcal{D}(\Omega_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*)$, our goal in this paragraph is to make sense of the following Fubini equality :

$$\begin{aligned} \left\langle 1, \langle G_\Lambda(f)^*(\delta_0^{n+1}), \Psi \rangle_{\mathring{T}^*U \times \mathbb{R}^*} \right\rangle_{\Omega_\Lambda} &= \langle G_\Lambda^*(\delta_0^{n+1}), \Psi \rangle_{\Omega_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*} \\ &= \left\langle 1, \langle G_\Lambda(y, \eta, t)^*(\delta_0^{n+1}), \Psi \rangle_{\Omega_\Lambda} \right\rangle_{\mathring{T}^*U \times \mathbb{R}^*}. \end{aligned}$$

The fact that we can “intertwine the order of integration” plays a central role in the proof of Proposition 4.8.

As was already explained in section 3, the fact that the distribution $G_\Lambda(f)^* \delta_0^{n+1}$ is well defined for f in Ω_Λ follows from the following theorem of Hörmander – Th. 8.2.4 in [39]:

Theorem 4.10. *Let $\Omega_1 \subset \mathbb{R}^{d_1}$ and $\Omega_2 \subset \mathbb{R}^{d_2}$ be two open subsets. For any smooth map $G : \Omega_1 \rightarrow \Omega_2$, the normal of G is defined as*

$$N_G := \{(G(x), \tau) \in \Omega_2 \times \mathbb{R}^{d_2^*} : \tau \circ d_x G = 0\},$$

*the pull-back operation extends uniquely to the distributions $u \in \mathcal{D}'(\Omega_2)$ whose wavefront $WF(u)$ does not intersect the normal N_G of G . Moreover, the wavefront set of G^*u is contained in the set*

$$G^*WF(u) := \{(x, \tau \circ d_x G) : (G(x), \tau) \in WF(u)\}.$$

Remark 4.11. For the definition of the wavefront set, we refer to [39, 14]. Recall that, in the case of δ_0^{n+1} , one has

$$WF(\delta_0^{n+1}) := \{(0, \tau) : \tau \neq 0\} \subset T^*\mathbb{R}^{n+1}.$$

In the following, we will apply this Theorem in three distinct situations. Recall that, for every f in \mathcal{H}_Λ , we defined the following map on $\mathcal{H}_\Lambda \times \phi(U) \times \mathbb{R}^n \times \mathbb{R}^*$:

$$G_\Lambda : (f, y, \eta, t) \in \mathcal{H}_\Lambda \times \phi(U) \times \mathbb{R}^n \times \mathbb{R}^* \longmapsto \left(f(y); \frac{t}{\Lambda} \partial_{y_1} f - \eta_1, \dots, \frac{t}{\Lambda} \partial_{y_n} f - \eta_n \right) \in \mathbb{R} \times \mathbb{R}^n,$$

and our idea is to think of $\mathcal{H}_\Lambda \times \phi(U) \times \mathbb{R}^n \times \mathbb{R}^*$ as the cartesian product of \mathcal{H}_Λ with $\phi(U) \times \mathbb{R}^n \times \mathbb{R}^*$ and to think of the global distribution $G_\Lambda^* \delta_0^{n+1}$ also as a distribution on \mathcal{H}_Λ depending smoothly on the parameters in $\phi(U) \times \mathbb{R}^n \times \mathbb{R}^*$ and conversely as a distribution on $\phi(U) \times \mathbb{R}^n \times \mathbb{R}^*$ depending smoothly on the parameters in \mathcal{H}_Λ . To be more precise, we will use Hörmander’s Theorem in the following three cases :

- (1) We fix f in Ω_Λ , and we consider the partial map $G_\Lambda(f) : \mathring{T}^*U \times \mathbb{R}^* \rightarrow \mathbb{R}^{n+1}$. As $d_y f \neq 0$ on $f^{-1}(0)$, one can verify that

$$N_{G_\Lambda(f)} \cap WF(\delta_0^{n+1}) = \emptyset.$$

It means that if we freeze f , the pull-back $G_\Lambda(f)^* \delta_0^{n+1}$ is a well-defined distribution on $\mathring{T}^*U \times \mathbb{R}^*$.

- (2) We consider the map $G_\Lambda : \mathcal{H}_\Lambda \times \mathring{T}^*U \times \mathbb{R}_+^* \rightarrow \mathbb{R}^{n+1}$. In order to apply Hörmander's result, we would need to verify

$$N_{G_\Lambda} \cap WF(\delta_0^{n+1}) = \emptyset,$$

which is slightly less obvious.

- (3) We fix (y, η, t) in $\mathring{T}^*U \times \mathbb{R}^*$, and we consider the partial map $G_\Lambda(y, \eta, t) : \mathcal{H}_\Lambda \rightarrow \mathbb{R}^{n+1}$. The situation in this case is also slightly different from the first case.

The second and the third cases are in some sense related to the notion of ampleness appearing for instance in [30, 45], and they are contained in the following lemma:

Lemma 4.12. *There exists $\Lambda_0 > 0$ such that, for every $\Lambda \geq \Lambda_0$,*

$$N_{G_\Lambda} \cap WF(\delta_0^{n+1}) = \emptyset,$$

and, for every (y, η, t) in $\mathring{T}^*U \times \mathbb{R}^*$, one has

$$N_{G_\Lambda(y, \eta, t)} \cap WF(\delta_0^{n+1}) = \emptyset.$$

In particular, $G_\Lambda^*(\delta_0^{n+1})$ is well defined as a distribution on $\mathcal{H}_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*$ and $G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})$ as a distribution on \mathcal{H}_Λ .

Remark 4.13. We note that in the second and in the third cases, the distributions are defined on the whole space \mathcal{H}_Λ , and not only on Ω_Λ . In particular, we can consider their restriction to the open subset Ω_Λ .

Before continuing our discussion on the properties of these pulled-back distributions, we start by giving the proof of the lemma.

Proof. Suppose that $(0, \tau)$ belongs to the normal set $N_{G_\Lambda(y, \eta, t)}$ (the other case works similarly). Then, for every f in \mathcal{H}_Λ , one has

$$\tau_0 f(y) + \sum_{j=1}^{N(\Lambda)} \tau_j \frac{t \partial_{y^j} f}{\Lambda} = 0.$$

Consider now a family of functions $f_0^\Lambda, f_1^\Lambda, \dots, f_n^\Lambda$ in \mathcal{H}_Λ , we write

$$\det \left(\left(f_l^\Lambda(y), \frac{t \partial_{y^1} f_l^\Lambda}{\Lambda}, \dots, \frac{t \partial_{y^n} f_l^\Lambda}{\Lambda} \right)_{0 \leq l \leq n} \right) = \frac{t^n}{\Lambda^n} \det \left((f_l^\Lambda(y), \partial_{y^1} f_l^\Lambda, \dots, \partial_{y^n} f_l^\Lambda)_{0 \leq l \leq n} \right).$$

If we are able to find a family $(f_l^\Lambda)_{0 \leq l \leq n}$ in \mathcal{H}_Λ such that the right-hand side of this equality does not vanish, we will deduce that $\tau = 0$. For that purpose, we note that we can find⁴ a family of smooth functions f_0, f_1, \dots, f_n on M such that the right hand side does not vanish for every y in U . Unfortunately, these functions do not a priori belong to \mathcal{H}_Λ but we can solve this issue since by Lemma A.1 proved in appendix, finite linear combinations of eigenfunctions are everywhere dense in $C^\infty(M)$. Said concretely, to every $0 \leq l \leq n$ corresponds a sequence $(f_l^\Lambda)_\Lambda$ in \mathcal{H}_Λ which converges to f_l in every \mathcal{C}^k norm (this

⁴We may have to pick a slightly smaller neighborhood U .

is just an approximation property like the Stone Weierstrass Theorem but on Riemannian manifolds). From this, we deduce the statement of the lemma for Λ large enough. \square

We refer the reader to the appendix where a similar method yields an unusual proof of the Whitney embedding Theorem.

As a direct consequence of the pull-back theorem of Hörmander, we can also describe the wavefront set of $G_\Lambda^*(\delta_0^{n+1})$ as follows:

Corollary 4.14. *For $\Lambda > 0$ large enough, the wavefront set of $G_\Lambda^*(\delta_0^{n+1})$ is included in the following subset of $T^*(\mathcal{H}_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*)$:*

$$\begin{aligned} WF(G_\Lambda^*(\delta_0^{n+1})) &\subset \{(y^j, \eta_j, t, f; \widehat{y}_j, \widehat{\eta}^j, \widehat{t}, \widehat{f}) \text{ such that } G_\Lambda(f, y, \eta, t) = (0, \dots, 0) \\ &\quad \widehat{y}_j = \tau_0 \partial_{y^j} f + \tau_1 \frac{t}{\Lambda} \partial_{y^j y^1}^2 f + \dots + \tau_n \frac{t}{\Lambda} \partial_{y^j y^n}^2 f, \widehat{\eta}_j = \tau_j, \\ &\quad \widehat{t} = \frac{1}{\Lambda} \sum_{j=1}^n \tau_j \partial_{y^j} f, \forall 1 \leq i \leq N(\Lambda), \widehat{f}_i = \tau_0 e_i + \sum_{j=1}^n \frac{t \tau_j}{\Lambda} \partial_{x^j} e_i \\ &\quad \text{for } (\tau_0, \tau_1, \dots, \tau_n) \neq (0, \dots, 0)\}. \end{aligned}$$

Remark 4.15. This corollary has the following consequence. The set $WF(G_\Lambda^*(\delta_0^{n+1}))$ does not contain any element of the form $(y^j, \eta_j, t, f; 0, 0, 0, \widehat{f})$, with $\widehat{f} \neq 0$ or $(y^j, \eta_j, t, f; \widehat{y}_j, \widehat{\eta}^j, \widehat{t}, 0)$ with $(\widehat{y}_j, \widehat{\eta}^j, \widehat{t}) \neq 0$. The first observation is more or less immediate if one uses Theorem 4.3 in the case $\alpha = \beta = 0$. The second one follows from the proof of lemma 4.12.

The relevance of such a remark is due to the following general result which can be found in [18, Proposition 1.3 p. 502] :

Proposition 4.16. *Let u be a distribution in $\mathcal{D}'(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})$. We denote by $(x, y; \xi, \eta) \in T^*(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})$ the coordinates in cotangent space. If $WF(u)$ contains no elements of the form $\{(x, y; \xi, 0) \text{ such that } \xi \neq 0\}$, then $u(x, y)$ is smooth in x with value distribution in y in the sense that for all test function $\varphi_2(y)$, $x \mapsto \langle u(x, \cdot), \varphi_2(\cdot) \rangle_{\mathbb{R}^{n_2}}$ is smooth in x .*

Thus, one finds that there exists $f \in \mathcal{H}_\Lambda \mapsto T_\Lambda(f) \in \mathcal{D}'(\mathring{T}^*U \times \mathbb{R}^*)$ such that, for every $\psi_1 \in \mathcal{D}(\mathcal{H}_\Lambda)$ and for every $\psi_2 \in \mathcal{D}(\mathring{T}^*U \times \mathbb{R}^*)$,

$$f \mapsto \langle T_\Lambda(f), \psi_2 \rangle_{\mathring{T}^*U \times \mathbb{R}^*} \in \mathcal{C}^\infty(\mathcal{H}_\Lambda),$$

and

$$(y, \eta, t) \mapsto \langle G_\Lambda(y, \eta, t)^* \delta_0^{n+1}, \psi_1 \rangle_{\Omega_\Lambda} \in \mathcal{C}^\infty(\mathring{T}^*U \times \mathbb{R}^*).$$

Moreover, one has the “following distributional version of the Fubini Theorem” :

$$\begin{aligned} \forall (\psi_1, \psi_2) \in \mathcal{D}(\mathcal{H}_\Lambda) \times \mathcal{D}(\mathring{T}^*U \times \mathbb{R}^*) \\ \langle \psi_1, \langle T_\Lambda(f), \psi_2 \rangle_{\mathring{T}^*U \times \mathbb{R}^*} \rangle_{\mathcal{H}_\Lambda} &= \langle G_\Lambda^*(\delta_0^{n+1}), \psi_1 \boxtimes \psi_2 \rangle_{\mathcal{H}_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*} \\ &= \left\langle \psi_2, \langle G_\Lambda(y, \eta, t)^* (\delta_0^{n+1}), \psi_1 \rangle_{\mathcal{H}_\Lambda} \right\rangle_{\mathring{T}^*U \times \mathbb{R}^*}. \end{aligned}$$

Remark 4.17. Note that $T_\Lambda(f)$ is well defined on the whole space \mathcal{H}_Λ while $G_\Lambda(f)^*(\delta_0^{n+1})$ was only defined on Ω_Λ by Theorem 4.10. For a fixed f in Ω_Λ , one has in fact $T_\Lambda(f) = G_\Lambda(f)^*(\delta_0^{n+1})$.

By density of the functions of the form $\psi_1(f) \boxtimes \psi_2(y, \eta, t)$ (where \boxtimes is the exterior tensor product) in $\mathcal{D}(\Omega_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*)$ – see Ch. IV in [53], given any test function $\Psi(f, y, \eta, t)$ in $\mathcal{D}(\Omega_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*)$, one has the following Fubini equality :

$$(30) \quad \begin{aligned} \langle 1, \langle T_\Lambda, \Psi \rangle_{\mathring{T}^*U \times \mathbb{R}^*} \rangle_{\mathcal{H}_\Lambda} &= \langle G_\Lambda^*(\delta_0^{n+1}), \Psi \rangle_{\mathcal{H}_\Lambda \times \mathring{T}^*U \times \mathbb{R}^*} \\ &= \left\langle 1, \langle G_\Lambda(y, \eta, t)^*(\delta_0^{n+1}), \Psi \rangle_{\mathcal{H}_\Lambda} \right\rangle_{\mathring{T}^*U \times \mathbb{R}^*}. \end{aligned}$$

Remark 4.18. We emphasize that all the distributions considered so far are constructed from the Dirac distribution δ_0^{n+1} and that they are positive distributions. In particular, according to [53] (Ch. I), *they can all be identified with positive Radon measures*. Therefore, it makes sense to test them against nonnegative measurable functions which are not necessarily integrable (or integrable function which are not necessarily smooth). This identification will be used several times in our proof of proposition 4.8 below.

Remark 4.19. We now make a final useful comment. We fix $(\psi_1, \psi_2) \in \mathcal{D}(\mathcal{H}_\Lambda) \times \mathcal{D}(\mathring{T}^*U \times \mathbb{R}^*)$. According to (30), one has

$$\begin{aligned} &\int_{\mathring{T}^*U \times \mathbb{R}^*} \psi_2(y, \eta, t) \left(\int_{\mathcal{H}_\Lambda} \psi_1(f) G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df) \right) d^n y d^n \eta dt \\ &= \int_{\mathcal{H}_\Lambda} \psi_1(f) \left(\int_{\mathring{T}^*U \times \mathbb{R}^*} \psi_2(y, \eta, t) T_\Lambda(d^n y, d^n \eta, dt) \right) df. \end{aligned}$$

From the above discussion, one knows that

$$f \mapsto \psi_1(f) \left(\int_{\mathring{T}^*U \times \mathbb{R}^*} \psi_2(y, \eta, t) T_\Lambda(d^n y, d^n \eta, dt) \right)$$

is a smooth compactly supported function. As $\Omega_\Lambda = \mathcal{H}_\Lambda \setminus D_\Lambda$ and D_Λ has zero Lebesgue measure, one has then

$$\begin{aligned} &\int_{\mathring{T}^*U \times \mathbb{R}^*} \psi_2(y, \eta, t) \left(\int_{\mathcal{H}_\Lambda} \psi_1(f) G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df) \right) d^n y d^n \eta dt \\ &= \int_{\Omega_\Lambda} \psi_1(f) \left(\int_{\mathring{T}^*U \times \mathbb{R}^*} \psi_2(y, \eta, t) G_\Lambda(f)^*(\delta_0^{n+1})(d^n y, d^n \eta, dt) \right) df. \\ &= \int_{\mathring{T}^*U \times \mathbb{R}^*} \psi_2(y, \eta, t) \left(\int_{\Omega_\Lambda} \psi_1(f) G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df) \right) d^n y d^n \eta dt, \end{aligned}$$

where the second equality follow from the fact that

$$G_\Lambda(f)^*(\delta_0^{n+1})(d^n y, d^n \eta, dt) df \text{ and } G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df) d^n y d^n \eta dt$$

define the same Radon measure on the space $\Omega_\Lambda \times \dot{T}^*U \times \mathbb{R}^*$. Note also that this equality holds for any smooth compactly supported function ψ_2 . In particular, one finds that, for a.e. (y, η, t) in $\dot{T}^*U \times \mathbb{R}^*$,

$$(31) \quad \int_{\mathcal{H}_\Lambda} \psi_1(f) G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df) = \int_{\Omega_\Lambda} \psi_1(f) G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df).$$

4.4.2. *Proof of proposition 4.8 – First step.* We now give the proof of Proposition 4.8. As a first step, we will give a crude upper bound on $|I_\Lambda^{(k)}(f, \omega, U)|$. Let $\tilde{\omega}$ be a smooth function compactly supported on $K \subset \dot{T}^*U$ such that, for every multiindices (α, β) satisfying $|\alpha| + |\beta| = n$, and for every $1 \leq p, q \leq n$, one has

$$\forall (y, \eta) \in \dot{T}^*U, |\omega_{\alpha, \beta}(y, \eta) \eta_p \eta_q| \leq \tilde{\omega}(y, \eta).$$

Moreover, we choose $\tilde{\omega}$ so that $\sup_{(y, \eta) \in \dot{T}^*U} \tilde{\omega}(y, \eta) \leq C \|\omega\|$ where $C > 0$ is a constant depending only on the support of ω , and where $\|\cdot\|$ is the norm on smooth forms defined at the beginning of paragraph 4.4. As was already mentioned, $G_\Lambda(f)^*(\delta_0^{n+1})$ is a positive distribution on $\dot{T}^*U \times \mathbb{R}^*$ hence a positive Radon measure $G_\Lambda(f)^*(\delta_0^{n+1})(d^n y, d^n \eta, dt)$ on $\dot{T}^*U \times \mathbb{R}^*$. In particular, one has

$$(32) \quad |I_\Lambda^{(k)}(f, \omega, U)| \leq \int_{\dot{T}^*U \times \mathbb{R}^*} \tilde{\omega}(y, \eta) |t|^{n-3-k} Q_k \left(\left(|\partial_{y^j y^l}^2 f / \Lambda^2| \right)_{j, l} \right) G_\Lambda(f)^*(\delta_0^{n+1})(d^n y, d^n \eta, dt),$$

for some homogeneous polynomial Q_k of degree $n-1-k$ with nonnegative coefficients and which is independent of $\omega, U, (y, \eta, t)$ and Λ . Then, according to (32), the integrability of $I_\Lambda^{(k)}(f, \omega, U)$ with respect to $d\mu_\Lambda$ will follow from the integrability of the map

$$(y, \eta, t, f) \in \dot{T}^*U \times \mathbb{R}^* \times \mathcal{H}_\Lambda \longmapsto \tilde{\omega}(y, \eta) |t|^{n-3-k} Q_k \left(\left(|\partial_{y^j y^l}^2 f / \Lambda^2| \right)_{j, l} \right) \in \mathbb{R}_+$$

with respect to the Radon measure $\mu_\Lambda(f) G_\Lambda^*(\delta_0^{n+1})(df, d^n y, d^n \eta, dt)$ with

$$\mu_\Lambda(f) := e^{-\frac{N(\Lambda) \|f\|^2}{2}} \left(\frac{N(\Lambda)}{2\pi} \right)^{\frac{N(\Lambda)}{2}}.$$

Thus, according to (32) and in order to prove the lemma, it remains to show that

$$\int_{\dot{T}^*U \times \mathbb{R}^* \times \mathcal{H}_\Lambda} \tilde{\omega}(y, \eta) |t|^{n-3-k} Q_k \left(\left(|\partial_{y^j y^l}^2 f / \Lambda^2| \right)_{j, l} \right) \mu_\Lambda(f) G_\Lambda^*(\delta_0^{n+1})(df, d^n y, d^n \eta, dt)$$

is uniformly bounded with respect to Λ . Let $\mathbf{j} := (j_1, j_2, \dots, j_{2(n-k)-3}, j_{2(n-k-1)})$ be an element in $\{1, \dots, n\}^{2(n-1-k)}$. As Q_k is an homogeneous polynomial of degree $n-1-k$, it is sufficient to prove that

$$(33) \quad \tilde{I}_\Lambda^{(k)}(\mathbf{j}, \omega, U) := \int_{\dot{T}^*U \times \mathbb{R}^* \times \mathcal{H}_\Lambda} \tilde{\omega}(y, \eta) |t|^{n-3-k} \prod_{p=1}^{n-1-k} \left| \frac{\partial_{y^{j_{2p-1}} y^{j_{2p}}}^2 f}{\Lambda^2} \right| \mu_\Lambda(f) G_\Lambda^*(\delta_0^{n+1})(df, d^n y, d^n \eta, dt)$$

is uniformly bounded with respect to Λ .

4.4.3. *Proof of proposition 4.8 – Step 2: Intertwining integrals.* We can now use (30), and integrate first with respect to the f variable. For that purpose, we define the following nonnegative Radon measure on Ω_Λ :

$$\nu_{\Lambda,1}^{(y,\eta,t)}(df) = \mu_\Lambda(f)G_\Lambda(y,\eta,t)^*(\delta_0^{n+1})(df),$$

and

$$\nu_{\Lambda,2}^{(y,\eta,t)}(df) = \prod_{p=1}^{n-1-k} \left| \frac{\partial^2 y^{j_{2p-1}} y^{j_{2p}} f(y)}{\Lambda^2} \right|^2 \mu_\Lambda(f)G_\Lambda(y,\eta,t)^*(\delta_0^{n+1})(df).$$

Then, combining (33) and the Fubini result (30) to Hölder inequality, one finds that :

$$\begin{aligned} \tilde{I}_\Lambda^{(k)}(\mathbf{j}, \omega, U) &= \int_{\dot{T}^*U \times \mathbb{R}^* \times \Omega_\Lambda} \tilde{\omega}(y, \eta) |t|^{n-3-k} \prod_{p=1}^{n-1-k} \left| \frac{\partial^2 y^{j_{2p-1}} y^{j_{2p}} f}{\Lambda^2} \right| \mu_\Lambda(f) G_\Lambda^*(\delta_0^{n+1})(df, d^n y, d^n \eta, dt) \\ &= \int_{\dot{T}^*U \times \mathbb{R}^*} \tilde{\omega}(y, \eta) |t|^{n-3-k} \left(\int_{\Omega_\Lambda} \prod_{p=1}^{n-1-k} \left| \frac{\partial^2 y^{j_{2p-1}} y^{j_{2p}} f}{\Lambda^2} \right| \mu_\Lambda(f) G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df) \right) d^n y d^n \eta dt \\ &\leq \int_{\dot{T}^*U \times \mathbb{R}^*} \tilde{\omega}(y, \eta) |t|^{n-3-k} \sqrt{\nu_{\Lambda,1}^{(y,\eta,t)}(\Omega_\Lambda)} \sqrt{\nu_{\Lambda,2}^{(y,\eta,t)}(\Omega_\Lambda)} d^n y d^n \eta dt. \end{aligned}$$

Finally, we get that :

$$(34) \quad \tilde{I}_\Lambda^{(k)}(\mathbf{j}, \omega, U) \leq \int_{\dot{T}^*U \times \mathbb{R}^*} \tilde{\omega}(y, \eta) |t|^{n-3-k} \sqrt{\nu_{\Lambda,1}^{(y,\eta,t)}(\Omega_\Lambda)} \sqrt{\nu_{\Lambda,2}^{(y,\eta,t)}(\Omega_\Lambda)} d^n y d^n \eta dt.$$

The problem is that we do not know a priori if $(y, \eta, t) \mapsto \nu_{\Lambda,1}^{(y,\eta,t)}(\Omega_\Lambda)$ and $(y, \eta, t) \mapsto \nu_{\Lambda,2}^{(y,\eta,t)}(\Omega_\Lambda)$ are integrable functions of (y, η, t) . In particular, the r.h.s. of 34 can a priori be infinite. We will now compute $\nu_{\Lambda,q}^{(y,\eta,t)}(\Omega_\Lambda)$ for $q = 1, 2$ and show that it is in fact uniformly bounded in terms of Λ by an integrable function. We note that the case $q = 1$ is a particular case of the case $q = 2$ (when $k = n - 1$). Thus, we only need to compute $\nu_{\Lambda,2}^{(y,\eta,t)}(\Omega_\Lambda)$. For that purpose, we introduce the following regularization of the Dirac distribution, for every fixed t in \mathbb{R}^* ,

$$\delta_{t, \frac{1}{m}}^{n+1}(f_0, f_1, \dots, f_n) := \int_{\mathbb{R}^{n+1}} e^{-\frac{|\tau_0|^2 - t^2 |\tau|^2}{m}} e^{-2i\pi(\tau_0, \tau) \cdot f} d\tau_0 d\tau.$$

Remark 4.20. We denote by $\Gamma = \{0\} \times (\mathbb{R}^{n+1})^*$ the wave front of δ_0^{n+1} . Using the conventions of paragraph 8.2 in [39], δ_0^{n+1} belongs to the space $\mathcal{D}'_\Gamma(\mathbb{R}^{n+1})$. One can verify that $(\delta_{t, \frac{1}{m}}^{n+1})_{m \geq 1}$ sequentially converges to δ_0^{n+1} for the topology on $\mathcal{D}'_\Gamma(\mathbb{R}^{n+1})$ for every fixed t in \mathbb{R}^* . Combining the sequential continuity of the pull-back operation (Th. 8.2.4 in [39], see also [15]) to lemma 4.12, we find that $(G_\Lambda(y, \eta, t)^*(\delta_{t, \frac{1}{m}}^{n+1}))_{m \geq 1}$ weakly converges to $G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})$ for the topology on $\mathcal{D}'(\mathbb{R}^{N(\Lambda)})$ and for every fixed (y, η, t) in $\dot{T}^*U \times \mathbb{R}^*$.

We can then define the following sequences of regularized positive Radon measures, for every $m \geq 1$:

$$\nu_{\Lambda,2,m}^{(y,\eta,t)}(df) = \prod_{p=1}^{n-1-k} \left| \frac{\partial_{y^{j_{2p-1}} y^{j_{2p}}}^2 f(y)}{\Lambda^2} \right|^2 \mu_{\Lambda}(f) G_{\Lambda}(y, \eta, t)^* (\delta_{\frac{1}{m}}^{n+1})(f) df,$$

We will now verify that $(\nu_{\Lambda,2,m}^{(y,\eta,t)}(\Omega_{\Lambda}))_{m \geq 1}$ is uniformly bounded in terms of m as follows:

$$(35) \quad \forall t > 0, \nu_{\Lambda,2,m}^{(y,\eta,t)}(\Omega_{\Lambda}) \leq \frac{C_0}{t^n} e^{-\frac{1}{C_0 t^2}},$$

where the constant C_0 is uniform for (y, η) in the support of $\tilde{\omega}$, for $0 < m \leq 1$ and $\Lambda > 0$ large enough. Then, combining this to remark 4.20, one can deduce that $\nu_{\Lambda,2}^{(y,\eta,t)}(\Omega_{\Lambda})$ is bounded by the same quantity and thanks to (34), we can conclude that $\tilde{I}_{\Lambda}^{(k)}(\mathbf{j}, \omega, U)$ is uniformly bounded in terms of Λ which was exactly what we were aiming for.

4.4.4. *Proof of proposition 4.8 – Final step: Gaussian integral bounds on $\nu_{\Lambda,2}^{(y,\eta,t)}(\Omega_{\Lambda})$.* It remains to prove (35). We fix (y, η, t) in $\hat{T}^*U \times \mathbb{R}^*$ and $m \geq 1$. We define the following map from \mathcal{H}_{Λ} to \mathbb{R}^{n^2+n+1} :

$$J_{\Lambda}^2(y) : f \mapsto F = \left(f(y); \left(\frac{\partial_{y^l} f(y)}{\Lambda} \right)_{1 \leq l \leq n}; \left(\frac{\partial_{y^r y^s}^2 f(y)}{\Lambda^2} \right)_{1 \leq r, s \leq n} \right).$$

For the coordinates in \mathbb{R}^{n^2+n+1} , we set

$$F = (F_0; F_1, \dots, F_n; F_{1,1}, \dots, F_{n,1}, F_{1,2}, \dots, F_{n,2}, \dots, F_{1,n}, F_{2,n}, \dots, F_{n,n}).$$

We now consider the pushforward of the Gaussian measure μ_{Λ} under the map $J_{\Lambda}^2(y)$. It induces the following Gaussian measure on \mathbb{R}^{n^2+n+1} :

$$(36) \quad d\tilde{\nu}_{\Lambda}^y(F) := d(J_{\Lambda}^2(y)_* \mu_{\Lambda})(F),$$

whose Fourier transform has the following expression:

$$(37) \quad \int_{\mathbb{R}^{n^2+n+1}} e^{-2i\pi T \cdot F} d\tilde{\nu}_{\Lambda}^y(F) = e^{-\frac{2\pi^2}{\text{vol}_g(M)} \langle T, A_{\Lambda}(y) T \rangle}.$$

Thanks to Remark 4.6, one can give an explicit expression for the matrix $A_{\Lambda}(y)$, namely

$$A_{\Lambda}(y) := A_0 + \mathcal{O}(\Lambda^{-1}) = \begin{pmatrix} A_{\Lambda}^{1,1}(y) & A_{\Lambda}^{1,2}(y) \\ A_{\Lambda}^{2,1}(y) & A_{\Lambda}^{2,2}(y) \end{pmatrix},$$

with the constant in the remainder uniformly bounded in terms of y and Λ , and

$$A_0 := \begin{pmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{pmatrix},$$

where the matrices $A_{i,j}$ do not depend on y and are defined as follows.

One has: $A_{1,1}$ is an $(n+1) \times (n+1)$ matrix, $A_{1,2}$ is an $(n+1) \times n^2$ matrix, $A_{2,1}$ is an $n^2 \times (n+1)$ matrix, and $A_{2,2}$ is an $n^2 \times n^2$ matrix. Their expressions are given by

$$A_{1,1} := \begin{pmatrix} 1 & 0_{1 \times n} \\ 0_{n \times 1} & \frac{1}{n+2} \text{Id}_n \end{pmatrix}, \quad A_{2,1} = A_{1,2}^T = -\frac{1}{n+2} \left((\delta_{r,s})_{1 \leq r,s \leq n}, 0_{n^2 \times n} \right),$$

and

$$A_{2,2} := \frac{1}{(n+2)(n+4)} (B_{s,s'})_{1 \leq s,s' \leq n},$$

with $B_{s,s'}$ an $n \times n$ matrix which is equal to

$$B_{s,s} := \text{Id} + 2 \text{diag}((\delta_{j,s})_{1 \leq j \leq n}) \text{ if } s = s',$$

and

$$B_{s,s'} = B_{s',s}^T := (\delta_{i,s} \delta_{j,s'} + \delta_{i,s'} \delta_{j,s})_{1 \leq i,j \leq n} \text{ if } s \neq s'.$$

With these conventions, one has

$$(38) \quad \nu_{\Lambda,2,m}^{(y,\eta,t)}(\Omega_\Lambda) \leq \int_{\mathbb{R}^{n+1}} e^{2i\pi\tau \cdot \eta} e^{-\frac{|\tau_0|^2 - t^2 |\tau|^2}{m}} \hat{\delta}_t^y(\tau_0, \tau) d\tau_0 d\tau,$$

where

$$(39) \quad \hat{\delta}_t^y(\tau_0, \tau) := \int_{\mathbb{R}^{n^2+n+1}} \prod_{p=1}^{n-1-k} F_{j_{2p-1}, j_{2p}}^2 e^{-2i\pi(\tau_0, t\tau) \cdot (F_0, \dots, F_n)} d\tilde{\nu}_\Lambda^y(F).$$

This quantity can also be rewritten more explicitly as

$$(40) \quad \hat{\delta}_t^y(\tau_0, \tau) = (-1)^{n-1-k} \prod_{p=1}^{n-1-k} \left(\frac{1}{2\pi} \frac{\partial}{\partial T_{j_{2p-1}, j_{2p}}} \right)^2 \left(e^{-\frac{2\pi^2 \langle T, A_\Lambda(y) T \rangle}{\text{Vol}_g(M)}} \right)_{|T=(\tau_0, t\tau, 0)},$$

or equivalently

$$\hat{\delta}_t^y(\tau_0, \tau) = R_{y,\Lambda}^{\mathbf{j}}(\tau_0, t\tau) e^{-\frac{2\pi^2}{\text{Vol}_g(M)} \langle (\tau_0, t\tau), A_\Lambda^{1,1}(y)(\tau_0, t\tau) \rangle},$$

for some polynom $R_{y,\Lambda}^{\mathbf{j}}$ depending only on y , Λ and \mathbf{j} and whose coefficients are uniformly bounded in terms of Λ and y . Combining this to (38), one finds that

$$(41) \quad \nu_{\Lambda,2,m}^{(y,\eta,t)}(\Omega_\Lambda) \leq \frac{1}{t^n \sqrt{\det B_\Lambda(y)}} R_{y,\Lambda}^{\mathbf{j}} \left(\frac{1}{2i\pi} \frac{\partial}{\partial \eta'_0}, \left(\frac{1}{2i\pi} \frac{\partial}{\partial \eta'_l} \right)_{1 \leq l \leq n} \right) \left(e^{-\pi \langle \eta', B_{\Lambda,m}(y) \eta' \rangle} \right)_{|\eta'=(0,\eta/t)},$$

where $B_{\Lambda,m}(y) = \frac{\pi}{\text{Vol}_g(M)} A_\Lambda^{1,1}(y) + \frac{1}{2\pi m}$. From this expression, we deduce that there exists a constant $C_0 > 0$, for every (y, η) in the support of $\tilde{\omega}$ (which is a compact subset of \mathring{T}^*U) and for every $m \geq 1$, one has

$$\forall t > 0, \quad \nu_{\Lambda,2,m}^{(y,\eta,t)}(\Omega_\Lambda) \leq \frac{C_0}{t^n} e^{-\frac{1}{C_0 t^2}},$$

which is exactly the upper bound (35) from which we get :

$$\begin{aligned} \tilde{I}_\Lambda^{(k)}(\mathbf{j}, \omega, U) &\leq \int_{\dot{T}^*U \times \mathbb{R}^*} \tilde{\omega}(y, \eta) |t|^{n-3-k} \sqrt{\nu_{\Lambda,1}^{(y,\eta,t)}(\Omega_\Lambda)} \sqrt{\nu_{\Lambda,2}^{(y,\eta,t)}(\Omega_\Lambda)} d^n y d^m \eta dt \\ &\leq C_K \|\omega\| \underbrace{\int_{\mathbb{R}^*} |t|^{n-3-k} \frac{1}{t^n} e^{-\frac{1}{C_K t^2}} dt}_{<+\infty}, \end{aligned}$$

for some constant C_K depending only on the support K of ω . The conclusion of the proposition 4.8 follows.

4.4.5. *A final remark.* We fix $k = 0$. We note that we have in fact proven something slightly stronger than what we claim in the proposition. Namely, we have shown that the map

$$(y, \eta, t, f) \mapsto t^{n-3} \eta_p \eta_{\sigma(p)} \left(\prod_{j \neq p} \frac{\partial_{y^j, y^{\sigma(j)}}^2 f}{\Lambda^2} \right) \mu_\Lambda(f) \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta)$$

belongs to $L^1(\dot{T}^*U \times \mathbb{R}^* \times \Omega_\Lambda, G_\Lambda^*(\delta_0^{n+1}))$ for every $1 \leq p \leq n$ and for every σ in S_n . Thus, regarding remark 4.9 and equality (30), it remains to compute, for every $1 \leq p \leq n$ and for every σ in S_n , the following expectation:

$$I_\Lambda(p, \sigma) := \int_{\dot{T}^*U \times \mathbb{R}^*} t^{n-3} \eta_p \eta_{\sigma(p)} \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) \left(\int_{\Omega_\Lambda} \mu_\Lambda(f) \prod_{j \neq p} \frac{\partial_{y^j, y^{\sigma(j)}}^2 f}{\Lambda^2} G_\Lambda(y, \eta, t)^*(\delta_0^{n+1})(df) \right) d^n y d^m \eta dt.$$

4.5. **Expectation of the leading term.** In the previous paragraph, we have verified that all the terms in the asymptotic expansion

$$\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) = - \sum_{k=0}^{n-1} \Lambda^{-k} I_\Lambda^{(k)}(f, \omega, U)$$

belong to $L^1(\mathcal{H}_\Lambda, d\mu_\Lambda)$ and that their L^1 norm is uniformly bounded with respect to Λ . We will now compute precisely the expectation of the leading term $I_\Lambda^{(0)}(f, \omega, U)$ from which the second part of Theorem 4.1 will follow.

Recall from remarks 4.9 and paragraph 4.4.5 that the expectation of the leading term decomposes as

$$(42) \quad \int_{\mathcal{H}_\Lambda} I_\Lambda^{(0)}(f, \omega, U) d\mu_\Lambda(f) = \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) I_\Lambda(p, \sigma).$$

In particular, it remains to compute the value of $I_\Lambda(p, \sigma)$.

4.5.1. *Regularization of $I_\Lambda(p, \sigma)$.* We fix $1 \leq p \leq n$ and σ in S_n . For that purpose, using the notations of the proof of proposition 4.8, we define the following regularization, for every fixed (y, η, t) in $\mathring{T}^*U \times \mathbb{R}^*$,

$$I_{\Lambda, p, \sigma}^{(m)}(y, \eta, t) = \int_{\mathcal{H}_\Lambda} \left(\prod_{j \neq p} \frac{\partial_{y^j, y^{\sigma(j)}}^2 f}{\Lambda^2} \right) \mu_\Lambda(f) G_\Lambda(y, \eta, t)^* (\delta_{t, \frac{1}{m}}^{n+1})(df).$$

Note that

$$f \mapsto \left(\prod_{j \neq p} \frac{\partial_{y^j, y^{\sigma(j)}}^2 f}{\Lambda^2} \right) \mu_\Lambda(f)^{\frac{1}{2}}$$

tends to 0 as $\|f\|$ tends to infinity; thus, it can be approximated (in the \mathcal{C}^0 topology) by a function $\psi_1(f)$ in $\mathcal{C}_c^\infty(\mathcal{H}_\Lambda)$. Moreover, mimicking the arguments of the proof of proposition 4.8, one can verify that the sequence

$$\left(\int_{\mathcal{H}_\Lambda} \mu_\Lambda(f)^{\frac{1}{2}} G_\Lambda(y, \eta, t)^* (\delta_{t, \frac{1}{m}}^{n+1})(df) \right)_{m \geq 1}$$

is uniformly bounded in terms of $m \geq 1$ (not necessarily in terms of $\Lambda, (y, \eta, t)$). Combining these two observations with (31), we find that, for every $1 \leq p \leq n$, for every σ in S_n , and for a.e. (y, η, t) in $\mathring{T}^*U \times \mathbb{R}^*$,

$$\lim_{m \rightarrow +\infty} I_{\Lambda, p, \sigma}^{(m)}(y, \eta, t) = \int_{\Omega_\Lambda} \mu_\Lambda(f) \prod_{j \neq p} \frac{\partial_{y^j, y^{\sigma(j)}}^2 f}{\Lambda^2} G_\Lambda(y, \eta, t)^* (\delta_0^{n+1})(df).$$

According to the proof of proposition 4.8, one also knows that there exists a constant $C_0 > 0$ such that, for every $m \geq 1$, for every $1 \leq p \leq n$, for every σ in S_n , one has

$$\forall (y, \eta, t) \in \text{supp}(\omega_{(0, \dots, 0), (1, \dots, 1)}) \times \mathbb{R}^*, \quad |I_{\Lambda, p, \sigma}^{(m)}(y, \eta, t)| \leq \frac{C_0}{t^n} e^{-\frac{1}{C_0 t^2}}.$$

In particular, from the dominated convergence theorem, we can deduce that

$$(43) \quad I_\Lambda(p, \sigma) = \lim_{m \rightarrow +\infty} \int_{\mathring{T}^*U \times \mathbb{R}^*} t^{n-3} \eta_p \eta_{\sigma(p)} \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) I_{\Lambda, p, \sigma}^{(m)}(y, \eta, t) d^n y d^n \eta dt.$$

Using Gaussian integration, we will now compute explicitly this limit as m tends to $+\infty$ and as $\Lambda \rightarrow +\infty$. Following the proof of proposition 4.8 – see e.g. equality (40), one has

$$I_{\Lambda, p, \sigma}^{(m)}(y, \eta, t) = (-1)^{\frac{n-1}{2}} \int_{\mathbb{R}^{n+1}} e^{2i\pi\tau \cdot \eta} e^{-\frac{|\tau_0|^2 - t^2 |\tau|^2}{m}} \prod_{j \neq p} \left(\frac{1}{2\pi} \frac{\partial}{\partial T_{j, \sigma(j)}} \right) \left(e^{-\frac{2\pi^2 \langle T, A_\Lambda(y) T \rangle}{\text{Vol}_g(M)}} \right) \Big|_{T=(\tau_0, t\tau, 0)} d\tau_0 d\tau.$$

Following the end of the proof of proposition 4.8, one can also verify that there exists a constant $C_0 > 0$, such that for every (y, η) in the support of $\tilde{\omega}$ and for every $m \geq 1$, one has

$$(44) \quad \forall t > 0, \quad |I_{\Lambda, p, \sigma}^{(m)}(y, \eta, t) - \tilde{I}_{\Lambda, p, \sigma}^{(m)}(\eta, t)| \leq \frac{C_0 \Lambda^{-1}}{t^n} e^{-\frac{1}{C_0 t^2}},$$

where

$$\tilde{I}_{\Lambda,p,\sigma}^{(m)}(\eta, t) = \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \int_{\mathbb{R}^{n+1}} e^{2i\pi\tau \cdot \eta} e^{-\frac{|\tau_0|^2 - t^2 |\tau|^2}{m}} \prod_{j \neq p} \left(\frac{\partial}{\partial T_{j,\sigma(j)}} \right) \left(e^{-\frac{2\pi^2 \langle T, A_0 T \rangle}{\text{Vol}_g(M)}} \right)_{T=(\tau_0, t\tau)} d\tau_0 d\tau.$$

In particular, provided that the limit $\lim_{m \rightarrow +\infty} \tilde{I}_{\Lambda,p,\sigma}^{(m)}(\eta, t)$ exists a.e., one has by the dominated convergence theorem

(45)

$$I_{\Lambda}(p, \sigma) = \int_{\hat{T}^*U \times \mathbb{R}^*} t^{n-3} \eta_p \eta_{\sigma(p)} \omega_{(0,\dots,0),(1,\dots,1)}(y, \eta) \left(\lim_{m \rightarrow +\infty} \tilde{I}_{\Lambda,p,\sigma}^{(m)}(\eta, t) \right) d^n y d^n \eta dt + \mathcal{O}(\Lambda^{-1}).$$

Remark 4.21. We note that, thanks to the particular form of the matrix A_0 , one has

$$\tilde{I}_{\Lambda,p,\sigma}^{(m)}(\eta, t) = \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \int_{\mathbb{R}^{n+1}} e^{2i\pi\tau \cdot \eta} e^{-\frac{|\tau_0|^2 - t^2 |\tau|^2}{m}} e^{-\frac{2\pi^2 t^2 |\tau|^2}{\text{Vol}_g(M)(n+2)}} H_{p,\sigma}(\tau_0) d\tau_0 d\tau,$$

where

$$H_{p,\sigma}(\tau_0) := \prod_{j \neq p} \left(\frac{\partial}{\partial T_{j,\sigma(j)}} \right) \left(e^{-\frac{2\pi^2}{\text{Vol}_g(M)} \langle (T_0; 0; (T_{i,j})), A_0(T_0; 0; (T_{i,j})) \rangle} \right)_{T=(\tau_0; 0; 0)}.$$

After integrating over the τ variable and taking the limit $m \rightarrow +\infty$, one finds

$$\lim_{m \rightarrow +\infty} \tilde{I}_{\Lambda,p,\sigma}^{(m)}(\eta, t) = \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \frac{1}{t^n} \left(\frac{(n+2)\text{Vol}_g(M)}{2\pi} \right)^{\frac{n}{2}} e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2}{2t^2}} \int_{\mathbb{R}} H_{p,\sigma}(\tau_0) d\tau_0.$$

4.5.2. *Computation of the leading term.* Recall that everything we want to calculate is the leading term $-\int_{\mathcal{H}_{\Lambda}} I_{\Lambda}^{(0)}(f, \omega, U) d\mu_{\Lambda}(f)$ of $\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})_{\Lambda}} \omega(y, dy, \eta, d\eta)$ which, thanks to Remark 4.21, is given by

$$\begin{aligned} \int_{\mathcal{H}_{\Lambda}} I_{\Lambda}^{(0)}(f, \omega, U) d\mu_{\Lambda}(f) &= \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) I_{\Lambda}(p, \sigma) \\ &= \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) \int_{\hat{T}^*U \times \mathbb{R}^*} t^{n-3} \eta_p \eta_{\sigma(p)} \omega_{(0,\dots,0),(1,\dots,1)}(y, \eta) \lim_{m \rightarrow +\infty} \left(\tilde{I}_{\Lambda,p,\sigma}^{(m)}(\eta, t) \right) d^n y d^n \eta dt \\ &= \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \left(\frac{(n+2)\text{Vol}_g(M)}{2\pi} \right)^{\frac{n}{2}} \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) \int_{\hat{T}^*U \times \mathbb{R}^*} \eta_p \eta_{\sigma(p)} \omega_{(0,\dots,0),(1,\dots,1)}(y, \eta) \frac{1}{t^3} \\ &\quad e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2}{2t^2}} \int_{\mathbb{R}} H_{p,\sigma}(\tau_0) d\tau_0 d^n y d^n \eta dt \end{aligned}$$

In order to alleviate notations, we introduce:

$$(46) \quad I_0(\eta) = \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) \eta_p \eta_{\sigma(p)} \int_{\mathbb{R}} H_{p,\sigma}(\tau_0) d\tau_0.$$

Then, we express everything in terms of $I_0(\eta)$:

$$\begin{aligned}
 & \int_{\mathcal{H}_\Lambda} I_\Lambda^{(0)}(f, \omega, U) d\mu_\Lambda(f) \\
 = & \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \left(\frac{(n+2)\text{Vol}_g(M)}{2\pi} \right)^{\frac{n}{2}} \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) \int_{T^*U \times \mathbb{R}^*} \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) \eta_p \eta_{\sigma(p)} \\
 & \frac{1}{t^3} e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2}{2t^2}} \int_{\mathbb{R}} H_{p, \sigma}(\tau_0) d\tau_0 d^n y d^n \eta dt \\
 = & \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \left(\frac{(n+2)\text{Vol}_g(M)}{2\pi} \right)^{\frac{n}{2}} \int_{T^*U \times \mathbb{R}^*} \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) \\
 & \frac{1}{t^3} e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2}{2t^2}} I_0(\eta) d^n y d^n \eta dt \text{ by definition of } I_0(\eta) \\
 = & \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \left(\frac{(n+2)\text{Vol}_g(M)}{2\pi} \right)^{\frac{n}{2}} \int_{T^*U} \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) I_0(\eta) d^n y d^n \eta \\
 & \times \left(\int_0^{+\infty} t^3 e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2 t^2}{2}} \frac{dt}{t^2} + \int_0^{-\infty} t^3 e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2 t^2}{2}} \frac{dt}{t^2} \right) \\
 = & \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \left(\frac{(n+2)\text{Vol}_g(M)}{2\pi} \right)^{\frac{n}{2}} \int_{T^*U} \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) I_0(\eta) d^n y d^n \eta \\
 & \int_0^{+\infty} e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2 t^2}{2}} 2t dt \\
 = & \frac{2(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \frac{((n+2)\text{Vol}_g(M))^{\frac{n}{2}-1}}{(2\pi)^{\frac{n}{2}}} \int_{T^*U} \omega_{(0, \dots, 0), (1, \dots, 1)}(y, \eta) \frac{I_0(\eta)}{|\eta|^2} d^n y d^n \eta.
 \end{aligned}$$

Therefore,

$$(47) \quad \int_{\mathcal{H}_\Lambda} \left(\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) \right) d\mu_\Lambda(f) = A_n \int_{\tilde{T}^*U} \frac{I_0(\eta)}{|\eta|^2} dy^1 \wedge \dots \wedge dy^n \wedge \omega(y, dy, \eta, d\eta) + \mathcal{O}(\Lambda^{-1}),$$

where

$$A_n := \frac{2(-1)^{\frac{n+1}{2}} ((n+2)\text{Vol}_g(M))^{\frac{n}{2}-1}}{(2\pi)^{n-1} (2\pi)^{\frac{n}{2}}}.$$

In order to conclude, it remains to compute $I_0(\eta)$. This combinatorial calculation will be the purpose of the next two paragraphs.

4.6. Computation of $I_0(\eta)$. Recall that $I_0(\eta)$ is given by the sum (46). We start by computing explicitly each term in the sum. We fix σ in S_n and $1 \leq p \leq n$. Moreover, we set

$$\tilde{T} := (T_0; 0; (T_{i,j})_{1 \leq i, j \leq n}) \in \mathbb{R}^{n^2+n+1},$$

which appears in the definition of $H_{p,\sigma}(\tau_0)$ – see Remark 4.21. Our first step will be to use the Faà di Bruno’s formula for partial derivatives [36] to give an explicit expression of

$$\prod_{j \neq p} \left(\frac{\partial}{\partial T_{j,\sigma(j)}} \right) \left(e^{-\frac{2\pi^2 \langle \tilde{T}, A_0 \tilde{T} \rangle}{\text{Vol}_g(M)}} \right).$$

Recall that this formula states that

$$\prod_{j \neq p} \left(\frac{\partial}{\partial T_{j,\sigma(j)}} \right) \left(e^{-\frac{2\pi^2 \langle \tilde{T}, A_0 \tilde{T} \rangle}{\text{Vol}_g(M)}} \right) = e^{-\frac{2\pi^2 \langle \tilde{T}, A_0 \tilde{T} \rangle}{\text{Vol}_g(M)}} \sum_{\mathfrak{a} \in \mathcal{P}_{\sigma,p}^{(0)}} \left(-\frac{2\pi^2}{\text{Vol}_g(M)} \right)^{|\mathfrak{a}|} \prod_{\mathfrak{a} \in \mathfrak{a}} \frac{\partial^{|\mathfrak{a}|} \langle \tilde{T}, A_0 \tilde{T} \rangle}{\prod_{j \in \mathfrak{a}} \partial T_{j,\sigma(j)}},$$

where $\mathcal{P}_{\sigma,p}^{(0)}$ is the set of partitions of $\{1 \leq j \leq n : j \neq p\}$. We now fix some notations:

$$\mathcal{P}_{\sigma,p} := \{ \mathfrak{a} \text{ partition of } \{1 \leq j \leq n : j \neq p\} : \forall \mathfrak{a} \in \mathfrak{a}, |\mathfrak{a}| \leq 2 \text{ and } \sigma(\mathfrak{a}) = \mathfrak{a} \}.$$

Remark 4.22. We note that $\mathcal{P}_{\sigma,p}$ is empty if $\sigma(p) \neq p$. Moreover, for every σ such that $\sigma(p) = p$, the above subset is empty if σ is not a product of disjoint 2-cycles.

For a given \mathfrak{a} in $\mathcal{P}_{\sigma,p}$, we define

- $\mathfrak{a}_1 := \{ \mathfrak{a} \in \mathfrak{a} : |\mathfrak{a}| = 1 \}$;
- $\mathfrak{a}_2 := \{ \mathfrak{a} \in \mathfrak{a} : |\mathfrak{a}| = 2 \text{ and } \sigma_{|\mathfrak{a}} = \text{id} \}$;
- $\mathfrak{a}_3 := \{ \mathfrak{a} \in \mathfrak{a} : |\mathfrak{a}| = 2 \text{ and } \sigma_{|\mathfrak{a}} \neq \text{id} \}$.

Using these conventions and the specific form of the matrix A_0 – see paragraph 4.4.4, one finds that

$$\begin{aligned} & \prod_{j \neq p} \left(\frac{\partial}{\partial T_{j,\sigma(j)}} \right) \left(e^{-\frac{2\pi^2 \langle T, A_0 T \rangle}{\text{Vol}_g(M)}} \right) \Big|_{T=(\tau_0;0;0)} \\ &= e^{-\frac{2\pi^2 \tau_0^2}{\text{Vol}_g(M)}} \sum_{\mathfrak{a} \in \mathcal{P}_{\sigma,p}} \left(-\frac{4\pi^2}{\text{Vol}_g(M)(n+2)} \right)^{|\mathfrak{a}|} \left(\frac{1}{n+4} \right)^{|\mathfrak{a}_2|+|\mathfrak{a}_3|} (-\tau_0)^{|\mathfrak{a}_1|}. \end{aligned}$$

Observe now that

$$\int_{\mathbb{R}} \tau_0^{|\mathfrak{a}_1|} e^{-\frac{2\pi^2 \tau_0^2}{\text{Vol}_g(M)}} d\tau_0 = \left(\frac{\text{Vol}_g(M)}{2\pi} \right)^{\frac{1}{2}} (|\mathfrak{a}_1| - 1)!! \left(\frac{\text{Vol}_g(M)}{4\pi^2} \right)^{\frac{|\mathfrak{a}_1|}{2}}$$

if $|\mathfrak{a}_1| \equiv 0 \pmod{2}$, and that it is equal to 0 otherwise. We now set

$$C_{\sigma,p} := \sum_{\mathfrak{a} \in \mathcal{P}_{\sigma,p} : |\mathfrak{a}_1| \equiv 0 \pmod{2}} \left(-\frac{1}{n+2} \right)^{|\mathfrak{a}|} (|\mathfrak{a}_1| - 1)!! \left(\frac{4\pi^2}{\text{Vol}_g(M)} \right)^{|\mathfrak{a}| - \frac{|\mathfrak{a}_1|}{2}} \left(\frac{1}{n+4} \right)^{|\mathfrak{a}_2|+|\mathfrak{a}_3|},$$

whenever $\sigma(p) = p$ and $C_{\sigma,p} := 0$ otherwise.

Remark 4.23. We note that the sum is empty as soon as n is even. In fact, requiring $|\mathfrak{a}_1| \equiv 0 \pmod{2}$ imposes that $n - 1$ is even.

From the above calculation, we deduce

$$(48) \quad I_0(\eta) = \sum_{p=1}^n \sum_{\sigma \in S_n} \varepsilon(\sigma) \eta_p \eta_{\sigma(p)} \int_{\mathbb{R}} H_{p,\sigma}(\tau_0) d\tau_0 = \left(\frac{\text{Vol}_g(M)}{2\pi} \right)^{\frac{1}{2}} \sum_{p=1}^n \eta_p^2 \sum_{\sigma \in S_n} \varepsilon(\sigma) C_{\sigma,p}.$$

In particular, from remark 4.23, one has $I_0(\eta) = 0$ if n is even. We also note that $\sum_{\sigma \in S_n} C_{\sigma,p}$ is independent of the choice of p . Thus, one finds, whenever n is odd,

$$(49) \quad I_0(\eta) = \left(\frac{\text{Vol}_g(M)}{2\pi} \right)^{\frac{1}{2}} |\eta|^2 \sum_{\sigma \in S_n} \varepsilon(\sigma) C_{\sigma,p}.$$

From (47), one finally obtains

$$(50) \quad \int_{\mathcal{H}_\Lambda} \left(\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, \eta, dy, d\eta) \right) d\mu_\Lambda(f) \\ = \left(\frac{\text{Vol}_g(M)}{2\pi} \right)^{\frac{1}{2}} A_n \left(\sum_{\sigma \in S_n} \varepsilon(\sigma) C_{\sigma,p} \right) \int_{\hat{T}^*U} dy^1 \wedge \dots \wedge dy^n \wedge \omega(y, dy, \eta, d\eta) + \mathcal{O}(\Lambda^{-1}),$$

where A_n was defined in (47).

4.7. Computation of $\sum_{\sigma \in S_n} C_{\sigma,p}$. From (50), it remains to compute $\sum_{\sigma \in S_n} C_{\sigma,p}$ which is independent of the choice of p . For that purpose, we fix $0 \leq k \leq \frac{n-1}{2}$ and we suppose that σ is the product of k disjoint 2-cycles $\sigma_1, \dots, \sigma_k$. Then, we compute the value of $C_{\sigma,p}$ in this case. We note that in this case $|\mathfrak{A}_3|$ is necessarily equal to k , and thus

$$C_{\sigma,p} = \left(\frac{1}{n+4} \right)^k \sum_{\mathfrak{A} \in \mathcal{P}_{\sigma,p}: |\mathfrak{A}_1| \equiv 0 \pmod{2}} \left(-\frac{1}{n+2} \right)^{|\mathfrak{A}|} (|\mathfrak{A}_1| - 1)!! \left(\frac{4\pi^2}{\text{Vol}_g(M)} \right)^{|\mathfrak{A}| - \frac{|\mathfrak{A}_1|}{2}} \left(\frac{1}{n+4} \right)^{|\mathfrak{A}_2|}.$$

We also note the following useful relations

$$n-1 = |\mathcal{A}_1| + 2|\mathcal{A}_2| + 2|\mathcal{A}_3| \quad \text{and} \quad \mathcal{A} = |\mathcal{A}_1| + |\mathcal{A}_2| + |\mathcal{A}_3|.$$

Recall now that the number of permutations in $\{1, 2, \dots, 2l\}$ with l cycles of length 2 is equal to $\frac{(2l)!}{l!2^l}$ – see Th. 6.9 in [13]. From this observation, we deduce that

$$C_{\sigma,p} = \left(\frac{4\pi^2}{(n+2)^2 \text{Vol}_g(M)} \right)^{\frac{n-1}{2}} \left(-\frac{n+2}{n+4} \right)^k \sum_{l=0}^{\frac{n-1}{2}-k} \frac{(2l)!}{l!2^l} C_{n-1-2k}^{2l} (n-2(k+l+1))!! \left(-\frac{n+2}{n+4} \right)^l,$$

and thus

$$C_{\sigma,p} = \left(\frac{2\pi^2}{2(n+2)^2 \text{Vol}_g(M)} \right)^{\frac{n-1}{2}} \left(-\frac{2(n+2)}{n+4} \right)^k \sum_{l=0}^{\frac{n-1}{2}-k} \frac{(n-1-2k)!}{l! \left(\frac{n-1}{2} - (k+l) \right)!} \left(-\frac{n+2}{n+4} \right)^l.$$

Thanks to our formula for $C_{\sigma,p}$ when σ is a product of k disjoint 2-cycles (thus $\varepsilon(\sigma) = (-1)^k$), we deduce that

$$\sum_{\sigma \in S_n} \varepsilon(\sigma) C_{\sigma,p} = \left(\frac{2\pi^2}{(n+2)^2 \text{Vol}_g(M)} \right)^{\frac{n-1}{2}} \sum_{k=0}^{\frac{n-1}{2}} C_{n-1}^{2k} \frac{(2k)!}{k! 2^k} 2^k \sum_{l=0}^{\frac{n-1}{2}-k} (-1)^l \frac{(n-1-2k)!}{l! 2^l \left(\frac{n-1}{2} - (k+l)\right)!} \left(\frac{n+2}{n+4}\right)^{l+k}.$$

After simplification, it gives

$$\sum_{\sigma \in S_n} \varepsilon(\sigma) C_{\sigma,p} = (n-1)! \left(\frac{2\pi^2}{(n+2)^2 \text{Vol}_g(M)} \right)^{\frac{n-1}{2}} \sum_{k=0}^{\frac{n-1}{2}} \left(\frac{n+2}{n+4}\right)^k \sum_{l=0}^{\frac{n-1}{2}-k} \frac{\left(-\frac{n+2}{n+4}\right)^l}{k! l! \left(\frac{n-1}{2} - (k+l)\right)!},$$

and thus

$$\sum_{\sigma \in S_n} \varepsilon(\sigma) C_{\sigma,p} = (n-1)! \left(\frac{2\pi^2}{(n+2)^2 \text{Vol}_g(M)} \right)^{\frac{n-1}{2}} \sum_{k=0}^{\frac{n-1}{2}} \frac{1}{k! \left(\frac{n-1}{2} - k\right)!} \left(\frac{n+2}{n+4}\right)^k \left(1 - \frac{n+2}{n+4}\right)^{\frac{n-1}{2}-k}.$$

Finally, we obtain

$$(51) \quad \sum_{\sigma \in S_n} \varepsilon(\sigma) C_{\sigma,p} = \frac{(n-1)!}{\left(\frac{n-1}{2}\right)!} \left(\frac{2\pi^2}{(n+2)^2 \text{Vol}_g(M)} \right)^{\frac{n-1}{2}}.$$

4.8. The conclusion. Regarding (50) and (51), we define

$$B_n := \frac{2(-1)^{\frac{n+1}{2}} (n-1)!}{(2\pi)^{n-1}} \frac{\pi^{\frac{n-3}{2}}}{\left(\frac{n-1}{2}\right)!} \frac{1}{2(n+2)^{\frac{n}{2}}} = \frac{(-1)^{\frac{n+1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})} \frac{2}{(n+2)^{\frac{n}{2}}}.$$

if $n \equiv 1 \pmod{2}$ and $B_n = 0$ otherwise. Finally, still thanks to (50), we conclude

$$\int_{\mathcal{H}_\Lambda} \left(\frac{1}{\Lambda^n} \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta) \right) d\mu_\Lambda(f) = B_n \int_{\dot{T}^*U} \omega_{(0,\dots,0),(1,\dots,1)}(y, \eta) d^n y d^n \eta + \mathcal{O}(\Lambda^{-1}),$$

which is exactly the content of the second part of Theorem 4.1.

5. FORMAL DERIVATION OF THE MAIN RESULT VIA BEREZIN INTEGRATION

The derivation of the main theorem (namely Theo. 4.1) is slightly technical due to the fact that we have to justify carefully the smallness of several terms and the inversion of several integrals but also due to some complicated combinatorics. The purpose of this last section is to show how the combinatorial aspects of the proof can be treated more easily at the expense of a little bit of abstraction by making use of the so-called Berezin integral [4, 33, 23, 57].

In this section, we will not pay too much attention to the inversion of integrals and to the size of the remainder terms, and we will mostly focus on the computation of the leading term of the asymptotics. Similar arguments as the ones presented in the previous section would in fact provide a rigorous treatment of the calculation developed in the present section. The reason for presenting this alternative approach in a formal (but non completely rigorous manner) is that we believe that it can be helpful (at least for the

reader familiar with Berezin integration) to understand and follow the main lines of our proof, especially its combinatorial aspects. It is also plausible that this kind of intuitive calculation could be used for other related questions.

5.1. A brief reminder on Berezin integration. Berezin integration is a convenient formalism that allows us to represent the wedge product appearing in our formulas as oscillatory integrals. This formalism was introduced by Berezin [4] and it is a more or less standard tool in quantum field theory where one aims at computing integrals over both bosonic (even) and fermionic (odd) variables. Basic introductions to this formalism are described by Disertori [23] or by Tao [57] with a view towards random matrix theory – see also the book of Guillemin and Sternberg [33] for considerations on Fourier transforms in this context. Following closely the presentation of these references, we give here a brief overview on it that should be sufficient for the purpose of our formal calculation.

5.1.1. Odd and even variables. Let $V = \mathbb{R}^d$ be a finite dimensional vector space. One can define the so called exterior algebra $\bigwedge(V) := \bigoplus_{k=0}^{+\infty} \bigwedge^k(V)$. For simplicity of notations, we will denote by $\Phi_1\Phi_2 := \Phi_1 \wedge \Phi_2$ the product on this algebra. This algebra can be splitted in two parts. One part is made of the so-called even (or bosonic) elements $\bigwedge(V)_{\text{even}} := \bigoplus_{k=0}^{+\infty} \bigwedge^{2k}(V)$, while the other is made of the so-called odd (or fermionic) elements $\bigwedge(V)_{\text{odd}} := \bigoplus_{k=0}^{+\infty} \bigwedge^{2k+1}(V)$. Any odd element Π commutes with every even element p , while it anticommutes with odd elements. From this observation, one can deduce that any element Φ in $\bigoplus_{k=1}^{+\infty} \bigwedge^k(V)$ is nilpotent.

Fix two nonnegative integers d_1 and d_2 . For every $i \leq d_1$, we fix a map

$$p_i : \mathbb{R}^{d_1} \rightarrow \bigwedge(V)_{\text{even}}, \quad x \mapsto x_i + n_i(x),$$

where $n_i(x)$ is nilpotent. Any “sufficiently nice” function f on \mathbb{R}^{d_1} can be extended as a function of even variables as follows:

$$f(p_1, \dots, p_{d_1}) := \sum_{k_i \geq 0: 1 \leq i \leq d_1} \frac{1}{k_1! \dots k_{d_1}!} \frac{\partial^{k_1 + \dots + k_{d_1}} f}{\partial x_1^{k_1} \dots \partial x_{d_1}^{k_{d_1}}} n_1(x)^{k_1} \dots n_{d_1}(x)^{k_{d_1}}.$$

In the following, we will in fact consider functions of even variables $p = (p_1, \dots, p_{d_1})$ and odd variables $\Pi = (\Pi_1, \dots, \Pi_{d_2})$, i.e.

$$f(p, \Pi) := f_0(p) + \sum_{k \geq 1} \sum_{i_1, \dots, i_k} f_{i_1, \dots, i_k}(p) \Pi_{i_1} \dots \Pi_{i_k}.$$

We say that such a function is defined on the space $\bigwedge(V)^{(d_1|d_2)}$.

5.1.2. Integration of functions of odd and even variables. Given a function f on $\bigwedge(V)^{(d_1|d_2)}$, one can define the Berezin integral of f as follows:

$$\int_{\mathbb{R}^{(d_1|d_2)}} f(p, \Pi) dp d\Pi := \int_{\mathbb{R}^{d_1}} \partial_{\Pi_1} \dots \partial_{\Pi_{d_2}} f dx_1 dx_2 \dots dx_{d_1}.$$

We refer the reader to [23] for basic properties on this integration procedure and for a review of simple examples. One can verify that any element Φ of $\Lambda(V)$ can be exponentiated as follows:

$$\exp(\Phi) := \sum_{k=0}^{+\infty} \frac{\Phi^k}{k!}.$$

Important properties of the exponential are that $\exp(\Pi) = 1 + \Pi$ for any odd element Π and that, for any even element p ,

$$(52) \quad \exp(p + \Phi) = \exp(p) \exp(\Phi) = \exp(\Phi) \exp(p).$$

Following [33] – Chapter 7, one can also define a notion of “super Fourier transform”, as follows:

$$\mathcal{F}(f)(q, Q) := \int_{\mathbb{R}^{(d_1|d_2)}} \exp(-2i\pi(q.p + Q.\Pi)) f(p, \Pi) dp d\Pi.$$

The prefix “super” is here to emphasize that we consider functions depending both on even (bosonic) and odd (fermionic) variables. We observe that $q.p + Q.\Pi$ is an even element. Note that an element Q in $\Lambda(V)_{\text{odd}}$ can be written as follows:

$$(53) \quad Q = -\frac{1}{2i\pi} \int_{\mathbb{R}^{(0|1)}} \exp(-2i\pi Q\Pi) d\Pi,$$

and more generally a product of elements Q_1, \dots, Q_{d_2} in $\Lambda(V)_{\text{odd}}$ as

$$(54) \quad Q_1 \dots Q_{d_2} = \frac{1}{(-2i\pi)^{d_2}} \int_{\mathbb{R}^{(0|d_2)}} \exp\left(-2i\pi \sum_{j=1}^{d_2} Q_j \Pi_j\right) d\Pi_1 \dots d\Pi_{d_2}.$$

This formula gives us a representation of products of odd elements as “oscillatory” integrals. This observation is the first key point which is at the heart of the calculation below.

5.1.3. Representation of integration currents as oscillatory integrals over even and odd variables. We consider a submanifold S in some larger manifold X given by equations $G = (g_i)_{1 \leq i \leq d} = 0$. Then it is a well known fact that the delta function $\delta_0^d \circ G$ supported on S can be represented as an oscillatory integral. So a natural question is how to represent the integration current $[S] = [\{G = 0\}]$ as an oscillatory integral? The answer involves the introduction of odd variables and the use of the Berezin integral.

Proposition 5.1. *Let X be a smooth oriented manifold with orientation $[\omega_1]$,*

$$G := (g_i)_{1 \leq i \leq d} : X \rightarrow \mathbb{R}^d$$

be a smooth function such that the differentials $d_x g_i$ are linearly independent for every $x \in X$ satisfying $g_1(x) = \dots = g_d(x) = 0$. Set S to be the submanifold in X defined by the regular system of equations $\{g_1 = \dots = g_d = 0\}$ and oriented by $[\omega_2]$. If $dg_1 \wedge \dots \wedge dg_d$ has orientation compatible with the pair of orientations $[\omega_1], [\omega_2]$ then the integration current $[S]$ is represented by the oscillatory integral

$$(55) \quad [S] = \frac{1}{(-2i\pi)^d} \int_{\mathbb{R}^{(d|d)}} e^{-2i\pi(\sum_{j=1}^d \tau_j g_j + \Pi_j dg_j)} d\tau_1 \dots d\tau_d d\Pi_1 \dots d\Pi_d.$$

Proof. The proof follows from a simple calculation, by Lemma 3.4, we find that :

$$[S] = G^* \delta_0^d dg_1 \wedge \cdots \wedge dg_d$$

then using the classical representation of the delta function by oscillatory integral

$$G^* \delta_0^d dg_1 \wedge \cdots \wedge dg_d = \int_{\mathbb{R}^d} e^{-2i\pi \sum_{j=1}^d \tau_j g_j} d\tau_1 \dots d\tau_d dg_1 \wedge \cdots \wedge dg_d$$

and using the representation of the exterior product $dg_1 \wedge \cdots \wedge dg_d$ in terms of the Berezin integral we end up with

$$[S] = \frac{1}{(-2i\pi)^d} \int_{\mathbb{R}^{(d|d)}} e^{-2i\pi(\sum_{j=1}^d \tau_j g_j + \Pi_j dg_j)} d\tau_1 \dots d\tau_d d\Pi_1 \dots d\Pi_d.$$

□

In fact, this oscillatory integral formula is already known in the literature since it appears in the work of Frenkel, Losev and Nekrasov on instantonic quantum field theory [27, p. 23].

5.1.4. *Fourier transform of Gaussian integrals.* We observe that the following holds:

Lemma 5.2. *Let $N \geq 1$. Let $p = (p_1, \dots, p_N)$ be an element in $\Lambda(V)_{\text{even}}^N$. Then, one has*

$$(56) \quad e^{-\frac{p^2}{2N}} = \left(\frac{N}{2\pi}\right)^{\frac{N}{2}} \int_{\mathbb{R}^N} e^{i\langle c, p \rangle} e^{-N\frac{\langle c, c \rangle}{2}} d^N c,$$

where $\langle c, p \rangle = \sum_{j=1}^N c_j p_j$, and $p^2 = \sum_{j=1}^N p_j^2$.

This lemma will be the second key point in the computation of the asymptotics via Berezin integration.

Proof. Let $\mathbb{C}[[X_1, \dots, X_N]]$ be the commutative algebra of formal power series in N indeterminates (X_1, \dots, X_N) . First, we note that

$$(57) \quad e^{-\frac{X^2}{2N}} \quad \text{and} \quad \left(\frac{N}{2\pi}\right)^{\frac{N}{2}} \int_{\mathbb{R}^N} e^{i\langle c, X \rangle} e^{-N\frac{\langle c, c \rangle}{2}} d^N c$$

are well defined as formal power series since the integral on the right hand side is defined by expanding the exponential function $e^{i\langle c, X \rangle}$ in powers of X under the integral sign where, for each (i_1, \dots, i_k) , the term $\int_{\mathbb{R}^N} (c_{i_1} \dots c_{i_k}) e^{-N\frac{\langle c, c \rangle}{2}} d^N c$ is a convergent integral that can be explicitly calculated by the Wick lemma. Then the equality between the two formal series in (57) follows from the classical result on the Fourier transform of the Gaussian measure replacing the formal indeterminate X by $y \in \mathbb{R}^N$ and by uniqueness of the Taylor series expansion in y at $y = 0$. □

Remark 5.3. Conceptually, we note that this proof shows that we can define the Fourier transform of a Gaussian measure in a purely algebraic language. We emphasize that the argument would work for more general symmetric positive definite matrices A than $N\text{Id}$.

5.2. Asymptotic expansion in a local chart. We fix a local geodesic normal coordinate chart (U, ϕ) and $\omega(y, dy, \eta, d\eta)$ an element in $\mathcal{D}_n(\mathring{T}^*U)$. Recall that we want to compute the expectation of

$$\mathcal{I}_\Lambda(f, \omega) := \int_{N^*(\{f=0\})} \omega(y, dy, \eta, d\eta)$$

with respect to the Gaussian measure $d\mu_\Lambda$. We can write formally

$$\mathcal{I}_\Lambda(f, \omega) = \int_{\mathring{T}^*U \times \mathbb{R}^*} \delta_0^{n+1} \left(f, \frac{td_y f}{\Lambda} - \eta \right) df \wedge \bigwedge_{j=1}^n d \left(\frac{t\partial_{y^j} f}{\Lambda} - \eta_j \right) \wedge \omega(y, dy, \eta, d\eta).$$

Recall that one of the reason why the calculation of the expectation against the Gaussian measure is possible is the fact that the Dirac distribution can be represented by an oscillatory integral over even variables, i.e.

$$\delta_0^{n+1} \left(f, \frac{td_y f}{\Lambda} - \eta \right) = \int_{\mathbb{R}^{(n+1|0)}} \exp \left(-2i\pi \left(\tau_0 f + \tau \cdot \left(\frac{td_y f}{\Lambda} - \eta \right) \right) \right) d\tau_0 d\tau.$$

Then, some of the difficulties in the proof come from the fact that the wedge product has not at first sight the same simple structure. By Proposition 5.1, we see that the advantage of the Berezin formalism described above is that it allows us to write the wedge product as an oscillatory integral over odd variables in $\mathbb{R}^{(0|n)}$. Before writing this oscillatory representation – see equation (59), we proceed to a few simplifications that will make the calculation slightly simpler. More precisely, we expand the wedge product, and we obtain that

$$\mathcal{I}_\Lambda(f, \omega) = \Lambda \sum_{r,s=1}^n (-1)^r \int_{\mathring{T}^*U \times \mathbb{R}^*} \frac{\eta_r \eta_s}{t^2} dy^s \wedge \mathcal{I}_\Lambda^{r,s}(f, t, y, dy, \eta, d\eta) \wedge \omega(y, dy, \eta, d\eta) \wedge dt,$$

where

$$\mathcal{I}_\Lambda^{r,s}(f, t, y, dy, \eta, d\eta) := \delta_0^{n+1} \left(f, \frac{td_y f}{\Lambda} - \eta \right) \bigwedge_{j \neq r} d \left(\frac{t\partial_{y^j} f}{\Lambda} - \eta_j \right).$$

For every t in \mathbb{R}^* and every f in \mathcal{H}_Λ , this defines an element in $\mathcal{D}'_{n+1}(\mathring{T}^*U)$. The main contribution in this integral will come from the terms involving derivatives of f – this is exactly the content of lemma 4.8 above. Thus, one can set

$$\mathcal{I}_{\Lambda,0}^{r,s}(f, t, y, dy, \eta) := \delta_0^{n+1} \left(f, \frac{td_y f}{\Lambda} - \eta \right) \bigwedge_{j \neq r} d \left(\frac{\partial_{y^j} f}{\Lambda^2} \right),$$

and then one finds

$$(58) \quad \mathcal{I}_\Lambda(f, \omega) = \Lambda^n \sum_{r,s=1}^n (-1)^r \int_{\mathring{T}^*U \times \mathbb{R}^*} t^{n-3} \eta_r \eta_s dy^s \wedge \mathcal{I}_{\Lambda,0}^{r,s}(f, t, y, dy, \eta) \wedge \omega(y, dy, \eta, d\eta) \wedge dt + \mathcal{O}(\Lambda^{n-1}).$$

In order to get the conclusion, everything boils down to computing the expectation of $\mathcal{I}_{\Lambda,0}^{r,s}(f, t, y, dy, \eta)$ with respect to $d\mu_\Lambda$ for a given (y, η, t) in $\mathring{T}^*U \times \mathbb{R}^*$. In order to alleviate notations, we will just write $\mathcal{I}_{\Lambda,0}^{r,s}(f)$ and re-establish the dependence in (y, η, t) after. Using

the Berezin Fourier transform exactly as in Proposition 5.1, one can write the following “oscillatory” integral:

(59)

$$\mathcal{I}_{\Lambda,0}^{r,s}(f) = \frac{(-1)^{\frac{n-1}{2}}}{(2\pi)^{n-1}} \int_{\mathbb{R}^{(n+1|n-1)}} \exp \left(-2i\pi \left(p_0 f + p \cdot \left(\frac{td_y f}{\Lambda} - \eta \right) + \sum_{j \neq r,k} \frac{\partial_{y^j y^k}^2 f}{\Lambda^2} \Pi_j dy^k \right) \right) dp_0 dp d\Pi.$$

Remark 5.4. The fact that $\mathcal{I}_{\Lambda,0}^{r,s}(f)$ has a simple representation as an oscillatory integral is the *central observation* that makes our formal calculation much simpler than the detailed combinatorial proof we gave before. Note that the element $\Pi_j dy^k$ is even since it is the product of two odd elements and thus, we are in a situation where we can apply lemma 5.2. As was already mentioned, we do not pay too much attention to the inversion of integrals and to the size of the remainders, and we mostly focus on the conceptual aspects that make our proof works.

Recall that we aim at computing the expectation of $\mathcal{I}_{\Lambda,0}^{r,s}(f)$ with respect to the Gaussian measure $d\mu_\Lambda$. For that purpose, we fix (τ_0, τ, Π) in $\mathbb{R}^{(n+1|n-1)}$, and we use lemma 5.2 to write

$$\begin{aligned} & \int_{\mathcal{H}_\Lambda} \exp \left(-2i\pi \left(\tau_0 f + \tau \cdot \left(\frac{td_y f}{\Lambda} \right) + \sum_{j \neq r,k} \frac{\partial_{y^j y^k}^2 f}{\Lambda^2} \Pi_j dy^k \right) \right) d\mu_\Lambda(f) \\ (60) \quad & = \exp \left(-\frac{2\pi^2}{\text{Vol}_g(M)} \langle (\tau_0, t\tau, (\Pi_j dy^k)_{j \neq r,k}), A_0(\tau_0, t\tau, (\Pi_j dy^k)_{j \neq r,k}) \rangle \right) (1 + \mathcal{O}(\Lambda^{-1})), \end{aligned}$$

where the covariance matrix A_0 was defined in paragraph 4.4.

We can now expand

$$\begin{aligned} \langle (\tau_0, t\tau, (\Pi_j dy^k)_{j \neq r,k}), A_0(\tau_0, t\tau, (\Pi_j dy^k)_{j \neq r,k}) \rangle & = \frac{1}{n+2} t^2 |\tau|^2 + \tau_0^2 - \frac{2}{n+2} \tau_0 \sum_{j \neq r} \Pi_j dy^j \\ & + \frac{2}{(n+2)(n+4)} \sum_{j,k \neq r} (\Pi_j dy^k \Pi_k dy^j + \Pi_j dy^j \Pi_k dy^k). \end{aligned}$$

Thanks to the anti-commutation rules for odd variables, one finds that the last term in this sum is in fact equal to 0. After integrating over the even variables (τ_0, τ) , one finds that the following equalities hold:

$$(61) \quad \int_{\mathbb{R}^{(n|0)}} e^{2i\pi\tau \cdot \eta} \exp \left(-\frac{2\pi^2 t^2 |\tau|^2}{(n+2)\text{Vol}_g(M)} \right) d\tau = \left(\frac{\text{Vol}_g(M)(n+2)}{2\pi t^2} \right)^{\frac{n}{2}} e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2}{2t^2}},$$

and

$$\begin{aligned} (62) \quad & \int_{\mathbb{R}^{(1|0)}} \exp \left(-\frac{2\pi^2}{\text{Vol}_g(M)} \left(\tau_0^2 - \frac{2}{n+2} \tau_0 \sum_{j \neq r} \Pi_j dy^j \right) \right) d\tau_0 \\ & = \left(\frac{\text{Vol}_g(M)}{2\pi} \right)^{\frac{1}{2}} \exp \left(\frac{2\pi^2}{\text{Vol}_g(M)(n+2)^2} \left(\sum_{j \neq r} \Pi_j dy^j \right)^2 \right). \end{aligned}$$

Remark 5.5. Observe that the last equality follows one more time from lemma 5.2.

It now remains to integrate (62) with respect to the odd variables $(\Pi_j)_{j \neq r}$. Regarding the definition of the Berezin integral, we have to compute that

$$\frac{\partial}{\partial \Pi_1} \cdots \frac{\hat{\partial}}{\partial \Pi_r} \cdots \frac{\partial}{\partial \Pi_n} \exp \left(\frac{2\pi^2}{\text{Vol}_g(M)(n+2)^2} \left(\sum_{j \neq r} \Pi_j dy^j \right)^2 \right).$$

One can verify that if n is even, this quantity vanishes, and then the total expectation vanishes. In the case where n is odd, one finds from the multinomial Newton formula that

$$\begin{aligned} \frac{\partial}{\partial \Pi_1} \cdots \frac{\hat{\partial}}{\partial \Pi_r} \cdots \frac{\partial}{\partial \Pi_n} \exp \left(\frac{2\pi^2}{\text{Vol}_g(M)(n+2)^2} \left(\sum_{j \neq r} \Pi_j dy^j \right)^2 \right) \\ = \frac{(n-1)!}{\left(\frac{n-1}{2}\right)!} \left(\frac{2\pi^2}{\text{Vol}_g(M)(n+2)^2} \right)^{\frac{n-1}{2}} dy^1 \wedge \cdots \hat{d}y^r \wedge \cdots dy^n. \end{aligned}$$

Combining this to (62), we obtain

$$\begin{aligned} (63) \quad \int_{\mathbb{R}^{(|n-1|)}} \exp \left(-\frac{2\pi^2}{\text{Vol}_g(M)} \left(\tau_0^2 - \frac{2}{n+2} \tau_0 \sum_{j \neq p, q} \Pi_j dy^j \right) \right) d\tau_0 [d\Pi] \\ = \left(\frac{\text{Vol}_g(M)}{2\pi} \right)^{\frac{1}{2}} \frac{(n-1)!}{\left(\frac{n-1}{2}\right)!} \left(\frac{2\pi^2}{\text{Vol}_g(M)(n+2)^2} \right)^{\frac{n-1}{2}} dy^1 \wedge \cdots \hat{d}y^r \wedge \cdots dy^n. \end{aligned}$$

From (61) and (63), we deduce that

$$\begin{aligned} \int_{\mathcal{H}_\Lambda} \mathcal{I}_{\Lambda,0}^{r,s}(f, t, y, dy, \eta) d\mu_\Lambda(f) \\ = \frac{(-1)^{\frac{n-1}{2}} (n-1)!}{(2\pi)^{n-1}} \frac{\pi^{\frac{n-3}{2}}}{\left(\frac{n-1}{2}\right)!} \frac{1}{2(n+2)^{\frac{n}{2}}} (n+2) \text{Vol}_g(M) e^{-\frac{(n+2)\text{Vol}_g(M)|\eta|^2}{2t^2}} dy^1 \wedge \cdots \hat{d}y^r \wedge \cdots dy^n (1 + \mathcal{O}(\Lambda^{-1})). \end{aligned}$$

We now use equality (58). Summing over r and s and integrating over t , one finally gets that, if n is odd,

$$\int_{\mathcal{H}_\Lambda} \mathcal{I}_\Lambda(f, \omega) d\mu_\Lambda(f) = \frac{2(-1)^{\frac{n+1}{2}}}{\pi \text{Vol}(\mathbb{S}^{n-1})} \left(\frac{\Lambda}{\sqrt{n+2}} \right)^n \int_{\dot{T}^*U} dy^1 \wedge \cdots \hat{d}y^r \wedge \cdots \omega(y, dy, \eta, d\eta) + \mathcal{O}(\Lambda^{n-1}).$$

APPENDIX A. A FUNCTIONAL ANALYTIC PROOF OF WHITNEY EMBEDDING THEOREM

In this short appendix, we would like to prove briefly the approximation property used in the proof of Lemma 4.12, and to illustrate it by giving a simple proof of the Whitney embedding Theorem. In fact, on a smooth compact manifold M , if we have enough smooth functions e_1, \dots, e_N so that their linear combinations approximates any smooth function with enough accuracy, then we will prove that we can embed the manifold M in \mathbb{R}^N for N sufficiently large by the simple map :

$$x \in M \mapsto (e_1(x), \dots, e_N(x)) \in \mathbb{R}^N.$$

A.1. Approximation of smooth functions. We denote by $H^s(M)$ the Sobolev space of functions or distributions of order s i.e. the set of all distributions t such that $\Delta_g^{\frac{s}{2}}t$ belongs to $L^2(M)$ with norm $\|\cdot\|_{H^s(M)} = \|(1 - \Delta_g)^{\frac{s}{2}}\cdot\|_{L^2(M)}$. Note by the spectral Theorem that

$$\|(1 - \Delta_g)^{\frac{s}{2}}t\|_{L^2(M)} = \left(\sum_{i=1}^{+\infty} (1 + \lambda_i^2)^{\frac{s}{2}} |\langle t, e_i \rangle_{L^2}|^2 \right)^{\frac{1}{2}},$$

where we use the conventions of the introduction. First, we show that eigenfunctions of the Laplace operator span a vector space which is everywhere dense in $C^\infty(M)$. This property was used to prove lemma 4.12.

Lemma A.1. *Let (M, g) be a smooth compact boundaryless Riemannian manifold. Then, finite combination of eigenfunctions $(e_i)_i$ of the Laplace-Beltrami operator Δ_g spans a dense subspace of $C^\infty(M)$. More precisely, for every C^k norm $\|\cdot\|_{C^k}$:*

$$\|f - \sum_{i=1}^N \langle f, e_i \rangle_{L^2} e_i\|_{C^k} \xrightarrow{N \rightarrow \infty} 0.$$

Proof. Eigenfunctions $(e_i)_i$ of the Laplace operator are smooth by elliptic regularity. Thus, any finite combination of them belongs to $C^\infty(M)$. By the Sobolev embedding Theorem, it suffices to show that for any $s > 0$,

$$\|f - \sum_{i=1}^N \langle f, e_i \rangle_{L^2} e_i\|_{H^s} \xrightarrow{N \rightarrow \infty} 0.$$

For that purpose, we write

$$\|f - \sum_{i=1}^N \langle f, e_i \rangle_{L^2} e_i\|_{H^s}^2 = \sum_{i=N+1}^{\infty} (1 + \lambda_i^2)^{\frac{s}{2}} |\langle f, e_i \rangle_{L^2}|^2,$$

which tends to 0 as f is smooth. □

A.2. Whitney embedding Theorem. We will now show how this approximation property can be used to prove the following version of Whitney embedding Theorem:

Theorem A.2. *Let (M, g) be a smooth compact Riemannian manifold without boundary. Let $(e_i)_i$ be the sequence of eigenfunctions of the Laplace-Beltrami operator as defined in the introduction. Then, there exists N_0 such that for any $N \geq N_0$, the map*

$$(64) \quad i_N : x \in M \mapsto (e_1(x), \dots, e_N(x)) \in \mathbb{R}^N$$

is an embedding.

Remark A.3. The above Theorem holds true for any sequence $(e_i)_i$ of smooth functions which spans a dense subspace of $C^\infty(M)$ and so that for any smooth function $f \in C^\infty(M)$, $f = \sum_0^\infty f_i e_i$ where the sequence $(f_i)_i$ is **uniquely** defined. In particular, it means that the $(e_i)_i$ are linearly independent.

Proof. We first prove that i_N is an immersion then we show it is injective.

Fix some N . To show that i_N is an immersion, we proceed as in the proof of lemma 4.12. It suffices to show that for all $x \in M$, $di_N(x)$ has maximal rank in other words

$$x \mapsto (de_1, \dots, de_N)$$

has rank $\dim M = n$. Equivalently, it means that near every $x_0 \in M$, there is a neighborhood U , n sequences $(a_i^1)_{1 \leq i \leq N}, \dots, (a_i^n)_{1 \leq i \leq N}$ such that for all $x \in U$, the covectors

$$\sum_{i=1}^N a_i^1 de_i(x), \dots, \sum_{i=1}^N a_i^n de_i(x)$$

are linearly independent. In other words, the n -form

$$\left(\sum_{i=1}^N a_i^1 de_i \right) \wedge \dots \wedge \left(\sum_{i=1}^N a_i^n de_i \right)$$

never vanishes on U . But this is easy to achieve since for every x_0 , we can always find some sufficiently small neighborhood U of x_0 and functions f_1, \dots, f_n so that $df_1 \wedge \dots \wedge df_n(x) \neq 0, \forall x \in U$. And since the family $(e_i)_i$ spans a dense subspace of $C^\infty(M)$, we can always find N_0 large enough so that the vector space spanned by $(e_i)_{i \leq N_0}$ is very close to f_1, \dots, f_n in the C^1 topology.

The last part consists in proving that the above map is injective for N large enough. By the previous part, we can fix N_0 large enough so that i_{N_0} is an immersion. Since an immersion is locally injective and M is compact, there is some $\varepsilon > 0$, so that for all $x \in M$, consider the ball $B(x, \varepsilon)$ centered around x of radius ε , the map $i_{N_0} : B(x, \varepsilon) \mapsto \mathbb{R}^{N_0}$ is injective. In other words if $i_{N_0}(x) = i_{N_0}(y)$ for some $(x, y) \in M^2$ then $x = y$ or $d(x, y) > \varepsilon$. We denote by Δ_M the diagonal in $M \times M$. Consider the map $i_{N_0} \times i_{N_0} : (M \times M) \setminus \Delta_M \mapsto \mathbb{R}^{N_0} \times \mathbb{R}^{N_0}$ which reaches the diagonal $\Delta \subset \mathbb{R}^{N_0} \times \mathbb{R}^{N_0}$ exactly at the pair of points (x, y) where i_{N_0} fails to be injective. First note that $(i_{N_0} \times i_{N_0})^{-1}(\Delta) = \{(x, y) \text{ such that } i_{N_0}(x) = i_{N_0}(y), d(x, y) \geq \varepsilon\}$ is a closed subset of $M \times M \setminus \Delta_M$. Since $M \times M \setminus \Delta_M$ is precompact, the closed subset $(i_{N_0} \times i_{N_0})^{-1}(\Delta)$ is compact. Note by definition of i_N that for all N , $(i_{N+1} \times i_{N+1})^{-1}(\Delta) \subset (i_N \times i_N)^{-1}(\Delta)$. Hence consider the sequence $((i_N \times i_N)^{-1}(\Delta))_N$ for $N \geq N_0$, it is a decreasing sequence of compact subsets in $M \times M \setminus \{(x, y) \in M^2 \text{ s.t. } d(x, y) < \varepsilon\}$. By contradiction, assume that for all $N \geq N_0$, the map i_N fails to be injective, then it means that $(i_N \times i_N)^{-1}(\Delta)$ is non empty for all N , then for all N there is a pair $(x_N, y_N) \in M \times M \setminus \{(x, y) \in M^2 \text{ s.t. } d(x, y) < \varepsilon\}$ such that $i_N(x_N) = i_N(y_N)$ for all N . By compactness of $M \times M \setminus \{(x, y) \in M^2 \text{ s.t. } d(x, y) < \varepsilon\}$, we may assume that the sequence $(x_N, y_N)_N$ converges to some element (x, y) in $M \times M \setminus \{(x, y) \in M^2 \text{ s.t. } d(x, y) < \varepsilon\}$. It follows by definition of i_N that $e_i(x_k) = e_i(y_k)$ for all $k \geq i$ and by continuity of the the functions $(e_i)_i$ and the fact that $x_k \rightarrow x, y_k \rightarrow y$, we also have that $e_i(x) = e_i(y)$ for all i .

On the other hand, we know that there is some smooth function f such that $f(x) \neq f(y)$. And since $f = \sum_{i=1}^{+\infty} \langle f, e_i \rangle e_i$, this implies that $f(x) = \sum_{i=1}^{+\infty} \langle f, e_i \rangle e_i(x) = \sum_{i=1}^{+\infty} \langle f, e_i \rangle e_i(y) = f(y)$ which contradicts the fact that $f(x) \neq f(y)$. \square

REFERENCES

- [1] S. Alesker, *Theory of valuations on manifolds II*, Adv. Math. 207 (2006), 420–454.
- [2] S. Alesker, *Theory of valuations on manifolds: a survey*, Geom. Funct. Anal. 17 (2007), no. 4, 1321–1341
- [3] P. Bérard, *Volume des ensembles nodaux des fonctions propres du laplacien*, Bony–Sjöstrand–Meyer seminar, 1984–1985 (Palaiseau), École polytechnique, 1985, Exp. No.XIV, pp. 1–9.
- [4] F.A. Berezin, *Introduction to Superanalysis*, Math. Phys. and Appl. Math., 9 D. Reidel Publishing Co., Dordrecht-Boston (1987).
- [5] A. Bernig, *The normal cycle of a compact definable set*, Israel Journal of Mathematics, 159(1) (2007), 373–411.
- [6] M. Berry, *Regular and irregular semiclassical wavefunctions*, J. Phys. A 10 (1977), no. 12, 2083–2091.
- [7] M. Berry, *Statistics of nodal lines and points in chaotic quantum billiards: perimeter corrections, fluctuations, curvature*, J. Phys. A 35 (2002), 3025–3038.
- [8] X. Bin, *Derivatives of the spectral function and Sobolev norms of eigenfunctions on a closed Riemannian manifold*, Ann. Global. Anal. and Geom. **26**, 231–252 (2004)
- [9] P. Bleher, B. Shiffman and S. Zelditch, *Universality and scaling of correlations between zeros on complex manifolds*. Invent. Math. 142 (2000), no. 2, 351–395.
- [10] P. Bleher, B. Shiffman and S. Zelditch, *Universality and scaling of zeros on symplectic manifolds*. Random matrix models and their applications, 31–69, Math. Sci. Res. Inst. Publ., 40, Cambridge Univ. Press, Cambridge, 2001.
- [11] G. Blum, S. Gnuzmann and U. Smilansky, *Nodal domain statistics—a criterion for quantum chaos*, Phys. Rev. Lett. 88 114101 (2002)
- [12] E. Bogomolny, R. Dubertrand and C. Schmit, *SLE description of the nodal lines of random wavefunctions*. J. Phys. A 40 (2007), no. 3, 381–395
- [13] M. Bóna, *A walk through combinatorics*, 2nd Edition, World Scientific Publishing, (2006)
- [14] C. Brouder, V. Dang, F. Helein, *A smooth introduction to the wavefront set*, to appear in Journal of Math. Physics, arXiv:1404.1778 (2014)
- [15] C. Brouder, V. Dang, F. Helein, *Boundedness and continuity of the fundamental operations on distributions having a specified wave front set. (with a counter example by Semyon Alesker)*, arXiv:1409.7662 (2014)
- [16] J. Brüning, *Über Knoten Eigenfunktionen des Laplace-Beltrami Operators*. Math. Z. **158** (1978), 15–21.
- [17] Y. Canzani and P. Sarnak, *On the topology of the zero sets of monochromatic random waves*, arXiv:1412.4437 (2014)
- [18] J. Chazarain, A. Piriou, *Introduction to the theory of linear differential equations*, Studies in mathematics and its applications, Vol. 14, North Holland Publishing (1982)
- [19] S.S. Chern, *A Simple Intrinsic Proof of the Gauss-Bonnet Formula for Closed Riemannian Manifolds*. Ann. of Math. (2) 45, (1944), 747–752.
- [20] S.S. Chern, *On the curvatura integra in a Riemannian manifold*. Ann. of Math. (2) 46, (1945), 674–684.
- [21] T. Colding and W. P. Minicozzi II, *Lower bounds for nodal sets of eigenfunctions*. Comm. Math. Phys. **306** (2011), no. 3, 777–784.
- [22] G. de Rham, *Differentiable manifolds: Forms, Currents, Harmonic Forms*, Springer (1980)
- [23] M. Disertori, *Modèles de matrices aléatoires et supersymétrie*, Habilitation à diriger des recherches, available at <http://lmrs.univ-rouen.fr/Persopage/Disertori/publications.html> (2011)
- [24] R. T. Dong, *Nodal sets of eigenfunctions on Riemann surfaces*. J. Differential Geom. **36** (1992), no. 2, 493–506.
- [25] H. Donnelly and C. Fefferman, *Nodal sets of eigenfunctions on Riemannian manifolds*. Invent. Math. **93** (1988), 161–183.

- [26] H. Donnelly and C. Fefferman, *Nodal sets for eigenfunctions of the Laplacian on surfaces*. J. Amer. Math. Soc. **3** (1990), 332–353.
- [27] E. Frenkel, A. Losev and N. Nekrasov, *Instantons beyond topological theory. I*, Journal of the Institute of Mathematics of Jussieu, 10(03) (2011), 463–565.
- [28] J. H. Fu, *Curvature measures of subanalytic sets*, American Journal of Mathematics (1994), 819–880.
- [29] D. Gayet and J.Y. Welschinger, *Expected topology of random real algebraic submanifolds*, Journal of the Institute of Mathematics of Jussieu, online (2014)
- [30] D. Gayet and J.Y. Welschinger, *Betti numbers of random nodal sets of elliptic pseudo-differential operators*, arXiv:1406.0934 (2014)
- [31] D. Gayet and J.Y. Welschinger, *Universal components of random nodal sets*, arXiv:1503.01582 (2015)
- [32] M. Giaquinta, G. Modica, J. Soucek, *Cartesian Currents in the Calculus of Variations I*, Springer-Verlag Berlin Heidelberg (1998)
- [33] V. Guillemin, S. Sternberg, *Supersymmetry and equivariant de Rham theory*, Springer-Verlag Berlin Heidelberg (1999)
- [34] Q. Han and F.H. Lin, *Nodal sets of solutions of elliptic differential equations*, book in preparation (online at <http://www.nd.edu/~qhan/nodal.pdf>).
- [35] R. Hardt and L. Simon, *Nodal sets for solutions of elliptic equations*. J. Differential Geom. 30 (1989), no. 2, 505–522
- [36] M. Hardy, *Combinatorics of partial derivatives*. Electronic Journal of Combinatorics 13 (1): R1 (2006)
- [37] H. Hezari and C. D. Sogge, *A natural lower bound for the size of nodal sets*. Anal. PDE **5** (2012), no. 5, 1133–1137.
- [38] L. Hörmander, *The spectral function of an elliptic operator*, Acta Math. 121 (1968), 193–218.
- [39] L. Hörmander, *The Analysis of Linear Partial Differential Operators I. Distribution Theory and Fourier Analysis*, second ed., Springer Verlag, Berlin, 1990
- [40] D. Jerison and G. Lebeau, *Nodal sets of sums of eigenfunctions*. Harmonic analysis and partial differential equations (Chicago, IL, 1996), 223–239, Chicago Lectures in Math., Univ. Chicago Press, Chicago, IL, 1999
- [41] M. Kac, *On the average number of real roots of a random algebraic equation*, Bull. AMS **49** (1943), 314–320
- [42] M. Kashiwara, *Index theorem for constructible sheaves*, Astérisque, 130 (1985), 193–209.
- [43] M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Springer Science
- [44] E. Kostlan, *On the distribution of roots of random polynomials*, From topology to computation: proceedings of the Smalefest (Berkeley, CA, 1990), Springer, New York (1993), 419–431
- [45] T. Letendre, *Expected volume and Euler characteristic of random submanifolds*, arXiv:1408.2107 (2014)
- [46] A. Losev and S. Slizovskiy, *New observables in topological instantonic field theories*, Journal of Geometry and Physics, 61(10) (2011), 1868–1880.
- [47] F. Nazarov and M. Sodin, *On the number of nodal domains of random spherical harmonics*, Amer. J. Math. 131 (2009), no. 5, 1337–1357
- [48] F. Nazarov and M. Sodin, *Asymptotic laws for the spatial distribution and the number of connected components of zero sets of Gaussian random functions*, preprint arXiv:1507.02017 (2015)
- [49] L. Nicolaescu, *Critical sets of random smooth functions on compact manifolds*, preprint arXiv:1101.5990, to appear in Asian J. Math. (2011)
- [50] H. Park, *Kinematic formulas for the real subspaces of complex space forms of dimension 2 and 3*, Phd Thesis (2002)
- [51] Z. Rudnick and I. Wigman, *On the volume of nodal sets for eigenfunctions of the Laplacian on the torus*, Annales Henri Poincaré **9** (2008), 109–130.
- [52] P. Sarnak and I. Wigman, *Topologies of nodal sets of random band limited functions*, arXiv:1312.7858 version 2 (2014).
- [53] L. Schwartz, *Théorie des distributions*, Hermann (1966)

- [54] B. Shiffman, S. Zelditch, *Distribution of zeros of random and quantum chaotic sections of positive line bundles*. Comm. Math. Phys. 200 (1999), no. 3, 661–683
- [55] M. Sodin, *Lectures on random nodal portraits*, Lecture notes for a mini-course given at the St. Petersburg Summer School in Probability and Statistical Physics (June, 2012), available at: <http://www.math.tau.ac.il/~sodin/SPB-Lecture-Notes.pdf> (2012)
- [56] C. D. Sogge and S. Zelditch, *Lower bounds on the Hausdorff measure of nodal sets II*. Math. Res. Lett. **19** (2012), no. 6, 1361–1364.
- [57] T. Tao, *Supercommutative gaussian integration, and the gaussian unitary ensemble*, available at <https://terrytao.wordpress.com/tag/berezin-integration/> (2013)
- [58] I. Wigman, *Fluctuations of the nodal length of random spherical harmonics*, Comm. Math. Phys. **298** (2010)
- [59] S.-T. Yau, *Seminar on Differential Geometry*, Ann. of Math. Stud., **102**, Princeton Univ. Press, Princeton, N.J., 1982.
- [60] S. Zelditch, *Real and complex zeros of Riemannian random waves*, Spectral analysis in geometry and number theory, Contemp. Math., vol. 484, Amer. Math. Soc., Providence, RI, 2009, pp. 321–342.
- [61] S. Zelditch, *Eigenfunctions and nodal sets*. Geometry and topology, 237–308, Surv. Differ. Geom., **18**, Int. Press, Somerville, MA, 2013.

LABORATOIRE PAUL PAINLEVÉ (U.M.R. CNRS 8524), U.F.R. DE MATHÉMATIQUES, UNIVERSITÉ
LILLE 1, 59655 VILLENEUVE D’ASCQ CEDEX, FRANCE

E-mail address: `nguyen-viet.dang@math.univ-lille1.fr`

E-mail address: `gabriel.riviere@math.univ-lille1.fr`