

### Image analysis of cheese microstructure and diffusion properties of solutes: An important link to understand cheese ripening mechanisms

Juliana Valle Costa Silva, Chantal Cauty, David Legland, Sylvie Lortal,

Juliane Floury

#### ► To cite this version:

Juliana Valle Costa Silva, Chantal Cauty, David Legland, Sylvie Lortal, Juliane Floury. Image analysis of cheese microstructure and diffusion properties of solutes: An important link to understand cheese ripening mechanisms. DREAM Project International Conference, Institut National de Recherche Agronomique (INRA). UAR Département Caractérisation et Elaboration des Produits Issus de l'Agriculture (1008)., Jun 2013, Nantes, France. hal-01189705

#### HAL Id: hal-01189705 https://hal.science/hal-01189705v1

Submitted on 3 Jun 2020  $\,$ 


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







"Image analysis of cheese microstructure and diffusion properties of solutes: An important link to understand cheese ripening mechanisms"




#### Juliana VALLE COSTA SILVA1, Chantal CAUTY1, David LEGLAND2, Sylvie LORTAL1, Juliane FLOURY1,3

1INRA, Science and Technology of Milk and Eggs, Rennes, France 2INRA, UMR782, Food Process Engineering and Microbiology, Thiverval-Grignon, France 3Agrocampus-Ouest, UMR1253, Science and Technology of Milk and Eggs, Rennes, France


### Context

### Cheese

- Kipening cheese making technology
- Complex and specific of each variety of cheese



### **Objective**



To enable a better understanding of the **ripening mechanisms**...

### **Strategy**




How can you understand and control the **ripening mechanisms** of...


Different cheeses...

Of different microstructures..




With different and complex steps of production and developme





#### Mothode

#### **UF-model cheese manufacture**




#### Mothode

Diffusion of solutes through the UF-model cheese and in the ultrafiltrate using the FRAP technique Prebleach Bleach Recovery Fluorescent molecule Bleached molecule **Fluorescent solutes:** Flash F FITC-dextrans (4 kDa to 2 Mba) Intensité émise F...

Flexible and neutral solutes

**FITC-labeled dairy proteins** (α-LA, β-LG and BSA)

Rigid and negatively charged solutes



6

Temps Typical recovery curve of fluorescence in function of time

F<sub>0</sub>

0

t = 0

temps de

retour, τ

Deff: effectif diffusion coefficient in the UF-model cheese (µm2/s)

Dag: diffusion coefficient in the aqueous phase (ultrafiltrate)

*wM:* profile width in the radial direction (*m*)

Design of the state of the stat

Fraction


mobile

#### Mothode

#### Study of the microstructure of the UF-model cheese using TEM

#### Freeze-substitution and low-temperature embedding protocol:

Protects the structure of dairy constituents



#### High-pressure freezing:

- Samples are frozen in liquid nitrogen (-196°C) at high-pressure (2100 bars)

- Advantage: at 2 100 bars the melting point of water decreases because the water becomes 1 500 times more viscous. The increase in viscosity reduces the ice crystal growth rate.

- The sample is immersed in a solvent
- Substitution of ice by the solvent

#### The solvent is replaced with a low-temperature resin:


- Ethanol 3:1 resin
- Ethanol 1:1 resin
- Ethanol 1:3 resin
- 100% resin

Thin sections **(90nm thick)** of embedded material are cut with a diamond knife using a Reichert ultramicrotome

#### Mothode

#### Quantitative analysis of the microstructure of the UF-Model Cheese

8



Shortest path containing a point x in the matrix, connecting two opposite edges of the image (Delarue et al. , 2003).


#### Dicqueeion

Influence of different physicochemical characteristics of solutes on diffusion in the UF-model cheese

Influence of the solute size...

FITC-dextrans (4 kDa to 2 MD

- Flexible and neutral solutes



**Figure 1.** Logarithmic plots of experimental diffusion coefficients of FITC-dextrans in the UF-model cheese ( $\Box$ ) and in the ultrafiltrate ( $\Box$ ) as a function of their molecular weight.

The **larger FITC-dextrans** were **no more hindered** by the structure of the UF-model cheese when compared to the smaller ones.

The mobility of FITC-dextrans in the UFmodel cheese was reduced by a **constant obstruction factor** (k = 0.42).

The **pore sizes** of this model cheese were believed to be sufficiently large when compared to the **size of the solute molecules**.


#### Dicqueeion

Influence of different physicochemical characteristics of solutes on diffusion in the UF-model cheese

Influence of the shape and charge of solutes...

**FITC-labeled dairy proteins** ( $\alpha$ -LA,  $\beta$ -LG and BSA)

- Rigid and negatively charged solutes



**Figure 2.** Logarithmic plots of experimental diffusion coefficients of FITC-dextrans ( $\Box$ ) and FITC-dairy proteins ( $\Box$ ) in the UF-model cheese as a function of their hydrodynamic radius.

**Rigid and negatively charged solutes** were **hindered more** than the **flexible and neutral solutes** in the UF-model cheese

The existence of steric and electrostatic interactions between the **protein matrix** of the UF-model cheese and the FITC-dairy proteins is suggested.


The diffusion through the protein aggregates could play an important role in the overall diffusion of the FITC-dairy proteins, retarding their movement when they diffuse in the UF-model cheese.




#### Dicouccion

Influence of the microstructure of the UF-model cheese on diffusion of solutes

Image Analysis of the UF-model cheese



TEM micrograph of the UF-model cheese



Segmented (binary) image of the UF-model cheese

### **Quantification of relevant features...**


Diffusion properties

Structure/ composition


#### Dicqueeion

Influence of the microstructure of the UF-model cheese on diffusion of solutes

Image Analysis of the UF-model cheese



 $\epsilon = 0.638 \pm 0.007$ 





Tortuosity distribution in the UF-model cheese. Blue zones: low tortuosity. Yellow/orange zones: high tortuosity

 $\tau = 1.085 \pm 0.008$ 

### Under way...

To establish the relationship between these **microstructural parameters** and **diffusion properties** by comparing results obtained in **model matrices** with **different microstructures**.

12

D


d1(x)

d,(x)

# Conclusions and Perspectives

- The physicochemical characteristics of solutes influence their diffusion in the UF-model cheese
- The diffusion of other solutes (flexible and negatively charged, rigid or flexible positively charged) may be studied to elucidate the solute-matrix interactions
- Studies of the diffusion and structure in other model cheese matrices are required to the development of predictive models of diffusion of solutes
- A better understanding of the influence of the matrix microstructure on the diffusion of solutes should allow a more generic view of cheese ripening kinetics for future innovations









# **Acknowledgements**

### My research team (MICRO): Juliane Floury

- Sylvie Lortal ٠
- Anne-Thierry •
- Sophie Jeanson •
- Valérie Gagnaire •
- **Chantal Cauty** •







14







**Platform MRic** 

٠

٠



### **David Legland**











# Thank you for your attention!

Juliana.ValleCostaSilva@rennes.inra.fr