

# Assessment on nitrogen flows in livestock farming system in France. Main issues and options

Jean-Louis Peyraud, Pierre Cellier, Catherine Donnars, Olivier Rechauchère, Frans Aarts, Fabrice Béline, Christian C. Bockstaller, Luc Delaby, Jean-Yves Dourmad, Pierre Dupraz, et al.

#### ▶ To cite this version:

Jean-Louis Peyraud, Pierre Cellier, Catherine Donnars, Olivier Rechauchère, Frans Aarts, et al.. Assessment on nitrogen flows in livestock farming system in France. Main issues and options. International Conference Planet under pressure: New knowledge, towards solutions, 2012, London, United Kingdom. 1 p. hal-01190254

HAL Id: hal-01190254

https://hal.science/hal-01190254

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Assessment on nitrogen flows in livestock farming system in France. Main issues and options.

Peyraud J.L.a, Cellier P.a, Donnars C.a, Rechauchère O.a, Aarts F.b, Béline F.c, Bockstaller C.a, Delaby L.a, Dourmad J.-Y.a, Dupraz P.a, Durand P.a, Faverdin P.a, Fiorelli J.-L.a, Gaigné C.a, Kuikman P.b, Langlais A.d, Le Goffe P.e, Lescoat P.a, Nicourt C.a, Rochette P.f, Vertes F.a, Veysset P.a, Bourblanc M.g, Morvan T.a, Parnaudeau V.a, Le Perchec S.a, Girard A.a, Guillaume F.a

<sup>a</sup> INRA, France, <sup>b</sup> WUR, Wageningen, The Netherlands, <sup>c</sup> Irstea, Rennes, France, <sup>d</sup> CNRS, Rennes, France, <sup>e</sup> AgroCampus Ouest, Rennes, France, <sup>f</sup> Agriculture and Agri-Food Canada, <sup>g</sup> Cirad, France

- 1. Background: Livestock consumes large amounts of agricultural products and is a major source of reactive nitrogen. Moreover different economic and societal factors contributed to a concentration of livestock in the West of France, leading to an increased pressure on the environment. The surplus of N causes pollution especially in the atmosphere, water ressources and coastal ecosystems.
- 2. Objectives: A collective expert assessment (Esco) was conducted whose objectives were to
  - make a synthesis of updated knowledge on N flows in livestock farming activities on, from the animal to the regional scale with a specific focus on the farming system. All forms of nitrogen (nitrate, ammonia, nitrous oxide, others) were considered as well as the link with impact.
  - make a comparison between different livestock systems.

This collective expert

by the French Ministries


of Ecology and Agriculture

- identify several possible actions in livestock breeding systems, e.g. improved techniques and management, change in the system, territorial and economic incentives.

#### 3. Methods

The cross-disciplinary assessment was based on an analysis of the literature by a panel of 22 scientists from a wide range of disciplines: animal sciences, agronomy, social sciences, biogeochemistry, etc.

- 1332 references were analyzed from 272 scientific journals; 82% after 1998.
- 3/4 of the references were on biophysical processes (of which half on N fluxes in livestock farming systems and approx. 20% on indicators), and ¼ in social sciences.
- The references were also analyzed according to the journal which they were originating from, the origin of the authors and the main topics of the assessment.



article


# 4 Results

This section makes a synthesis of the main findings.

### A framework to analyse livestock farming system

Choices in N management are taken by farmers. But they involve:

- (i) Herd unit, manure management, soils & growing crops.
- (ii) Groups of farms or local organisations where deals may open new opportunity for N flows control.
- (iii) Environmental issues linked to farming activity (iv) Stakeholders with their policies and social debate.



> Livestock farming provides major part of N flows.

and  $N_2O$  (>70%) and accounts for more than 50% of nitrate.

> N efficiency per animal is low but it increases at the

Efficiency

Medium

(40%)

Low input and organic farming show the highest N efficiency.

Drivers

Farming system:

rotation

farm scale thanks to internal recycling

results from complex interactions and thus

It decreases when N input increases.

an improvement at one stage can be cancelled

by a bad management at a previous or next stage.

Grazing reduces N emissions because there is no

storage and grass absorbs N directlyfrom excreta.

meat. Limited

made with feed

recommendations.

Margins are higher

for farms. The N balance

progress can still be

nitrate. However, livestock farming has a prominent place in the


#### > Specialisation and concentration amplify problems.

From the 50<sup>th</sup>, the agricultural system moved from mixed farming to specialized systems and increased concentration in animal husbandry due to economic rationality and agroindustry pressure. West of France presently concentrates half of hogs and poultry and 20% of cattle on only 6% of farmland. Consequently, livestock farming has no longer sufficient land for providing feed and spreading manure. Scientific literature emphasizes that this technical and economic logic locks the system because firms and farmers are strictly connected in their activities.

#### > A complicated legal frame did not allow to achieve the objectives.

The French N regulations framework is characterized by a overabundance of plans that make it difficult to find a compromise between agreeing with the European directives, maintaining the production and reaching environmental aims. The problem of pollutions progressively

entered into the public debate in the 90<sup>th</sup> thereby questioning the livestock farming practices and making the farmers more difficult in some cases.



# been considered from the point of view of water contamination by whole N cycle: it is the main provider of atmospheric ammonia (90%) From Citepa (2011) Efficiency **Drivers** Knowledge

A minority of N input is exported in products: eggs, milk or Production yields (10-40%)Feeding practices Knowledge Farm High **Production intensity Animal** Herd grassland, legumes, crop Soils and Manure crops Knowledge **Drivers** Losses Globally, N efficiency in livestock farming is around 40-50%. Very variable Housing > grazing Housing> spreading > Large uncertainty (20-80%) NH<sub>3</sub>: 25-55% storage on emissions

Manure treatments

#### > N farm budgets are convenient methods for improving nitrogen management. The budget has to take into acount all the flows related to the farm in order to screen every source of waste and identify a range of progress.

10%


Budget (kg N/ha) **Average N flows** Farming system system **Dairy farm** 238 148 Input + crops **177** Output 80 ha UAA 82 LU Crops = 25 haBalance 61 West of France Variations of N into the soil ↓ Dissolved organic N 48 and NO<sub>3</sub>-

Organic

Pig farm

+ crops

dairy farm



+ crops  160 ha UAA  107 LU  Crops = 65 ha	Output	21	93
	Balance	46	41
East of France			


Input

134

199

**67** 

952




it	84 ha UAA 400 sows Breeding and fattening Slurry treatment (60%) West of France	Output	730	140
		Balance	222	59

Input

## > The impact of nitrogen losses also depends on landscape sensitivity.

That is the capacity of the agroecosystem and the environment (surrounding ecosystems, soils) to use or to transform the excess N.

Animal density, fraction of farmlands and grassland in the area influence nitrate and ammonia losses. This shows the need to better account for the territorial vulnerability when setting up environmental policies. Then a critical load of N might be allowed for a given area ,whose amount depends on the local environmental sensitivity.



### A range of options for reducing the impact of excess nitrogen in livestock farming has been defined:

- Applying more extensively measures for decreasing ammonia emissions, which has not been considered very thoroughly in France up to now.
- \* Manure treatment for exporting them to remote places with fewer nitrogen issues.
- Improving nitrogen balance on farm instead only in fields, considering thus all the flows and losses of nitrogen in the system. Yet it is difficult to implement due to the lack of data to estimate llocal flows.
- Relocating a fraction of livestock farms, or mixing crop and livestock systems could be envisaged on a case by case basis, between regions or between adjacent areas with different nitrogen loads. Inside critical areas, this might require reorganizing the farming system dramatically. This can only be done with cooperation between local partners.
- \* Environmental policies face with the problem of the mostly diffuse sources of nitrogen sources and with the difficulty to give a precise value to most N-related damages. Penalties based on estimating nitrogen surplus seems to be the best compromise.

These options must consider interactions with other environmental issues (energy budget, phosphorus, pollution swapping) and economic issues (farmers income, global production).

# > Territorial options and deals between areas

