

Trees feel mechanical strain: from genes expression to cambium activity modulation

Eric Badel, Ludovic Martin, Nathalie Leblanc-Fournier, Jean-Louis J.-L. Julien, Catherine Coutand, Mélanie M. Decourteix, Catherine Lenne, Bruno Moulia

▶ To cite this version:

Eric Badel, Ludovic Martin, Nathalie Leblanc-Fournier, Jean-Louis J.-L. Julien, Catherine Coutand, et al.. Trees feel mechanical strain: from genes expression to cambium activity modulation. International Symposium on Wood Structure in Plant Biology and Ecology, Apr 2013, Naples, Italy. hal-01190306

HAL Id: hal-01190306 https://hal.science/hal-01190306v1

Submitted on 6 Jun 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Trees acclimation to strains induced by wind: from genes expression to stem structure

¹L. Martin, ²E. Badel, ¹N. Leblanc-Fournier, ¹M. Decourteix, ¹C. Lenne, ²C. Coutand, ²B. Moulia and ¹J.L. Julien

1 Université BLAISE PASCAL, UMR547 PIAF, F-63177 AUBIERE 2 INRA, UMR547 PIAF, F-63100 CLERMONT-FERRAND




0.2 0.1 Daily 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 Time (days)

Response of secondary growth to repeated daily bending. Open circles represent growth response to one single bending (1B). Closed circles represent growth response to 9 successive bendings at 1-day intervals (9B-1d). Dash squares, model of additive effects (linear time integration, 9x1B). Open squares, model with a sensitivity shift after 3 daily bendings (3x1B; modation)

In nature, mechanical stimuli do not occur as a single bending. In this experiment, successive bendings were separated at day scale, mimicking the alternance between windy or quiet weather.


As soon as a second bending was applied, a diminution of growth and molecular responses to subsequent bending were observed. Our results show that plants acclimate rapidly to mechanical loadings and a desensitization period of a few days occurs after a single transitory bending. This acclimation process provides a basis for a mechanistic analysis of response sensitivity to mechanical loadings such as wind (4).




Mechanoresponsive genes expression after repeated daily

Mechanoresponsive genes expression after repeated daily bending. C : control (no load). 1B : one single bending. xB-yd : x bendings each separated by y days. *PtaZFP2* : Populus tremula*alba Zinc Finger Proteing gene, *PtaTCH4* : Populus tremula*alba Touch4 xyloglucan endotransglucosylase/hydrolyse gene. Significant differences (*P* < 0.05) of responses are indicated by different

Multiple mechanical stimuli affect anatomical pattern and secondary growth rate



Stem anatomical modifications induced by multiple bendings. (a) to (d) Control plants. (e) to (h) Plants subjected to 6 bendings, 2 successive bendings separated by 1 day, 3 days without sollicitations between each set of bendings. (a, b, e, f) Toluidine blue staining. (c, d, g, f) Blue astra-safratine staining c cambium, c to us stamma, to u, g, r) nove astra-sarratime staffing. c: cambium, cp: cortical parenchyma, fix: flexure wood, gl: G layer, ph: phloem, pi: pith, scl: sclerenchyma, x: xyelm, xf: xylem fiber, xr: xylem ray, xv: xylem vessel.



Cambial activity induced by local strains applied to the stem: experimentations and modelling. Plants were subjected to 3 successive daily bendings per week during 4 months. (a) Transversal section of a stimulated plant (h) Relative stimulated plant. (b) Relative ovalisation induced by bending. (c) Strain field for living tissues. (d) Weekly growth rate measurements (blue lines) and growth rate modeling (red lines).

 $\Delta R(i,t) = \Delta C(i,t)(1 + \alpha \, \varepsilon_{LL}(i))$

In response to bendings, cambium activity is impacted by the strain. Growth rate is identically stimulated according to the strain level both in the stretched part and in the compressed part of the stem. Wood differentiation is modulated according to the type of mechanical loading (tension and compression).

(1) Coutand C. and Moulia B. (2000). Biomechanical study of the effect of a controlled bending on tomato stem elongation: local strain sensing and spatial integration of the signal. Journal of Experimental Botany 51(352): 1825-1842. (2) Coutand C.*, Martin L.*, Leblanc-Fournier N., Decourteix M., Julien J.L. and Moulia B. (2009). Strain mechanosensing quantitatively controls diameter growth and PlaZFP2 gene expression in poplar. Plant Physiology 151: 1-10. (3) Moulia B., Der Loughian C., Bastien R., Martin L., Rodriguez M., Gourcilleau D., Barbacci A., Badel E., Franchel J., Lenne C., Roeckel-Drevet P., Allain J.M., Frachisse J.M., de Langre E., Coutand C., Leblanc-Fournier N., and Julien J.L. (2011). Integrative mechanicbiology of growth and architectural development in changing mechanical environments. In P. Wojtaszek [ed.], Mechanical Integration of Plant Cells and Plants, Signaling and Communication in Plants 9, 269-302. Springler-Vieran Ratin Heindeharra