

All cows are worth to be genotyped!

Didier Boichard, Romain Dassonneville, Sophie Mattalia-Elie, Vincent Ducrocq, Sebastien Fritz

▶ To cite this version:

Didier Boichard, Romain Dassonneville, Sophie Mattalia-Elie, Vincent Ducrocq, Sebastien Fritz. All cows are worth to be genotyped!. 64th Annual Meeting European Association for Animal Production (EAAP), Aug 2013, Nantes, France. pp.184. hal-01190501

HAL Id: hal-01190501

https://hal.science/hal-01190501

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Session 10 Theatre 2

All cows are worth to be genotyped!

D. Boichard¹, R. Dassonneville^{1,2}, S. Mattalia², V. Ducrocq¹, S. Fritz³

¹INRA, UMR1313 GABI, 78350 Jouy en Josas, France, ²Institut de l'Elevage, 149 rue de Bercy, 75595 Paris, France, ³UNCEIA, 149 rue de Bercy, 75595 Paris, France

In dairy cattle, genomic evaluation based on male reference populations has the same accuracy for males and females and a very similar accuracy for all traits. This provides a new opportunity to efficiently implement within herd selection. Much more than before, the farmer can customize his breeding goal due to the larger panel of available bulls and the accurate evaluation of females for all traits. The proportion of genotyped females is very sensitive to the evaluation cost. It is anticipated that, in European conditions, this technology can be generalized to a large proportion of the female population if the cost is below 40€. The genetic interest and the profitability of this choice are highly dependent on the selection intensity which can be applied within herd. With conventional reproduction, most females are needed for replacement and the interest is limited. With sexed semen, a technology expected to strongly develop, the upper half of the herd (based on the breeding objective of the farmer) can be oriented to produce female calves for replacement whereas the remaining part does not contribute to replacement and can be mated for other purposes, especially in crossbreeding. In practice, it is worth to start genotyping at least all young animals during their first year of life, allowing to gradually increase the proportion of genotyped animals in the herd. In addition to selection, genotyping females provides useful information for matings (today genetic defects, tomorrow optimal matings based on inbreeding minimization, QTL pyramiding, non additive effects...). Finally, it should be emphasized that these females will contribute the reference population of the future and there is a strong general interest to increase the proportion of genotyped cows and decrease the genotyping cost.

EAAP – 64th Annual Meeting, Nantes

184