



**HAL**  
open science

## The "Alaas" and the stock farming in Yakutia

Gilles Selleron, Jean-Paul Métaillé

► **To cite this version:**

Gilles Selleron, Jean-Paul Métaillé. The "Alaas" and the stock farming in Yakutia. Wetlands as archives of the cultural landscape: from research to management, Jan 2009, Gênes, Italy. 2009. hal-01190752

**HAL Id: hal-01190752**

**<https://hal.science/hal-01190752>**

Submitted on 1 Sep 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# THE "ALAAS" AND THE STOCK FARMING IN YAKUTIA

*A sub-arctic pastoral system based upon cryogenic wetlands*

Selleron G., Métailié J.-P. University of Toulouse, GEODE-CNRS, France



The agriculture of Yakutia (Sakha republic) is characterized by a breeding of cows and horses in extreme climatic conditions. The pastoral zones are located between 60°N and 65°N, in an area of Siberia with very strong continental subarctic climate. At Yakutsk (62° 05' N, 129°45'E; 103 m a.s.l.), the mean annual temperature is -10.0°C, and mean monthly temperatures in January and July are -41.2°C and 18.7°C, respectively. Mean annual precipitation is 237 mm (National Astronomical Observatory, 2001), of which 150mm in summer.

Permafrost is present in all the territory and thickness can reach 500 m. near Yakutsk ; in central Yakutia the melting of permafrost in summer can reach a depth of 2-3 meters in sunny conditions, but only 50 cm in cold places (northern expositions and under forests). The geology of central Yakutia is mainly of quaternary loess deposits covering the alluvial terraces of Lena and Aldan rivers.

Yakutsk : meteorological data

months	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Maxima (°C)	-6	-2	8	21	31	33	33	27	21	3	-4	-8	38
Mean maximal temperature (°C)	-40	-31	-14	0	12	23	25	23	12	-4	-21	-37	-6
Mean annual temperature (°C)	-41	-36	-22	6	7	15	18	15	6	-9	-29	-39	-8
Mean minimal temperature (°C)	-46	-41	-30	-14	0	8	12	9	1	-12	-33	-43	-16
Minima (°C)	-43	-44	-55	-41	-19	-7	-2	-6	-14	-41	-45	-50	-64
Mean rainfall (mm)	9	4	6	10	18	37	39	37	29	20	16	12	240


Source : MétéoFrance

The vegetation of alaaes is closely dependent of thermokarstic morphology and climatic conditions; the presence of lakes and stagnant water, and the fluctuation of the water level create concentric zones of herbaceous vegetation : in the center of the alaaes or around the lakes, temporary marshes dominate with *Carex* and *Scirpus* ; a second ring corresponds to the rich meadows which can be mown and provide the great quantity of hay necessary for the winter. In the peripheral parts, the soils are dry and saline, and the vegetation is dominated by the *Artemisia* steppe, which can be only grazed. In the sunny woods of larch (*Larix Gmelinii*) and birches, the cattle can also find some grass in the underwood and in the burned stands ; on the opposite, in the northern exposures and in the dense forests, the permafrost remains superficial and the soil is covered only by lichens.

LANDSAT-ETM 2001 ("pseudo true colors")  
 Dark blue : deep waters of the duede and lakes. Light blue: surface waters  
 Dark pink : wet pastures. Clearly pink: dry pastures and steppe.  
 Light green : marshes. Dark green: Larch woodlands



The alaaes are thermokarstic depressions distributed throughout the central Yakutia. They are formed by thermal disturbance of the ice-rich permafrost surface and an increase in depth of the active layer (annually freezingthawing layer). The alaaes, of circular forms, range in size from ca. 0.1 to 20 km in diameter, and 3 to 20 m in depth ; the first stage of alaaes process, a very active small lake depression, is locally named *duede*. The duede are very numerous in the forest areas. In the lowland of central Yakutia, approximately 20% of land area is covered with alaaes but the density is very variable.





In the valley of the Taata river, the classification of the Landsat image show the great number and extension of the alaaes, which allows a good disponibility of meadows and hay. This region has been still today one of the main centers of the Yakut stockbreeding.



Sakhas, or Yakuts, are stockbreeders of horses and cows ; they are of Turco-Mongolian origin and would have colonized Yakutia during the Middle Ages, towards XIVth-XVth centuries, by transplanting a stockbreeding adapted to less cold areas. From central Yakutia, they colonized more septentrional areas (Vilyui, Verkhoyansk), probably during 17th and 18th centuries. Traditionnaly, yakut stockbreeding is based upon stabulation of the cows during the long and very cold winter (8-9 monthes), and summer transhumance at short distance in the nearby pastures. The horses, very robust, stand outside during winter.

Untill the Revolution, the yakuts lived in family hamlets in the middle of the pastures of the clan. The summer habitat was a "tepee" of birch bark. During the soviet period, they were gathered in kholkoz and sovkhozes, with great collective cowsheds and rustic Yakut cows were replaced by imported races (Simmental, Holstein) . Agriculture and ploughing were intensified in the dry parts of the alaaes and terraces, causing the degradation of the fragile soils. From the fall of communism, the yakuts reconstituted family farms in the villages and the alaaes were allotted according to the number of family members. The interannual fluctuation of water level in the alaaes does not allow the precise delimitation of an exploitable surface.



Legend for Landsat-ETM 2001 classification:  
 Larch (green), Larch-birches (light green), Degraded forest (fires) (brown), Degraded forest (insects) (dark brown), Wet meadows (orange), Good meadows (yellow), Marshes (purple), Lakes, waters (blue), Baren soil (grey).



References :  
 Crute, Susan A. 2008. "Eating Hay": The Ecology, Economy and Culture of Vilyui Sakha Smallholders of Northeastern Siberia. Human Ecology, vol. 36-2, pp.161-174  
 Brouckov, A. ; Fukuda, M. ; Fedorov, A. ; Konstantinov, P. ; Iwahana, G. 2004. Thermokarst as a short-term permafrost disturbance, Central Yakutia. Permafrost and periglacial processes, vol. 15, n°1, pp. 81-87  
 Gavrilov, P.P. ; Efremov, P.V. 2003. Effects of cryogenic processes on Yakutian landscapes under climate warming. In : Springman & Arenson (eds.) Permafrost. Proceedings of the 8th international conference on permafrost, pp. 277-281  
 Katamura F., Fukuda M., Bosikov N.P., Desyatkin R.V. 2009. Forest fires and vegetation during the Holocene in central Yakutia, eastern Siberia, J. For. Res. 14, pp. 30-36  
 Katamura F., Fukuda M., Bosikov N.P., Desyatkin R.V., Nakamura T., Morizumi J. 2008. Thermokarst Formation and Vegetation Dynamics Inferred from a Palynological Study in Central Yakutia, Eastern Siberia, Russia. Arctic, Antarctic, and Alpine Research, Vol. 38, No. 4, pp. 561-570  
 Métailié J.-P., Selleron G., Blanc F., Galop D. 2007. Au pays des alas. In Crubezy E. (dir) Chamane. Kyys, jeune fille des glaces. Editions Erance - 119-126.  
 Mirkin, B.M. ; Gogoleva, P.A. ; Kononov, K.E. 1985. The vegetation of central yakutian alaaes. Folia geobotanica et phytotaxonomica, 20, pp.

Because of this connection between thermokarstic process and dynamics of the vegetation, it is probable that the fodder resource of the alaaes preexisted on arrival of the first Sakhas. One can make the hypothesis that the presence of this resource was one of the factors of the installation of these stockbreeders, which transported their cattle of regions distant from several thousands of kilometers. Indeed, the principal areas of historical Sakha settlement (like the valley of Taata river, 100 km in the east of the Lena) are characterized by the abundance of the alaaes and the associated pastures. However, the studies attest of an increase in herbaceous vegetation during the last centuries, showing an impact of the yakut stockbreeders and an extension of the pastures in the surroundings of the natural alaaes.