



Characterization of orphan sesquiterpene biosynthesis genes in *Botrytis cinerea*

Javier Barua, Michelli Massaroli, Josefina Aleu, Muriel Viaud, Isidro G Collado

► To cite this version:

Javier Barua, Michelli Massaroli, Josefina Aleu, Muriel Viaud, Isidro G Collado. Characterization of orphan sesquiterpene biosynthesis genes in *Botrytis cinerea*. First biennial meeting Chemical biology group, RSEQ, Mar 2012, Santiago de Compostela, Spain. hal-01191225

HAL Id: hal-01191225


<https://hal.science/hal-01191225>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FIRST BIENNIAL MEETING CHEMICAL BIOLOGY GROUP REAL SOCIEDAD ESPAÑOLA DE QUÍMICA




Characterization of Orphan Sesquiterpene Biosynthesis Genes in *Botrytis Cinerea*

Javier Barúa,² Michelli Massaroli,² Josefina Aleu,² Muriel Viaud,¹ Isidro G. Collado²

¹ INRA, BIOGER, 78350 Thiverval-Grignon, France. ² Organic Chemistry Department, University of Cádiz, Campus Ciencia y Tecnología, 11510 Puerto Real, Cádiz, Spain.

Advances in genomic studies, in combination with increased access to DNA sequencing, are providing a wealth of information about how natural products are assembled, mechanisms by which natural product gene clusters can be manipulated to yield new product diversity, and the genetic potential of individual organisms.

Botrytis cinerea is a well known pathogen of a number of commercial crops and produces many structurally diverse metabolites. *Botrytis* produces two series of phytotoxic metabolites: a family of characteristic sesquiterpene metabolites which contain the basic botryane skeleton, principally botrydial (**1**) and two polyketide lactones types: botcinins (**2**, **3**) and botrylactone (**4**).¹ Additionally, abscisic acid (a phytohormone) has been isolated and reported.²



However, genomic data revealed that *B. cinerea* has 43 key enzymes involved in the biosynthesis of secondary metabolites, including many polyketides, peptides and terpenes, some of them are specific of this phytopathogen. The study displayed six genes coding for putative sesquiterpene cyclases (STC) in *B. cinerea*.³

So, we have undertaken a chemical genetic project oriented to characterization of these orphan biosynthetic genes and their corresponding cryptic metabolites from *B. cinerea*, by a molecular- and chemical-based epigenetic approach, using mutant strains which have silenced genes coding for five STCs. As a result the sesquiterpene cyclases STC1 and STC4 have been characterized and some aspect about the role played by STC1, STC2 and STC4 in the pathogenicity of *B. cinerea* determined.

- 1 J. Moraga, C. Pinedo, R. Durán-Patrón, I. G. Collado, R. Hernández-Galán, *Tetrahedron*, 2011, **67**, 8583.
- 2 V. Siewers, L. Kokkelink, J. Smedsgaard, P. Tudzynski, *Appl Environ Microbiol.*, 2006, **72**, 4619-4626.
- 3 J. Amselem, C. Cuomo, J. van Kan, M. Viaud, et al., *PLoS Genet.*, 2011, **7**, e1002230.