

The functional characterization of BcBOT3-BcBOT5 genes involved in the biosynthesis of the toxins botrydial

Javier Moraga, Berengere Dalmais-Lenaers, Josefina Aleu, Muriel Viaud, Isidro.G. Collado

▶ To cite this version:

Javier Moraga, Berengere Dalmais-Lenaers, Josefina Aleu, Muriel Viaud, Isidro.G. Collado. The functional characterization of BcBOT3-BcBOT5 genes involved in the biosynthesis of the toxins botrydial. Bio-communication- Phytochemical Society of Europe., Sep 2012, Puerto Real, Cadiz, Spain. Francisco A. Macias, 2012. hal-01191226


HAL Id: hal-01191226

https://hal.science/hal-01191226

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Book of abstract Biocom12 Congress. @ 2012 Francisco A. Macias Editor

ISBN 84-695-4508-6.(Spain)

Editor-in-Chief:

Francisco A. Macias, Department of Organic Chemistry, Faculty of Sciences, University of Cádiz, Spain

Phone: 34-956012770 Puerto real (Cadiz)

E-mail: famacias@uca.es

Associate Editors:

Jose A. Alvarez, University of Cadiz, Spain Ana M. Simonet, University of Cadiz, Spain Ascension Torres, University of Cadiz, Spain Rosa M. Varela, University of Cadiz, Spain

First Edition: Copisteria San Rafael, S.L.,

Logo Design: Estudio Crealia www.e-crealia.com **Cover Design**: David Chinchilla and Belen Macias **Electronic Layout**: Estudio Crealia www.e-crealia.com

Cover Illustration: Manuel Espinosa

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of traslation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction and storage in data banks. Duplication of this publication or parts thereof is only permitted from the publishers.

Printed on acid free and low chlorine paper.

Printed in Spain.



Waters P_5

The functional characterization of BcBOT3-BcBOT5 genes involved in the biosynthesis of the toxins botrydial

<u>Javier Moraga</u> (javier.moraga@uca.es) ⁽¹⁾, B. Dalmais ⁽²⁾, J. Aleu ⁽¹⁾, M. Viaud ⁽²⁾, I. G. Collado ⁽¹⁾

1 University of Cadiz | Organic Chemistry Department, Campus Ciencia y Tecnología, 11510, Puerto Real, SPAIN 2 INRA, BIOGER | , 78850, Thiverval-Grignon, FRANCE

Botrytis cinerea is a well-known pathogen affecting a number of commercial crops, which produces a variety of structurally diverse metabolites. There is no evidence for the production of host-specific toxins by this fungus, which is in accordance with the broad host range of this pathogen. B. cinerea produces two series of phytotoxic metabolites: a family of sesquiterpenes which contain the basic botriane skeleton and a family of polyketides.

Advances in genomic studies, in combination with increased access to DNA sequencing, are providing a wealth of information about how natural products are assembled, mechanisms by which natural product gene clusters can be manipulated to yield new product diversity, and the genetic potential of individual organisms.

In previous works we provided the functional characterization of BcBOT1 and BcBOT2 genes, which encode a P450 monooxygenase and a sesquiterpene cyclase, respectively. The region surrounding the previously characterized BcBOT1 gene was identified as the botrydial biosynthetic gene cluster: BcBOT 2 to BcBOT 5 genes. Wherein BcBOT3 and BcBOT4 genes encode two P450 monooxygenases and BcBOT5 shows high homology to known acetyl transferases.

In this communication we report the functional characterisation of the BcBOT3-5 genes and the encoded key enzymes which catalyze intermediate steps in the biosynthesis pathway leading to botrydial and its derivati.



