



## Structural and functional comparison of pyrrolnitrin- and iprodione-induced modifications in the class III histidine-kinase Bos1 of [i]Botrytis cinerea[/i]

Sabine, Helma Fillinger-David, Sakhr Ajouz, Philippe C. Nicot, Pierre Leroux, Marc Bardin

### ► To cite this version:

Sabine, Helma Fillinger-David, Sakhr Ajouz, Philippe C. Nicot, Pierre Leroux, Marc Bardin. Structural and functional comparison of pyrrolnitrin- and iprodione-induced modifications in the class III histidine-kinase Bos1 of [i]Botrytis cinerea[/i]. 11. European conference on fungal genetics, Mar 2012, Marburg, Germany. hal-01191323

**HAL Id: hal-01191323**

**<https://hal.science/hal-01191323>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



11TH EUROPEAN CONFERENCE ON FUNGAL GENETICS

PHILIPPS-UNIVERSITÄT MARBURG, GERMANY

MARCH 30 - APRIL 2 2012

ECFG11



MEETING ABSTRACTS

## PS7.6

### Structural and Functional Comparison of Pyrrolnitrin- and Iprodione-induced Modifications in the Class III histidine-kinase Bos1 of *Botrytis cinerea*

Sabine Fillinger<sup>[1]</sup> Sakhr Ajouz<sup>[2]</sup> Philippe Nicot<sup>[2]</sup> Pierre Leroux<sup>[1]</sup> Marc Bardin<sup>[2]</sup>

<sup>1</sup> INRA UR 1290, BIOGER CPP, BP01, F-78850 Thiverval-Grignon, France <sup>2</sup> INRA, UR407, Plant Pathology Unit, Domaine St Maurice, F-84140 Montfavet, France

Dicarboximides and phenylpyrroles are commonly used fungicides against plant pathogenic ascomycetes. Although their effect on fungal osmosensing systems has been shown in many studies, their modes-of-action still remain unclear. Laboratory- or field-mutants of fungi resistant to either or both fungicide categories generally harbour point mutations in the sensor histidine kinase of the osmotic signal transduction cascade. We compared the mechanisms of resistance to the dicarboximide iprodione and to pyrrolnitrin, a structural analogue of phenylpyrroles, in *Botrytis cinerea*. Pyrrolnitrin-induced mutants and iprodione-induced mutants of *B. cinerea* were produced *in vitro*. For the pyrrolnitrin-induced mutants, high level of resistance to pyrrolnitrin was associated with a high level of resistance to iprodione. For the iprodione-induced mutants, the high level of resistance to iprodione generated variable levels of resistance to pyrrolnitrin and phenylpyrroles. All selected mutants showed hypersensitivity to high osmolarity and regardless of their resistance levels to phenylpyrroles, they showed strongly reduced fitness parameters (sporulation, mycelial growth, aggressiveness on plants) compared to the parental phenotypes. The sequences of the osmosensing class III histidine kinase encoding gene *bos1* showed different mutations in both types of mutants. All of them affected the HAMP-domains of the histidine-kinase showing that each of the six HAMP domains is important for signal-transduction. Structure modelling of the HAMP domains revealed that the replacements of hydrophobic residues within the HAMP domains generally affected their helical structure, probably abolishing signal transduction. The mutation of residues E529, T581, or E692 – without consequences on HAMP structure – highlighted their involvement in signal transduction.