



Estimation of genetic trends from 1977 to 2000 for stress-responsive systems in French Large White and Landrace pig populations using frozen semen

Aline Foury, Thierry Tribout, Christophe Bazin, Yvon Billon, Marcel Bouffaud, Jean-Marcel Gogu , Jean Pierre Bidanel, Pierre Morm de

► To cite this version:

Aline Foury, Thierry Tribout, Christophe Bazin, Yvon Billon, Marcel Bouffaud, et al.. Estimation of genetic trends from 1977 to 2000 for stress-responsive systems in French Large White and Landrace pig populations using frozen semen. *Animal*, 2009, 3 (12), pp.1681-1687. 10.1017/S1751731109990504 . hal-01193458

HAL Id: hal-01193458

<https://hal.science/hal-01193458>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin e au d p t et   la diffusion de documents scientifiques de niveau recherche, publi s ou non,  manant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv s.

Estimation of genetic trends from 1977 to 2000 for stress-responsive systems in French Large White and Landrace pig populations using frozen semen

A. Foury¹, T. Tribout², C. Bazin³, Y. Billon⁴, M. Bouffaud⁵, J. M. Gogué⁶, J. P. Bidanel² and P. Mormède^{1†}

¹Université Victor Segalen Bordeaux 2, PsyNuGen, INRA UMR1286, CNRS UMR5226, F-33076 Bordeaux, France; ²INRA UMR1313 Génétique Animale et Biologie Intégrative, F-78352 Jouy-en-Josas, France; ³IFIP, Pôle Génétique, F-35651 Le Rheu, France; ⁴INRA, UE967 Génétique Expérimentale en Productions Animales F-17700 Surgères, France; ⁵INRA, UE450 Testage de Porcs F35650 Le Rheu, France; ⁶INRA, UE332 Domaine Expérimental de Bourges, F-18300 Osmoy, France

(Received 30 January 2009; Accepted 16 June 2009; First published online 12 August 2009)

An experimental design aiming at analysing the consequences of genetic selection from 1977 to 1998–2000 on the evolution of stress-responsive systems in the French Large White (LW) and Landrace (LR) pig populations was conducted by INRA and IFIP-Institut du Porc. Large White sows were inseminated with semen from LW boars born in 1977 (frozen semen) or in 1998 and their second-generation offspring were station-tested. Landrace sows were inseminated with semen from LR boars born in 1977 (frozen semen) or in 1999 to 2000, and their progeny was station-tested. Urinary concentration of stress hormones (cortisol and catecholamines) and traits related to carcass composition (estimated carcass lean content (ECLC) and global adiposity) and meat quality (pH 24 h) were measured. For the two populations, selection carried out since 1977 led to an increase in ECLC and a decrease in carcass adiposity. Between 1977 and 1998 to 2000, urinary concentrations of stress hormones were unchanged in the LR breed, but were decreased in the LW breed. Moreover, for the animals generated from LW boars born in 1977 and in 1998, urinary cortisol levels were negatively correlated with ECLC. Therefore, in the LW breed, selection carried out for higher ECLC resulted in a decrease in cortisol production, as well as a reduction of catecholamine production that may be responsible for the lower ultimate pH of meat. Therefore, selection carried out for increased carcass lean content led, in this breed, to large modifications in the functioning of the stress-responsive systems, thereby influencing a large range of physiological regulations and technical properties such as carcass composition and meat pH, which remained however in the normal range for acceptable meat quality.

Keywords: catecholamines, cortisol, fatness, genetic trend, stress

Implications

During the last 30 years, the reduction of carcass fat content in pigs was one of the main objectives of genetic selection. It resulted in an important genetic increase in carcass lean content in French pig breeds. Due to the influence of cortisol and catecholamines on energy metabolism, the down-regulation of neuroendocrine stress-responsive systems may have contributed to this trend in the Large White breed. However, considering the importance of these hormones on several major production traits such as meat quality and newborn survival, this genetically driven down-regulation of cortisol and catecholamine production may also be responsible for the observed worsening of these traits. A better knowledge of

the balance between favourable and unfavourable effects of these hormones, and of their genetic regulation, should help designing future selection strategies to jointly improve the level of performance, carcass composition, product quality and animal welfare.

Introduction

The two main stress-responsive neuroendocrine systems play a critical role in the regulation of energy fluxes. The hypothalamic–pituitary–adrenocortical (HPA) axis influences feeding behaviour, pancreatic hormone secretion, energy expenditure and the protein/lipid balance (Dallman *et al.*, 1993). Cortisol, the main hormone of the axis, favours the accretion of fat at the expense of proteins (Devenport *et al.*, 1989). On the other

[†] E-mail: Pierre.Mormede@bordeaux.inra.fr

hand, catecholamines (adrenaline and noradrenaline) released by the sympathetic nervous system increase the use of energy stores (glycogen and lipids) (Scheurink and Steffens, 1990) and exert anabolic effects on protein metabolism (Navegantes *et al.*, 2002). Sympathetic activation by stress before slaughter reduces muscle glycogen content and *post-mortem* acidification, leading to dark, firm and dry type meat (Fernandez and Tornberg, 1991; Monin, 2003).

It is now well established that genetic factors influence individual variations in stress behavioural and neuroendocrine responses (Morm de *et al.*, 2002). In a study using a F2 segregating population between the Duroc and Large White pig breeds, we demonstrated that cortisol levels measured in urine collected in the bladder after slaughter correlated negatively to lean content of the carcass, and that urinary catecholamine levels correlated positively to ultimate pH of the meat (Foury *et al.*, 2005). In the context of a larger study to explore the genetic components of meat quality (Plastow *et al.*, 2005), the activity of neuroendocrine stress systems and the relationship of neuroendocrine characteristics with carcass composition were studied in five genetic lines representing a significant proportion of European pig production (Foury *et al.*, 2007). The Large White, Landrace, Duroc and Meishan breeds show basal urinary cortisol levels negatively correlated to carcass lean content. Genetic selection on stress reactivity traits could improve both animal welfare and product quality. The mapping of chromosome regions involved in genetic variations opens the way to the identification of the molecular mechanisms involved in individual differences in stress responses (D saut s *et al.*, 2002; Ousova *et al.*, 2004).

During the last 30 years, the reduction of carcass fat content was one of the main objectives of genetic selection, together with the increase in growth rate and, more recently, the prolificacy of sows (Legault, 1998; Renand *et al.*, 2003). It resulted in an important genetic increase in

the carcass lean content of French Large White (LW) and French Landrace (LR) populations (Bazin *et al.*, 2003; Tribout *et al.*, 2004). Selection for lean content also led to modifications in fibre type composition, which resulted in a higher proportion of glycolytic fibres and fibre hypertrophy (Rahelic and Puac, 1981; Weiler *et al.*, 1995). Because of a reduced capillary density and an impaired nuclear control of cellular processes, fibre hypertrophy seems to be associated with a lower capacity of the fibres to adapt to activity-induced demands (Rehfeldt *et al.*, 2000). Therefore, the aim of this study was to analyse the consequences of genetic selection on the evolution of the stress-responsive neuroendocrine systems of pigs issued from LW and LR boars born either in 1977 or between 1998 and 2000.

Material and methods

Animals and experimental design

The present study is part of a vast experimental design aiming at estimating realized genetic trends in the French maternal porcine populations between 1977 and 1998 (LW) or between 1977 and 1999 to 2000 (LR) for a large number of growth, reproduction, body composition and carcass quality traits. Two different groups of animals (referred to as G77 and G98 for French LW and L77 and L99 for French LR) have been produced, by inseminating French LW and LR sows with semen from boars of the same breed born either in 1977 (frozen kept) or between 1998 and 2000. The experimental design is schematized in Figure 1.

For the LW crossbreeding, randomly chosen males and females from each of the two F1 groups generated from boars born either in 1977 or in 1998 were kept for breeding and produced, respectively, G77 and G98 pigs by *inter se* mating. This part of the experiment took place in two INRA experimental herds: Le Magneraud (Charente-Maritime, France) and Bourges (Cher, France). Further details on the




Figure 1 Experimental design.

experimental design are given in Tribout *et al.* (2004). At a live weight of approximately 106 kg, after 18 h of fasting and 4.5 h of transport, 236 pigs (castrated males and females) were slaughtered after electrical stunning in two INRA experimental abattoirs: Jouy-en-Josas (Yvelines, France) and Saint-Gilles (Ille-et-Vilaine, France).

The French Pig Institute (IFIP) was in charge of the LR crossbreeding. French LR sows were inseminated with semen from boars born either in 1977 or in 1999 and 2000 in seven farms and produced respectively L77 and L99 pigs. They were transferred in two testing stations for fattening: Maureon (Morbihan, France) and Le Rheu (Ille-et-Vilaine, France). Further details on the experimental design are given in Bazin *et al.* (2003). At a live weight of approximately 105 kg, after 18 h of fasting and 0.5 h of transport, 189 pigs (castrated males and females) were slaughtered after electrical stunning in the Cooperl-Industrie abattoir (Montfort-sur-Meu, Ille-et-Vilaine, France).

Sampling procedure and chemical analyses

After evisceration, urine samples of 136 LW pigs and 126 LR pigs were collected from the bladder to measure stress hormone levels (cortisol and catecholamines). A preservative (ethylene diamine tetra acetic acid (EDTA) 10%, 1 ml/40 ml) was added and samples were frozen until analysis.

Urinary cortisol was assayed using a solid-phase extraction procedure on C18 cartridges followed by HPLC with UV absorbance detection (254 nm), as previously described (Hay and Mormède, 1997b). The intra- and inter-assay coefficients of variation (%) were 7.4 and 10.6, respectively.

Urinary catecholamines (adrenaline and noradrenaline) were assayed using an ion-exchange purification procedure followed by HPLC with electrochemical detection, as previously described (Hay and Mormède, 1997a). The intra- and inter-assay coefficients of variation (%) were 7.0 and 7.1 for adrenaline, and 6.5 and 11.6 for noradrenaline, respectively.

Concentrations of hormones in urine were expressed as their ratio to creatinine content (ng/mg creatinine), to correct for the variable dilution of urine related to water intake (Crockett *et al.*, 1993). Creatinine levels were determined using a colorimetric quantitative reaction (Creatinine; BIO-LABO, Fismes, France).

Carcass composition measurements

The day after slaughter, the backfat thickness at the last cervical vertebra (neck), last dorsal vertebra (back) and last lumbar vertebra (rump) at the sectioned edge of the carcass were recorded with a ruler. Ultimate pH was measured 24 h after slaughter in the *semi-membranosus* muscle (pH24). Estimated carcass lean content (ECLC) was calculated from three joint weights expressed as a proportion of the half-carcass weight using the following equation (Métayer and Daumas, 1998):

$$\text{ECLC} = 5.684 + 1.197\% \text{ ham} + 1.076\% \text{ loin} - 1.059\% \text{ backfat.}$$

Statistical analyses

Because the LR and LW populations are totally independent, the data were analysed separately with the Statistical Analysis Systems (SAS) Institute software (Statistical Analysis Systems Institute, 2000). Hormone levels in urine were transformed into their logarithmic scores (ln) for data normalization. Experimental group (G77 v. G98 or L77 v. L99), sex, farrowing batch and station, abattoir and slaughter batch were included as fixed effects and live weight as a co-variable in the MIXED procedure of SAS when significant (Table 1). Least-squares means were generated by the LSMEANS statement. Pearson correlations were calculated between hormone concentrations, carcass composition (lean meat content and mean backfat thickness) and pH24 with the CORR procedure.

The realised genetic trends from 1977 to 1998 (ΔG) and their standard errors (s.e. ΔG) were estimated for each

Table 1 Fixed between subjects effects considered in the models of analysis for each trait

Effects	Cortisol urine	Adrenaline urine	Noradrenaline urine	ECLC	Backfat	pH24
LW						
Experimental group	*	**	*	****	****	*
Sex	ns	**	**	****	*	ns
Farrowing batch	*	ns	ns	*	ns	ns
Slaughter batch	ns	****	****	ns	**	ns
Abattoir	****	ns	***	****	****	****
Live weight (co-variable)	ns	ns	ns	ns	ns	ns
LR						
Experimental group	ns	ns	ns	****	****	ns
Sex	ns	*	ns	****	****	*
Station	**	****	*	**	ns	***
Slaughter batch	*	ns	ns	ns	ns	ns
Live weight (co-variable)	ns	ns	ns	****	****	ns

ECLC = estimated carcass lean content; Backfat = average of the backfat thickness measured at the neck, back and rump levels; pH24 = ultimate pH in the *semi-membranosus* muscle; LW = French Large White; LR = French Landrace.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$; ns: non-significant.

considered trait as proposed by Smith (1977):

$$\Delta G = 2 \times (\text{G98 least-squares mean} - \text{G77 least-squares mean}),$$

and s.e. $\Delta G = 2 \times \text{s.e. (G98 least-squares mean - G77 least-squares mean)}$.

Results

Genetic group least-squares means and estimated genetic trends for stress hormones are shown in Tables 2 and 3 for LW and LR crossings, respectively. For the LW breed, urinary levels of cortisol, adrenaline and noradrenaline were lower in 1998 than in 1977 (−24.4%, −26.7% and −18.9% respectively). ECLC increased between 1977 and 1998 and carcass adiposity dropped significantly. Ultimate pH was lower in 1998 than in 1977. In the LR breed, ECLC and carcass adiposity followed the same trends as in LW, but ultimate pH, urinary levels of cortisol, adrenaline and noradrenaline did not significantly differ from 1977 to 1999.

Pearson correlations between hormone concentrations and carcass composition were calculated. No significant correlations were found for the pigs generated from LR boars. Significant correlations were found for the pigs generated from LW boars born in 1977: urinary cortisol and adrenaline levels were negatively correlated with ECLC ($r = -0.27$, $P < 0.05$ and $r = -0.36$, $P < 0.01$, respectively), and urinary adrenaline concentrations were also

positively correlated with mean backfat thickness ($r = 0.33$, $P < 0.05$). For the pigs generated from LW boars born in 1998, urinary cortisol levels were negatively correlated with ECLC ($r = -0.23$, $P = 0.05$). Correlations between levels of cortisol and ECLC for the LW population are presented in Figure 2. No significant correlations were found between stress hormones and pH24.

Discussion

The aim of this study was to explore the consequences of this selection on the stress-responsive neuroendocrine systems of pigs issued from LW and LR boars born in 1977 or between 1998 and 2000. The results obtained here for ECLC, carcass adiposity and ultimate pH are in accordance with those obtained by Tribout *et al.* (2004) and Bazin *et al.* (2003) with the complete set of animals from the experiment. This shows that our sample of animals is representative of the whole experimental populations.

The generations studied in the two populations (F1 in LR and F2 in LW) and farming and slaughter conditions were different between LW and LR populations. As the results obtained may depend on all these factors, the two breeds were analyzed independently.

In the LW population, the consequences of selection are obviously a strong increase in carcass lean content and an important decrease in carcass adiposity, but also a decrease in cortisol and catecholamines levels. Cortisol favours the storage of lipids (in presence of insulin) at the expense of

Table 2 Least-squares means and estimated genetic trends (ΔG) for endocrine, carcass composition and meat quality traits of G77 and G98 pigs (Large White breed)

Traits	G77 <i>n</i>	G98 <i>n</i>	G77 LSM (s.e.)	G98 LSM (s.e.)	ΔG (s.e.)	$P > t $ $H_0: \Delta G = 0$
Cortisol urine (ng/mg creatinine), ln	63	70	2.97 (0.07)	2.69 (0.07)	−0.56 (0.17)	**
Adrenaline urine (ng/mg creatinine), ln	57	58	1.96 (0.10)	1.65 (0.10)	−0.63 (0.22)	**
Noradrenaline urine (ng/mg creatinine), ln	58	63	2.60 (0.08)	2.39 (0.08)	−0.42 (0.18)	*
ECLC (%)	116	120	52.20 (0.30)	56.00 (0.29)	7.59 (0.82)	****
Mean backfat thickness (mm)	116	120	24.51 (0.28)	22.13 (0.28)	−4.76 (0.80)	***
pH24	116	120	5.84 (0.03)	5.76 (0.03)	−0.17 (0.07)	*

LSM = least-squares means; *n* = number of observations in each group; $\Delta G = 2 \times (\text{G98 LSM} - \text{G77 LSM})$; ECLC = estimated carcass lean content; Mean backfat thickness = average of the backfat thickness measured at the neck, back and rump levels; pH24 = ultimate pH in the *semi-membranosus* muscle.

$P > |t|$ = *t*-test probability for $\Delta G \neq 0$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$.

Table 3 Least-squares means and estimated genetic trends (ΔG) for endocrine, carcass composition and meat quality traits of L77 and L99 pigs (Landrace breed)

Traits	L77 <i>n</i>	L99 <i>n</i>	L77 LSM (s.e.)	L99 LSM (s.e.)	ΔG (s.e.)	$P > t $ $H_0: \Delta G = 0$
Cortisol urine (ng/mg creatinine), ln	59	58	3.93 (0.08)	3.97 (0.08)	0.06 (0.22)	ns
Adrenaline urine (ng/mg creatinine), ln	58	58	2.25 (0.07)	2.28 (0.08)	0.07 (0.17)	ns
Noradrenaline urine (ng/mg creatinine), ln	58	59	2.67 (0.07)	2.68 (0.07)	0.04 (0.18)	ns
ECLC (%)	91	98	48.62 (0.33)	51.16 (0.32)	5.07 (0.92)	****
Mean backfat thickness (mm)	91	98	27.87 (0.36)	23.83 (0.35)	−8.06 (1.02)	****
pH24	91	98	5.68 (0.02)	5.69 (0.02)	0.02 (0.06)	ns

LSM = least-squares means; *n* = number of observations in each group; $\Delta G = 2 \times (\text{L99 LSM} - \text{L77 LSM})$; ECLC = estimated carcass lean content; Mean backfat thickness = average of the backfat thickness measured at the neck, back and rump levels; pH24 = ultimate pH in the *semi-membranosus* muscle.

$P > |t|$ = *t*-test probability for $\Delta G \neq 0$; **** $P < 0.0001$; ns: non-significant.




Figure 2 Scatter plot and regression line between estimated carcass lean content (ECLC) and concentration of cortisol in urine (ng/mg creatinine, ln scale) in the Large White (LW) breed.

proteins, via peripheral catabolism and hepatic neoglucogenesis (Devenport *et al.*, 1989; Dallman *et al.*, 1993); it is not surprising that selection on carcass lean content results in the selection of animals with lower levels of cortisol in the LW population. Nevertheless, the hypothesis of a genetic drift of hormone production independent of the selection traits cannot be completely discarded. The negative correlation between cortisol and lean content for the G77 and G98 pigs illustrates the relationship between cortisol and fatness. It also shows that selection did not reduce variability in the LW population between 1977 and 1998. The relationship found between catecholamines and fatness in the G77 pigs is more difficult to understand. As the activation of β -adrenergic receptors increases mobilization of fat and reduces protein catabolism (Navegantes *et al.*, 2002), we could have expected the reverse result. Part of this relationship probably results from the coupling between adrenaline and cortisol secretions (Foury *et al.*, 2005). Indeed, the synthesis of adrenaline is under control of cortisol that induces phenylethanolamine-*N*-methyltransferase, the enzyme converting noradrenaline into adrenaline in the adrenal medulla (Ciaranello, 1978). The expected correlation between cortisol and ECLC was confirmed in G98 pigs; however, no relationship between adrenaline and carcass composition measures was found.

On the same experimental design, other studies have estimated genetic trends for traits measured at farrowing in

the LW breed (Canario *et al.*, 2007a and 2007b). The results suggest that the improvement in lean growth rate and in sow prolificacy from 1977 to 1998 has resulted in a lower maturity of piglets at birth. Leenhouwers *et al.* (2002) showed that the degree of physiological maturity of piglets increases with increasing circulating cortisol concentrations at birth. Therefore, the selection of animals with a lower production of cortisol can lead to a lower maturity of piglets.

In the LW population, selection on lean tissue growth was also accompanied by a decrease in catecholamines and ultimate pH levels. Indeed, sympathetic activation before slaughter increases muscle glycogenolysis and therefore reduces lactic acid production *post-mortem* and meat acidification (Fernandez and Tornberg, 1991). Therefore a reduced *pre-mortem* sympathetic activation, as observed in Large White pigs, spares muscle glycogen and, as a consequence, decreases ultimate pH. However, ultimate pH remains in the normal range for acceptable meat quality.

In the LR breed, selection did not affect cortisol levels. A hypothesis to explain this fact could be that genes responsible for the variability of cortisol secretion are not segregating in the LR population. However, this hypothesis could not be checked in the present population and no estimate of the heritability of blood and urinary cortisol concentrations is, to our knowledge, available in the literature. The stability of cortisol levels could be an explanation to the more limited gain in lean content for the LR

breed in comparison with the LW breed, even if mean backfat thickness shows a greater decrease in LR than in LW pigs. Indeed, lean content may be more informative than carcass adiposity of the influence of cortisol that changes the proteins/lipids balance (Devenport *et al.*, 1989). This more limited gain in the LR breed can also be explained by the fact that one of the main selection goals in the 1980s and 1990s was the eradication of the halothane sensitivity gene (Renand *et al.*, 2003), which led to a reduction in the selection pressure on carcass composition traits. The eradication of this gene probably did not have consequences on cortisol levels since the halothane sensitivity mutation does not influence HPA axis activity (Morm de and Dantzer, 1978; Terlouw *et al.*, 2001).

The present data show that, due to the influence of cortisol and catecholamines on energy metabolism, selection performed from 1977 to 1998 in the LW population to increase the carcass lean content has resulted in the selection of low cortisol and catecholamines level animals. Moreover, this indirect selection on stress-responsive systems may be responsible for negative trends on other traits such as meat quality (Lonergan *et al.*, 2001) or newborn survival (Canario *et al.*, 2007a and 2007b; Herpin *et al.*, 1993). A better understanding of the molecular and physiological mechanisms involved in the relationships between endocrine systems (cortisol and catecholamines) and different production traits would help to optimize selection to improve together the level of production, carcass composition, product quality and animal welfare.

Acknowledgements

We thank the staff of Le Magneraud, Bourges and Le Rheu experimental units and of Mauron testing stations for their careful data management, and Francis Bussi res from the INRA Pig and Goat Insemination experimental unit for collecting, conditioning and stocking the semen of 1977 boars 30 years ago.

This work was supported by the INRA, the French Pig Institute (IFIP) and the 'Actions Innovantes' from the French Ministry of Agriculture.

References

Bazin C, Tiger E, Tribout T, Bouffaud M, Madigand G, Boulard J, Deschodt G, Fl  o J-Y, Gueblez R, Maignel L and Bidanel JP 2003. Estimation, par utilisation de semence congel  e en   levage de s  lection, du progr  s g  n  tique r  alis   entre 1977 et 2000 dans les races Large White et Landrace Fran  ais pour les caract  res de croissance, de carcasse et de qualit   de la viande. *Journ  es de la Recherche Porcine en France* 35, 277–284.

Canario L, P  re MC, Tribout T, Thomas F, David C, Gogu  J, Herpin P, Bidanel JP and Le Dividich J 2007a. Estimation of genetic trends from 1977 to 1998 of body composition and physiological state of Large White pigs at birth. *Animal* 1, 1409–1413.

Canario L, Rydhmer L, Gogu  J, Tribout T and Bidanel JP 2007b. Estimation of genetic trends from 1977 to 1998 for farrowing characteristics in the French Large White breed using frozen semen. *Animal* 1, 929–938.

Ciaranello RD 1978. Regulation of phenylethanolamine N-methyltransferase. *Biochemical Pharmacology* 27, 1895–1897.

Crockett CM, Bowers CL, Sackett GP and Bowden DM 1993. Urinary cortisol responses of longtailed macaques to 5 cage sizes, tethering, sedation, and room change. *American Journal of Primatology* 30, 55–74.

Dallman MF, Strack AM, Akana SF, Bradbury MJ, Hanson ES, Scribner KA and Smith M 1993. Feast and famine: critical role of glucocorticoids with insulin in daily energy flow. *Frontiers in Neuroendocrinology* 14, 303–347.

D  saut  s C, Bidanel J, Milan D, Iannuccelli N, Amigues Y, Bourgeois F, Caritez J, Renard C, Chevalet C and Morm de P 2002. Genetic linkage mapping of quantitative trait loci for behavioral and neuroendocrine stress response traits in pigs. *Journal of Animal Science* 80, 2276–2285.

Devenport L, Knehans A, Sundstrom A and Thomas T 1989. Corticosterone's dual metabolic actions. *Life Sciences* 45, 1389–1396.

Fernandez X and Tornberg E 1991. A review of the causes of variation in muscle glycogen content and ultimate pH in pigs. *Journal of Muscle Foods* 2, 209–235.

Foury A, Devillers N, Sanchez MP, Griffon H, Le Roy P and Morm de P 2005. Stress hormones, carcass composition and meat quality in Large White   Duroc pigs. *Meat Science* 69, 703–707.

Foury A, Geverink N, Gil M, Gispert M, Hortos M, Furnols M, Carri n D, Blott S, Plastow G and Morm de P 2007. Stress neuroendocrine profiles in five pig breeding lines and the relationship with carcass composition. *Animal* 1, 973–982.

Hay M and Morm de P 1997a. Determination of catecholamines and methoxycatecholamines excretion patterns in pig and rat urine by ion-exchange liquid chromatography with electrochemical detection. *Journal of Chromatography B* 703, 15–23.

Hay M and Morm de P 1997b. Improved determination of urinary cortisol and cortisone, or corticosterone and 11-dehydrocorticosterone by high-performance liquid chromatography with ultraviolet absorbance detection. *Journal of Chromatography B* 702, 33–39.

Herpin P, Dividich J and Amaral N 1993. Effect of selection for lean tissue growth on body composition and physiological state of the pig at birth. *Journal of Animal Science* 71, 2645–2653.

Leenhouders JI, Knol E, Groot P, Vos H and Lende T 2002. Fetal development in the pig in relation to genetic merit for piglet survival. *Journal of Animal Science* 80, 1759–1770.

Legault C 1998. G  n  tique et prolificit   chez la truie: la voie hyperprolif  rique et la voie sino-europ  enne. *Productions Animales* 11, 214–218.

Lonergan SM, Huff-Lonergan E, Rowe LJ, Kuhlers DL and Jungst SB 2001. Selection for lean growth efficiency in Duroc pigs influences pork quality. *Journal of Animal Science* 79, 2075–2085.

M  tayer A and Dumas G 1998. Estimation of the lean meat content of pig carcasses using the measurement of different cuts. *Journ  es de la Recherche Porcine en France* 30, 7–11.

Monin G 2003. Abattage des porcs et qualit  s des carcasses et des viandes. *Productions Animales* 16, 251–262.

Morm de P, Courvoisier H, Ramos A, Marissal-Arvy N, Ousova O, D  saut  s C, Duclos M, Chaouloff F and Moisan MP 2002. Molecular genetic approaches to investigate individual variations in behavioral and neuroendocrine stress responses. *Psychoneuroendocrinology* 27, 563–583.

Morm de P and Dantzer R 1978. Behavioral and pituitary-adrenal characteristics of pigs differing by their susceptibility to the malignant hyperthermia syndrome induced by halothane anesthesia. 2. pituitary-adrenal-function. *Annales de Recherches V  t  rinaires* 9, 569–576.

Navegantes L, Migliorini R and Kettelhut I 2002. Adrenergic control of protein metabolism in skeletal muscle. *Current Opinion in Clinical Nutrition and Metabolic Care* 5, 281–286.

Ousova O, Guyonnet-Duperat V, Iannuccelli N, Didanel JP, Milan D, Genet C, Llamas B, Yerle M, Gellin J, Chardon P, Emptoz-Bonneton A, Pugeat M, Morm de P and Moisan MP 2004. Corticosteroid binding globulin: a new target for cortisol-driven obesity. *Molecular Endocrinology* 18, 1687–1696.

Plastow GS, Carri n D, Gil M, Garc  a-Regueiro JA, Font i Furnols M, Gispert M, Oliver MA, Velarde A, Gu  rdia MD, Hort  s M, Rius MA, S  rraga C, D    s I, Valero A, Sosnicki A, Klont R, Dorman S, Wilkinson JM, Evans G, Sargent C, Davey G, Connolly D, Houeix B, Maltin CM, Hayes HE, Anandavijayan V, Foury A, Geverink N, Cairns M, Tilley RE, Morm de P and Blott SC 2005. Quality pork genes and meat production. *Meat Science* 70, 409–421.

Rahelic S and Puac S 1981. Fiber types in *Longissimus dorsi* from wild and highly selected pig breeds. *Meat Science* 5, 439–450.

Rehfeldt C, Fiedler I, Dietl G and Ender K 2000. Myogenesis and postnatal skeletal muscle cell growth as influenced by selection. *Livestock Production Science* 66, 177–188.

Genetic trends from 1977 to 2000 for stress-responsive systems in pigs

- Renand G, Larzul C, Le Bihan-Duval E and Le Roy P 2003. L'amélioration génétique de la qualité de la viande dans les différentes espèces: situation actuelle et perspectives à court et moyen terme. *Productions Animales* 16, 159–173.
- Scheurink AJ and Steffens AB 1990. Central and peripheral control of sympathoadrenal activity and energy metabolism in rats. *Physiology and Behavior* 48, 909–920.
- Smith C 1977. Use of stored frozen semen and embryos to measure genetic trends in farm livestock. *Zeitschrift für Tierzüchtung und Züchtungsbiologie* 94, 119–130.
- Statistical Analysis Systems Institute 2000. *Statistics Version 8.02*. SAS Institute Inc., Cary, NC, USA.
- Terlouw C, Ludriks A, Schouten W, Vaessen S, Fernandez X, Andanson S and Père MC 2001. Response of pigs to slaughter stress: predominance of the halothane sensitivity allele. *Viandes et Produits Carnés* 22, 127–136.
- Tribout T, Caritez JC, Gogué J, Gruand J, Bouffaud M, Billon Y, Péry C, Griffon H, Brénot S, Le Tiran M-H, Bussièrès F, Le Roy P and Bidanel JP 2004. Estimation, par utilisation de semence congelée, du progrès génétique réalisé en France entre 1977 et 1998 dans la race porcine Large White: résultats pour quelques caractères de production et de qualité des tissus gras et maigres. *Journées de la Recherche Porcine en France* 36, 275–282.
- Weiler U, Appell HJ, Kremser M, Hofacker S and Glaus R 1995. Consequences of selection on muscle composition – a comparative-study on Gracilis muscle in wild and domestic pigs. *Anatomia Histologia Embryologia* 24, 77–80.