



HAL
open science

Biomarkers of beef tenderness, moving towards analytical tools

Brigitte Picard, Didier Micol, Isabelle Cassar-Malek, C. Denoyelle, Gilles Renand, Jean-François J.-F. Hocquette, C. Capel, L. Journaux

► **To cite this version:**

Brigitte Picard, Didier Micol, Isabelle Cassar-Malek, C. Denoyelle, Gilles Renand, et al.. Biomarkers of beef tenderness, moving towards analytical tools. 64. Annual Meeting of the European Federation of Animal Science (EAAP), European Association for Animal Production (EAAP). ITA., Aug 2013, Nantes, France. hal-01193813

HAL Id: hal-01193813

<https://hal.science/hal-01193813v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biomarkers of beef tenderness, moving towards analytical tools

B. Picard¹, D. Micol¹, I. Cassar-Malek¹, C. Denoyelle², G. Renand³, J.F. Hocquette¹, C. Capel² and L. Journaux²

¹INRA-VetAgro Sup, UMRH 1213, Theix, 63122, France, ²Institut de l'Elevage, 149 rue de Bercy, 75012 Paris, France, ³INRA, UMR 1313, 78352 Jouy en Josas, France; picard@clermont.inra.fr

Variation in beef tenderness is a major problem for the beef industry. Currently, there are no accurate commercial techniques available to evaluate the tenderness of carcasses, and this capacity is even poorer for live animals. The beef industry needs commercially viable tenderness measurements for progress to occur. For years, functional genomic programs have been conducted to search for biomarkers of this quality. The main results showed that, in the Longissimus thoracis of the tenderest animals, the muscle proteins associated with the fast glycolytic type (phosphoglucosmutase, lactate dehydrogenase B, triphosphate isomerase, glyceraldehyde 3-phosphate dehydrogenase, isoforms of fast troponin T, β enolase ...) were less abundant. Several proteins involved in calcium metabolism such as parvalbumin were also identified as positive markers of tenderness, in agreement with the role of calcium in post-mortem ageing. Proteins of heat shock (Hsp) family have been revealed as markers of tenderness at mRNA and protein levels in different experiments (Hsp40, Hsp27, Hsp20, α -B crystallin, Hsp70). This is consistent with a theory that the anti-apoptotic activity of Hsp could slow down the process of cell death early post-mortem and consequently improve tenderness. Other proteins involved in oxidative stress, such as superoxide dismutase 1 or peroxiredoxin 6, have been related negatively with tenderness. Currently, the abundance of these proteins is being measured in different muscles within a representative sample of the French production systems. The final step is to provide a tool for the beef industry to enable it to measure the abundance of these tenderness markers. Its use will provide objective data on the potential tenderness of individuals pre or post slaughter allowing production to be tailored to maximise tenderness and/or decide the appropriate market.

