



**HAL**  
open science

## Digestion and genetics in broilers

Sandrine Grasteau, Agnès Narcy, Cécile Arnould, Aline Bertin, Marie Yvonne Boscher, Fabien Brossier, Fanny Calenge, Irène Gabriel, Edouard Guitton, Mélynda Hassouna, et al.

► **To cite this version:**

Sandrine Grasteau, Agnès Narcy, Cécile Arnould, Aline Bertin, Marie Yvonne Boscher, et al.. Digestion and genetics in broilers. Towards the chicken of the future, Royal Veterinary College - University of London. GBR., Apr 2014, Londres, United Kingdom. hal-01193902

**HAL Id: hal-01193902**

**<https://hal.science/hal-01193902>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Towards the Chicken of the Future Symposium**  
**23-24 April, 2014 at The Royal Veterinary College**  
**Official Programme**

**RVC**  
Royal  
Veterinary  
College  
University of London



**MANCHESTER**  
1824  
The University of Manchester

**Towards the chicken of the future...**  
**A BBSRC funded research project**



[thechickenofthefuture.com](http://thechickenofthefuture.com)

Although our findings point to a metabolic origin of the pathogenesis of these conditions, it is important to stress that our observations do not negate the importance of bacterial infection in the evolution of the lesions. However, we submit that the pathogens play a critical role only in the terminal stages of lesion development.

### **2.3 Digestion and genetics in broilers**

Sandrine Mignon-Grasteau<sup>1</sup>, Agnès Narcy<sup>1</sup>, Cécile Arnould<sup>2</sup>, Aline Bertin<sup>2</sup>, Marie-Yvonne Boscher<sup>3</sup>, Fabien Brossier<sup>4,5</sup>, Fanny Calenge<sup>6</sup>, Irène Gabriel<sup>1</sup>, Edouard Guitton<sup>7</sup>, Mélynda Hassouna<sup>8</sup>, Anne-Christine Lalmanach<sup>4,5</sup>, Maria-Céleste Le Bourhis<sup>9</sup>, Bertrand Méda<sup>1</sup>, Pascale Quéré<sup>4,5</sup>, Nicole Rideau<sup>1</sup>, Catherine Schouler<sup>4,5</sup>, Bernard Carré<sup>1</sup>

<sup>1</sup>UR083 Recherches Avicoles, INRA, 37380 Nouzilly, France;

<sup>2</sup>UMR085 PRC, INRA, 37380 Nouzilly, France;

<sup>3</sup>INRA, LABOGENA, Domaine de Vilvert, F-78352 Jouy en Josas cedex, France

<sup>4</sup>UMR1282 Infectiologie et Santé Publique, INRA, 37380 Nouzilly, France ;

<sup>5</sup>Université François Rabelais de Tours, UMR1282 Infectiologie et Santé Publique, F-37000 Tours, France

<sup>6</sup>INRA UMR1313 Génétique Animale et Biologie Intégrative, 78352 Jouy-en-Josas

<sup>7</sup>UE1277 PFIE Plateforme d'Infectiologie Expérimentale, INRA, 37380 Nouzilly, France

<sup>8</sup>UMR INRA/Agrocampus 1069 SAS, 65, rue de St-Brieuc, CS 84215, 35042 Rennes Cedex – France

<sup>9</sup>UE1295 PEAT Pôle d'Expérimentation Avicole de Tours, INRA, 37380 Nouzilly, France  
[sandrine.grasteau@tours.inra.fr](mailto:sandrine.grasteau@tours.inra.fr)

Poultry production will increase in the future to meet the feed demand of the growing human population. This will increase the pressure on land use to produce cereals and proteins for poultry diets and also the environmental pressure due to the increase of poultry manure discharge. Selecting birds able to digest more varied and less optimal diets could be a way to minimize these negative impacts of poultry production. Projects developed at INRA to reach this goal were focused on three different axes.

We first demonstrated that selection for digestive efficiency was possible by creating divergent lines selected for high (D+) or low (D-) digestive efficiency using a diet difficult to digest. We also showed that the success of the selection is diet-dependent. We also identified several genomic regions having an effect on digestive efficiency that should soon be confirmed by gene expression analyses.

In a second step, we studied the physiological consequences of selection on digestibility to understand which mechanisms were affected. The development of the digestive tract is a key factor in differences observed between the D+ and D-. Major differences have also been observed on feed retention time, excretion traits and microbiota composition. D+ birds expressed higher behavioural inhibition in novel environments but reduced food neophobia compared to D- birds.

In a third step, we tried to assess the impact of this selection on a wide range of traits related to the economic, social and environmental pillars of sustainability. The D+ birds have a positive impact on economic and environmental traits, with better efficiency, lower consumption, and lower manure production. The two lines were not different on general activity and health traits, apart from a better bone yield and resistance to colibacillosis in D+ line. The D- birds were more sensitive to the diet than D+ birds.