

# To develop with or without the prion protein

Sophie Halliez, Bruno Passet, Séverine Martin-Lannerée, Julia Hernandez-Rapp, Hubert Laude, Sophie Mouillet–richard, Jean-Luc Vilotte, Vincent Béringue

## ► To cite this version:

Sophie Halliez, Bruno Passet, Séverine Martin-Lannerée, Julia Hernandez-Rapp, Hubert Laude, et al.. To develop with or without the prion protein. Frontiers in Cell and Developmental Biology, 2014, 2, pp.1-10. 10.3389/fcell.2014.00058 . hal-01193977

## HAL Id: hal-01193977 https://hal.science/hal-01193977

Submitted on 27 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## To develop with or without the prion protein

Sophie Halliez<sup>1</sup>\*, Bruno Passet<sup>2</sup>, Séverine Martin-Lannerée<sup>3,4</sup>, Julia Hernandez-Rapp<sup>3,4</sup>, Hubert Laude<sup>1</sup>, Sophie Mouillet-Richard<sup>3,4</sup>, Jean-Luc Vilotte<sup>2</sup> and Vincent Béringue<sup>1</sup>\*

<sup>1</sup> Institut National de la Recherche Agronomique, U892 Virologie et Immunologie Moléculaires, Jouy-en-Josas, France

<sup>2</sup> Institut National de la Recherche Agronomique, UMR1313 Génétique Animale et Biologie Intégrative, Jouy-en-Josas, France

<sup>3</sup> Institut National de la Santé et de la Recherche Médicale, UMR-S1124, Paris, France

<sup>4</sup> Université Paris Descartes, Sorbonne Paris Cité, UMR-S1124, Paris, France

#### Edited by:

Kim Newton, Genentech, Inc., USA

#### Reviewed by:

Peter Christian Kloehn, University College London UK Joseph W. Lewcock, Genentech, Inc., USA

#### \*Correspondence:

Sophie Halliez and Vincent Béringue, Institut National de la Recherche Aaronomique, U892 Virologie et Immunologie Moléculaires, Equipe MAP2, Bâtiment 440, 78350 Jouy-en-Josas, France e-mail: shalliez@jouy.inra.fr; vincent.beringue@jouy.inra.fr

The deletion of the cellular form of the prion protein (PrPC) in mouse, goat, and cattle has no drastic phenotypic consequence. This stands in apparent contradiction with PrP<sup>C</sup> quasi-ubiquitous expression and conserved primary and tertiary structures in mammals, and its pivotal role in neurodegenerative diseases such as prion and Alzheimer's diseases. In zebrafish embryos, depletion of PrP ortholog leads to a severe loss-of-function phenotype. This raises the question of a potential role of PrP<sup>C</sup> in the development of all vertebrates. This view is further supported by the early expression of the PrP<sup>C</sup> encoding gene (Prnp) in many tissues of the mouse embryo, the transient disruption of a broad number of cellular pathways in early  $Prnp^{-/-}$  mouse embryos, and a growing body of evidence for PrP<sup>C</sup> involvement in the regulation of cell proliferation and differentiation in various types of mammalian stem cells and progenitors. Finally, several studies in both zebrafish embryos and in mammalian cells and tissues in formation support a role for PrP<sup>C</sup> in cell adhesion, extra-cellular matrix interactions and cytoskeleton. In this review, we summarize and compare the different models used to decipher PrP<sup>C</sup> functions at early developmental stages during embryo- and organo-genesis and discuss their relevance.

Keywords: prion protein, development, neural development, stem cells, cell adhesion, extra-cellular matrix, cytoskeleton

#### **INTRODUCTION**

Prion diseases are a group of fatal and transmissible neurodegenerative diseases affecting a broad range of mammals including humans. The causative agent (the prion) is primarily composed of abnormally folded and aggregated forms of a hostencoded protein, the cellular prion protein (PrP<sup>C</sup>). PrP<sup>C</sup> is a glycosyl-phosphatidyl-inositol anchored cell surface sialoglycoprotein associated with lipid rafts (Taylor et al., 2009). PrP primary sequence is highly conserved among mammals (Wopfner et al., 1999) and PrP putative functional domains are structurally conserved between mammals, avians, and fish (Wopfner et al., 1999; Rivera-Milla et al., 2006). PrP<sup>C</sup> is widely expressed in nearly all the organism, albeit at highest levels in the adult nervous system (Bendheim et al., 1992). Together, these data lend support for an essential role of PrP<sup>C</sup> in mammals and possibly in vertebrates in general. However, the production of mice, goat, and cattle lacking PrP did not lead to any obvious phenotype (Bueler et al., 1992; Manson et al., 1994a; Richt et al., 2007; Yu et al., 2009) except, for mice, a resistance to experimental prion diseases (Bueler et al., 1993; Prusiner et al., 1993; Manson et al., 1994b; Mallucci et al., 2003). Additionally, goats, naturally devoid of PrP<sup>C</sup> due to a nonsense mutation, do not seem to present any abnormal phenotype (Benestad et al., 2012). Subtle behavioral and oxidative stressrelated alterations have been then reported in adult mice devoid of PrP<sup>C</sup> (see Table 1) (Tobler et al., 1996; Wong et al., 2001; Roucou et al., 2004; Meotti et al., 2007; Sanchez-Alavez et al., 2008; Le Pichon et al., 2009; Gadotti et al., 2012) although some of them

are debated (Steele et al., 2007) and some may be related to Prnpflanking genes rather than to PrP<sup>C</sup> absence itself (Nuvolone et al., 2013). However, none of them seems, at first glance, so important to justify the conserved structure and broad expression of PrP<sup>C</sup>. The physiological role of PrP<sup>C</sup> still remains highly uncertain despite more than two decades of research and numerous proposed functions (Nicolas et al., 2009; Martins et al., 2010). To conciliate these discrepant data, it has been hypothesized that PrP<sup>C</sup> function is either dispensable or redundant with that of other proteins. Yet, recent advances, notably in the developmental biology, shed a new light on PrP<sup>C</sup> functions and suggest that, perhaps, the quest for PrP<sup>C</sup> functions has been made at the wrong place and/or at the wrong period of time.

## EARLY DEVELOPMENTAL EXPRESSION OF THE PrP GENES **IN SELECTED VERTEBRATES**

The apparent lack of a phenotype in mice invalidated for PrP<sup>C</sup> sounds at odds with the very early and quasi-ubiquitous expression pattern of the protein at embryonic and postnatal stages, respectively. Expression of the gene encoding PrP-Prnp-is detected in post-implantation embryo from embryonic day (E) 6.5 in extra-embryonic regions (Manson et al., 1992) and from E7.5 (late allantoic bud stage) in cardiac mesoderm (Hidaka et al., 2010). Prnp expression is then observed in the developing central nervous system and heart around E8 before extending rapidly almost to the entire embryo (Tremblay et al., 2007). Prnp may even be expressed earlier on as PrP mRNAs have been

## Table 1 | Phenotypes associated to PrP invalidation/ectopic activation.

Life period/developmental stage	Type of manipulation	Phenotype(s)	Comments	References
Early embryo (zebrafish)	<i>PrP-1</i> knockdown	Lethal – gastrulation arrest due to impaired cell adhesion	<ul> <li>partial rescue of the morphants by the injection of PrP mRNA including mammalian sequence</li> </ul>	Malaga-Trillo et al., 2009
Early embryo (mouse)	Prnp knockout embryos	<b>Moderately severe</b> – transcriptomic analysis shows differential expression of genes from multiple cell pathways	– transient phenotype	Khalife et al., 2011; Passet et al., 2012
Early embryo (mouse)	PrP and Shadoo co-invalidation obtained by: <i>Sprn</i> knockdown and <i>Prnp</i> knockout Double knockout	Controversial <ul> <li>impaired trophoblast</li> <li>development, lethality</li> <li>no phenotype reported but no</li> <li>assessment at embryonic stage</li> </ul>	– discussed in Makhzami et al. (2014)	Young et al., 2009; Passet et al., 2012 Daude et al., 2012
Pharyngula stage (zebrafish)	<i>PrP-2</i> knockdown <i>PrP-2</i> knockout	Controversial – differential expression of genes involved in multiple cell pathways – no obvious developmental abnormalities but no transcriptomic analysis performed	– no phenotype rescue of the morphants	Nourizadeh-Lillabadi et al., 2010 Fleisch et al., 2013
Larva (zebrafish)	<i>PrP-2</i> knockdown <i>PrP-2</i> knockout	Controversial – head malformations, missing neuronal structures – no obvious abnormalities at larval stage	– no phenotype rescue of the morphants	Malaga-Trillo et al., 2009; Nourizadeh-Lillabadi et al., 2010 Fleisch et al., 2013
Late embryo/newborn (mouse)	<i>Prnp</i> knockout embryos	<ul> <li>Minor</li> <li>increased proliferation and maturation delay of the oligodendrocyte precursor cell population in the brain</li> <li>earlier formation of dentin and enamel in the developing tooth</li> </ul>	<ul> <li>no brain abnormalities or myelin defect</li> <li>no enamel defect but reduced hardness of dentin at adult stage</li> </ul>	Bribian et al., 2012 Zhang et al., 2011
Juvenile (mouse)	Prnp knockout mice	Moderately severe – functional abnormalities and persisting cell proliferation in the cerebellum, impaired locomotor abilities	– transient phenotype	Prestori et al., 2008
Adult Brain (mouse)	<i>Prnp</i> knockout mice	<ul> <li>Minor</li> <li>transcriptomic and proteomic analysis revealed no important differences</li> <li>increased protein oxidation, protein ubiquitination and lipid peroxidation</li> <li>increased proliferation and maturation delay of the oligodendrocyte precursor cells</li> </ul>	– no brain abnormalities or myelin defect associated	Crecelius et al., 2008; Chadi et al., 2010 Wong et al., 2001 Bribian et al., 2012

#### Table 1 | Continued

Life period/developmental stage	Type of manipulation	Phenotype(s)	Comments	References
		<ul> <li>decreased cell proliferation in the dentate gyrus (adult neurogenic region)</li> <li>functional abnormalities in the hippocampus</li> </ul>		Steele et al., 2006 Collinge et al., 1994
Adult brain (mouse)	Prnp overexpressing mice	Minor – increased cell proliferation in the subventricular zone (adult neurogenic region) – shorten astrocyte maturation phase	– no brain abnormalities associated	Steele et al., 2006 Hartmann et al., 2013
Adult extraneural tissues (mouse)	<i>Prnp</i> knockout mice	<ul> <li>Minor</li> <li>delayed mineralization of the continuously erupting incisors</li> <li>slower regeneration of muscle fibers</li> <li>shortening of intestinal villi, cell cycle alterations in intestinal crypts and reduced size of desmosomes in intestinal epithelium</li> </ul>	– regeneration and renewing tissues	Zhang et al., 2011; Stella et al., 2010 Morel et al., 2008
Adult (mouse)	<i>Prnp</i> knockout mice	Minor and/or partially controversial – depending of the study, no phenotype observed to minor alterations such as altered olfactory behavior	<ul> <li>few studies were carried out using distinct genetic background</li> </ul>	Bueler et al., 1992; Collinge et al., 1994; Tobler et al., 1996; Walz et al., 1999; Nico et al., 2005; Meotti et al., 2007; Sanchez-Alavez et al., 2008; Le Pichon et al., 2009; Gadotti et al., 2012; Rial et al., 2014
Adult (zebrafish)	<i>PrP-2</i> knockout	Minor – increased susceptibility to a convulsant drug – kinetics alteration of NMDA receptors		Fleisch et al., 2013
Adult (zebrafish)	PrP-1 knockout	Unknown	– no transgenic (inducible) line established	
Aged animal (mouse)	Prnp knockout mice	Minor – behavior alterations		Rial et al., 2009; Massimino et al., 2013
Aged animal nerves (mouse)	<i>Prnp</i> knockout and conditional <i>Prnp</i> knockout	Minor – increased myelin abnormalities in peripheral nerves	– several genetic background analyzed	Bremer et al., 2010

found maternally expressed in the zygotes of *Xenopus laevis* and zebrafish (*Danio rerio*) (van Rosmalen et al., 2006; Malaga-Trillo et al., 2009).

Two PrP homologs, PrP-1, PrP-2 and a more divergent form, PrP-3, have been identified in zebrafish (Cotto et al., 2005;

Rivera-Milla et al., 2006). The expression patterns of the corresponding genes (*PrP-1*, *PrP-2*, and *PrP-3*, respectively) in zebrafish embryos are partially conflicting (Cotto et al., 2005; Malaga-Trillo et al., 2009) although both studies agree on their spatio-temporal complementarity. Both studies describe early PrP gene expression from the mid-blastula stage although it is *PrP-2* according to the Cotto et al. study and *PrP-1* according to Malaga-Trillo et al. At pharyngula stage (24–48 h postfertilization), both studies show a strong and broad expression of *PrP-2* in the developing central nervous system. Additionally, the Cotto et al. study reports the expression of *PrP-1* in the floor plate, an essential organizing center of the central nervous system and in the peripheral nervous system around the same developmental period. The expression of *PrP-2* and *PrP-3* in the developing neuromasts (sensory organs) is also described at the larval stage. PrP-encoding genes are also expressed in developing organs or tissues outside the nervous system such as the heart (*PrP-2* and *PrP-3*), the kidney (*PrP-2*), the pectoral fins (*PrP-2* and *PrP-3*) and the intestinal epithelium (*PrP-2*) (Cotto et al., 2005).

## THE EMBRYO SPILLS THE BEANS

In sharp contrast with the situation in mammals, knockdown (KD) of PrP-1 or PrP-2 by morpholino injection led to severe biological phenotypes in zebrafish (see Table 1): gastrulation arrest for PrP-1 KD (Malaga-Trillo et al., 2009) and malformations of the brain and the eyes associated to a reduced number of peripheral neurons for PrP-2 KD (Malaga-Trillo et al., 2009; Nourizadeh-Lillabadi et al., 2010). However, PrP-2 KDinduced phenotype is subject to caution as PrP-2 knockout (KO) by zinc finger nuclease-induced targeted mutation did not lead to any obvious developmental phenotype (Fleisch et al., 2013). Additional and aspecific effects due to morpholino injection may have occurred. There is less concern about PrP-1 KD phenotype as partial rescue was obtained by the injection of PrP-1 mRNA, PrP-2 mRNA and, remarkably, mouse Prnp mRNA (Malaga-Trillo et al., 2009), strongly suggesting that zebrafish and mammalian PrPs may share conserved functions.

PrP inactivation studies in zebrafish allow investigating PrP<sup>C</sup> functions. They also favor a more critical role of PrP<sup>C</sup> in early development rather than during postnatal stages (see Table 1). Accordingly, transcriptomic analyses (by RNAseq) of  $Prnp^{-/-}$  versus wild-type mouse embryos identified numerous genes differentially expressed (73 and 263 genes at E6.5 and E7.5, respectively) (Khalife et al., 2011), while only 1 gene was found differentially expressed in adult brains (Chadi et al., 2010). Proteomic analysis also failed to identify major alterations (Crecelius et al., 2008) although many cellular stress markers such as protein oxidation, protein ubiquitination, and lipid peroxidation were reportedly activated in the brains of  $Prnp^{-/-}$  versus wild-type mice (Wong et al., 2001). This suggests either that PrP<sup>C</sup> is quite dispensable in the adult brain or that, in a  $Prnp^{-/-}$ context, alternative mechanisms are activated during the brain development to compensate the absence of PrP<sup>C</sup>. Supporting the second hypothesis, the number of genes differentially expressed in the adult  $Prnp^{-/-}$  mouse brain was substantially higher upon postnatal depletion in neurons than embryonically (Chadi et al., 2010), albeit to a less extent than in the embryo. The genes differentially expressed in the embryo covered various cell functions including adhesion, apoptosis and proliferation and are related to heart formation and blood vessel development, vascular diseases, immune response, nervous system development, and prion diseases (Khalife et al., 2011). Interestingly and in line with the idea of evolutionary conserved functions of PrPs, similar biological functions were impacted by *PrP-2* KD in zebrafish, as revealed by transcriptomic analyses: development including that of the cardiovascular and nervous systems specifically, cell death, cellular assembly, cell cycle, immunological, and neurological diseases (Nourizadeh-Lillabadi et al., 2010).

Finally, the KD of the *Sprn* gene encoding a PrP-related protein, Shadoo resulted in embryonic lethality between E8 and E11 specifically in  $Prnp^{-/-}$  embryos, supporting a role for  $PrP^{C}$ and Shadoo in mouse embryogenesis, notably cell adhesion and placenta development, and a possible functional redundancy between the two proteins (Young et al., 2009; Passet et al., 2012). However, the recent generation of viable  $Prnp^{-/-}$ ;  $Sprn^{-/-}$  animals is challenging this view (Daude et al., 2012). These apparently conflicting observations are discussed in Makhzami et al. (2014).

Thus, studying the consequences of PrP inactivation during developmental stages rather than in adult animals could be a more efficient, although not easier, strategy to decipher PrP<sup>C</sup> functions. Additionally, studying PrP<sup>C</sup> implications in regenerative processes and in renewing tissues could also be an informative approach (see **Table 1**).

## THE CURRENT LIMITS OF IN VIVO STUDIES IN MAMMALS

As PrP<sup>C</sup> is mediating neuronal dysfunction and degeneration in prion diseases and reportedly in Alzheimer's disease (Lauren et al., 2009; Klyubin et al., 2014), it has been logically suspected to play a role in neuronal homeostasis and during neural development. As PrP<sup>C</sup> functions in mature neurons are not under the scope of the present review, we will not discuss them and only focus on developmental aspects. In vivo direct evidence supporting a role of PrP<sup>C</sup> in neural development is quite elusive and PrP<sup>C</sup> expression pattern remains one of the strongest pro arguments. Indeed, PrP<sup>C</sup> expression increases as neurons mature, although it is not detected in mitotically active neural progenitor cells (Steele et al., 2006; Peralta et al., 2011). Other experimental evidence was obtained from the comparative study of WT and  $Prnp^{-/-}$ and/or Prnp overexpressing animals (see Table 1). The levels of PrP<sup>C</sup> were found to positively regulate the proliferation in neurogenic regions in adult mice (Steele et al., 2006; Bribian et al., 2012) and the maturation of astrocytes in the developing brain (Arantes et al., 2009; Hartmann et al., 2013). Conversely, the absence of PrP<sup>C</sup> was associated with an increased proliferation in oligodendrocyte precursor cells both in embryos and adults (Bribian et al., 2012). However, no brain morphology abnormalities or myelin defects could be observed in  $Prnp^{-/-}$  animals, casting doubt on a crucial role of PrP<sup>C</sup> in cell proliferation and maturation during brain development or, at least, suggesting the occurrence of compensatory mechanisms. Yet, electrophysiological recordings on hippocampus from  $Prnp^{-/-}$  mice (Collinge et al., 1994) and on cerebellum from juvenile  $Prnp^{-/-}$  mice (Prestori et al., 2008) revealed functional alterations. However, no hippocampus-related behavioral or learning alterations were observed in Prnp<sup>-/-</sup> mice (Bueler et al., 1992) suggesting the hippocampus functional alterations are minor or at least compensated. Conversely, cerebellum functional alterations were found associated to a failure of juvenile  $Prnp^{-/-}$  mice in motor control

test and to protracted cell proliferation in the cerebellum granular layer at the third week of postnatal development although all these abnormalities vanished in older animals (Prestori et al., 2008). Taken together, these arguments suggest a delay in the maturation of the cerebellar granule cells in  $Prnp^{-/-}$  mice.

Further dissecting the underlying mechanisms responsible for the subtle phenotypes in  $Prnp^{-/-}$  mice is a particularly arduous task *in vivo*. For example, determining whether the effects are cellautonomous would require the use of several transgenic lines or genetic chimeras to limit the expression of  $PrP^{C}$  to a subset of cells. And this question is far from being irrelevant as  $PrP^{C}$  is a cell surface protein. Such topology could elicit specific cell responses in neighboring cells. Additionally,  $PrP^{C}$  may act at distance since secreted forms of have been described both *in vitro* and *in vivo* (Borchelt et al., 1993; Harris et al., 1993).

### CONTRIBUTIONS OF *IN VITRO* AND *EX VIVO* MODELS TO UNDERSTANDING THE ROLE OF PrP<sup>C</sup> IN NEURAL DEVELOPMENT

The use of in vitro or ex vivo models has generated a wealth of data with regard to the involvement of PrP<sup>C</sup> in neural development and, importantly, has comforted the putative functions suggested by the in vivo approaches. Moreover they often allow pinpointing more precisely the cellular processes and signaling pathways involving PrP<sup>C</sup>. Collectively, these experimental data argue that PrP<sup>C</sup> positively regulates (1) the differentiation of pluripotent progenitors cells toward the neural lineage (Peralta et al., 2011), (2) the proliferation of neural progenitor cells (Prodromidou et al., 2014) (3) the self-renewal of neural progenitor cells (Santos et al., 2011; Prodromidou et al., 2014) (4) the neuronal differentiation in terms of choice of cell fate (Steele et al., 2006) and (5) acquisition of neuronal features (Graner et al., 2000). However, how PrP<sup>C</sup> fulfills these different actions is only partially understood. The current idea is that PrP<sup>C</sup> is part of multiprotein signaling complexes, able to bind various partners, and participate to signal transduction events along different cellular pathways depending on the cellular context (Caughey and Baron, 2006; Watts et al., 2009; Stuermer, 2011; Hirsch et al., 2014). Non-exhaustively, the interaction between PrP<sup>C</sup> and one of its identified ligand, Stress inducible protein 1 (STI1), has been shown to promote the self-renewal of the neural progenitor cells (Santos et al., 2011). Besides, interaction with the neural cell adhesion molecule (NCAM) can induce their neuronal differentiation (Prodromidou et al., 2014). In cultured neurons, STI1 binding to PrP<sup>C</sup> increases protein synthesis (Roffe et al., 2010), STI1 and Laminin-γ1 binding promotes intracellular Ca<sup>2+</sup> increase and axonogenesis (Santos et al., 2013), and vitronectin binding, axonal growth (Hajj et al., 2007). Experimental evidence also favors a role of PrP<sup>C</sup> in glia development since the lack of PrP<sup>C</sup> promotes the proliferation of oligodendrocyte precursor cells and delays their differentiation in culture (Bribian et al., 2012). Studies in astrocyte primary cultures revealed that the involvement of PrP<sup>C</sup> in astrocytes maturation is dependent upon its interaction with STI1 and the downstream activation of ERK1/2 (Arantes et al., 2009). Of note, PrP<sup>C</sup> action does not necessarily rely on PrP<sup>C</sup> expression solely in the cell population considered. For instance, proper neuritogenesis has been shown

to require  $PrP^{C}$  expression in both neurons and surrounding astrocytes in co-cultures experiments (Lima et al., 2007). Beyond a role in the interactions between both populations,  $PrP^{C}$  serves as a neuronal receptor for laminin (Graner et al., 2000) and is also involved in the organization of the extracellular matrix by the astrocytes, especially through laminin deposition (Lima et al., 2007).

It should also be stressed that  $PrP^{C}$  partners and the cellular context could tightly determine the final consequences of  $PrP^{C}$  stimulation. For example  $PrP^{C}$  interaction with contactinassociated protein (Caspr) protects Caspr from proteolysis allowing it to inhibit more efficiently neurite outgrowth (Devanathan et al., 2010). On the opposite,  $PrP^{C}$  and NCAM interactions allow stabilization of NCAM in lipid rafts allowing Fyn kinase activation and promoting neurite outgrowth (Santuccione et al., 2005).

*In vitro*, PrP<sup>C</sup> is thereby found involved in all the major steps of neural development. Since it notably regulates the self-renewal and differentiation processes in neural progenitor cells, the question arises as to whether this role can be extended to other cell types.

## WHAT THE STUDY OF STEM CELLS REVEALS

The role of PrP<sup>C</sup> is far from being restricted to neural development and a growing body of evidence support its involvement in stem cell biology see also Martin-Lannerée et al. (2014). Whether PrP<sup>C</sup> is expressed in embryonic stem cells (ESC) at basal state remains unclear (Lee and Baskakov, 2010; Miranda et al., 2011), possibly due to limitations in the sensitivity of the detection methods, choice of the target (i.e., the gene or the protein itself) and methods of production and culture of the cells. Nevertheless, there is a consensus that PrP<sup>C</sup> expression progressively increases during stem cell differentiation (Miranda et al., 2011; Lee and Baskakov, 2013). To understand the role played by PrP<sup>C</sup> during early embryogenesis, the consequences of Prnp KD or overexpression were studied in human ESC (Lee and Baskakov, 2013). Forcing PrP<sup>C</sup> expression under self-renewal conditions was found to alter cell cycle regulation and to promote ESC differentiation. Blocking PrP<sup>C</sup> expression in differentiating ESC also impacts on cell cycle regulation and inhibits the differentiation toward ectodermal lineages, in line with PrP<sup>C</sup> functions in neural differentiation (Peralta et al., 2011) and the reduced expression of Nestin, an ectodermal marker, observed in ESC in the absence of PrP<sup>C</sup> (Miranda et al., 2011; Peralta et al., 2011). Mesodermal and endodermal differentiations were, however, not affected. Finally, overexpressing PrP<sup>C</sup> in differentiating ESC promotes proliferation and inhibits differentiation toward all the lineages (Lee and Baskakov, 2013).

Further supporting  $PrP^C$  function in stem cell biology,  $PrP^C$  is expressed in hematopoietic stem cells and promotes their selfrenewal (Zhang et al., 2006).  $PrP^C$  is therefore involved in the self-renewal and differentiation of stem cells although its role and requirement seem to vary along the differentiation process and the cell lineage considered. Identifying the molecular mechanisms at play is a particularly challenging task since self-renewal, differentiation and proliferation are highly intertwined processes impacting each other.

## THE ROLE OF PrP<sup>C</sup> IN DEVELOPING AND RENEWING TISSUES OUTSIDE THE NERVOUS SYSTEM

Looking outside the central nervous system in newborn to full-grown animals allowed pinpointing subtle phenotypes in the absence of  $PrP^C$  (see **Table 1**), in perfect consistency with  $PrP^C$  widespread expression.  $PrP^C$  is expressed in developing human and murine teeth and study of dental cell cultures and teeth from  $Prnp^{-/-}$  animals revealed multiple alterations during tooth formation (Schneider et al., 2007; Zhang et al., 2011). *In vitro*, embryonic dental mesenchymal cells from  $Prnp^{-/-}$  embryos proliferate more rapidly and differentiate earlier and coherently,  $Prnp^{-/-}$  newborn mice exhibit earlier formation of mesenchymally- and epithelially-derived dentin and enamel, respectively (Zhang et al., 2011).

 $PrP^{C}$  is also abundantly expressed in muscles and, although no abnormalities could be observed in a physiological context, it is involved in the regeneration of skeletal muscles in adult mice. Regeneration of locally damaged muscle fibers occurs at slower pace in  $Prnp^{-/-}$  animals. Yet total recovery is finally achieved (Stella et al., 2010). This is associated with a longer phase of proliferation and a delayed maturation of muscle precursors, likely due to a reduced level of released myogenic factors (Stella et al., 2010). However, the identity of the releasing cells is difficult to establish, showing again the limits of the *in vivo* paradigm.

Another extra-neural tissue where  $PrP^{C}$  depletion induces alterations is the intestinal epithelium, a constantly renewing tissue.  $PrP^{C}$  is expressed by human enterocytes (Morel et al., 2004) and  $Prnp^{-/-}$  animals exhibit shorter intestinal villi associated to an increased number of mitotic cells (Morel et al., 2008).

Collectively, these observations support a role of PrP<sup>C</sup> in the regulation of cell cycle and cell differentiation extraneurally. But how can these functions be exerted?

## PrP<sup>C</sup> REGULATES CELLULAR ADHESION, EXTRA-CELLULAR MATRIX INTERACTIONS, AND CYTOSKELETON REMODELING

PrP-1 KD-induced gastrulation arrest in zebrafish embryos is due to impaired morphogenetic cell movements. This defect is, at least partially, caused by the disruption of cellular adhesion (Malaga-Trillo et al., 2009). The study revealed that PrP-1 normally accumulates at cell-cell contacts at early embryonic stages and regulates cell adhesion, including E-cadherin processing and/or storage, in a cell-autonomous way. Such functions may be conserved in mammals as injection of mouse mRNA Prnp partially rescued the PrP-1 KD-induced phenotype (Malaga-Trillo et al., 2009). Consistent with this idea, a number of experimental evidence have linked PrP<sup>C</sup> to cell adhesion and intimately associated processes such as cytoskeleton remodeling and interactions with extra-cellular matrix (Petit et al., 2013): (i) the expression pattern of PrP<sup>C</sup>. PrP<sup>C</sup> is expressed at the cell-to-cell contacts, growth cone, and other extending process tips in cultured neural progenitors and neurons (Santuccione et al., 2005; Devanathan et al., 2010; Miyazawa et al., 2010), cell-to-cell junctions in human intestinal epithelium villi (Morel et al., 2004), cell-to-cell contacts in primary cultured endothelial cells (Viegas et al., 2006), focal adhesions in HeLa cell lines (Schrock et al., 2009) and the apical face of ameloblasts of developing teeth (Zhang et al.,

2011); (ii) relevant PrP<sup>C</sup> partners and ligands, such as cell junction proteins including integrins, the cytoskeleton protein actin and components of the extra-cellular matrix (Graner et al., 2000; Schmitt-Ulms et al., 2001; Morel et al., 2004, 2008; Nieznanski et al., 2005; Hajj et al., 2007; Watts et al., 2009); (iii) alterations in cell adhesion, cytoskeleton dynamics, and extra-cellular matrix interactions upon ectopic expression or deletion of PrP<sup>C</sup> in different cell systems. Prnp KD leads to alterations in the actin cytoskeleton, remodeling of focal adhesions and over-production of fibronectin in neuronal progenitors (Loubet et al., 2012), to disruption of adherens junctions in carcinoma cells (Solis et al., 2012) and to miss-localization of cell junction-related proteins in enterocytes (Morel et al., 2008). As for Prnp overexpression, it promotes aggregation and filopodia formation as well as alterations of the focal adhesion dynamics in neuroblastoma N2a cells (Mange et al., 2002; Schrock et al., 2009). Subtle changes were even noticed in vivo such as a decrease in the length of desmosomes in the intestinal epithelium of  $Prnp^{-/-}$  mice (Morel et al., 2008).

Interestingly, the alterations induced by the deletion of PrP<sup>C</sup> can be partially rescued by neutralizing antibodies to specific integrins: use of  $\beta$ 1 and  $\beta$ 5 integrins was found to restore neuritogenesis in differentiating PrP-deficient neuronal progenitors (Loubet et al., 2012) and *Nanog* expression in differentiating PrP-deficient ESC (Miranda et al., 2011), respectively. Moreover, modulation of integrin activity was suspected to be a spontaneous compensatory mechanism occurring in the absence of PrP<sup>C</sup> (Hajj et al., 2007). This last observation supports an involvement of PrP<sup>C</sup> in extra-cellular matrix interactions and suggests a possible adaptive mechanism to the absence of PrP<sup>C</sup>, which may account for the lack of drastic phenotypes in *Prnp<sup>-/-</sup>* animals.

Taken together, these observations demonstrate a clear involvement of  $PrP^{C}$  in the regulation of cell adhesion, the interactions with the extra-cellular matrix and the cytoskeleton dynamics. The links between these functions and the role of  $PrP^{C}$  in the regulation of cell proliferation and differentiation have been the focus of few studies (Miranda et al., 2011; Loubet et al., 2012). Are they under-estimated? And are these functions still ongoing in adult organism?

## **STUDYING PrP<sup>C</sup> FUNCTIONS DURING AGING**

Little attention has been devoted to the potential functions of PrP<sup>C</sup> during aging (see Table 1 and also Gasperini and Legname, 2014). It is commonly accepted that the cellular machinery deteriorates or, at least, evolves with aging. As the potential development of adaptive mechanisms to the absence of PrP<sup>C</sup> is assumed to complicate the study of  $PrP^{C}$  functions in  $Prnp^{-/-}$  mice, the study of aged animals could be a way to bypass such difficulty: because of the age-related alterations accumulation, the tolerance to PrP<sup>C</sup> depletion could partially break. In line with this scenario, behavioral differences between WT and Prnp<sup>-/-</sup> mice seem more pronounced when performed on aged animals (Rial et al., 2009; Massimino et al., 2013). Age-related physiological traits can also be found amplified in absence of PrP<sup>C</sup>. Myelin abnormalities in the peripheral nervous system accumulate with aging (Verdu et al., 2000) and sciatic nerves from moderately aged mice devoid of PrP<sup>C</sup> (in selected or all cell populations) showed

highly precocious and increased demyelination in the absence of neuronal  $PrP^C$  (Bremer et al., 2010). If the axons-glia interactions are known to play a key role in myelination during development (Sherman and Brophy, 2005), their role in myelin maintenance in adulthood is less well-understood. However, myelin maintenance is clearly established as an active process (Bremer et al., 2011). The early demyelination observed in the absence of neuronal  $PrP^C$  suggests axonal  $PrP^C$  participates in myelin maintenance and is likely required, directly or indirectly, in axons-Schwann cells communication.

Thus, aging may perhaps be the ultimate stage to unravel the mystery of  $PrP^{C}$  function(s). However, it has to be considered that aging could also impact on  $PrP^{C}$  and/or its function(s). For instance, the age-related change in the cholesterol/sphingolipids ratio in lipid rafts is suspected to alter  $PrP^{C}$  compartmentalization (Agostini et al., 2013), which could ultimately impact on its functions. A functional study of  $PrP^{C}$  during aging could therefore be difficult to analyse due to the numerous changes occurring at the same time.

#### **CONCLUDING REMARKS**

A major function of PrP<sup>C</sup>, consistent with its cell surface expression consists in the modulation of signaling pathways in response to various cues. The panel of putative PrP<sup>C</sup> ligands is quite large, including soluble ligands, extra-cellular matrix components, and adhesion molecules. Their modes of interaction with PrP<sup>C</sup> are variable, including cooperation (Santos et al., 2013) and both cis- and trans-interactions for a same ligand (Santuccione et al., 2005). The cellular context can strongly influence the composition of the PrP<sup>C</sup>-multiprotein complexes and the sub-cellular localization of PrP<sup>C</sup> is highly dynamic: it can cycle rapidly from the plasma membrane to internal membranes (Griffiths et al., 2007) possibly in response to a given stimulus (Lee et al., 2001; Brown and Harris, 2003; Caetano et al., 2008). All these elements taken together may allow a specific response to a given signal with regard to the cell type and differentiation state. This may explain why PrP<sup>C</sup> appears to be involved in a multitude of functions. Regarding development, this view perfectly fits the requirement of different stem and progenitor cells by eliciting appropriate cell responses to adapt efficiently to different microenvironments. But how PrP<sup>C</sup> precisely impacts on stemness maintenance and differentiation processes remains unresolved. The potential implication of PrP<sup>C</sup> in cell polarization and cytoskeleton remodeling in differentiating stem cells is a very attractive hypothesis and should obviously deserve further investigation. Cell polarity and cytoskeleton dynamics are life-long important cell processes, notably during cell fate determination. They have been studied more particularly during drosophilia neurogenesis and in mammalian neuroepithelial cells (Fietz and Huttner, 2011), but they are fundamental processes for all sort of stem and progenitor cells such as hematopoietic progenitor cells (Bullock et al., 2007). Our unpublished data (Halliez et al., in preparation) reveal a polarized expression of PrP<sup>C</sup> in neural and cardiovascular progenitor cells of the developing mouse embryo and therefore are compatible with such idea.

If PrP<sup>C</sup> is involved in so many cell lineages, understanding the lack of drastic phenotype in PrP<sup>C</sup>-depleted animals remains an

ongoing issue. Yet, a modulation of the cell maturation phase (generally delayed but also shortened) is consistently observed in different models (Prnp KO but also KD and overexpression) and may therefore be the general price to pay for the loss of PrP<sup>C</sup> (see **Table 1**). This may have a more selective impact on free-range than on laboratory animals (for example acquisition of perfect motor control) and justify the high degree of PrP conservation through evolution. Moreover, building an organism is a highly complex task requiring spatial and temporal finetuning of a broad number of biological processes via multiple regulatory networks. Cells, tissues and organs do not develop separately along each other but rely on multiple interactions to develop in a concerted, interconnected, and time-scheduled manner. Partial redundancies into and between signaling pathways ensure robustness to the system. That two or more proteins may exert overlapping functions during development and therefore allow tolerance to the loss of one of them is not unprecedented (Yoon et al., 2005; Santamaria and Ortega, 2006; Nicolae et al., 2007). Moreover, it is not unheard-of, notably in plants where environment adaptation is largely studied, that redundant genes in a regular context allow survival under environmental modifications and stress (Wu et al., 2002). Interestingly, PrP KO animals seem to react differentially than WT animals to stressful situations (Nico et al., 2005) and to alcohol (Rial et al., 2014) and show increased susceptibility to convulsant agents (Walz et al., 1999; Fleisch et al., 2013) (see Table 1). The general function of PrP<sup>C</sup> could be to facilitate responses to external stimulus/factors at the cell- and at the organism-scale. Yet, laboratory animals are raised in standardized and protective environment and are often homogenous genetically. In such conditions, the time window to apprehend the cell specific functions of PrP<sup>C</sup> may be restricted to the developmental and perinatal phases, with so much information provided and so much transformation occurring at the same time.

## REFERENCES

- Agostini, F., Dotti, C. G., Perez-Canamas, A., Ledesma, M. D., Benetti, F., and Legname, G. (2013). Prion protein accumulation in lipid rafts of mouse aging brain. *PLoS ONE* 8:e74244. doi: 10.1371/journal.pone.0074244
- Arantes, C., Nomizo, R., Lopes, M. H., Hajj, G. N., Lima, F. R., and Martins, V. R. (2009). Prion protein and its ligand stress inducible protein 1 regulate astrocyte development. *Glia* 57, 1439–1449. doi: 10.1002/glia.20861
- Bendheim, P. E., Brown, H. R., Rudelli, R. D., Scala, L. J., Goller, N. L., Wen, G. Y., et al. (1992). Nearly ubiquitous tissue distribution of the scrapie agent precursor protein. *Neurology* 42, 149–156. doi: 10.1212/WNL.42.1.149
- Benestad, S. L., Austbo, L., Tranulis, M. A., Espenes, A., and Olsaker, I. (2012). Healthy goats naturally devoid of prion protein. *Vet. Res.* 43, 87. doi: 10.1186/1297-9716-43-87
- Borchelt, D. R., Rogers, M., Stahl, N., Telling, G., and Prusiner, S. B. (1993). Release of the cellular prion protein from cultured cells after loss of its glycoinositol phospholipid anchor. *Glycobiology* 3, 319–329. doi: 10.1093/glycob/ 3.4.319
- Bremer, J., Baumann, F., Tiberi, C., Wessig, C., Fischer, H., Schwarz, P., et al. (2010). Axonal prion protein is required for peripheral myelin maintenance. *Nat. Neurosci.* 13, 310–318. doi: 10.1038/nn.2483
- Bremer, M., Frob, F., Kichko, T., Reeh, P., Tamm, E. R., Suter, U., et al. (2011). Sox10 is required for Schwann-cell homeostasis and myelin maintenance in the adult peripheral nerve. *Glia* 59, 1022–1032. doi: 10.1002/glia.21173
- Bribian, A., Fontana, X., Llorens, F., Gavin, R., Reina, M., Garcia-Verdugo, J. M., et al. (2012). Role of the cellular prion protein in oligodendrocyte precursor cell proliferation and differentiation in the developing and adult mouse CNS. *PLoS ONE* 7:e33872. doi: 10.1371/journal.pone.0033872

- Brown, L. R., and Harris, D. A. (2003). Copper and zinc cause delivery of the prion protein from the plasma membrane to a subset of early endosomes and the Golgi. J. Neurochem. 87, 353–363. doi: 10.1046/j.1471-4159.2003.01996.x
- Bueler, H., Aguzzi, A., Sailer, A., Greiner, R. A., Autenried, P., Aguet, M., et al. (1993). Mice devoid of PrP are resistant to scrapie. *Cell* 73, 1339–1347. doi: 10.1016/0092-8674(93)90360-3
- Bueler, H., Fischer, M., Lang, Y., Bluethmann, H., Lipp, H. P., DeArmond, S. J., et al. (1992). Normal development and behaviour of mice lacking the neuronal cell-surface PrP protein. *Nature* 356, 577–582. doi: 10.1038/356577a0
- Bullock, T. E., Wen, B., Marley, S. B., and Gordon, M. Y. (2007). Potential of CD34 in the regulation of symmetrical and asymmetrical divisions by hematopoietic progenitor cells. *Stem Cells* 25, 844–851. doi: 10.1634/stemcells.2006-0346
- Caetano, F. A., Lopes, M. H., Hajj, G. N., Machado, C. F., Pinto Arantes, C., Magalhaes, A. C., et al. (2008). Endocytosis of prion protein is required for ERK1/2 signaling induced by stress-inducible protein 1. J. Neurosci. 28, 6691–6702. doi: 10.1523/JNEUROSCI.1701-08.2008
- Caughey, B., and Baron, G. S. (2006). Prions and their partners in crime. *Nature* 443, 803–810. doi: 10.1038/nature05294
- Chadi, S., Young, R., Le Guillou, S., Tilly, G., Bitton, F., Martin-Magniette, M. L., et al. (2010). Brain transcriptional stability upon prion protein-encoding gene invalidation in zygotic or adult mouse. *BMC Genomics* 11:448. doi: 10.1186/1471-2164-11-448
- Collinge, J., Whittington, M. A., Sidle, K. C., Smith, C. J., Palmer, M. S., Clarke, A. R., et al. (1994). Prion protein is necessary for normal synaptic function. *Nature* 370, 295–297. doi: 10.1038/370295a0
- Cotto, E., Andre, M., Forgue, J., Fleury, H. J., and Babin, P. J. (2005). Molecular characterization, phylogenetic relationships, and developmental expression patterns of prion genes in zebrafish (*Danio rerio*). *FEBS J.* 272, 500–513. doi: 10.1111/j.1742-4658.2004.04492.x
- Crecelius, A. C., Helmstetter, D., Strangmann, J., Mitteregger, G., Frohlich, T., Arnold, G. J., et al. (2008). The brain proteome profile is highly conserved between Prnp-/- and Prnp+/+ mice. *Neuroreport* 19, 1027–1031. doi: 10.1097/WNR.0b013e3283046157
- Daude, N., Wohlgemuth, S., Brown, R., Pitstick, R., Gapeshina, H., Yang, J., et al. (2012). Knockout of the prion protein (PrP)-like Sprn gene does not produce embryonic lethality in combination with PrP(C)-deficiency. *Proc. Natl. Acad. Sci. U.S.A.* 109, 9035–9040. doi: 10.1073/pnas.1202130109
- Devanathan, V., Jakovcevski, I., Santuccione, A., Li, S., Lee, H. J., Peles, E., et al. (2010). Cellular form of prion protein inhibits Reelin-mediated shedding of Caspr from the neuronal cell surface to potentiate Caspr-mediated inhibition of neurite outgrowth. J. Neurosci. 30, 9292–9305. doi: 10.1523/JNEUROSCI.5657-09.2010
- Fietz, S. A., and Huttner, W. B. (2011). Cortical progenitor expansion, self-renewal and neurogenesis-a polarized perspective. *Curr. Opin. Neurobiol.* 21, 23–35. doi: 10.1016/j.conb.2010.10.002
- Fleisch, V. C., Leighton, P. L., Wang, H., Pillay, L. M., Ritzel, R. G., Bhinder, G., et al. (2013). Targeted mutation of the gene encoding prion protein in zebrafish reveals a conserved role in neuron excitability. *Neurobiol. Dis.* 55, 11–25. doi: 10.1016/j.nbd.2013.03.007
- Gadotti, V. M., Bonfield, S. P., and Zamponi, G. W. (2012). Depressive-like behaviour of mice lacking cellular prion protein. *Behav. Brain Res.* 227, 319–323. doi: 10.1016/j.bbr.2011.03.012
- Gasperini, L., and Legname, G. (2014). Prion protein and aging. Front. Cell Dev. Biol. 2:44. doi: 10.3389/fcell.2014.00044
- Graner, E., Mercadante, A. F., Zanata, S. M., Forlenza, O. V., Cabral, A. L., Veiga, S. S., et al. (2000). Cellular prion protein binds laminin and mediates neuritogenesis. *Brain Res. Mol. Brain Res.* 76, 85–92. doi: 10.1016/S0169-328X(99) 00334-4
- Griffiths, R. E., Heesom, K. J., and Anstee, D. J. (2007). Normal prion protein trafficking in cultured human erythroblasts. *Blood* 110, 4518–4525. doi: 10.1182/blood-2007-04-085183
- Hajj, G. N., Lopes, M. H., Mercadante, A. F., Veiga, S. S., da Silveira, R. B., Santos, T. G., et al. (2007). Cellular prion protein interaction with vitronectin supports axonal growth and is compensated by integrins. *J. Cell Sci.* 120, 1915–1926. doi: 10.1242/jcs.03459
- Harris, D. A., Huber, M. T., van Dijken, P., Shyng, S. L., Chait, B. T., and Wang, R. (1993). Processing of a cellular prion protein: identification of N- and C-terminal cleavage sites. *Biochemistry* 32, 1009–1016. doi: 10.1021/bi00055a003

- Hartmann, C. A., Martins, V. R., and Lima, F. R. (2013). High levels of cellular prion protein improve astrocyte development. *FEBS Lett.* 587, 238–244. doi: 10.1016/j.febslet.2012.11.032
- Hidaka, K., Shirai, M., Lee, J. K., Wakayama, T., Kodama, I., Schneider, M. D., et al. (2010). The cellular prion protein identifies bipotential cardiomyogenic progenitors. *Circ. Res.* 106, 111–119. doi: 10.1161/CIRCRESAHA.109. 209478
- Hirsch, T. Z., Hernandez-Rapp, J., Martin-Lannerée, S., Launay, J. M., and Mouillet-Richard, S. (2014). PrP signalling in neurons: from basics to clinical challenges. *Biochimie*. 104, 2–11. doi: 10.1016/j.biochi.2014.06.009
- Khalife, M., Young, R., Passet, B., Halliez, S., Vilotte, M., Jaffrezic, F., et al. (2011). Transcriptomic analysis brings new insight into the biological role of the prion protein during mouse embryogenesis. *PLoS ONE* 6:e23253. doi: 10.1371/journal.pone.0023253
- Klyubin, I., Nicoll, A. J., Khalili-Shirazi, A., Farmer, M., Canning, S., Mably, A., et al. (2014). Peripheral administration of a humanized anti-PrP antibody blocks Alzheimer's disease abeta synaptotoxicity. *J. Neurosci.* 34, 6140–6145. doi: 10.1523/JNEUROSCI.3526-13.2014
- Lauren, J., Gimbel, D. A., Nygaard, H. B., Gilbert, J. W., and Strittmatter, S. M. (2009). Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. *Nature* 457, 1128–1132. doi: 10.1038/nature 07761
- Lee, K. S., Magalhaes, A. C., Zanata, S. M., Brentani, R. R., Martins, V. R., and Prado, M. A. (2001). Internalization of mammalian fluorescent cellular prion protein and N-terminal deletion mutants in living cells. *J. Neurochem.* 79, 79–87. doi: 10.1046/j.1471-4159.2001.00529.x
- Lee, Y. J., and Baskakov, I. V. (2010). Treatment with normal prion protein delays differentiation and helps to maintain high proliferation activity in human embryonic stem cells. *J. Neurochem.* 114, 362–373. doi: 10.1111/j.1471-4159.2010.06601.x
- Lee, Y. J., and Baskakov, I. V. (2013). The cellular form of the prion protein is involved in controlling cell cycle dynamics, self-renewal, and the fate of human embryonic stem cell differentiation. J. Neurochem. 124, 310–322. doi: 10.1111/j.1471-4159.2012.07913.x
- Le Pichon, C. E., Valley, M. T., Polymenidou, M., Chesler, A. T., Sagdullaev, B. T., Aguzzi, A., et al. (2009). Olfactory behavior and physiology are disrupted in prion protein knockout mice. *Nat. Neurosci.* 12, 60–69. doi: 10.1038/nn.2238
- Lima, F. R., Arantes, C. P., Muras, A. G., Nomizo, R., Brentani, R. R., and Martins, V. R. (2007). Cellular prion protein expression in astrocytes modulates neuronal survival and differentiation. *J. Neurochem.* 103, 2164–2176. doi: 10.1111/j.1471-4159.2007.04904.x
- Loubet, D., Dakowski, C., Pietri, M., Pradines, E., Bernard, S., Callebert, J., et al. (2012). Neuritogenesis: the prion protein controls beta1 integrin signaling activity. *FASEB J.* 26, 678–690. doi: 10.1096/fj.11-185579
- Makhzami, S., Passet, B., Halliez, S., Castille, J., Moazami-Goudarzi, K., Duchesne, A., et al. (2014). The prion protein family: a view from the placenta. *Front. Cell Dev. Biol.* 2:35. doi: 10.3389/fcell.2014.00035
- Malaga-Trillo, E., Solis, G. P., Schrock, Y., Geiss, C., Luncz, L., Thomanetz, V., et al. (2009). Regulation of embryonic cell adhesion by the prion protein. *PLoS Biol.* 7:e55. doi: 10.1371/journal.pbio.1000055
- Mallucci, G., Dickinson, A., Linehan, J., Klohn, P. C., Brandner, S., and Collinge, J. (2003). Depleting neuronal PrP in prion infection prevents disease and reverses spongiosis. *Science* 302, 871–874. doi: 10.1126/science.1090187
- Mange, A., Milhavet, O., Umlauf, D., Harris, D., and Lehmann, S. (2002). PrPdependent cell adhesion in N2a neuroblastoma cells. *FEBS Lett.* 514, 159–162. doi: 10.1016/S0014-5793(02)02338-4
- Manson, J. C., Clarke, A. R., Hooper, M. L., Aitchison, L., McConnell, I., and Hope, J. (1994a). 129/Ola mice carrying a null mutation in PrP that abolishes mRNA production are developmentally normal. *Mol. Neurobiol.* 8, 121–127. doi: 10.1007/BF02780662
- Manson, J. C., Clarke, A. R., McBride, P. A., McConnell, I., and Hope, J. (1994b). PrP gene dosage determines the timing but not the final intensity or distribution of lesions in scrapie pathology. *Neurodegeneration* 3, 331–340.
- Manson, J., West, J. D., Thomson, V., McBride, P., Kaufman, M. H., and Hope, J. (1992). The prion protein gene: a role in mouse embryogenesis? *Development* 115, 117–122.
- Martin-Lannerée, S., Hirsch, T. Z., Hernandez-Rapp, J., Halliez, S., Vilotte, J.-L., Launay, J.-M., et al. (2014). PrP<sup>C</sup> from stem cells to cancer. *Front. Cell Dev. Biol.* 2:55. doi: 10.3389/fcell.2014.00055

- Martins, V. R., Beraldo, F. H., Hajj, G. N., Lopes, M. H., Lee, K. S., Prado, M. A., et al. (2010). Prion protein: orchestrating neurotrophic activities. *Curr. Issues Mol. Biol.* 12, 63–86.
- Massimino, M. L., Redaelli, M., Bertoli, A., Sorgato, M. C., and Mucignat-Caretta, C. (2013). Altered behavioral aspects of aged mice lacking the cellular prion protein. *Physiol. Behav.* 119, 86–91. doi: 10.1016/j.physbeh.2013.06.006
- Meotti, F. C., Carqueja, C. L., Gadotti Vde, M., Tasca, C. I., Walz, R., and Santos, A. R. (2007). Involvement of cellular prion protein in the nociceptive response in mice. *Brain Res.* 1151, 84–90. doi: 10.1016/j.brainres.2007.03.024
- Miranda, A., Pericuesta, E., Ramirez, M. A., and Gutierrez-Adan, A. (2011). Prion protein expression regulates embryonic stem cell pluripotency and differentiation. *PLoS ONE* 6:e18422. doi: 10.1371/journal.pone.0018422
- Miyazawa, K., Emmerling, K., and Manuelidis, L. (2010). Proliferative arrest of neural cells induces prion protein synthesis, nanotube formation, and cell-to-cell contacts. J. Cell. Biochem. 111, 239–247. doi: 10.1002/jcb.22723
- Morel, E., Fouquet, S., Chateau, D., Yvernault, L., Frobert, Y., Pincon-Raymond, M., et al. (2004). The cellular prion protein PrPc is expressed in human enterocytes in cell-cell junctional domains. *J. Biol. Chem.* 279, 1499–1505. doi: 10.1074/jbc.M308578200
- Morel, E., Fouquet, S., Strup-Perrot, C., Pichol Thievend, C., Petit, C., Loew, D., et al. (2008). The cellular prion protein PrP(c) is involved in the proliferation of epithelial cells and in the distribution of junction-associated proteins. *PLoS ONE* 3:e3000. doi: 10.1371/journal.pone.0003000
- Nico, P. B., de-Paris, F., Vinade, E. R., Amaral, O. B., Rockenbach, I., Soares, B. L., et al. (2005). Altered behavioural response to acute stress in mice lacking cellular prion protein. *Behav. Brain Res.* 162, 173–181. doi: 10.1016/j.bbr.2005.02.003
- Nicolae, C., Ko, Y. P., Miosge, N., Niehoff, A., Studer, D., Enggist, L., et al. (2007). Abnormal collagen fibrils in cartilage of matrilin-1/matrilin-3-deficient mice. *J. Biol. Chem.* 282, 22163–22175. doi: 10.1074/jbc.M610994200
- Nicolas, O., Gavin, R., and del Rio, J. A. (2009). New insights into cellular prion protein (PrPc) functions: the "ying and yang" of a relevant protein. *Brain Res. Rev.* 61, 170–184. doi: 10.1016/j.brainresrev.2009.06.002
- Nieznanski, K., Nieznanska, H., Skowronek, K. J., Osiecka, K. M., and Stepkowski, D. (2005). Direct interaction between prion protein and tubulin. *Biochem. Biophys. Res. Commun.* 334, 403–411. doi: 10.1016/j.bbrc.2005.06.092
- Nourizadeh-Lillabadi, R., Seilo Torgersen, J., Vestrheim, O., Konig, M., Alestrom, P., and Syed, M. (2010). Early embryonic gene expression profiling of zebrafish prion protein (Prp2) morphants. *PLoS ONE* 5:e13573. doi: 10.1371/journal.pone.0013573
- Nuvolone, M., Kana, V., Hutter, G., Sakata, D., Mortin-Toth, S. M., Russo, G., et al. (2013). SIRPalpha polymorphisms, but not the prion protein, control phagocytosis of apoptotic cells. J. Exp. Med. 210, 2539–2552. doi: 10.1084/jem.20131274
- Passet, B., Young, R., Makhzami, S., Vilotte, M., Jaffrezic, F., Halliez, S., et al. (2012). Prion protein and Shadoo are involved in overlapping embryonic pathways and trophoblastic development. *PLoS ONE* 7:e41959. doi: 10.1371/journal.pone.0041959
- Peralta, O. A., Huckle, W. R., and Eyestone, W. H. (2011). Expression and knockdown of cellular prion protein (PrPC) in differentiating mouse embryonic stem cells. *Differentiation* 81, 68–77. doi: 10.1016/j.diff.2010.09.181
- Petit, C. S., Besnier, L., Morel, E., Rousset, M., and Thenet, S. (2013). Roles of the cellular prion protein in the regulation of cell-cell junctions and barrier function. *Tissue Barriers* 1:e24377. doi: 10.4161/tisb.24377
- Prestori, F., Rossi, P., Bearzatto, B., Laine, J., Necchi, D., Diwakar, S., et al. (2008). Altered neuron excitability and synaptic plasticity in the cerebellar granular layer of juvenile prion protein knock-out mice with impaired motor control. J. Neurosci. 28, 7091–7103. doi: 10.1523/JNEUROSCI.0409-08.2008
- Prodromidou, K., Papastefanaki, F., Sklaviadis, T., and Matsas, R. (2014). Functional cross-talk between the cellular prion protein and the neural cell adhesion molecule is critical for neuronal differentiation of neural stem/precursor cells. *Stem Cells* 32, 1674–1687. doi: 10.1002/stem.1663
- Prusiner, S. B., Groth, D., Serban, A., Koehler, R., Foster, D., Torchia, M., et al. (1993). Ablation of the prion protein (PrP) gene in mice prevents scrapie and facilitates production of anti-PrP antibodies. *Proc. Natl. Acad. Sci. U.S.A.* 90, 10608–10612. doi: 10.1073/pnas.90.22.10608
- Rial, D., Duarte, F. S., Xikota, J. C., Schmitz, A. E., Dafre, A. L., Figueiredo, C. P., et al. (2009). Cellular prion protein modulates age-related behavioral and neurochemical alterations in mice. *Neuroscience* 164, 896–907. doi: 10.1016/j.neuroscience.2009.09.005

- Rial, D., Pandolfo, P., Bitencourt, R. M., Pamplona, F. A., Moreira, K. M., Hipolide, D., et al. (2014). Cellular prion protein (PrP(C)) modulates ethanol-induced behavioral adaptive changes in mice. *Behav. Brain Res.* 271, 325–332. doi: 10.1016/j.bbr.2014.05.067
- Richt, J. A., Kasinathan, P., Hamir, A. N., Castilla, J., Sathiyaseelan, T., Vargas, F., et al. (2007). Production of cattle lacking prion protein. *Nat. Biotechnol.* 25, 132–138. doi: 10.1038/nbt1271
- Rivera-Milla, E., Oidtmann, B., Panagiotidis, C. H., Baier, M., Sklaviadis, T., Hoffmann, R., et al. (2006). Disparate evolution of prion protein domains and the distinct origin of Doppel- and prion-related loci revealed by fish-to-mammal comparisons. *FASEB J.* 20, 317–319. doi: 10.1096/fj.05-4279fje
- Roffe, M., Beraldo, F. H., Bester, R., Nunziante, M., Bach, C., Mancini, G., et al. (2010). Prion protein interaction with stress-inducible protein 1 enhances neuronal protein synthesis via mTOR. *Proc. Natl. Acad. Sci. U.S.A.* 107, 13147–13152. doi: 10.1073/pnas.1000784107
- Roucou, X., Gains, M., and LeBlanc, A. C. (2004). Neuroprotective functions of prion protein. J. Neurosci. Res. 75, 153–161. doi: 10.1002/jnr.10864
- Sanchez-Alavez, M., Criado, J. R., Klein, I., Moroncini, G., and Conti, B. (2008). Hypothalamic-pituitary-adrenal axis disregulation in PrPC-null mice. *Neuroreport* 19, 1473–1477. doi: 10.1097/WNR.0b013e32830f1e90
- Santamaria, D., and Ortega, S. (2006). Cyclins and CDKS in development and cancer: lessons from genetically modified mice. *Front. Biosci.* 11, 1164–1188. doi: 10.2741/1871
- Santos, T. G., Beraldo, F. H., Hajj, G. N., Lopes, M. H., Roffe, M., Lupinacci, F. C., et al. (2013). Laminin-gamma1 chain and stress inducible protein 1 synergistically mediate PrPC-dependent axonal growth via Ca2+ mobilization in dorsal root ganglia neurons. J. Neurochem. 124, 210–223. doi: 10.1111/jnc.12091
- Santos, T. G., Silva, I. R., Costa-Silva, B., Lepique, A. P., Martins, V. R., and Lopes, M. H. (2011). Enhanced neural progenitor/stem cells self-renewal via the interaction of stress-inducible protein 1 with the prion protein. *Stem Cells* 29, 1126–1136. doi: 10.1002/stem.664
- Santuccione, A., Sytnyk, V., Leshchyns'ka, I., and Schachner, M. (2005). Prion protein recruits its neuronal receptor NCAM to lipid rafts to activate p59fyn and to enhance neurite outgrowth. J. Cell Biol. 169, 341–354. doi: 10.1083/jcb.200409127
- Schmitt-Ulms, G., Legname, G., Baldwin, M. A., Ball, H. L., Bradon, N., Bosque, P. J., et al. (2001). Binding of neural cell adhesion molecules (N-CAMs) to the cellular prion protein. *J. Mol. Biol.* 314, 1209–1225. doi: 10.1006/jmbi. 2000.5183
- Schneider, K., Korkmaz, Y., Addicks, K., Lang, H., and Raab, W. H. (2007). Prion protein (PrP) in human teeth: an unprecedented pointer to PrP's function. *J. Endod.* 33, 110–113. doi: 10.1016/j.joen.2006.11.010
- Schrock, Y., Solis, G. P., and Stuermer, C. A. (2009). Regulation of focal adhesion formation and filopodia extension by the cellular prion protein (PrPC). *FEBS Lett.* 583, 389–393. doi: 10.1016/j.febslet.2008.12.038
- Sherman, D. L., and Brophy, P. J. (2005). Mechanisms of axon ensheathment and myelin growth. Nat. Rev. Neurosci. 6, 683–690. doi: 10.1038/nrn1743
- Solis, G. P., Schrock, Y., Hulsbusch, N., Wiechers, M., Plattner, H., and Stuermer, C. A. (2012). Reggies/flotillins regulate E-cadherin-mediated cell contact formation by affecting EGFR trafficking. *Mol. Biol. Cell* 23, 1812–1825. doi: 10.1091/mbc.E11-12-1006
- Steele, A. D., Emsley, J. G., Ozdinler, P. H., Lindquist, S., and Macklis, J. D. (2006). Prion protein (PrPc) positively regulates neural precursor proliferation during developmental and adult mammalian neurogenesis. *Proc. Natl. Acad. Sci. U.S.A.* 103, 3416–3421. doi: 10.1073/pnas.0511290103
- Steele, A. D., Lindquist, S., and Aguzzi, A. (2007). The prion protein knockout mouse: a phenotype under challenge. *Prion* 1, 83–93. doi: 10.4161/pri.1.2.4346
- Stella, R., Massimino, M. L., Sandri, M., Sorgato, M. C., and Bertoli, A. (2010). Cellular prion protein promotes regeneration of adult muscle tissue. *Mol. Cell. Biol.* 30, 4864–4876. doi: 10.1128/MCB.01040-09
- Stuermer, C. A. (2011). Reggie/flotillin and the targeted delivery of cargo. *J. Neurochem.* 116, 708–713. doi: 10.1111/j.1471-4159.2010.07007.x
- Taylor, D. R., Whitehouse, I. J., and Hooper, N. M. (2009). Glypican-1 mediates both prion protein lipid raft association and disease isoform formation. *PLoS Pathog.* 5:e1000666. doi: 10.1371/journal.ppat.1000666
- Tobler, I., Gaus, S. E., Deboer, T., Achermann, P., Fischer, M., Rulicke, T., et al. (1996). Altered circadian activity rhythms and sleep in mice devoid of prion protein. *Nature* 380, 639–642. doi: 10.1038/380639a0

- Tremblay, P., Bouzamondo-Bernstein, E., Heinrich, C., Prusiner, S. B., and DeArmond, S. J. (2007). Developmental expression of PrP in the postimplantation embryo. *Brain Res.* 1139, 60–67. doi: 10.1016/j.brainres.2006. 12.055
- van Rosmalen, J. W., Born, J. M., and Martens, G. J. (2006). Prion protein mRNA expression in *Xenopus laevis*: no induction during melanotrope cell activation. *Brain Res.* 1075, 20–25. doi: 10.1016/j.brainres.2005.12.105
- Verdu, E., Ceballos, D., Vilches, J. J., and Navarro, X. (2000). Influence of aging on peripheral nerve function and regeneration. J. Peripher. Nerv. Syst. 5, 191–208. doi: 10.1046/j.1529-8027.2000.00026.x
- Viegas, P., Chaverot, N., Enslen, H., Perriere, N., Couraud, P. O., and Cazaubon, S. (2006). Junctional expression of the prion protein PrPC by brain endothelial cells: a role in trans-endothelial migration of human monocytes. *J. Cell Sci.* 119, 4634–4643. doi: 10.1242/jcs.03222
- Walz, R., Amaral, O. B., Rockenbach, I. C., Roesler, R., Izquierdo, I., Cavalheiro, E. A., et al. (1999). Increased sensitivity to seizures in mice lacking cellular prion protein. *Epilepsia* 40, 1679–1682. doi: 10.1111/j.1528-1157.1999. tb01583.x
- Watts, J. C., Huo, H., Bai, Y., Ehsani, S., Jeon, A. H., Shi, T., et al. (2009). Interactome analyses identify ties of PrP and its mammalian paralogs to oligomannosidic N-glycans and endoplasmic reticulum-derived chaperones. *PLoS Pathog.* 5:e1000608. doi: 10.1371/annotation/9eb11869-6acb-49b0-978eabedc3cc545a
- Wong, B. S., Liu, T., Li, R., Pan, T., Petersen, R. B., Smith, M. A., et al. (2001). Increased levels of oxidative stress markers detected in the brains of mice devoid of prion protein. *J. Neurochem.* 76, 565–572. doi: 10.1046/j.1471-4159.2001.00028.x
- Wopfner, F., Weidenhofer, G., Schneider, R., von Brunn, A., Gilch, S., Schwarz, T. F., et al. (1999). Analysis of 27 mammalian and 9 avian PrPs reveals high conservation of flexible regions of the prion protein. *J. Mol. Biol.* 289, 1163–1178. doi: 10.1006/jmbi.1999.2831
- Wu, Z., Liang, F., Hong, B., Young, J. C., Sussman, M. R., Harper, J. F., et al. (2002). An endoplasmic reticulum-bound Ca(2+)/Mn(2+) pump, ECA1, supports plant growth and confers tolerance to Mn(2+) stress. *Plant Physiol.* 130, 128–137. doi: 10.1104/pp.004440

- Yoon, B. S., Ovchinnikov, D. A., Yoshii, I., Mishina, Y., Behringer, R. R., and Lyons, K. M. (2005). Bmpr1a and Bmpr1b have overlapping functions and are essential for chondrogenesis *in vivo. Proc. Natl. Acad. Sci. U.S.A.* 102, 5062–5067. doi: 10.1073/pnas.0500031102
- Young, R., Passet, B., Vilotte, M., Cribiu, E. P., Béringue, V., Le Provost, F., et al. (2009). The prion or the related Shadoo protein is required for early mouse embryogenesis. *FEBS Lett.* 583, 3296–3300. doi: 10.1016/j.febslet.2009.09.027
- Yu, G., Chen, J., Xu, Y., Zhu, C., Yu, H., Liu, S., et al. (2009). Generation of goats lacking prion protein. *Mol. Reprod. Dev.* 76, 3. doi: 10.1002/mrd.20960
- Zhang, C. C., Steele, A. D., Lindquist, S., and Lodish, H. F. (2006). Prion protein is expressed on long-term repopulating hematopoietic stem cells and is important for their self-renewal. *Proc. Natl. Acad. Sci. U.S.A.* 103, 2184–2189. doi: 10.1073/pnas.0510577103
- Zhang, Y., Kim, S. O., Opsahl-Vital, S., Ho, S. P., Souron, J. B., Kim, C., et al. (2011). Multiple effects of the cellular prion protein on tooth development. *Int. J. Dev. Biol.* 55, 953–960. doi: 10.1387/ijdb.113348yz

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 28 July 2014; accepted: 22 September 2014; published online: 13 October 2014.

Citation: Halliez S, Passet B, Martin-Lannerée S, Hernandez-Rapp J, Laude H, Mouillet-Richard S, Vilotte J-L and Béringue V (2014) To develop with or without the prion protein. Front. Cell Dev. Biol. **2**:58. doi: 10.3389/fcell.2014.00058

This article was submitted to Cell Death and Survival, a section of the journal Frontiers in Cell and Developmental Biology.

Copyright © 2014 Halliez, Passet, Martin-Lannerée, Hernandez-Rapp, Laude, Mouillet-Richard, Vilotte and Béringue. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.