



HAL
open science

**Three-year breeding cycle of rainbow trout
(*Oncorhynchus mykiss*) fed a plant-based diet, totally
free of marine resources: consequences for reproduction,
fatty acid composition and progeny survival**

Viviana Lazzarotto, Geneviève Corraze, Amandine Leprevost, Edwige Quillet,
Mathilde Dupont-Nivet, Françoise Médale

► **To cite this version:**

Viviana Lazzarotto, Geneviève Corraze, Amandine Leprevost, Edwige Quillet, Mathilde Dupont-Nivet, et al.. Three-year breeding cycle of rainbow trout (*Oncorhynchus mykiss*) fed a plant-based diet, totally free of marine resources: consequences for reproduction, fatty acid composition and progeny survival. PLoS ONE, 2015, 10 (2), 17 p. 10.1371/journal.pone.0117609 . hal-01194043

HAL Id: hal-01194043

<https://hal.science/hal-01194043>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Three-Year Breeding Cycle of Rainbow Trout (*Oncorhynchus mykiss*) Fed a Plant-Based Diet, Totally Free of Marine Resources: Consequences for Reproduction, Fatty Acid Composition and Progeny Survival

Viviana Lazzarotto¹, Geneviève Corraze¹, Amandine Leprevost¹, Edwige Quillet², Mathilde Dupont-Nivet², Françoise Médale^{1,2*}

1 INRA, UR 1067 "Nutrition, Métabolisme, Aquaculture", Aquapôle, 64310 Saint-Pée-sur-Nivelle, France, **2** INRA, UMR 1313 "Génétique Animale et Biologie Intégrative", 78352 Jouy-en-Josas, France

* medale@st-pee.inra.fr



CrossMark
click for updates

OPEN ACCESS

Citation: Lazzarotto V, Corraze G, Leprevost A, Quillet E, Dupont-Nivet M, Médale F (2015) Three-Year Breeding Cycle of Rainbow Trout (*Oncorhynchus mykiss*) Fed a Plant-Based Diet, Totally Free of Marine Resources: Consequences for Reproduction, Fatty Acid Composition and Progeny Survival. PLoS ONE 10(2): e0117609. doi:10.1371/journal.pone.0117609

Academic Editor: Jose Luis Balcazar, Catalan Institute for Water Research (ICRA), SPAIN

Received: September 26, 2014

Accepted: December 29, 2014

Published: February 6, 2015

Copyright: © 2015 Lazzarotto et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was supported by the FP7 European project ARRANA (n° 288925) and by the FUI project VegeAqua. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Terrestrial plant resources are increasingly used as substitutes for fish meal and fish oil in fish feed in order to reduce the reliance of aquaculture on marine fishery resources. Although many studies have been conducted to assess the effects of such nutritional transition, no whole breeding cycles of fish fed diets free from marine resources has been reported to date. We therefore studied the reproductive performance of trout after a complete cycle of breeding while consuming a diet totally devoid of marine ingredients and thus of n-3 long chain polyunsaturated fatty acids (n-3 LC-PUFAs) that play a major role in the formation of ova. Two groups of female rainbow trout were fed from first feeding either a commercial diet (C, marine and plant ingredients), or a 100% plant-based diet (V, blend of plant proteins and vegetable oils). Livers, viscera, carcasses and ova were sampled at spawning and analyzed for lipids and fatty acids. Although the V-diet was devoid of n-3 LC-PUFAs, significant amounts of EPA and DHA were found in livers and ova, demonstrating efficient bioconversion of linolenic acid and selective orientation towards the ova. Some ova were fertilized to assess the reproductive performance and offspring survival. We observed for the first time that trout fed a 100% plant-based diet over a 3-year breeding cycle were able to produce ova and viable alevins, although the ova were smaller. The survival of offspring from V-fed females was lower (-22%) at first spawning, but not at the second. Our study showed that, in addition to being able to grow on a plant-based diet, rainbow trout reared entirely on such a diet can successfully produce ova in which neo-synthesized n-3 LC-PUFAs are accumulated, leading to viable offspring. However, further adjustment of the feed formula is still needed to optimize reproductive performance.

Introduction

Aquaculture is currently the fastest growing animal production sector and has been expanding continuously for the last 25 years to meet the increasing demands in terms of fish as food that fisheries cannot meet. Aquaculture now contributes to 40% of the total fish production for human consumption and is expected to reach 60% by 2020 [1,2]. This supposes a growth rate of aquaculture that cannot be achieved by using fish meal (FM) and fish oil (FO) as major dietary sources. Their use in fish production is more than ever limited because of their decreasing availability and high cost. Carnivorous fish species, such as salmonid species, remain among the highest consumers of fish oil (51% of the market share) and the third highest of fish meal (19.5% of the market share) [3]. Among salmonids, rainbow trout (*Oncorhynchus mykiss*) farming represents the main freshwater production in Europe [3], and intense research efforts are being focused on replacing marine feedstuffs (FM and FO) with more easily available plant-based ingredients [4] in their diets.

Replacement of FM by alternative terrestrial plant products has been explored for more than twenty years [4–6]. Although substantial reduction in dietary levels of FM can be achieved, there are still several difficulties to be overcome with regard to total replacement of FM by plant ingredients, even in salmonids. Some previous studies have shown that the complete replacement of FM by plant sources of protein leads to lower growth performance in rainbow trout [5], possibly linked to decreased feed intake [7]. Moreover, poorer digestibility of carbohydrates [8], that are abundant in plant-based diets, and the presence of antinutritional factors [9] can also be responsible for poorer growth performance. With regard to FO, several studies carried out in salmonids (rainbow trout, brown trout-*Salmo trutta*-, Atlantic salmon-*Salmo salar*) have shown that it is possible to replace FO totally by individual or mixtures of vegetable oils, without affecting growth or feed efficiency [10,11], provided that n-3 LC-PUFA requirements are met by the lipids contained in fish meal. A major problem when replacing both FM and FO by plant sources is the lack of n-3 long chain polyunsaturated fatty acids (n-3 LC-PUFAs), such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). These fatty acids (FA) are important in the fish life-cycle, most notably for their roles in flesh quality [12] and reproduction, egg quality and offspring development [13,14]. In both salmonids and many cultured fish, sexual maturation and gonad development occur at the expense of stored energy, including lipids [15]. Previous studies in rainbow trout have shown that during sexual maturation lipids are initially mobilized from visceral adipose tissue [16], and mobilization from secondary storage sites (e.g. muscle) occurs only in the long term [17]. Moreover, visceral adipose tissue appears to be the primary source of energy for vitellogenesis [18], a crucial step in the female reproductive cycle. After mobilization of adipose fatty acids, this process ensures subsequent hepatic synthesis and export of lipoproteins (vitellogenin) which, together with lipids and vitamins, is finally taken up by the ova through endocytosis, providing energy reserves for ovum and offspring development. During this process, n-3 LC-PUFAs such as EPA and DHA are preferentially incorporated into ova, typically at a ratio of 2:1 [19]. EPA and DHA, as well as n-6 PUFA arachidonic acid (ARA), are recognized as determining factors in egg quality of several species [13,20–25] and in offspring development [26]. In addition to meeting energy requirements of early ontogenesis, one specific role of n-3 LC-PUFAs, in particular DHA, is incorporation into forming membranes and maintaining their fluidity [27]. Salmonids fed a diet lacking in n-3 LC-PUFAs are capable of biosynthesizing DHA via the desaturation and elongation of linolenic acid (available in plant ingredients, such as linseed oil) [17]. However, it has also been demonstrated that such bioconversion is not sufficient to compensate for the lack of dietary n-3 LC-PUFAs, resulting in a significant reduction of these FA in fish tissues [28]. While several studies have examined the effects of a plant-based diet on growth, metabolism

and fish flesh quality, the consequences of supplying a plant-based diet devoid of FM and FO across an entire life-cycle on reproductive performance and offspring survival have not been studied to date. This constitutes a significant gap in knowledge, since successful reproduction, egg quality and offspring development are key elements in fish farming [29].

We investigated whether rainbow trout reared on a plant-based diet from first feeding to reproduction were capable of synthesizing sufficient amounts of n-6 and n-3 LC-PUFAs from precursors to survive and reproduce. Furthermore, in addition to reproductive performance the incorporation of n-3 and n-6 PUFAs into ova and the subsequent trans-generational effects on offspring survival were also studied, in order to assess whether female trout exclusively reared on a plant-based diet can produce viable ova and offspring.

Materials and Methods

Ethics

The experiment was carried out in strict accordance with EU legal frameworks relating to the protection of animals used for scientific purposes (Directive 2010/63/EU) and guidelines of the French legislation governing the ethical treatment of animals (Decree no. 2001-464, May 29th, 2001). It was approved by the ethics committee of INRA (INRA 2002-36, April 14, 2002). The INRA experimental facility is certified for animal services under the permit number B29-277-02 by the French veterinary services, which is the competent authority and the scientist in charge of the experimentation received training and personal authorization (N°B64 10 003). During the experiment absence or occasional presence of dead (opaque, whitish) or unfertilized (transparent) eggs was daily checked, as well as the eggs shape and size. Dead, unfertilized or not normally shaped/sized eggs were carefully removed with a pipette by competent person in charge of the experiment fish rearing. As well as for the eggs, fish were daily monitored during the study. If any clinical symptoms (i.e. morphological abnormality, restlessness or uncoordinated movements) were observed, fish were sedated by immersion in 2% benzocaine solution and then euthanized by immersion in a 6% benzocaine solution (anesthetic overdose) during 3 minutes, to be certain that death was achieved.

Diets

The experiment was conducted with two different diets: a commercial (C) diet from “*Le Gouessant*” (Lamballe, France) containing a mix of FM, FO and plant ingredients, and an experimental plant-based diet (V), completely free from FM and FO, which were replaced by a blend of plant ingredients. The latter diet was formulated by UR NuMeA (INRA, Saint- Pée sur Nivelles, France) and manufactured at the INRA experimental facilities (Donzacq, France). Ingredients and compositions of the diets are presented in [Table 1](#). In the C-diet 45% of FM and 50% of FO were replaced by plant ingredients. During the feeding trial, the producer of the commercial diet (*Le Gouessant*) changed the origin of the ingredients (mainly FO), so that fish of the C group received diets (C₁ and C₂) with slightly different FA composition over the three breeding years. The V-diet contained only plant protein sources and a blend of vegetable oils (50% rapeseed oil, 30% linseed oil, 20% palm oil). This blend was chosen in order to provide an overall amount of FA classes closely resembling the proportions of FA classes found in fish oil. No n-6 or n-3 long chain poly-unsaturated fatty acids (LC-PUFAs), such as eicosapentaenoic acid (EPA), docosahexaenoic acid (EPA) or arachidonic acid (ARA), were present in the V diet, whereas it contained high levels of 18:3 n-3 (alpha-linolenic acid, ALA) and 18:2 n-6 (linoleic acid, LA) compared to the C diet. The fatty acid composition of the two diets is provided in [Table 2](#).

Table 1. Ingredients and composition of diets.

Diets	C	V
Ingredients (g/Kg)		
Fish meal *	434	0
Corn gluten	0	170
Soybean meal	163	200
Wheat gluten	0	250
Durum wheat	100	49.8
White lupin	0	57.2
Dehulled peas	86	30
Fish oil **	105	0
Soybean oil	105	0
Rapeseed oil	0	62
Linseed oil	0	37
Palm oil	0	24
Soy lecithin	0	20
L-lysine	0	15
L-arginine	0	10
CaHPO ₄ .2H ₂ O (18%P)	0	35
Binder	0	20
Min.-Vit. Premix	7	20
Composition (% DM)		
Crude protein	40	44.8
Crude fat	28	23.3
Energy kJ/g DM	24.5	23.6

*Origin co-fishery products—all species

** Origin co-fishery products—sardines

C: commercial diet *Le Gouessant*, V: experimental 100% plant-based diet

doi:10.1371/journal.pone.0117609.t001

Fish and experimental design

Female rainbow trout were produced at the INRA fish facilities (PEIMA, Sizun, France). Throughout the experiment, fish were reared under natural photoperiod and temperature conditions. Fish were randomly divided into two groups that were reared either on the commercial diet (C) or on the plant-based FM/FO-free diet (V). The dietary treatment was applied from first feeding until the end of the trial. Female fish of the V-diet group were fed twice a day, until apparent satiety. In order to avoid wide differences in body weight at the time of analysis, feed ration of C-group was adjusted to that of V-group, because it is known that feeding rainbow trout with plant-based diets free of marine resources leads to reduced feed intake [5,7], which results in reduced weight gain in large size rainbow trout [30].

At the moment of first spawning (2 years old females), ten females from each dietary treatment were sacrificed by benzocaine overdose, weighed and measured. The ova, liver, digestive tract (intestine with perivisceral adipose tissue) and carcasses (whole gutted fish) were collected from each female and weighed. Three pools of 50 ova from each female were weighed and the average weight (mg) of a single ovum was calculated. Absolute fecundity was measured as number of ova/female and gonadosomatic index (GSI) was calculated as: (gonad weight/total female body weight) x 100.

Table 2. Fatty acid composition (% of total fatty acids) of diets.

Diets	C ₁	C ₂	V
Fatty acid			
Saturated	23.7	25.8	16.1
MUFA	22.5	38.3	39.8
Σ n-6	8.1	10.0	21.9
18:2 n-6 (LA)	5.7	8.5	21.7
20:2 n-6	0.2	0.4	0.05
20:3 n-6	0.2	0.2	0.0
20:4 n-6 (ARA)	1.3	0.6	0.0
22:2 n-6	0.4	0.1	0.2
22:4 n-6	0.1	nd	0.0
Σ n-3	36.3	20.2	20.1
18:3 n-3 (ALA)	1.5	2.7	20.1
18:4 n-3	2.5	1.9	0.0
20:3 n-3	0.1	0.2	0.0
20:4 n-3	0.9	0.7	0.0
20:5 n-3 (EPA)	17.6	6.9	0.0
22:4 n-3	0.3	nd	0.0
22:5 n-3	1.9	1.1	0.0
22:6 n-3 (DHA)	11.5	6.4	0.0

C₁₋₂: commercial diet *Le Gouessant*, V: experimental 100% plant-based diet

C₁: fed until year-2 spawning; C₂: fed between year-2 and year-3 spawning

MUFA: monounsaturated fatty acids

nd: not detected

doi:10.1371/journal.pone.0117609.t002

Rearing of the remaining fish continued with the two dietary treatments (C₂ and V) up to the next reproduction (3 years old females) when ova were sampled following the same protocol as described for the first.

At each spawning approximately 400 of the collected ova per female were fertilized with a pool of sperm collected from males fed a commercial diet and the survival rate of progeny was measured at the eyed stage, at hatching, and at the swim-up fry stage (before 1st feeding).

All collected tissues, ova and swim-up fry sampled were stored at -80°C until analysis.

Lipid and fatty acid analysis

Total lipid content of female tissues, ova and swim-up fry was quantified gravimetrically after extraction by dichloromethane/methanol (2:1, v/v), containing 0.01% of butylated hydroxytoluene (BHT) as antioxidant, according to Folch *et al.* [31]. Neutral (NL) and polar lipid (PL) fractions were separated on silica cartridges (Sep-Pak, Waters, Ireland), according to Juaneda and Roquelin [32].

Fatty acid methyl esters (FAME) were prepared by acid-catalyzed transmethylation, using boron trifluoride according to Shantha & Ackman [33]. FAME were then analyzed in a Varian 3900 gas chromatograph equipped with a fused silica DB Wax capillary column (30m x 0.25 mm internal diameter, film thickness 0.25 μm; JW Alltech, France). Injection volume was 1 μl, using helium as carrier gas (1 ml/min). The temperatures of the injector and the flame ionization detector were 260°C and 250°C, respectively. The thermal gradient was as follows: 100–180°C at 8°C/min, 180–220°C at 4°C/min and a constant temperature of 220°C for 20min.

Fatty acids were identified with reference to a known standard mixture (Sigma, St Louis, MO, USA) and peaks were integrated using Varian Star Chromatography Software (Star Software, version 5). The results for individual FA were expressed as percentage of total identified FA methyl esters and as quantities (g/100g tissue) for ARA, EPA and DHA.

Statistical analysis

Data were analyzed statistically using the R software version 2.14.0 and the Rcmdr package. The normality of distribution and the homogeneity of variance of the variables were tested with Shapiro-Wilk's and Levene's test, respectively. When both conditions were satisfied, an independent sample t-test was performed to assess the effects of the dietary treatment; the variables with non-parametric distribution were either normalized with an arcsin transformation or, if the criteria were still not met (some fatty acids), a non-parametric test (paired Wilcoxon test) was used for analysis.

Results

Biometric parameters and reproductive performance

Biometric parameters (Table 3) of fish of both groups were measured at the spawning of year-2 and year-3. At the year-2 spawning the body weights of females fed the V-diet were higher (+18%) than that of females fed the C-diet, indicating that limitation of food supplied to the C-fed group may have been too restrictive, but no such significant difference was found at the year-3 spawning. Female body lengths were similar at both spawnings irrespective of the dietary treatment. In terms of reproductive parameters (Table 3), total spawn weight from both C and V-fed females increased at the year-3 spawning, compared to the year-2. At the first spawning, the absolute fecundity (ova/female) of the V-fed females was higher (+17%) than that of the C-fed females. Absolute fecundity was higher at the year-3 spawning than at the year-2 for both the C-fed and V-fed females, and at that time no significant difference was detected between the two groups. Average ovum weight was significantly lower in V-fed fish at both spawnings (year-2: -17%, year-3: -12%). One female among the fish fed the V-diet did not

Table 3. Biometric parameters, reproduction performance and survival rates.

Diets	Year-2 spawning			Year-3 spawning		
	C	V	p-value	C	V	p-value
Biometric parameters						
Fish weight (g)	1185 ± 253	1446 ± 205	<0.05	3453 ± 727	3166 ± 539	ns
Fish length (mm)	395 ± 41	406 ± 20	ns	570 ± 51	558 ± 43	ns
Whole spawn weight (g)	224 ± 45	222 ± 43	ns	495 ± 140	349 ± 84	<0.05
Ova weight (mg)	53 ± 7	44 ± 7	<0.05	65 ± 9	57 ± 6	<0.05
Reproduction and survival performance						
Gonadosomatic index (%)	18.9 ± 2	15.3 ± 1.4	<0.05	14.3 ± 2.3	11.1 ± 2.4	<0.05
Absolute fecundity (ova female ⁻¹)	4243 ± 589	5113 ± 994	<0.05	7680 ± 2538	6334 ± 1700	ns
Eyed stage survival (%)	91 ± 4	69 ± 30	<0.05	90 ± 7	84 ± 12	ns
Hatching survival (% of eyed)	90 ± 5	56 ± 29	<0.05	87 ± 10	82 ± 14	ns
Swim-up fry survival (% of hatched)	85 ± 5	50 ± 32	<0.05	84 ± 7	78 ± 13	ns

Data are presented as mean ± SD. *p-values* were produced by independent sample t-test. *ns*: not significant

doi:10.1371/journal.pone.0117609.t003

produce any viable ova at the year-2 spawning. The gonadosomatic index of the V-fed females was significantly lower than that of the C-fed females, at both spawnings (-19% year-2 spawning; -28.8% year-3 spawning). A lower gonadosomatic index was observed for both groups at the year-2 spawning compared to the year-3.

Rates of progeny survival (Table 3) were assessed as percentage of fertilized ova that reached the eyed stage and that were then able to hatch and to develop into swim-up fry. The survival rates for progeny from the first spawning of V-fed females were lower than for progeny of broodstock fed the C-diet, and wide inter-individual variability was observed for the V-fed group (Fig. 1). However, at the year-3 spawning, no such significant diet-induced differences were detected between alevins from the two treatment groups (Table 3).

Tissue lipid composition

Fish fed the V-diet exhibited significantly lower lipid content in the liver (-38%), while higher lipid content was found in the digestive tract (+51%) and carcass (+44%) (Fig. 2). The lipid content in the carcass and digestive tract was mainly composed of neutral lipids (NL) for both V- and C-fed groups (>90% and >88%, respectively). However, the lipid content of livers was principally composed of polar lipids (PL) in both V- and C-fed females (65% and 55%, respectively).

In ova from the year-2 and the year-3 spawning (Fig. 2) no significant differences were observed in total lipid content (11%) or in the proportions of NL and PL, which represented 62–65% and 38–35%, respectively, for both female diets.

Similarly, no significant differences were detected in terms of total lipid content (5% i.e. lower than ova), or neutral or polar lipid fractions (68–69% and 31–32%, respectively) in swim-up fry from either treatment group (Fig. 2).




Fig 1. Individual variability of progeny survival rates at different stages from females fed the C and the V diet (in bold).

doi:10.1371/journal.pone.0117609.g001




Fig 2. Total lipid content in female tissues, ova and swim-up fry. Data are presented as mean \pm SD. p-values were produced by independent sample t-test. Different letters indicates significant differences (p -value < 0.05).

doi:10.1371/journal.pone.0117609.g002

Fatty acid profile

NL and PL fractions of the tissues collected from sexually mature females, ova from the year-2 and year-3 spawning and alevins were analyzed for fatty acid composition. The proportions of FAs of all the tissues analyzed are presented in [Table 4](#) and [Table 5](#). A significant effect of the dietary treatment was observed for all fatty acids analyzed.

Saturated fatty acids (SAT) were mainly present in the PL fraction in all maternal tissues and in ova and swim-up fry, with lower values for the V-fed group ([Table 4](#) and [5](#)). In ova from the year-2 spawning and swim-up fry, no significant difference was detected in the PL fraction in terms of SAT content ([Table 5](#)), whereas percentages of SAT of the NL fraction were lower in the V-fed group than in the C-fed group (-24% and -21% in ova and swim-up fry, respectively), as well as in all the others tissues analyzed ([Table 4](#)).

With regard to monounsaturated fatty acid (MUFA) content, feeding the V-diet led to higher percentages of these FA in all tissues analyzed (+40%), in ova (+31–38%) and swim-up fry (+34%) in both the PL and NL fractions, with a more pronounced effect in the NL fraction. MUFAs were the most abundant FA class in the NL fraction of all maternal tissues, and ova and swim-up fry from V-fed fish.

Total n-6 PUFAs were higher in maternal tissues, and in ova and swim-up fry from the V-fed group (+10–15%). Linoleic acid (LA, 18:2 n-6) was mostly recovered in the NL fraction in all the tissues analyzed from both C- and V-fed groups, with significantly higher values when fed the V-diet. A higher proportion of both n-6 PUFA and LA was found in ova from C-fed females at the year-3 compared to the year-2, both in PL and NL fractions.

ARA was mainly present in the PL fraction of all tissues, and in ova and swim-up fry, irrespective of diet, with higher percentages observed for the V-fed group, except in the digestive tract. Lower percentages of ARA were found in the NL fraction of all maternal tissues with the V-diet ([Table 4](#)). Similar ARA content (g/100g tissue) was found in the carcasses and digestive tracts of

Table 4. Fatty acid composition (% of total FA) of livers, carcasses and digestive tracts (polar and neutral lipid fractions).

Diets	Polar lipids			Neutral lipids		
	C	V	p-value	C	V	p-value
Liver						
Saturated	26.6 ± 2.2	21.9 ± 2.4	<0.05	22.1 ± 1.9	14.1 ± 1.5	<0.05
MUFA	13.8 ± 1.4	16.5 ± 2.4	<0.05	23.3 ± 4.0	38.5 ± 5.5	<0.05
n-6 PUFAs	10.8 ± 0.8	20.1 ± 1.0	<0.05	10.6 ± 0.5	21.9 ± 0.6	<0.05
18: 2 n-6	3.0 ± 0.3	8.2 ± 0.9	<0.05	6.4 ± 0.9	16.8 ± 1.3	<0.05
ARA	6.9 ± 0.7	7.4 ± 1.0	ns	3.2 ± 0.7	1.8 ± 0.7	<0.05
n-3 PUFAs	47.1 ± 3.8	40.9 ± 4.7	<0.05	37.9 ± 6.1	23.5 ± 4.7	<0.05
18: 3 n-3	0.5 ± 0.1	3.4 ± 0.9	<0.05	1.5 ± 0.3	8.8 ± 0.8	<0.05
EPA	12.0 ± 1.3	6.1 ± 1.4	<0.05	12.6 ± 2.1	2.7 ± 1.1	<0.05
DHA	29.0 ± 3.8	24.8 ± 3.4	<0.05	16.2 ± 3.7	6.7 ± 3.3	<0.05
Carcass						
Saturated	28.3 ± 1.0	23.6 ± 2.0	<0.05	25.1 ± 0.9	15.5 ± 0.8	<0.05
MUFA	15.7 ± 1.7	20.7 ± 2.0	<0.05	30.2 ± 1.6	47.5 ± 0.7	<0.05
n-6 PUFAs	6.2 ± 0.5	21.2 ± 0.8	<0.05	8.5 ± 0.4	21.7 ± 0.4	<0.05
18: 2 n-6	3.1 ± 0.3	14.0 ± 1.3	<0.05	7.1 ± 0.4	18.9 ± 0.3	<0.05
ARA	2.3 ± 0.1	3.9 ± 0.5	<0.05	1.1 ± 0.1	0.3 ± 0.1	<0.05
n-3 PUFAs	45.1 ± 2.8	32.1 ± 2.3	<0.05	31.4 ± 2.1	14.3 ± 0.7	<0.05
18: 3 n-3	1.0 ± 0.1	7.5 ± 0.7	<0.05	1.9 ± 0.1	9.9 ± 0.3	<0.05
EPA	11.9 ± 0.6	6.1 ± 0.8	<0.05	10.1 ± 1.0	0.7 ± 0.1	<0.05
DHA	27.1 ± 2.5	13.9 ± 2.2	<0.05	12.5 ± 0.9	0.9 ± 0.3	<0.05
Digestive tract						
Saturated	31.7 ± 1.2	17.9	nc	17.9 ± 1.9	11.7 ± 0.8	<0.05
MUFA	26.8 ± 1.5	35.6	nc	30.9 ± 2.0	48.4 ± 0.9	<0.05
n-6 PUFAs	9.2 ± 0.4	21.4	nc	11.0 ± 0.6	23.0 ± 0.8	<0.05
18: 2 n-6	5.4 ± 0.7	15.2	nc	7.9 ± 0.6	19.7 ± 0.7	<0.05
ARA	2.5 ± 0.5	2.2	nc	1.4 ± 0.2	0.4 ± 0.1	<0.05
n-3 PUFAs	27.1 ± 2.1	22.6	nc	32.9 ± 3.5	15.4 ± 1.0	<0.05
18: 3 n-3	1.1 ± 0.2	6.1	nc	1.6 ± 0.3	8.9 ± 0.6	<0.05
EPA	6.5 ± 1.0	2.4	nc	7.5 ± 0.9	0.6 ± 0.1	<0.05
DHA	14.3 ± 1.7	8.5	nc	13.1 ± 1.7	1.7 ± 0.3	<0.05

Data are presented as mean ± SD; *p-values* were produced by independent sample t-test or equivalent non-parametric test (two-sample Wilcoxon test)

MUFA: monounsaturated fatty acids

nc: *p-value* not produced, PL for the V group pooled for practical reasons

doi:10.1371/journal.pone.0117609.t004

females from the two groups (Table 6), as well as in the ova at the year-3 spawning (Table 7). Significantly higher (or equal) percentages of ARA were detected in both NL and PL lipid fractions in ova and swim-up fry of the V-fed group compared to the C-fed group (Table 5).

Percentages of n-3 PUFAs were higher in the PL fraction compared to the NL fraction. It was the most highly represented FA class in maternal tissues (muscle, liver), ova and swim-up fry. Lower percentages of n-3 PUFAs were observed in all tissues for the V-fed group (-5 to -15%). Higher proportions of linolenic acid (ALA, 18:3 n-3) were detected in the V-fed group, particularly in the NL fraction. Percentages of EPA (20:5 n-3) and DHA (22:6 n-3) were lower in all maternal tissues, ova and swim-up fry from V-fed females (Table 5). Lower levels of EPA and DHA

Table 5. Fatty acid composition (% of total FA) of ova and swim-up fry (polar and neutral lipid fractions).

Diets	Polar lipids			Neutral lipids		
	C	V	p-value	C	V	p-value
Ova (year-2 spawning)						
Saturated	29.5 ± 3.1	28.0 ± 2.9	ns	17.0 ± 2.4	12.9 ± 0.8	<0.05
MUFA	11.3 ± 1.1	14.2 ± 1.1	<0.05	21.7 ± 1.6	34.7 ± 1.6	<0.05
n-6 PUFAs	5.9 ± 0.4	16.6 ± 0.8	<0.05	8.3 ± 0.5	19.9 ± 0.4	<0.05
18: 2 n-6	1.2 ± 0.1	6.0 ± 0.7	<0.05	5.1 ± 0.4	13.1 ± 0.7	<0.05
ARA	3.3 ± 0.2	5.2 ± 0.8	<0.05	1.9 ± 0.1	1.9 ± 0.3	ns
n-3 PUFAs	51.0 ± 3.9	39.6 ± 4.2	<0.05	46.4 ± 3.8	29.6 ± 1.7	<0.05
18: 3 n-3	0.3 ± 0.0	1.8 ± 0.3	<0.05	1.8 ± 0.2	7.7 ± 0.4	<0.05
EPA	12.2 ± 0.8	9.0 ± 1.2	<0.05	17.1 ± 1.7	4.2 ± 0.4	<0.05
DHA	33.3 ± 3.7	22.7 ± 3.8	<0.05	18.2 ± 2.1	9.7 ± 1.4	<0.05
Ova (year-3 spawning)						
Saturated	33.4 ± 2.2	31.3 ± 1.0	<0.05	20.5 ± 1.3	15.1 ± 0.8	<0.05
MUFA	12.2 ± 0.5	15.3 ± 0.7	<0.05	27.1 ± 1.4	39.4 ± 1.1	<0.05
n-6 PUFAs	9.1 ± 0.6	18.1 ± 1.2	<0.05	15.9 ± 0.9	19.7 ± 0.7	<0.05
18: 2 n-6	3.0 ± 0.4	6.2 ± 1.2	<0.05	12.5 ± 0.8	13.8 ± 1.0	<0.05
ARA	3.2 ± 0.1	6.1 ± 1.4	<0.05	1.3 ± 0.1	1.9 ± 0.6	<0.05
n-3 PUFAs	43.3 ± 1.9	33.6 ± 1.8	<0.05	32.1 ± 2.2	23.5 ± 0.8	<0.05
18: 3 n-3	0.4 ± 0.0	1.4 ± 0.3	<0.05	2.5 ± 0.2	7.4 ± 1.0	<0.05
EPA	10.1 ± 0.3	8.7 ± 0.8	<0.05	9.4 ± 1.1	3.4 ± 0.6	<0.05
DHA	29.0 ± 1.4	19.7 ± 1.8	<0.05	14.3 ± 1.4	6.9 ± 1.1	<0.05
Swim-up fry (year-3 spawning)						
Saturated	31.1 ± 2.5	28.3 ± 1.5	ns	18.7 ± 1.2	14.7 ± 0.7	<0.05
MUFA	13.0 ± 0.4	15.1 ± 0.6	<0.05	24.1 ± 0.7	36.4 ± 0.7	<0.05
n-6 PUFAs	8.2 ± 0.4	13.7 ± 0.6	<0.05	15.5 ± 0.2	20.3 ± 0.3	<0.05
18: 2 n-6	3.2 ± 0.1	4.3 ± 0.2	<0.05	11.4 ± 0.3	13.0 ± 0.2	<0.05
ARA	3.3 ± 0.2	6.4 ± 0.6	<0.05	1.7 ± 0.1	2.4 ± 0.2	<0.05
n-3 PUFAs	44.4 ± 2.7	38.6 ± 2.6	<0.05	37.1 ± 1.9	25.9 ± 1.0	<0.05
18: 3 n-3	0.5 ± 0.0	1.3 ± 0.4	<0.05	2.3 ± 0.2	6.6 ± 0.1	<0.05
EPA	9.0 ± 0.7	7.7 ± 0.5	<0.05	9.9 ± 0.6	3.8 ± 0.2	<0.05
DHA	32.5 ± 2.4	28.5 ± 2.8	ns	18.3 ± 1.1	9.6 ± 0.7	<0.05

Data are presented as mean ± SD. *p-values* were produced by independent sample t-test or equivalent non-parametric test (two-sample Wilcoxon test)

MUFA: monounsaturated fatty acids

ns: not significant

doi:10.1371/journal.pone.0117609.t005

(g/100g tissue) were found in the digestive tracts, carcasses and ova of females fed the V-diet compared to those of the C-fed females (Table 6 and Table 7). In swim-up fry, the DHA content of the PL fraction was in the same range, irrespective of the broodstock dietary treatment (Table 5).

Discussion

The results of this study prove, for the first time to our knowledge, that rainbow trout can achieve a 3-year breeding cycle including two spawnings events, despite being fed a plant-based diet totally free from marine resources. Our study focused on characterization of the dietary effects on the lipid content and fatty acid profile of female tissues, ova and progeny across

Table 6. ARA, EPA and DHA content (g/100g tissue) in carcasses and digestive tracts of C- and V-fed females.

	Diets		p-value
	C	V	
<i>Carcass</i>			
ARA	0.12 ± 0.03	0.11 ± 0.03	ns
EPA	1.05 ± 0.3	0.22 ± 0.1	<0.05
DHA	1.44 ± 0.4	0.31 ± 0.1	<0.05
<i>Digestive tract</i>			
ARA	0.01 ± 0.01	0.02 ± 0.01	ns
EPA	0.06 ± 0.03	0.03 ± 0.01	<0.05
DHA	0.10 ± 0.05	0.07 ± 0.03	<0.05

Data are presented as mean ± SD. p-values were produced by independent sample t-test
 ns: not significant

doi:10.1371/journal.pone.0117609.t006

two spawning events, because these parameters are known to be key determinants of reproductive success and progeny survival [17, 19].

Several studies have been carried out to investigate the possibility of using plant-based diets in rearing fish but very few of them challenged fish with diets in which both FM and FO were substituted [34]. The use of plant-based diets in aquaculture-raised fish species has revealed limitations, such as reduction in feed-intake [35–38]. The concomitant replacement of dietary FM and FO is known to result in lower growth performance [5,39], probably linked to a combination of lower feed intake and lower feed efficiency [5]. This poorer growth effect of a totally plant-based diet is believed to be mainly related to the replacement of FM and not to the FO substitution in rainbow trout [40,41], European seabass [42] and gilthead sea bream [43]. Pereira *et al.* observed that the replacement of FM by plant proteins both reduced feed intake and resulted in poorer reproductive performance in rainbow trout [7], but it was difficult to be certain that the reproductive performance was impaired by FM replacement or by the reduced feed intake. In the present study we adjusted the feed intake of the C-group to that of the V-group, in order to avoid differences in growth and the potential subsequent effects on spawn [44]. Although at the first spawning this feed restriction was slightly too high, we reached our goal at the spawning of the year after.

Table 7. ARA, EPA and DHA content (g/100g ova) in ova from second- and third-year spawning.

Diets	Year-2 spawning			Year-3 spawning		
	C	V	p-value	C	V	p-value
<i>Fatty acid</i>						
ARA	0.6 ± 0.1	0.8 ± 0.2	<0.05	1.2 ± 0.2	1.3 ± 0.5	ns
EPA	3.9 ± 1.1	1.6 ± 0.3	<0.05	5.8 ± 1.4	2.1 ± 0.5	<0.05
DHA	6.1 ± 1.0	3.8 ± 0.8	<0.05	6.2 ± 1.8	2.7 ± 0.7	<0.05

Data are presented as mean ±SD. p-values were produced by independent sample t-test
 ns: not significant

doi:10.1371/journal.pone.0117609.t007

V-diet induces perivisceral and carcass lipid deposition

In teleost fish, stored fat supplies energy in periods of low feed availability and, more specifically, in periods of particularly high energy demand, such as reproduction and smoltification [17]. Depending on the species, fish are able to accumulate fat in different tissues [45,46], e.g. the liver in Atlantic cod [38] and perivisceral adipose tissue and flesh in Atlantic salmon and rainbow trout [5,38]. In the present study, females from both groups had significantly higher lipid content in the digestive tract (Fig. 2) than in the carcass and liver. This was in accordance with the fact that perivisceral adipose tissue is the main lipid storage site in rainbow trout [47], while the liver does not function as a fat store, despite its important role in FA metabolism [48].

The lipid content in the digestive tract was significantly higher in trout fed the V-diet than in those fed the C-diet, mainly due to the NL fraction. This result confirmed results from previous nutritional studies carried out on salmonids [5,38], thus indicating that feeding a 100% plant-based diet leads to greater lipid deposition in perivisceral fat. Previous studies also demonstrated that maternal dietary lipids and adipose tissue lipids mobilized during previtellogenic and vitellogenic development are transported to oocytes via the circulatory system [27,49,50]. Interestingly, in the present study the total lipid content of ova was similar in both dietary groups (11%) at both spawnings and, furthermore, no significant difference in lipid content was found between swim-up fry (5%) originating from the two dietary treatments. The decrease in terms of total lipid content (from 11% in ova to 5% in swim-up fry) is due to the fact that prior to exogenous feeding the embryo utilizes nutrients derived from the ovum, including lipids [51]. In conclusion, the fat content of ova and fry was not related to the stored maternal body fat.

V-diet affects FA profile in maternal tissues, ova and offspring

One of the main problems of the concomitant replacement of FM and FO by plant sources in fish feed is modification of the fatty acid (FA) profile, that results in decrease in n-3 PUFAs in the fish body [52–54]. Indeed, vegetable oils do not contain EPA and DHA, and the FA profile of fish tissue is known to mirror the FA composition of the diet [36,38]. Our results were consistent with these previous findings; we observed higher concentrations of MUFA, linolenic acid and n-6 PUFA in NL and PL fractions of all maternal fish tissues investigated, reflecting their abundance in the V-diet. In particular, feeding fish the latter diet led to similar or even higher concentrations of ARA in the PL fraction of maternal tissues, although the dietary intake of ARA was zero. When calculating the quantities of ARA (g ARA/100g tissue) we found higher (or equal) values for the V-fed group in the carcasses, digestive tracts and ova from both spawnings. These results can be explained by the higher proportion of n-6 PUFA in the V-diet (22%) compared to the C-diet (8% and 10%), mainly due to the high linoleic acid content, the ARA precursor. On the other hand, the levels of EPA and DHA, a characteristic component of dietary FO, were lower in all tissues of fish fed the V-diet, as were the concentrations of saturated FAs. The bioconversion of 18:2 n-6 into ARA in the V-fed group seems therefore to be greater than the bioconversion of 18:3 n-3 into EPA and DHA. However, measurable concentrations of both EPA and DHA were found in NL and PL fractions of all maternal tissues of trout reared on the V-diet that was devoid of these FAs. Several previous studies reported that the synthesis of n-3 LC-PUFAs, such as EPA and DHA, as measured by the desaturation and elongation of linolenic acid 18:3n-3, was increased in salmonids fed diets in which dietary FO was replaced by vegetable oils [55–57], although the replacement of FO by vegetable oils resulted in reduced levels of EPA irrespective of the species [36]. The present study is the first in which rainbow trout were fed across the whole life cycle (3 years) without dietary intake of

EPA and DHA (concomitant replacement of FM and FO). The fact that non-negligible quantities of these two FA were found, indicated that female rainbow trout are capable of synthesizing EPA and DHA from dietary linolenic acid and to store these two fatty acids in both NL (reserve) and PL (membrane) fractions. However, the EPA and DHA content of tissues, ova and swim-up fry was significantly lower in V-fed trout than in those fed the C-diet (Table 4 and 5), meaning that the biosynthesis of EPA and DHA was not enough to overcome the total absence of these two FAs in the diet.

The degree of modification of the FA composition as a result of dietary FO replacement was significantly different between lipid storage sites (adipose tissue, muscle) and other tissues. Irrespective of the diet, EPA and DHA were mainly stored in the PL fraction of all maternal tissues, confirming selective retention of n-3 PUFAs, particularly DHA, in the membranes [58,59].

Broodstock nutrition is known to play a major role in quality of eggs and larvae, and lipid and fatty acid composition has in particular been identified as the main factor that determines successful reproduction and survival of offspring [60]. Moreover, the lipids derived directly from the dietary intake of broodstock in the period preceding gonadogenesis determine the essential fatty acids vital for early survival and development of newly hatched progeny [27].

With regard to n-6 PUFAs in ova, higher or equal concentrations of ARA were found in NL and PL fractions of ova and alevins from V-fed trout and C-fed trout, suggesting that neo-synthesized ARA from the 18:2 n-6 precursor (LA) is stored in ova. This is important, since ARA plays a major role in the reproductive process [13,61]. Moreover ARA is considered to be the major precursor of eicosanoids in fish [62] and eicosanoids play an important role in ovulation and are probably involved in embryogenesis, hatching and early larval development [63,64].

Our study also revealed increased percentages of n-3 LC-PUFA, EPA and DHA in ova compared to maternal tissues in both C-fed and V-fed fish, suggesting active maternal transfer of these neosynthesized FAs from linolenic acid to the ova. The amounts of EPA and DHA found per 100g of tissue strongly suggested preferential mobilization from perivisceral lipid reserves, rather than from the carcass. As mentioned above, perivisceral fat storage was increased in females fed the V-diet so that sufficient stores were available for the vitellogenic process. The amounts of EPA and DHA found in ova supported the preferential incorporation of these FAs in ova.

EPA and DHA were stored in the reserves (NL fraction) in ova and swim-up fry in addition to the membranes (PL fraction), which are the main sites of deposition. Increased levels of EPA and DHA have previously been described in fish eggs [60,61], supporting the important role that DHA plays in egg and subsequent offspring development. n-3 PUFAs are stored in the reserves for potential use during development [17] or to be catabolized for energy after hatching [65]. For example, DHA is mainly incorporated into the phospholipid-rich vitellogenin which is synthesized in the liver, then transferred via the serum to the developing ova [27]. The accumulation of DHA in the PL fraction of ova has a major biological role as structural phospholipids [27] assuring the fluidity of cell membranes. Our results also showed a greater increase in DHA content from ova to alevins for both groups. This indicates that rainbow trout alevins are able to retain DHA selectively, mainly in the polar lipid fraction.

Measurements of absolute fecundity indicate that the V-diet does not diminish the quantity of ova produced, though ova from V-fed females were smaller. Moreover, with regard to offspring development, feeding the V-diet resulted in a significant decrease (>22%) in survival rate at all stages at first spawning (year-2), confirming previous studies [26,66]. Interestingly, no such significant difference in survival occurred in alevins developing from eggs from the year-3 spawning. It is well known that bigger broodstock produce greater number of eggs and also bigger sized eggs [67]. In our study, comparing the two spawning events, we observed an

increase of female weight at the second spawning and a concomitant increase of the ova weight for both females groups. The lower reproductive performance observed for the V-fed group at the first spawning, is possibly linked to the smaller size of eggs, in comparison with those from females fed the C-diet. Indeed, at the year-3 spawning, V-fed females produced ova with a weight comparable with that of ova produced by the C-fed females at the first spawning. We also initially hypothesized that differential accumulation of total lipids or specific changes in the FA profile of ova might be responsible for the differences in survival rate observed between the two spawns of the V-fed females. In this group, ova spawned by 3-years old females had higher ARA (1.3 vs 0.8 g/100g ova) and EPA (2.1 vs 1.6 g/100g ova) content than those spawned by the 2-years old females. Given the importance of these FAs in reproduction performance [68,69], the higher survival rates observed for offsprings developing from ova spawned by 3-years old females might be linked to the greater ARA and EPA content in ova. However, we also found a lower DHA content, in ova from V-fed females at the year-3 spawning compared to the year-2 spawning. Taking into account the crucial role of DHA in reproduction and embryo development [60], it seems that the higher survival rate observed for offsprings from V-fed females at the year-3 spawning is mostly linked to the egg size and energy reserves than to differences in FA profile of ova.

In conclusion, trout which were reared on plant-based diets completely devoid of long chain PUFAs throughout the entire life cycle (3 years) were able to produce viable ova and progeny. This was probably linked to the fact that the female rainbow trout were remarkably able to synthesize ARA, EPA and DHA from dietary precursors. These neo-synthesized LC-PUFAs were preferentially incorporated into ova, which in fish represent the main sources of nutrients utilized by the embryo prior to exogenous feeding [51]. However, further adjustments of the feed-formula are still needed to optimize reproductive performance, especially at the first spawning.

Acknowledgments

We thank P. Maunas and N. Turonnet from INRA experimental facilities of Lées Athas (Pyrénées Atlantique, France) and L. Labbé and T. Kerneis from INRA PEIMA facilities for fish rearing, and F. Sandres and F. Terrier for manufacturing the experimental diet (INRA, Donzaq, France). We thank L. Larroquet for technical assistance.

Author Contributions

Conceived and designed the experiments: FM MDN EQ. Performed the experiments: VL AL FM GC MDN EQ. Analyzed the data: VL AL GC FM. Wrote the paper: VL GC FM.

References

1. Spoden K, Nierenberg D (2013) Aquaculture Tries to Fill World's Insatiable Appetite for Seafood. *Vital Signs*: Springer. pp. 57–61.
2. FAO (2011) Aquaculture Department. 2013. Global Aquaculture Production Statistics for the year.
3. FAO (2009) The State of World Fisheries and Aquaculture: 2008.
4. Gatlin DM, Barrows FT, Brown P, Dabrowski K, Gaylord TG, et al. (2007) Expanding the utilization of sustainable plant products in aquafeeds: a review. *Aquaculture Research* 38: 551–579.
5. Panserat S, Hortopan G, Plagnes-Juan E, Kolditz C, Lansard M, et al. (2009) Differential gene expression after total replacement of dietary fish meal and fish oil by plant products in rainbow trout (*Oncorhynchus mykiss*) liver. *Aquaculture* 294: 123–131.
6. De Francesco M, Parisi G, Pérez-Sánchez J, Gomez-Réqueni P, Medale F, et al. (2007) Effect of high-level fish meal replacement by plant proteins in gilthead sea bream (*Sparus aurata*) on growth and body/fillet quality traits. *Aquaculture Nutrition* 13: 361–372.

7. Pereira J, Reis-Henriques MA, Sanchez J, Costa J (1998) Effect of protein source on the reproductive performance of female rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquaculture Research* 29: 751–760.
8. Enes P, Panserat S, Kaushik S, Oliva-Teles A (2011) Dietary carbohydrate utilization by European sea bass (*Dicentrarchus labrax* L.) and gilthead sea bream (*Sparus aurata* L.) juveniles. *Reviews in Fisheries Science* 19: 201–215.
9. Francis G, Makkar HP, Becker K (2001) Antinutritional factors present in plant-derived alternate fish feed ingredients and their effects in fish. *Aquaculture* 199: 197–227.
10. Bell JG, McGhee F, Campbell PJ, Sargent JR (2003) Rapeseed oil as an alternative to marine fish oil in diets of post-smolt Atlantic salmon (*Salmo salar*): changes in flesh fatty acid composition and effectiveness of subsequent fish oil “wash out”. *Aquaculture* 218: 515–528.
11. Richard N, Kaushik S, Larroquet L, Panserat S, Corraze G (2006) Replacing dietary fish oil by vegetable oils has little effect on lipogenesis, lipid transport and tissue lipid uptake in rainbow trout (*Oncorhynchus mykiss*). *British Journal of Nutrition* 96: 299–309. PMID: [16923224](#)
12. Rosenlund G, Corraze G, Izquierdo M, Torstensen BE (2011) The effects of fish oil replacement on nutritional and organoleptic qualities of farmed fish. In: Turchini GM, Ng WK, Tocher DR, editors. *Fish oil replacement and alternative lipid sources in aquaculture feeds*. Boca Raton: CRC Press, Taylor and Francis. pp. 487–522.
13. Salze G, Tocher DR, Roy WJ, Robertson DA (2005) Egg quality determinants in cod (*Gadus morhua* L.): egg performance and lipids in eggs from farmed and wild broodstock. *Aquaculture Research* 36: 1488–1499.
14. Sargent J, McEvoy L, Estevez A, Bell G, Bell M, et al. (1999) Lipid nutrition of marine fish during early development: current status and future directions. *Aquaculture* 179: 217–229.
15. Manor ML, Weber GM, Cleveland BM, Kenney PB (2014) Effects of feeding level and sexual maturation on fatty acid composition of energy stores in diploid and triploid rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 418: 17–25.
16. Nassour I, Léger CL (1989) Deposition and mobilisation of body fat during sexual maturation in female trout (*Salmo gairdneri* Richardson). *Aquatic Living Resources* 2: 153–159.
17. Tocher DR (2003) Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* 11: 107–184.
18. Manor ML, Weber GM, Salem M, Yao J, Aussanasuwannakul A, et al. (2012) Effect of sexual maturation and triploidy on chemical composition and fatty acid content of energy stores in female rainbow trout, *Oncorhynchus mykiss*. *Aquaculture* 364: 312–321.
19. Bromage NR, Roberts RJ (1995) *Broodstock management and egg and larval quality*. Oxford (UK): Blackwell Science Ltd. 424 p. PMID: [25144093](#)
20. Cerdá J, Zanuy S, Carrillo M, Ramos J, Serrano R (1995) Short-and long-term dietary effects on female sea bass *Dicentrarchus labrax*: seasonal changes in plasma profiles of lipids and sex steroids in relation to reproduction. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology* 111: 83–91.
21. Abi-Ayad A, Mélard C, Kestemont P (1997) Effect of n-3 fatty acids in Eurasian perch broodstock diet on egg fatty acid composition and larva stress resistance. *Aquaculture International* 5.
22. Navas J, Bruce M, Thrush M, Farndale B, Bromage N, et al. (1997) The impact of seasonal alteration in the lipid composition of broodstock diets on egg quality in the European sea bass. *Journal of Fish Biology* 51: 760–773.
23. Takeuchi T (1997) Essential fatty acid requirements of aquatic animals with emphasis on fish larvae and fingerlings. *Reviews in Fisheries Science* 5: 1–25.
24. Rodriguez C, Cejas J, Martin M, Badia P, Samper M, et al. (1998) Influence of n-3 highly unsaturated fatty acid deficiency on the lipid composition of broodstock gilthead seabream (*Sparus aurata* L.) and on egg quality. *Fish Physiology and Biochemistry* 18: 177–187.
25. Harel M, Tandler A, Kissil GW, Applebaum SW (1994) The kinetics of nutrient incorporation into body tissues of gilthead seabream (*Sparus aurata*) females and the subsequent effects on egg composition and egg quality. *British Journal of Nutrition* 72: 45–58. PMID: [7918328](#)
26. Leray C, Nonnotte G, Roubaud P, Leger C (1985) Incidence of (n-3) essential fatty acid deficiency on trout reproductive processes. *Reproduction, Nutrition, Development* 25: 567. PMID: [4023399](#)
27. Sargent J (1995) Origins and functions of egg lipids: nutritional implications. In: Bromage NR, Roberts RJ, editors. *Broodstock management and egg and larval quality*. Oxford (UK): Blackwell Science Ltd. pp. 353–372.

28. Turchini GM, Francis DS (2009) Fatty acid metabolism (desaturation, elongation and β -oxidation) in rainbow trout fed fish oil-or linseed oil-based diets. *British Journal of Nutrition* 102: 69–81. doi: [10.1017/S0007114508137874](https://doi.org/10.1017/S0007114508137874) PMID: [19123959](https://pubmed.ncbi.nlm.nih.gov/19123959/)
29. Hamre K, Yúfera M, Rønnestad I, Boglione C, Conceição LE, et al. (2013) Fish larval nutrition and feed formulation: knowledge gaps and bottlenecks for advances in larval rearing. *Reviews in Aquaculture* 5: S26–S58.
30. De Francesco M, Parisi G, Médale F, Lupi P, Kaushik SJ, et al. (2004) Effect of long-term feeding with a plant protein mixture based diet on growth and body/fillet quality traits of large rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 236: 413–429.
31. Folch J, Lees M, Sloane-Stanley G (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226: 497–509. PMID: [13428781](https://pubmed.ncbi.nlm.nih.gov/13428781/)
32. Juaneda P, Rocquelin G (1985) Rapid and convenient separation of phospholipids and non phosphorus lipids from rat heart using silica cartridges. *Lipids* 20: 40–41. PMID: [2982073](https://pubmed.ncbi.nlm.nih.gov/2982073/)
33. Shantha N, Ackman R (1990) Nervonic acid versus tricosanoic acid as internal standards in quantitative gas chromatographic analyses of fish oil longer-chain n—3 polyunsaturated fatty acid methyl esters. *J Chromatogr B: Biomed Sci Appl* 533: 1–10.
34. Le Boucher R, Quillet E, Vandeputte M, Lecalvez JM, Goardon L, et al. (2011) Plant-based diet in rainbow trout (*Oncorhynchus mykiss* Walbaum): Are there genotype-diet interactions for main production traits when fish are fed marine vs plant-based diets from the first meal? *Aquaculture* 321: 41–48.
35. Lim C, Webster CD, Lee C-S (2008) *Alternative protein sources in aquaculture diets*. New York (USA): The Haworth Press PMID: [25506952](https://pubmed.ncbi.nlm.nih.gov/25506952/)
36. Turchini GM, Torstensen BE, Ng WK (2009) Fish oil replacement in finfish nutrition. *Reviews in Aquaculture* 1: 10–57.
37. Médale F, Kaushik S (2009) Protein sources in feed for farmed fish. *Cahiers Agricultures* 18 103–111.
38. Corraze G, Kaushik S (2009) Lipid nutrition and fish oil replacement by vegetable oils in pisciculture. *Cahiers Agricultures* 18: 112–118.
39. Drew MD, Ogunkoya AE, Janz DM, Van Kessel AG (2007) Dietary influence of replacing fish meal and oil with canola protein concentrate and vegetable oils on growth performance, fatty acid composition and organochlorine residues in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 267: 260–268.
40. Kaushik S, Cravedi J, Lalles J, Sumpter J, Fauconneau B, et al. (1995) Partial or total replacement of fish meal by soybean protein on growth, protein utilization, potential estrogenic or antigenic effects, cholesterolemia and flesh quality in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture* 133: 257–274.
41. Panserat S, Kolditz C, Richard N, Plagnes-Juan E, Piumi F, et al. (2008) Hepatic gene expression profiles in juvenile rainbow trout (*Oncorhynchus mykiss*) fed fishmeal or fish oil-free diets. *British Journal of Nutrition* 100: 953–967. doi: [10.1017/S0007114508981411](https://doi.org/10.1017/S0007114508981411) PMID: [18439330](https://pubmed.ncbi.nlm.nih.gov/18439330/)
42. Kaushik S, Coves D, Dutto G, Blanc D (2004) Almost total replacement of fish meal by plant protein sources in the diet of a marine teleost, the European seabass, *Dicentrarchus labrax*. *Aquaculture* 230: 391–404.
43. Gómez-Requeni P, Mingarro M, Calduch-Giner J, Médale F, Martin S, et al. (2004) Protein growth performance, amino acid utilisation and somatotrophic axis responsiveness to fish meal replacement by plant protein sources in gilthead sea bream (*Sparus aurata*). *Aquaculture* 232: 493–510.
44. Bromage N, Jones J, Randall C, Thrush M, Davies B, et al. (1992) Broodstock management, fecundity, egg quality and the timing of egg production in the rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 100: 141–166.
45. Hardy RW, Scott TM, Harrell LW (1987) Replacement of herring oil with menhaden oil, soybean oil, or tallow in the diets of Atlantic salmon raised in marine net-pens. *Aquaculture* 65: 267–277.
46. Sargent J, Tocher D, Bell J (2002) The lipids. In: Halver J, Hardy R, editors. *Fish Nutrition*. London (UK): Academic Press. pp. 181–257. doi: [10.1054/nepr.2002.0062](https://doi.org/10.1054/nepr.2002.0062) PMID: [19036296](https://pubmed.ncbi.nlm.nih.gov/19036296/)
47. Corraze G, Kaushik S (1999) Les lipides des poissons marins et d'eau douce. *Oléagineux, Corps Gras, Lipides* 6: 111–115. doi: [10.1002/jms.3517](https://doi.org/10.1002/jms.3517) PMID: [25601692](https://pubmed.ncbi.nlm.nih.gov/25601692/)
48. Peragén J, Barroso JB, García-Salguero L, de la Higuera M, Lupiáñez JA (2000) Dietary alterations in protein, carbohydrates and fat increase liver protein-turnover rate and decrease overall growth rate in the rainbow trout (*Oncorhynchus mykiss*). *Molecular and Cellular Biochemistry* 209: 97–104. PMID: [10942206](https://pubmed.ncbi.nlm.nih.gov/10942206/)
49. Tyler C, Sumpter J (1996) Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries* 6: 287–318.
50. Patiño R, Sullivan CV (2002) Ovarian follicle growth, maturation, and ovulation in teleost fish. *Fish Physiology and Biochemistry* 26: 57–70.

51. Brooks S, Tyler CR, Sumpter JP (1997) Egg quality in fish: what makes a good egg? *Reviews in Fish Biology and Fisheries* 7: 387–416.
52. Benedito-Palos L, Navarro JC, Sitjà-Bobadilla A, Gordon Bell J, Kaushik S, et al. (2008) High levels of vegetable oils in plant protein-rich diets fed to gilthead sea bream (*Sparus aurata* L.): growth performance, muscle fatty acid profiles and histological alterations of target tissues. *British Journal of Nutrition* 100: 992–1003. doi: [10.1017/S0007114508966071](https://doi.org/10.1017/S0007114508966071) PMID: [18377678](https://pubmed.ncbi.nlm.nih.gov/18377678/)
53. Torstensen BE, Espe M, Stubhaug I, Lie Ø (2011) Dietary plant proteins and vegetable oil blends increase adiposity and plasma lipids in Atlantic salmon (*Salmo salar* L.). *British Journal of Nutrition* 106: 633–647. doi: [10.1017/S0007114511000729](https://doi.org/10.1017/S0007114511000729) PMID: [21535902](https://pubmed.ncbi.nlm.nih.gov/21535902/)
54. Matos E, Gonçalves A, Bandarra N, Colen R, Nunes ML, et al. (2012) Plant proteins and vegetable oil do not have detrimental effects on post-mortem muscle instrumental texture, sensory properties and nutritional value of gilthead seabream. *Aquaculture* 358: 205–212.
55. Tocher DR, Bell JG, Dick JR, Sargent JR (1997) Fatty acyl desaturation in isolated hepatocytes from Atlantic salmon (*Salmo salar*): Stimulation by dietary borage oil containing γ -linolenic acid. *Lipids* 32: 1237–1247. PMID: [9438233](https://pubmed.ncbi.nlm.nih.gov/9438233/)
56. Tocher DR, Bell JG, Dick JR, Henderson RJ, McGhee F, et al. (2000) Polyunsaturated fatty acid metabolism in Atlantic salmon (*Salmo salar*) undergoing parr-smolt transformation and the effects of dietary linseed and rapeseed oils. *Fish Physiology and Biochemistry* 23: 59–73.
57. Tocher DR, Bell JG, MacGlaughlin P, McGhee F, Dick JR (2001) Hepatocyte fatty acid desaturation and polyunsaturated fatty acid composition of liver in salmonids: effects of dietary vegetable oil. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 130: 257–270. PMID: [11544096](https://pubmed.ncbi.nlm.nih.gov/11544096/)
58. Izquierdo M (1996) Essential fatty acid requirements of cultured marine fish larvae. *Aquaculture Nutrition* 2: 183–191.
59. Nassour I (1988) Depot et mobilisation des lipides corporels au cours du cycle sexuel chez la truite arc-en-ciel: effets d'une carence en acides gras essentiels sur la composition en acides gras des différents tissus [PhD Thesis]: University Paris Diderot-Paris VII.
60. Izquierdo M, Fernandez-Palacios H, Tacon A (2001) Effect of broodstock nutrition on reproductive performance of fish. *Aquaculture* 197: 25–42.
61. Tocher DR, Sargent JR (1984) Analyses of lipids and fatty acids in ripe roes of some northwest European marine fish. *Lipids* 19: 492–499. doi: [10.1007/BF02534481](https://doi.org/10.1007/BF02534481) PMID: [21344278](https://pubmed.ncbi.nlm.nih.gov/21344278/)
62. Bell JG, Tocher DR, Sargent JR (1994) Effect of supplementation with 20: 3 (n–6), 20: 4 (n–6) and 20: 5 (n–3) on the production of prostaglandins E and F of the 1-, 2- and 3-series in turbot (*Scophthalmus maximus*) brain astroglial cells in primary culture. *Biochimica et Biophysica Acta - Lipids and Lipid Metabolism* 1211: 335–342.
63. Mustafa T, Srivastava K (1989) Prostaglandins (eicosanoids) and their role in ectothermic organisms. *Advances in Comparative and Environmental Physiology*: Springer. pp. 157–207.
64. Furuita H, Tanaka H, Yamamoto T, Suzuki N, Takeuchi T (2002) Effects of high levels of n–3 HUFA in broodstock diet on egg quality and egg fatty acid composition of Japanese flounder, *Paralichthys olivaceus*. *Aquaculture* 210: 323–333.
65. Rønnestad I, Koven W, Tandler A, Harel M, Fyhn HJ (1998) Utilisation of yolk fuels in developing eggs and larvae of European sea bass (*Dicentrarchus labrax*). *Aquaculture* 162: 157–170.
66. Watanabe T (1985) Importance of the study of broodstock nutrition for further development of aquaculture. In: Cowey CB, Mackie AM, Bell JG, editors. *Nutrition and Feeding in Fish*. London (UK): Academic Press. pp. 395–414.
67. Sargent RC, Taylor PD, Gross MR (1987) Parental care and the evolution of egg size in fishes. *American Naturalist*: 32–46.
68. Støttrup J, Jacobsen C, Tomkiewicz J, Jarlbæk H (2013) Modification of essential fatty acid composition in broodstock of cultured European eel *Anguilla anguilla* L. *Aquaculture Nutrition* 19: 172–185.
69. Fernández-Palacios H, Izquierdo MS, Robaina L, Valencia A, Salhi M, Vergara J (1995) Effect of n-3 HUFA level in broodstock diets on egg quality of gilthead seabream (*Sparus aurata* L.). *Aquaculture* 132: 325–337.