



HAL
open science

Higher Sobolev regularity for the fractional p-Laplace equation in the superquadratic case

Lorenzo Brasco, Erik Lindgren

► **To cite this version:**

Lorenzo Brasco, Erik Lindgren. Higher Sobolev regularity for the fractional p-Laplace equation in the superquadratic case. *Advances in Mathematics*, 2017, 304, pp.300 - 354. 10.1016/j.aim.2016.03.039 . hal-01198777

HAL Id: hal-01198777

<https://hal.science/hal-01198777>

Submitted on 14 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HIGHER SOBOLEV REGULARITY FOR THE FRACTIONAL p -LAPLACE EQUATION IN THE SUPERQUADRATIC CASE

LORENZO BRASCO AND ERIK LINDGREN

ABSTRACT. We prove that for $p \geq 2$ solutions of equations modeled by the fractional p -Laplacian improve their regularity on the scale of fractional Sobolev spaces. Moreover, under certain precise conditions, they are in $W_{loc}^{1,p}$ and their gradients are in a fractional Sobolev space as well. The relevant estimates are stable as the fractional order of differentiation s reaches 1.

CONTENTS

| | |
|---|----|
| 1. Introduction | 1 |
| 1.1. Aim of the paper | 1 |
| 1.2. Some expedient definitions | 3 |
| 1.3. Main results | 4 |
| 1.4. Comments | 6 |
| 1.5. Plan of the paper | 7 |
| 2. Preliminaries | 8 |
| 2.1. Notation | 8 |
| 2.2. Besov-type spaces | 8 |
| 2.3. Gagliardo seminorms and finite differences | 10 |
| 2.4. Special spaces | 11 |
| 3. Basic estimates | 12 |
| 3.1. Caccioppoli-type inequality | 13 |
| 3.2. Improving Lemma | 21 |
| 4. Proof of Theorem 1.4 | 23 |
| 4.1. The general scheme | 23 |
| 4.2. Case $t + s p \leq (p - 1)$ | 26 |
| 4.3. Case $t + s p > (p - 1)$ | 27 |
| 4.4. Robust estimate for $s \nearrow 1$ | 30 |
| 4.5. A note on more general lower order terms | 30 |
| Appendix A. Proof of Proposition 2.4 | 31 |
| Appendix B. Pointwise inequalities | 33 |
| References | 34 |

1. INTRODUCTION

1.1. **Aim of the paper.** Let $2 \leq p < \infty$ and let $\Omega \subset \mathbb{R}^N$ be an open set, consider a local weak solution u of the p -Laplace equation

$$-\Delta_p u = 0, \quad \text{in } \Omega.$$

This means that $u \in W_{loc}^{1,p}(\Omega)$ and verifies

$$\int_{\Omega} \langle |\nabla u|^{p-2} \nabla u, \nabla \varphi \rangle dx = 0,$$

Date: August 24, 2015.

2010 Mathematics Subject Classification. 35B65, 35J70, 35R09.

Key words and phrases. Fractional p -Laplacian, nonlocal elliptic equations, Besov regularity.

for every $\varphi \in W^{1,p}$ with compact support in Ω . Thus the operator $-\Delta_p$ arises from the first variation of the $W^{1,p}$ Sobolev seminorm. A classical regularity result by Uhlenbeck asserts that (see [29, Lemma 3.1])

$$|\nabla u|^{\frac{p-2}{2}} \nabla u \in W_{loc}^{1,2}(\Omega).$$

This in turn implies the following higher differentiability for the gradient itself

$$(1.1) \quad \nabla u \in W_{loc}^{\tau,p}(\Omega), \quad \text{for every } 0 < \tau < \frac{2}{p},$$

see also [23, Proposition 3.1] for a more comprehensive result.

In this paper we want to tackle this regularity issue for weak solutions of nonlocal and nonlinear equations like the *fractional p -Laplace equation*

$$(1.2) \quad (-\Delta_p)^s u = 0,$$

and prove the analogue of (1.1). Here $0 < s < 1$ is given. In order to clarify the content of this paper, it is useful to recall that various different definitions of fractional (or nonlocal) p -Laplacian have been recently proposed (see for example [3], [8] and [26]). The definition considered in this paper is the variational one. That is, if for every open set $E \subset \mathbb{R}^N$ we define the $W^{s,p}$ Gagliardo seminorm

$$[u]_{W^{s,p}(E)} := \left(\int_E \int_E \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{1}{p}},$$

then the operator $(-\Delta_p)^s$ arises as the first variation of

$$u \mapsto [u]_{W^{s,p}(\mathbb{R}^N)}^p.$$

This is in analogy with the case of $-\Delta_p$, which formally corresponds to the case $s = 1$. Operators of this type were, to best of our knowledge, first considered in [2] and [15]. A weak solution u of (1.2) verifies

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) dx dy = 0,$$

for every $\varphi \in W^{s,p}$ with compact support. The reader worried about the sloppiness of this definition is invited to jump to Definition 1.3 below. There one may find the precise description of the equation and the definition of weak solution.

We point out that for ease of readability for the moment we just focus on the operator $(-\Delta_p)^s$. But indeed we will treat more general operators, where the singular kernel $(x, y) \mapsto |x - y|^{-N-sp}$ is replaced by some slight generalizations of the latter.

Very recently the operator $(-\Delta_p)^s$ has been much studied and the low regularity of solutions is now quite well understood. The first important paper on the subject is [10] by Di Castro, Kuusi and Palatucci. There local Hölder regularity for solutions of (1.2) is proved by building De Giorgi-type techniques for the nonlocal and nonlinear setting, in a similar spirit as it was first done for the case $p = 2$ by Kassman in [16]. In the companion paper [11], the same authors also proved the Harnack inequality for solutions of the homogeneous equation. As for the inhomogeneous equation

$$(1.3) \quad (-\Delta_p)^s u = f,$$

it is unavoidable to mention the impressive paper [18] by Kuusi, Mingione and Sire, where very refined pointwise estimates of potential type are proved. These lead for example to local continuity of the solution under sharp assumptions on f (see [18, Corollary 1.2]). It is worth mentioning that [18] considers a general measure datum f , not necessarily belonging to the natural dual Sobolev space. In this case, the concept of solution has to be carefully defined. Finally, Iannizzotto, Mosconi and Squassina in [14] (see also [13]) succeeded in proving global Hölder regularity for the solution of the Dirichlet problem

$$\begin{cases} (-\Delta_p)^s u = f, & \text{in } \Omega, \\ u = 0, & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases}$$

under appropriate assumptions on the data f and Ω (see [14, Theorem 1.1]).

On the contrary, as for higher differentiability of solutions, the picture is less clear. Some results on this subject are contained in the recent papers [9, 17] and [25] (see also [19]). We postpone comments on these papers, let us now proceed to present our main result.

1.2. Some expedient definitions. In order to neatly state our contribution, we need some definitions. The first one is very similar to that of *nonlocal tail* of a function, introduced in [10]. Since the two definitions differ slightly, we prefer to introduce a different notation and terminology. In what follows, the writing $F \Subset E$ means that both F and E are open sets of \mathbb{R}^N , such that the closure of F is a compact set contained in E .

Definition 1.1. Let $1 < p < \infty$, $0 < s < 1$ and $\psi \in L^p_{loc}(\mathbb{R}^N)$. For every open and bounded set $E \subset \mathbb{R}^N$, we set

$$\text{Snail}(\psi; x, E) := \left[|E|^{\frac{s-p}{N}} \int_{\mathbb{R}^N \setminus E} \frac{|\psi(y)|^p}{|x-y|^{N+sp}} dy \right]^{\frac{1}{p}}, \quad x \in E.$$

In the definition it is intended that $\text{Snail}(\psi; x, E) = +\infty$ if the integral is not finite.

Definition 1.2 (Special spaces). Let $1 < p < \infty$, $0 < s < 1$ and $\Omega \subset \mathbb{R}^N$ be an open set. Given $\psi \in L^p_{loc}(\mathbb{R}^N)$, for every $F \Subset E \Subset \Omega$ we define

$$\langle \psi \rangle_{\mathcal{X}_s^p(F;E)} := \left(\int_E |\psi|^p dx + \int_F \text{Snail}(\psi; x, E)^p dx \right)^{\frac{1}{p}}.$$

For $0 \leq t \leq 1$, we also define the associated Nikol'skii-type quantity

$$(1.4) \quad \langle \psi \rangle_{\mathcal{Y}_s^{t,p}(F;E)} := \sup_{0 < |h| < \frac{1}{2} d(F,E)} \left(\int_F \text{Snail} \left(\frac{\delta_h \psi}{|h|^t}; x, E \right)^p dx \right)^{\frac{1}{p}},$$

where we set $d(F, E) := \text{dist}(F, \mathbb{R}^N \setminus E)$. Accordingly, we define the vector spaces

$$\mathcal{X}_s^p(\Omega) = \left\{ \psi \in L^p_{loc}(\mathbb{R}^N) : \langle \psi \rangle_{\mathcal{X}_s^p(F;E)} < +\infty, \text{ for every } F \Subset E \Subset \Omega \right\},$$

and

$$\mathcal{Y}_s^{t,p}(\Omega) = \left\{ \psi \in \mathcal{X}_s^p(\Omega) : \langle \psi \rangle_{\mathcal{Y}_s^{t,p}(F;E)} < +\infty, \text{ for every } F \Subset E \Subset \Omega \right\}.$$

In (1.4) it is intended that $|h|^0 = 1$, so that for $t = 0$ we have $\mathcal{Y}_s^{0,p}(\Omega) = \mathcal{X}_s^p(\Omega)$.

With the symbol $W_0^{s,p}(\Omega)$ we denote the completion of $C_0^\infty(\Omega)$ with respect to the norm

$$\psi \mapsto \|\psi\|_{L^p(\Omega)} + [u]_{W^{s,p}(\Omega)}.$$

Definition 1.3 (Operator and local weak solutions). Let $1 < p < \infty$ and $0 < s < 1$. We consider a measurable function $K : \mathbb{R}^N \rightarrow [0, +\infty)$ satisfying

$$(1.5) \quad \frac{1}{\Lambda} |z|^{N+sp} \leq K(z) \leq \Lambda |z|^{N+sp} \quad \text{for all } z \in \mathbb{R}^N,$$

for some $\Lambda \geq 1$. Given $f \in L^{p'}_{loc}(\Omega)$, we say that $u \in W_0^{s,p}(\Omega) \cap \mathcal{X}_s^p(\Omega)$ is a *local weak solution* of

$$(1.6) \quad (-\Delta_{p,K})^s u = f \quad \text{in } \Omega,$$

if

$$(1.7) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{K(x-y)} (\varphi(x) - \varphi(y)) dx dy = \int_{\Omega'} f \varphi dx,$$

for every $\Omega' \Subset \Omega$ and every $\varphi \in W_0^{s,p}(\Omega')$. It is intended that the test functions φ are extended by 0 outside Ω' in (1.7). The assumptions on u and K guarantee that the double integral in the left-hand side of (1.7) is absolutely convergent.

In the case $K(z) = |z|^{N+sp}$, we will simply write $(-\Delta_p)^s$ in place of $(-\Delta_{p,K})^s$.

1.3. Main results. The following is our main result. The parameter t below measures the degree of differentiability of the solution “at infinity”. The value $t = 0$ is admitted as well, thus the differentiability “at infinity” is not necessary to improve the local one. The case of the p -Laplacian formally corresponds to taking $s = t = 1$ in (1.11) below. In this case, the result boils down to the aforementioned one (1.1).

Theorem 1.4 (Higher differentiability). *Let $p \geq 2$, $0 < s < 1$ and $0 \leq t \leq s$. Let $u \in W_{loc}^{s,p}(\Omega) \cap \mathcal{Y}_s^{t,p}(\Omega)$ be a local weak solution of*

$$(-\Delta_{p,K})^s u = f, \quad \text{in } \Omega,$$

with $f \in W_{loc}^{s,p'}(\Omega)$ and K verifying (1.5). For every ball $B_R \Subset \Omega$ we define

$$(1.8) \quad \begin{aligned} \mathcal{A}_R(u, f) := & \left(R^{sp} [u]_{W^{s,p}(B_R)}^p + \frac{1}{s(1-s)} \|u\|_{L^p(B_R)}^p \right) \\ & + \frac{1}{s} \langle u \rangle_{\mathcal{X}_s^p(B_{\frac{3}{4}R}, B_R)}^p + R^{tp} \langle u \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{3}{4}R}, B_{\frac{7}{8}R})}^p \\ & + R^{sp'} \left(R^{s p'} [(1-s)f]_{W^{s,p'}(B_R)}^{p'} + \frac{1}{s(1-s)} \|(1-s)f\|_{L^{p'}(B_R)}^{p'} \right). \end{aligned}$$

Then we have:

i) if $\boxed{t + sp \leq p - 1}$

$$u \in W_{loc}^{\tau,p}(\Omega), \quad \text{for every } s \leq \tau < \frac{t + sp}{p - 1},$$

and for every ball $B_R \Subset \Omega$ there holds the scaling invariant estimate

$$(1.9) \quad [u]_{W^{\tau,p}(B_{R/2})}^p \leq \frac{C_1}{R^{\tau p}} \mathcal{A}_R(u, f),$$

for some $C_1 = C_1(N, p, s, \Lambda, t, \tau) > 0$;

ii) if $\boxed{t + sp > p - 1}$ we set

$$\Gamma := \frac{1 + t + sp}{p},$$

then

$$u \in W_{loc}^{1,p}(\Omega) \quad \text{and} \quad \nabla u \in W_{loc}^{\tau,p}(\Omega), \quad \text{for every } \tau < \Gamma - 1,$$

and for every ball $B_R \Subset \Omega$ there hold the scaling invariant estimates

$$(1.10) \quad \|\nabla u\|_{L^p(B_{R/2})}^p \leq \frac{C_2}{R^p} \mathcal{A}_R(u, f),$$

and

$$(1.11) \quad [\nabla u]_{W^{\tau,p}(B_{R/2})}^p \leq \frac{(2 - \Gamma)^{-p} (\Gamma - 1)^{-p}}{(\Gamma - 1 - \tau) \tau} \frac{C_3}{R^{p(1+\tau)}} \mathcal{A}_R(u, f).$$

for some $C_2 = C_2(N, p, s, \Lambda, t) > 0$ and $C_3 = C_3(N, p, s, \Lambda, t) > 0$.

Remark 1.5 (Behaviour of the constants). Let us fix $\ell_0 > p$, then for every $0 \leq t \leq s < 1$ such that

$$t + s(p + 1) \geq \ell_0,$$

estimates (1.10) and (1.11) can be replaced by

$$(1.12) \quad \|\nabla u\|_{L^p(B_{R/2})}^p \leq \frac{C}{(\ell_0 - p)^p} (1 - s) \frac{\mathcal{A}_R(u, f)}{R^p},$$

and

$$(1.13) \quad [\nabla u]_{W^{\tau,p}(B_{R/2})}^p \leq \frac{(2 - \Gamma)^{-p} (\Gamma - 1)^{-p}}{(\Gamma - 1 - \tau) \tau} \frac{C}{(\ell_0 - p)^p} (1 - s) \frac{\mathcal{A}_R(u, f)}{R^{p(1+\tau)}},$$

with $C > 0$ depending on N, p and Λ only. In particular, the estimates are stable for $s \nearrow 1$.

Remark 1.6 (Hölder continuity via embedding). By using Morrey-type embeddings for fractional Sobolev spaces (see [1, Theorem 7.57]), we get that a local weak solution $u \in W_{loc}^{s,p}(\Omega) \cap \mathcal{Y}_s^{t,p}(\Omega)$ is locally Hölder continuous for $p \geq 2$, $0 < s < 1$ and $0 \leq t \leq s$ such that

$$t + sp > \frac{p-1}{p} N, \quad \text{if } t + sp \leq p - 1,$$

or

$$t + sp > N - 1, \quad \text{if } t + sp > p - 1.$$

For example, in dimension $N = 2$ this is always the case if $p \geq 2$ and $s > (p-1)/p$.

Before proceeding further, let us illustrate some particular cases of the previous result. We start with the case where our solution u is a priori known to be globally bounded, a situation that is quite natural if u is constructed through viscosity methods (see [20]).

Corollary 1.7 (Bounded solutions). *Let $p \geq 2$ and $0 < s < 1$. Let $u \in W_{loc}^{s,p}(\Omega) \cap L^\infty(\mathbb{R}^N)$ be a local weak solution of (1.6), with $f \in W_{loc}^{s,p'}(\Omega)$ and K verifying (1.5). Then*

$$i) \text{ if } \boxed{s \leq (p-1)/p}$$

$$u \in W_{loc}^{\tau,p}(\Omega), \quad \text{for every } \tau < \frac{sp}{p-1},$$

$$ii) \text{ if } \boxed{s > (p-1)/p}$$

$$u \in W_{loc}^{1,p}(\Omega) \quad \text{and} \quad \nabla u \in W_{loc}^{\tau,p}(\Omega), \quad \text{for every } \tau < s - \frac{p-1}{p}.$$

Proof. The result follows from the simple observation that

$$L^\infty(\mathbb{R}^N) \subset \mathcal{X}_s^p(\Omega) = \mathcal{Y}_s^{0,p}(\Omega),$$

see (2.12) below. Thus we can apply Theorem 1.4 with $t = 0$. \square

An important case is that of nonlocal Dirichlet boundary value problems for the operator $(-\Delta_p)^s$. Indeed, since the “boundary datum” g is imposed on the whole complement $\mathbb{R}^N \setminus \Omega$, the solution u naturally inherits differentiability properties “at infinity” from g . We can tune the parameter t accordingly and improve the result. As in [6], we use the notation $\widetilde{W}_0^{s,p}(\Omega)$ to denote the completion of $C_0^\infty(\Omega)$ with respect to the norm

$$\psi \mapsto [\psi]_{W^{s,p}(\mathbb{R}^N)} + \|\psi\|_{L^p(\Omega)}.$$

Corollary 1.8 (Dirichlet problems). *Let $p \geq 2$ and $0 < s < 1$. Let $\Omega \subset \mathbb{R}^N$ be an open and bounded set. Given $f \in W^{s,p'}(\Omega)$, $g \in W^{s,p}(\mathbb{R}^N)$ and K verifying (1.5), we consider the (unique) solution $u \in W^{s,p}(\mathbb{R}^N)$ of the problem*

$$\begin{cases} (-\Delta_{p,K})^s u = f, & \text{in } \Omega, \\ u = g, & \text{in } \mathbb{R}^N \setminus \Omega. \end{cases}$$

This means that u coincides with g in $\mathbb{R}^N \setminus \Omega$ and verifies (1.7) for every test function $\varphi \in \widetilde{W}_0^{s,p}(\Omega)$. Then we have:

$$i) \text{ if } \boxed{s \leq (p-1)/(p+1)}$$

$$u \in W_{loc}^{\tau,p}(\Omega), \quad \text{for every } s \leq \tau < s \frac{p+1}{p-1};$$

$$ii) \text{ if } \boxed{s > (p-1)/(p+1)}$$

$$u \in W_{loc}^{1,p}(\Omega) \quad \text{and} \quad \nabla u \in W_{loc}^{\tau,p}(\Omega), \quad \text{for every } \tau < s \frac{p+1}{p} - \frac{p-1}{p}.$$

Proof. It is sufficient to observe that

$$W^{s,p}(\mathbb{R}^N) \subset \mathcal{Y}_s^{s,p}(\Omega),$$

see (2.13) below. Thus we can apply Theorem 1.4 with $t = s$. \square

1.4. **Comments.** Some comments are in order, we start with some words on the proof of Theorem 1.4.

- (*About the proof*) The starting point of the proof of Theorem 1.4 is standard, we differentiate equation (1.7) in a discrete sense. Then by testing the equation against fractional derivatives of the solution, i.e. quantities like

$$(1.14) \quad \frac{u(x+h) - u(x)}{|h|^\vartheta},$$

we establish a Caccioppoli-type inequality for finite differences of the solution (see Proposition 3.1). For the p -Laplacian this is a “one shot” estimate, i.e. by taking ϑ to be the exponent dictated by the hypothesis $u \in W_{loc}^{1,p}$ we directly reach (1.1) from this Caccioppoli-type inequality. On the contrary, in the nonlocal case this estimate may in general be iterated. The number of possible iterations depends of course on s , namely on how close it is to 1. Then the initial information $u \in W_{loc}^{s,p}$ can be recursively improved. At each step the differentiability gain is on a “hybrid scale”, which mixes two different ways of measuring fractional derivatives. Roughly speaking, at each step we are estimating the $W^{s,p}$ seminorm (i.e. s derivatives on the Gagliardo scale) of a finite difference (1.14) (i.e. ϑ derivatives on the Nikol’skii scale). The main point of the iteration is to identify the resulting quantity as the norm of $s + \vartheta$ derivatives of the solution, measured again on the Nikol’skii scale. We point out that this is a genuine Besov-type estimate (see Lemma 3.3).

- (*The right-hand side*) As for the right-hand side f , the hypothesis $W_{loc}^{s,p'}$ is certainly too strong and we could improve the differentiability of the solution under weaker assumptions. On the other hand, we prefer to avoid further complications in the statement (and the proof) of Theorem 1.4, thus for the moment we do not try to relax it.

It is natural to expect that a suitable variant of Theorem 1.4 holds true also for very weak solutions with measure data, by using perturbative and approximation arguments as in [22, Section 6].

- (*Previous results*) Let us now make some comments on the aforementioned papers [9, 17, 19] and [25]. Let us start with the linear case, corresponding to the choice $p = 2$. In [17] and [19], the authors consider general linear elliptic nonlocal equations like

$$(1.15) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))}{\mathcal{K}(x, y)} (\varphi(x) - \varphi(y)) dx dy = \int f \varphi, \quad \text{for every } \varphi,$$

where

$$f \in L^{\frac{2N}{N+2} + \delta_0} \quad \text{and} \quad \frac{1}{\Lambda} |x - y|^{N+2s} \leq \mathcal{K}(x, y) \leq \Lambda |x - y|^{N+2s}.$$

They prove that a solution $u \in W^{s,2}(\mathbb{R}^N)$ is indeed in $W^{s+\delta,2+\delta}(\mathbb{R}^N)$ for some

$$0 < \delta = \delta(N, s, \delta_0, \Lambda) < 1 - s,$$

see [17, Theorem 1.1]. This result is based on different techniques, namely it is obtained by means of a suitable *fractional Gehring Lemma* (see [17, Theorem 1.2]). We may notice that as a consequence of Theorem 1.4, in our case as well we can improve both the differentiability and the integrability exponent, just by using a standard interpolation argument.

In [9] it is still considered the equation (1.15), under the additional assumptions

$$f \in L^2 \quad \text{and} \quad \left| \frac{1}{\mathcal{K}(x+h, y+h)} - \frac{1}{\mathcal{K}(x, y)} \right| \leq C \frac{|h|^s}{|x-y|^{N+2s}}.$$

Observe that the previous condition on \mathcal{K} covers for example the case of kernels of the type $\mathcal{K}(x, y) = K(x - y)$. Then [9, Theorem 2.2] shows that the solution gains “almost” s -derivatives, i.e. $u \in$

$W_{loc}^{2s-\tau,2}$ for every $\tau > 0$. The proof relies on differentiating twice the equation in discrete sense. Though limited to linear equations, we may notice that the result of [9] is stronger than our Theorem 1.4 in the case $p = 2$. Indeed, if we consider Theorem 1.4 for $p = 2$ and $s > 1/2$ and we do not assume differentiability “at infinity” of the solution, i.e. we taking $t = 0$, then we obtain $u \in W_{loc}^{s+1/2-\tau,2}$, for every $\tau > 0$.

As for the general case $p \geq 2$, in [25] the author considers a “regional” version of (1.3), i.e. the equation

$$\int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) dx dy = \int_{\Omega} f \varphi, dx \quad \text{for every } \varphi,$$

where f belongs to the dual space of $W^{s-\varepsilon(p-1),p}(\Omega)$, for some $\varepsilon > 0$. In [25, Theorem 1.3] it is proved that there exists $\varepsilon_0 = \varepsilon_0(s, p, \Omega)$ such that for $\varepsilon < \varepsilon_0$, a solution $u \in W^{s,p}(\Omega)$ is indeed in $W_{loc}^{s+\varepsilon,p}(\Omega)$.

- (*Limit as $s \nearrow 1$*) Finally, we conclude this list of comments by stressing that estimates (1.12) and (1.13) display the correct dependence on the parameter s , at least in the asymptotical regime $s \nearrow 1$. Indeed, we recall that for a function $u \in W_{loc}^{1,p}$ we have the pointwise convergence

$$\lim_{s \nearrow 1} (1-s) \int_{B_R} \int_{B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy = C_{N,p} \int_{B_R} |\nabla u|^p dx,$$

see [4, 5]. Moreover, we also have the Γ -convergence of the two functionals displayed above with respect to the strong L^p topology, see [7] and [24]. Thus, in the standard case $K(z) = |z|^{N+sp}$, the estimates of Theorem 1.4 can be used to prove that solutions of the fractional p -Laplace equation converge strongly in $W_{loc}^{1,p}$ to solutions of the usual p -Laplace equation as $s \nearrow 1$, under suitable assumptions. For example, let $\Omega \subset \mathbb{R}^N$ be an open and bounded set, we note u_s the unique solution of

$$(-\Delta_p)^s u_s = f_s := \frac{f}{1-s}, \quad \text{in } \Omega, \quad u = 0, \quad \text{in } \mathbb{R}^N \setminus \Omega.$$

By using (1.12) and (1.13) it is possible to show that u_s converges strongly in $L^p(\Omega) \cap W_{loc}^{1,p}(\Omega)$ to the unique solution of

$$-\Delta_p u = f, \quad \text{in } \Omega, \quad u = 0, \quad \text{on } \partial\Omega.$$

1.5. Plan of the paper. In Section 2 we introduce all the definitions and the functional analytic stuff that will be needed throughout the whole paper. The core of the paper is Section 3, where the fundamental estimates are settled down. These are the Caccioppoli-type inequality of Proposition 3.1 and the Besov-Nikol’skii differentiability improvement of Lemma 3.3. The proof of Theorem 1.4 is then contained in Section 4. In the same section we also briefly comment the case of more general equations of the type

$$(-\Delta_p)^s u = f + \lambda |u|^{q-2} u,$$

see Subsection 4.5. We then conclude the paper with a couple of appendices: while the material of Appendix B is standard, Appendix A contains the proof of an embedding property of Besov-type spaces (Proposition 2.4), which is crucially exploited in the proof Theorem 1.4.

Acknowledgements. We thank Assia Benabdallah for a useful discussion on heat kernels. An informal discussion with Sunra Mosconi and Marco Squassina has lead to an improvement of Theorem 1.4, we wish to thank them. We are grateful to Matteo Cozzi for pointing out to us his paper [9]. Part of this work has been done during a visit of the first author to Stockholm and of the second author to Marseille. The following institutions and their facilities are kindly acknowledged: Department of Mathematics of KTH (Stockholm), FRUMAM (Marseille) and CPT (Marseille). The second author has been supported by the Swedish Research Council, grant no. 2012-3124.

2. PRELIMINARIES

2.1. Notation. Let $1 \leq p < \infty$ and $0 < \alpha < 1$. For an open set $\Omega \subset \mathbb{R}^N$, we denote by $W^{\alpha,p}(\Omega)$ the usual fractional Sobolev space defined as the set of functions such that

$$\|\psi\|_{W^{\alpha,p}(\Omega)} := [\psi]_{W^{\alpha,p}(\Omega)} + \|\psi\|_{L^p(\Omega)} < +\infty.$$

The quantity $[\cdot]_{W^{\alpha,p}}$ is the $W^{\alpha,p}$ Gagliardo seminorm, i.e.

$$[\psi]_{W^{\alpha,p}(\Omega)} = \left(\int_{\Omega} \int_{\Omega} \frac{|\psi(x) - \psi(y)|^p}{|x - y|^{N+\alpha p}} dx dy \right)^{\frac{1}{p}}.$$

The local variant $W_{loc}^{\alpha,p}(\Omega)$ is defined in a straightforward manner. Given $h \in \mathbb{R}^N \setminus \{0\}$, for a measurable function $\psi : \mathbb{R}^N \rightarrow \mathbb{R}$ we introduce the notation

$$\psi_h(x) := \psi(x + h) \quad \text{and} \quad \delta_h \psi(x) = \psi_h(x) - \psi(x).$$

We recall that for every pair of functions φ, ψ we have

$$(2.1) \quad \delta_h(\varphi \psi) = (\delta_h \varphi) \psi + \varphi_h (\delta_h \psi).$$

We also remind the notation δ_h^2 for the second order differences of a function, i.e.

$$(2.2) \quad \delta_h^2 \psi(x) = \delta_h(\delta_h \psi(x)) = \psi(x + 2h) - 2\psi(x + h) + \psi(x).$$

2.2. Besov-type spaces. The following spaces defined in terms of second order differences will be important¹.

Definition 2.1 (Besov-Nikol'skii spaces). Let $1 \leq p < \infty$ and $0 < \alpha < 2$. We say that $\psi \in \mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)$ if

$$\int_{\mathbb{R}^N} |\psi|^p dx < +\infty \quad \text{and} \quad [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} := \sup_{|h|>0} \int_{\mathbb{R}^N} \left| \frac{\delta_h^2 \psi}{|h|^{\alpha}} \right|^p dx < +\infty.$$

In this case, we set

$$\|\psi\|_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} := \|\psi\|_{L^p(\mathbb{R}^N)} + [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)}.$$

We now need a couple of simple preliminary result for $\mathcal{B}_{\infty}^{\alpha,p}$. The first one states that it is indeed sufficient to control second order difference quotients for small translations. This is not surprising, we omit the proof.

Lemma 2.2 (Reduction to small translations). *Let $1 \leq p < \infty$ and $0 < \alpha < 2$. If $\psi \in \mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)$ then for every $h_0 > 0$*

$$(2.3) \quad [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} \leq \left[\sup_{0 < |h| < h_0} \left\| \frac{\delta_h^2 \psi}{|h|^{\alpha}} \right\|_{L^p(\mathbb{R}^N)} + 3 h_0^{-\alpha} \|\psi\|_{L^p(\mathbb{R}^N)} \right].$$

In the case $0 < \alpha < 1$, second order difference quotients control first order ones². This is the content of the next result.

¹We recall that it is possible to consider the more general Besov space $\mathcal{B}_q^{\alpha,p}(\mathbb{R}^N)$, built up of L^p functions such that

$$\left(\int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \left| \frac{\delta_h^2 \psi}{|h|^{\alpha}} \right|^p dx \right)^{\frac{q}{p}} \frac{dh}{|h|^N} \right)^{\frac{1}{q}} < +\infty.$$

For $q = p$, we obtain the usual fractional Sobolev space $W^{\alpha,p}(\mathbb{R}^N)$. Also observe that our notation for Besov spaces is not the standard one: we prefer to adopt this in order to be consistent with that of $W^{\alpha,p}$.

²Actually, it is easy to see that they are equivalent in this range. Since we do not need the other estimate, we omit it.

Lemma 2.3. *Let $1 \leq p < \infty$ and $0 < \alpha < 1$. If $\psi \in \mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)$ then*

$$(2.4) \quad \sup_{|h|>0} \left\| \frac{\delta_h \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} \leq \frac{C}{1-\alpha} \left[[\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} + \|\psi\|_{L^p(\mathbb{R}^N)} \right],$$

for some universal constant $C > 0$. For every $h_0 > 0$, we also get

$$(2.5) \quad \sup_{0 < |h| < h_0} \left\| \frac{\delta_h \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} \leq \frac{C}{1-\alpha} \left[\sup_{0 < |h| < h_0} \left\| \frac{\delta_h^2 \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} + (h_0^{-\alpha} + 1) \|\psi\|_{L^p(\mathbb{R}^N)} \right].$$

Proof. We will deduce the required estimate (2.4) by using some elementary manipulations, see also [28, Chapter 2.6]. We start by observing that for every measurable function ψ we have

$$\delta_h \psi(x) = \frac{1}{2} \left(\delta_{2h} \psi(x) - \delta_h^2 \psi(x) \right).$$

Thus for every $h \in \mathbb{R}^N \setminus \{0\}$ we get

$$(2.6) \quad \left\| \frac{\delta_h \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} \leq \frac{1}{2} \left\| \frac{\delta_{2h} \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} + \frac{1}{2} \left\| \frac{\delta_h^2 \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)},$$

and observe that the second term on the right-hand side is uniformly bounded by the hypothesis. For the first one, we observe that if we set $h' = 2h$

$$\begin{aligned} \left\| \frac{\delta_{2h} \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} &= 2^\alpha \left\| \frac{\delta_{h'} \psi}{|h'|^\alpha} \right\|_{L^p(\mathbb{R}^N)} \leq 2^\alpha \sup_{0 < |h'| < \frac{1}{2}} \left\| \frac{\delta_{h'} \psi}{|h'|^\alpha} \right\|_{L^p(\mathbb{R}^N)} + 2^\alpha \sup_{\frac{1}{2} \leq |h'|} \left\| \frac{\delta_{h'} \psi}{|h'|^\alpha} \right\|_{L^p(\mathbb{R}^N)} \\ &\leq 2^\alpha \sup_{0 < |h'| < \frac{1}{2}} \left\| \frac{\delta_{h'} \psi}{|h'|^\alpha} \right\|_{L^p(\mathbb{R}^N)} + 2 \cdot 4^\alpha \|\psi\|_{L^p(\mathbb{R}^N)}. \end{aligned}$$

By using this estimate in (2.6), we get

$$\sup_{0 < |h| < \frac{1}{2}} \left\| \frac{\delta_h \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} \leq \frac{1}{2} \sup_{0 < |h| < \frac{1}{2}} \left\| \frac{\delta_h^2 \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)} + 4^\alpha \|\psi\|_{L^p(\mathbb{R}^N)} + 2^{\alpha-1} \sup_{0 < |h'| < \frac{1}{2}} \left\| \frac{\delta_{h'} \psi}{|h'|^\alpha} \right\|_{L^p(\mathbb{R}^N)}.$$

By recalling that $\alpha < 1$, the last term can be absorbed in the left-hand side and thus we get (2.4).

Finally, estimate (2.5) can be obtained by combining (2.4) and (2.3). \square

The following result on Besov spaces will play a crucial role. For the reader's convenience, we give a proof of this result in Appendix A. The proof is essentially taken from Stein's book [27] and is based on the so-called *thermic extension characterization* of Besov spaces (see [28, Chapter 2.6] for such a characterization).

Proposition 2.4. *Let $1 \leq p < \infty$ and $1 < \alpha < 2$. We have the continuous embedding $\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N) \hookrightarrow W^{1,p}(\mathbb{R}^N)$. In particular, for every $\psi \in \mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)$ we have $\nabla \psi \in L^p(\mathbb{R}^N)$, with the following estimate*

$$(2.7) \quad \|\nabla \psi\|_{L^p(\mathbb{R}^N)} \leq C \|\psi\|_{L^p(\mathbb{R}^N)} + \frac{C}{(\alpha-1)} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)},$$

for some constant $C = C(N, p) > 0$. Moreover, we also have

$$(2.8) \quad \sup_{|h|>0} \left\| \frac{\delta_h \psi}{|h|^{\alpha-1}} \right\|_{L^p(\mathbb{R}^N)} \leq \frac{C}{(2-\alpha)(\alpha-1)} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)},$$

still for some $C = C(N, p) > 0$.

Remark 2.5. The previous result is false for the borderline case $\alpha = 1$, see [27, Example page 148] for a counterexample.

2.3. Gagliardo seminorms and finite differences. We still need a couple of basic facts on fractional order Sobolev spaces. The following results are well-known, but here we want to stress the explicit dependence of the constants on the differentiability index.

Proposition 2.6. *Let $1 \leq p < \infty$ and $0 < \alpha < 1$.*

- (Global case) *For every $\psi \in W^{\alpha,p}(\mathbb{R}^N)$ there holds*

$$(2.9) \quad \sup_{|h|>0} \left\| \frac{\delta_h \psi}{|h|^\alpha} \right\|_{L^p(\mathbb{R}^N)}^p \leq C(1-\alpha) [\psi]_{W^{\alpha,p}(\mathbb{R}^N)}^p,$$

for a constant $C = C(N, p) > 0$;

- (Local case) *Let $\Omega \subset \mathbb{R}^N$ be an open set. Let $\psi \in W_{loc}^{\alpha,p}(\Omega)$, then for every ball $B_R \Subset \Omega$ and every $0 < h_0 \leq \text{dist}(B_R, \partial\Omega)/2$ we have*

$$(2.10) \quad \sup_{0 < |h| < h_0} \left\| \frac{\delta_h \psi}{|h|^\alpha} \right\|_{L^p(B_R)}^p \leq C \left[(1-\alpha) [\psi]_{W^{\alpha,p}(B_{R+h_0})}^p + \left(\frac{h_0^{-\alpha p}}{\alpha} + \frac{(R+h_0)^{(1-\alpha)p}}{h_0^p} \right) \|\psi\|_{L^p(B_{R+h_0})}^p \right],$$

for a constant $C = C(N, p) > 0$.

Proof. An elementary proof of (2.9) can be found for example in [6, Lemma A.1]. In order to prove (2.10), we first take a standard Lipschitz cut-off function η such that

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \text{ on } B_R, \quad \eta \equiv 0 \text{ on } \mathbb{R}^N \setminus B_{R+\frac{h_0}{2}}, \quad |\nabla \eta| \leq \frac{2}{h_0}.$$

Then we observe that $\psi \eta \in W^{\alpha,p}(\mathbb{R}^N)$, thus by using the discrete Leibniz rule (2.1), (2.9) and the properties of η we get

$$\begin{aligned} \sup_{0 < |h| < h_0} \left\| \frac{\delta_h \psi}{|h|^\alpha} \right\|_{L^p(B_R)}^p &\leq 2^{p-1} \sup_{0 < |h| < h_0} \left\| \frac{\delta_h(\psi \eta)}{|h|^\alpha} \right\|_{L^p(B_R)}^p + 2^{p-1} \sup_{0 < |h| < h_0} \left\| \frac{\delta_h \eta}{|h|^\alpha} \psi_h \right\|_{L^p(B_R)}^p \\ &\leq C(1-\alpha) [\psi \eta]_{W^{\alpha,p}(\mathbb{R}^N)}^p + C h_0^{-\alpha p} \|\psi\|_{L^p(B_{R+h_0})}^p. \end{aligned}$$

We now decompose the Gagliardo seminorm as follows

$$\begin{aligned} [\psi \eta]_{W^{\alpha,p}(\mathbb{R}^N)}^p &= [\psi \eta]_{W^{\alpha,p}(B_{R+h_0})}^p + 2 \int_{B_{R+h_0}} \int_{\mathbb{R}^N \setminus B_{R+h_0}} \frac{|\psi(x)|^p \eta(x)^p}{|x-y|^{N+\alpha p}} dx dy \\ &\leq C [\psi]_{W^{\alpha,p}(B_{R+h_0})}^p + C \int_{B_{R+h_0}} \int_{B_{R+h_0}} \frac{|\eta(x) - \eta(y)|^p}{|x-y|^{N+\alpha p}} |\psi(y)|^p dx dy \\ &\quad + 2 \int_{B_{R+\frac{h_0}{2}}} \int_{\mathbb{R}^N \setminus B_{R+h_0}} \frac{|\psi(x)|^p}{|x-y|^{N+\alpha p}} dx dy. \end{aligned}$$

By using the Lipschitz character of η , we can now easily get (2.10). \square

Proposition 2.7. *Let $1 \leq p < \infty$ and $0 < \alpha < \beta \leq 1$. Let $\psi \in L^p(\mathbb{R}^N)$ be such that for some $h_0 > 0$ we have*

$$\sup_{0 < |h| < h_0} \left\| \frac{\delta_h \psi}{|h|^\beta} \right\|_{L^p(\mathbb{R}^N)}^p < +\infty.$$

Then there holds

$$(2.11) \quad [\psi]_{W^{\alpha,p}(\mathbb{R}^N)}^p \leq C \left(\frac{h_0^{(\beta-\alpha)p}}{\beta-\alpha} \sup_{0 < |h| < h_0} \left\| \frac{\delta_h \psi}{|h|^\beta} \right\|_{L^p(\mathbb{R}^N)}^p + \frac{h_0^{-\alpha p}}{\alpha} \|\psi\|_{L^p(\mathbb{R}^N)}^p \right),$$

for some constant $C = C(N, p) > 0$.

Proof. The proof is elementary, we give it for completeness. Let us assume that the right-hand side of (2.11) is finite, otherwise there is nothing to prove. We have

$$\begin{aligned} [\psi]_{W^{\alpha,p}(\mathbb{R}^N)}^p &= \int_{\{|h|<h_0\}} \int_{\mathbb{R}^N} \frac{|\delta_h \psi(x)|^p}{|h|^{N+\alpha p}} dx dh + \int_{\{|h|\geq h_0\}} \int_{\mathbb{R}^N} \frac{|\delta_h \psi(x)|^p}{|h|^{N+\alpha p}} dx dh \\ &\leq \int_{\{|h|<h_0\}} \left(\int_{\mathbb{R}^N} \frac{|\delta_h \psi(x)|^p}{|h|^{\beta p}} dx \right) \frac{dh}{|h|^{N-(\beta-\alpha)p}} + 2^{p-1} \|\psi\|_{L^p(\mathbb{R}^N)}^p \int_{\{|h|\geq h_0\}} \frac{1}{|h|^{N+\alpha p}} dh \\ &\leq \frac{C h_0^{(\beta-\alpha)p}}{(\beta-\alpha)} \sup_{0<|h|<h_0} \left\| \frac{\delta_h \psi}{|h|^\beta} \right\|_{L^p(\mathbb{R}^N)}^p + \frac{C h_0^{-\alpha p}}{\alpha} \|\psi\|_{L^p(\mathbb{R}^N)}^p. \end{aligned}$$

The constant C above depends on N and p only. This concludes the proof. \square

2.4. Special spaces. In this subsection, we present some basic properties of the spaces $\mathcal{X}_s^p(\Omega)$ and $\mathcal{Y}_s^{t,p}(\Omega)$ we introduced in Definition 1.2. We recall the notation

$$d(F, E) := \text{dist}(F, \mathbb{R}^N \setminus E).$$

Lemma 2.8 (Inclusions). *Let $1 \leq p < \infty$, $0 < s < 1$ and $\Omega \subset \mathbb{R}^N$ an open set. Then we have the following inclusions*

$$(2.12) \quad L^q(\mathbb{R}^N) \subset \mathcal{X}_s^p(\Omega), \quad \text{for every } q \geq p,$$

$$(2.13) \quad W^{s,p}(\mathbb{R}^N) \subset \mathcal{Y}_s^{t,p}(\Omega), \quad \text{for every } 0 \leq t \leq s,$$

$$(2.14) \quad W_{loc}^{s,p}(\Omega) \cap \mathcal{Y}_s^{s,p}(\Omega) \subset W_{loc}^{\tau,p}(\mathbb{R}^N), \quad \text{for every } 0 < \tau < s.$$

Proof. The first inclusion (2.12) stems from the simple observation that for every $F \Subset E \Subset \Omega$ and $\psi \in L^q(\mathbb{R}^N)$, by Jensen inequality we have

$$\int_F \text{Snail}(\psi; x, E)^p dx \leq |F|^{\frac{p}{q}} |E|^{\frac{sp}{N}} \|\psi\|_{L^q(\mathbb{R}^N \setminus E)}^p \left(\int_F \int_{\mathbb{R}^N \setminus E} |x-y|^{-\frac{q}{q-p}(N+sp)} dx dy \right)^{\frac{q-p}{q}} < +\infty.$$

Similarly, for the second inclusion (2.13) we observe that for every $F \Subset E \Subset \Omega$

$$\sup_{0<|h|<\frac{1}{2}d(F,E)} \int_F \text{Snail} \left(\frac{\delta_h \psi}{|h|^s}; x, E \right)^p dx \leq \frac{|F| |E|^{\frac{sp}{N}}}{d(F,E)^{N+sp}} \sup_{0<|h|<\frac{1}{2}d(F,E)} \int_{\mathbb{R}^N \setminus E} \left| \frac{\delta_h \psi}{|h|^s} \right|^p dy,$$

and the last term is bounded by the $W^{s,p}$ seminorm of u , thanks to (2.9).

Finally, we prove (2.14). Let $\psi \in W_{loc}^{s,p}(\Omega) \cap \mathcal{Y}_s^{s,p}(\Omega)$. We take an open ball $B_R \Subset \Omega$, then by (2.10) we have

$$(2.15) \quad \sup_{0<|h|<h_0} \left\| \frac{\delta_h \psi}{|h|^s} \right\|_{L^p(B_R)}^p < +\infty, \quad \text{where } h_0 \leq \frac{1}{2} \text{dist}(B_R, \partial\Omega).$$

Also, by using definition (1.4) with $B_{R/2} \Subset B_R$ we get

$$\sup_{0<|h|<\frac{R}{4}} \int_{B_{\frac{R}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left| \frac{\delta_h \psi(y)}{|h|^s} \right|^p \frac{1}{|x-y|^{N+sp}} dy dx < +\infty,$$

thus in particular for every open and bounded set $\Omega \subset \mathcal{O} \subset \mathbb{R}^N$ we have

$$(2.16) \quad \frac{C R^N}{(\text{diam } \mathcal{O})^{N+sp}} \sup_{0<|h|<\frac{R}{4}} \left\| \frac{\delta_h \psi}{|h|^s} \right\|_{L^p(\mathcal{O} \setminus B_R)}^p < +\infty.$$

By joining (2.15) and (2.16), we get the conclusion from Proposition 2.7. \square

The following monotonicity properties will be needed in the proof of Theorem 1.4.

Lemma 2.9 (Monotonicity). *Let $1 \leq p < \infty$ and $0 < s < 1$. For an open set $\Omega \subset \mathbb{R}^N$, we consider two pairs of sets $F_1 \Subset E_1 \Subset \Omega$ and $F_2 \Subset E_2 \Subset \Omega$ such that*

$$F_1 \subset F_2 \quad \text{and} \quad E_1 \subset E_2.$$

Then for every $\psi \in \mathcal{X}_s^p(\Omega)$ we have

$$(2.17) \quad \int_{F_1} \text{Snail}(\psi; x, E_1)^p dx \leq \left(\frac{|E_1|}{|E_2|} \right)^{\frac{s p}{N}} \left[\int_{F_2} \text{Snail}(\psi; x, E_2)^p dx + \frac{|F_1| |E_2|^{\frac{s p}{N}}}{d(F_1, E_1)^{N+s p}} \int_{E_2 \setminus E_1} |\psi|^p dy \right].$$

In particular, we get

$$(2.18) \quad \langle \psi \rangle_{\mathcal{X}_s^p(F_1; E_1)}^p \leq \left(\frac{|E_1|}{|E_2|} \right)^{\frac{s p}{N}} \left(1 + \frac{|F_1| |E_2|^{\frac{s p}{N}}}{d(F_1, E_1)^{N+s p}} \right) \langle \psi \rangle_{\mathcal{X}_s^p(F_2; E_2)}^p,$$

Proof. The proof of (2.17) is elementary. We have

$$\begin{aligned} \int_{F_1} \text{Snail}(\psi; x, E_1)^p dx &= |E_1|^{\frac{s p}{N}} \int_{F_1} \int_{\mathbb{R}^N \setminus E_1} \frac{|\psi(y)|^p}{|x-y|^{N+s p}} dx dy \\ &= \left(\frac{|E_1|}{|E_2|} \right)^{\frac{s p}{N}} |E_2|^{\frac{s p}{N}} \int_{F_1} \int_{\mathbb{R}^N \setminus E_2} \frac{|\psi(y)|^p}{|x-y|^{N+s p}} dx dy \\ &\quad + \left(\frac{|E_1|}{|E_2|} \right)^{\frac{s p}{N}} |E_2|^{\frac{s p}{N}} \int_{F_1} \int_{E_2 \setminus E_1} \frac{|\psi(y)|^p}{|x-y|^{N+s p}} dx dy \\ &\leq \left(\frac{|E_1|}{|E_2|} \right)^{\frac{s p}{N}} \int_{F_2} \text{Snail}(\psi; x, E_2)^p dx \\ &\quad + \left(\frac{|E_1|}{|E_2|} \right)^{\frac{s p}{N}} \frac{|F_1| |E_2|^{\frac{s p}{N}}}{d(F_1, E_1)^{N+s p}} \int_{E_2 \setminus E_1} |\psi|^p dy. \end{aligned}$$

With some standard manipulations we get (2.18) as well. \square

3. BASIC ESTIMATES

Throughout the whole section, we denote by $u \in W_{loc}^{s,p}(\Omega) \cap \mathcal{Y}_s^{t,p}(\Omega)$ a local weak solution of (1.6), with right-hand side $f \in W_{loc}^{s,p'}(\Omega)$ and K satisfying (1.5). Thus for every $\Omega' \Subset \Omega$ and any $\varphi \in W_0^{s,p}(\Omega')$, the function u satisfies (1.7). For notational simplicity, we will set

$$(3.1) \quad d\mu = \frac{1}{K(x-y)} dx dy,$$

where K verifies the hypotheses of Theorem 1.4. We also set

$$J_p(t) = |t|^{p-2} t \quad \text{and} \quad V_p(t) = |t|^{\frac{p-2}{2}} t,$$

and then define the nonlinear function of the solution $\mathcal{V}_p : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$ by

$$(3.2) \quad \mathcal{V}_p(x, y) = V_p(u(x) - u(y)) = |u(x) - u(y)|^{\frac{p-2}{2}} (u(x) - u(y)).$$

By a slight abuse of notation, for every $h \in \mathbb{R}^N \setminus 0$ we will use the following convention

$$\delta_h \mathcal{V}_p(x, y) = (\mathcal{V}_p)_h(x, y) - \mathcal{V}_p(x, y) = V_p(u_h(x) - u_h(y)) - V_p(u(x) - u(y)).$$

3.1. Caccioppoli-type inequality. We start with the following general estimate containing a free parameter of differentiability γ . This is an iterative scheme which improves the differentiability of u . We notice that the case $s = t = \gamma = 1$ formally corresponds to the result (1.1) for the p -Laplacian.

Proposition 3.1 (Differentiability scheme). *Let $p \geq 2$, $0 < s < 1$ and $0 \leq t \leq s$. We take $B_r \Subset B_R \Subset \Omega$ a pair of concentric balls and fix*

$$0 < h_0 < \frac{1}{4} \min \left\{ \text{dist}(B_R; \partial\Omega), R - r, 1 \right\}.$$

We take η a standard C^2 cut-off function such that

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \quad \text{on } B_r, \quad \eta \equiv 0 \quad \text{on } \mathbb{R}^N \setminus B_{\frac{R+r}{2}}, \quad |\nabla \eta| \leq \frac{c_N}{R-r}, \quad |D^2 \eta| \leq \frac{c_N}{(R-r)^2}.$$

For every $h \in \mathbb{R}^N \setminus \{0\}$ such that $|h| < h_0$ and every $s \leq \gamma \leq 1$ we have

$$(3.3) \quad \begin{aligned} \left[\frac{\delta_h(u\eta)}{|h|^{\frac{\gamma+t}{p}}} \right]_{W^{s,p}(B_R)}^p &\leq \left(\frac{R}{R-r} \right)^p \frac{C}{(1-s)s} \frac{1}{(R-r)^{sp}} \left\| \frac{\delta_h u}{|h|^\gamma} \right\|_{L^p(B_R)}^p \\ &+ C \left(\frac{R}{R-r} \right)^p h_0^{-\gamma-t} \left([u]_{W^{s,p}(B_{R+h_0})}^p + \frac{1}{s(1-s)R^{sp}} \|u\|_{L^p(B_{R+h_0})}^p \right) \\ &+ \frac{C}{s} \left(\frac{R+r}{R-r} \right)^N \left(\frac{R+1}{R-r} \right)^{sp} \frac{1}{(R-s)^{sp}} \langle u \rangle_{\mathcal{X}_s^p(B_{\frac{R+r}{2}+h_0}, B_{R+h_0})}^p \\ &+ \frac{C}{R^{sp}} \langle u \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{R+r}{2}}; B_R)}^p + C(1-s)^{\frac{1}{p-1}} R^{sp'} \left\| \frac{\delta_h f}{|h|^s} \right\|_{L^{p'}(B_R)}^{p'} \end{aligned}$$

for a constant $C = C(N, p, \Lambda) > 0$.

Proof. We take a test function $\varphi \in W_0^{s,p}(B_{(R+r)/2})$. By testing (1.7) with φ_{-h} for $h \in \mathbb{R}^N \setminus \{0\}$ with $|h| < h_0$ and then changing variables, we get

$$(3.4) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \left(J_p(u_h(x) - u_h(y)) \right) \left(\varphi(x) - \varphi(y) \right) d\mu(x, y) = \int_{\Omega} f_h \varphi dx.$$

We recall that μ is the singular measure defined in (3.1). We now subtract (1.7) from (3.4), thus we get

$$(3.5) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \left(J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right) \left(\varphi(x) - \varphi(y) \right) d\mu(x, y) = \int_{\Omega} \delta_h f \varphi dx,$$

for every $\varphi \in W_0^{s,p}(B_{(R+r)/2})$. Finally, we insert in (3.5) the test function

$$\varphi = \frac{\delta_h u}{|h|^{\gamma+t}} \eta^p,$$

where η is the cut-off function of the statement. We now divide the double integral in (3.5) in three pieces:

$$\mathcal{I}_1 := \int_{B_R} \int_{B_R} \frac{\left(J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right)}{|h|^{\gamma+t}} \left(\delta_h u(x) \eta(x)^p - \delta_h u(y) \eta(y)^p \right) d\mu(x, y),$$

$$\mathcal{I}_2 := \int_{B_R} \int_{\mathbb{R}^N \setminus B_R} \frac{\left(J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right)}{|h|^{\gamma+t}} \delta_h u_h(x) \eta(x)^p d\mu(x, y),$$

and

$$\mathcal{I}_3 := - \int_{\mathbb{R}^N \setminus B_R} \int_{B_R} \frac{\left(J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right)}{|h|^{\gamma+t}} \delta_h u(y) \eta(y)^p d\mu(x, y),$$

We estimate each term separately.

Estimate of \mathcal{I}_1 . For the first, we start observing that

$$\begin{aligned} \delta_h u(x) \eta(x)^p - \delta_h u(y) \eta(y)^p &= \frac{(\delta_h u(x) - \delta_h u(y))}{2} (\eta(x)^p + \eta(y)^p) \\ &\quad + \frac{(\delta_h u(x) + \delta_h u(y))}{2} (\eta(x)^p - \eta(y)^p). \end{aligned}$$

Thus we get

$$\begin{aligned} & \left(J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right) \left((\delta_h u(x)) \eta(x)^p - (\delta_h u(y)) \eta(y)^p \right) \\ & \geq \left(J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right) (\delta_h u(x) - \delta_h u(y)) \left(\frac{\eta(x)^p + \eta(y)^p}{2} \right) \\ & \quad - \left| J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right| \left(|\delta_h u(x)| + |\delta_h u(y)| \right) \left| \frac{\eta(x)^p - \eta(y)^p}{2} \right|. \end{aligned}$$

The first term has a positive sign and we will keep it on the left-hand side. For the negative term, we proceed as follows: we use (B.2), the definition (3.2) of \mathcal{V}_p , Young inequality and (B.1) to get

$$\begin{aligned} & \left| J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right| \left(|\delta_h u(x)| + |\delta_h u(y)| \right) \left| \frac{\eta(x)^p - \eta(y)^p}{2} \right| \\ & \leq 2 \frac{p-1}{p} \left(|u_h(x) - u_h(y)|^{\frac{p-2}{2}} + |u(x) - u(y)|^{\frac{p-2}{2}} \right) \\ & \quad \times |\delta_h \mathcal{V}_p(x, y)| \left(|\delta_h u(x)| + |\delta_h u(y)| \right) \frac{\eta(x)^{\frac{p}{2}} + \eta(y)^{\frac{p}{2}}}{2} \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right| \\ & \leq \frac{C}{\varepsilon} \left(|u_h(x) - u_h(y)|^{\frac{p-2}{2}} + |u(x) - u(y)|^{\frac{p-2}{2}} \right)^2 \\ & \quad \times (|\delta_h u(x)|^2 + |\delta_h u(y)|^2) \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right|^2 \\ & \quad + C \varepsilon |\delta_h \mathcal{V}_p(x, y)|^2 (\eta(x)^p + \eta(y)^p) \\ & \leq \frac{C}{\varepsilon} \left(|u_h(x) - u_h(y)|^{\frac{p-2}{2}} + |u(x) - u(y)|^{\frac{p-2}{2}} \right)^2 (|\delta_h u(x)|^2 + |\delta_h u(y)|^2) \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right|^2 \\ & \quad + C \varepsilon \left(J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right) (\delta_h u(x) - \delta_h u(y)) (\eta(x)^p + \eta(y)^p), \end{aligned}$$

where $C = C(p) > 0$. By putting all the estimates together and choosing ε sufficiently small, we then get

$$\begin{aligned} \mathcal{I}_1 & \geq \frac{1}{C} \int_{B_R} \int_{B_R} \frac{J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y))}{|h|^{\gamma+t}} (\delta_h u(x) - \delta_h u(y)) (\eta(x)^p + \eta(y)^p) d\mu(x, y) \\ & \quad - C \int_{B_R} \int_{B_R} \left(|u_h(x) - u_h(y)|^{\frac{p-2}{2}} + |u(x) - u(y)|^{\frac{p-2}{2}} \right)^2 \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right|^2 \\ & \quad \times \frac{|\delta_h u(x)|^2 + |\delta_h u(y)|^2}{|h|^{\gamma+t}} d\mu(x, y), \end{aligned}$$

for some constant $C = C(p) > 0$. We can further estimate from below the positive term by using (B.4). This leads us to

$$\begin{aligned}
(3.6) \quad \mathcal{I}_1 &\geq \frac{1}{C} \int_{B_R} \int_{B_R} \left| \frac{\delta_h u(x)}{|h|^{\frac{\gamma+t}{p}}} - \frac{\delta_h u(y)}{|h|^{\frac{\gamma+t}{p}}} \right|^p (\eta(x)^p + \eta(y)^p) d\mu(x, y) \\
&\quad - C \int_{B_R} \int_{B_R} \left(|u_h(x) - u_h(y)|^{\frac{p-2}{2}} + |u(x) - u(y)|^{\frac{p-2}{2}} \right)^2 \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right|^2 \\
&\quad \times \frac{|\delta_h u(x)|^2 + |\delta_h u(y)|^2}{|h|^{\gamma+t}} d\mu(x, y).
\end{aligned}$$

We now observe that if we set for simplicity

$$A = \frac{\delta_h u(x)}{|h|^{\frac{\gamma+t}{p}}} \quad \text{and} \quad B = \frac{\delta_h u(y)}{|h|^{\frac{\gamma+t}{p}}},$$

then by using the convexity of $\tau \mapsto \tau^p$, we have

$$\begin{aligned}
|A\eta(x) - B\eta(y)|^p &= \left| (A-B) \frac{\eta(x) + \eta(y)}{2} + (A+B) \frac{\eta(x) - \eta(y)}{2} \right|^p \leq 2^{p-2} |A-B|^p (\eta(x)^p + \eta(y)^p) \\
&\quad + 2^{p-2} (|A|^p + |B|^p) |\eta(x) - \eta(y)|^p.
\end{aligned}$$

Thus from (3.6) together with the assumption (1.5) on K , we get the following lower bound for \mathcal{I}_1

$$\begin{aligned}
(3.7) \quad \mathcal{I}_1 &\geq \frac{1}{C} \left[\frac{\delta_h u}{|h|^{\frac{\gamma+t}{p}}} \eta \right]_{W^{s,p}(B_R)}^p \\
&\quad - C \int_{B_R} \int_{B_R} \left(|u_h(x) - u_h(y)|^{\frac{p-2}{2}} + |u(x) - u(y)|^{\frac{p-2}{2}} \right)^2 \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right|^2 \\
&\quad \times \frac{|\delta_h u(x)|^2 + |\delta_h u(y)|^2}{|h|^{\gamma+t}} d\mu(x, y) \\
&\quad - C \int_{B_R} \int_{B_R} \left(\frac{|\delta_h u(x)|^p}{|h|^{\gamma+t}} + \frac{|\delta_h u(y)|^p}{|h|^{\gamma+t}} \right) \frac{|\eta(x) - \eta(y)|^p}{|x-y|^{N+sp}} dx dy,
\end{aligned}$$

where $C = C(p, \Lambda) > 0$. We need to estimate the last two integrals. For the first one, by using Hölder's inequality, again the assumption (1.5) on K , the Lipschitz character of η and some simple manipulations we

get

$$\begin{aligned}
& \int_{B_R} \int_{B_R} \left(|u_h(x) - u_h(y)|^{\frac{p-2}{2}} + |u(x) - u(y)|^{\frac{p-2}{2}} \right)^2 \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right|^2 \\
& \quad \times \frac{|\delta_h u(x)|^2 + |\delta_h u(y)|^2}{|h|^{\gamma+t}} d\mu(x, y) \\
& \leq C \left[\int_{B_R} \int_{B_R} (|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2})^{\frac{p}{p-2}} d\mu(x, y) \right]^{\frac{p-2}{p}} \\
& \quad \times \left[\int_{B_R} \int_{B_R} \frac{|\delta_h u(x)|^p + |\delta_h u(y)|^p}{|h|^{(\gamma+t)\frac{p}{2}}} \left| \eta(x)^{\frac{p}{2}} - \eta(y)^{\frac{p}{2}} \right|^p d\mu(x, y) \right]^{\frac{2}{p}} \\
& \leq C [u]_{W^{s,p}(B_{R+h_0})}^{p-2} \left[\int_{B_R} \left| \frac{\delta_h u(x)}{|h|^{\frac{\gamma+t}{2}}} \right|^p \left(\int_{B_R} \frac{1}{|x-y|^{N+p(s-1)}} dy \right) dx \right]^{\frac{2}{p}} \\
& \leq \frac{C}{R^{2s}} \left(\frac{R}{R-r} \right)^2 [u]_{W^{s,p}(B_{R+h_0})}^{p-2} \left[\frac{1}{1-s} \int_{B_R} \left| \frac{\delta_h u}{|h|^{\frac{\gamma+t}{2}}} \right|^p dx \right]^{\frac{2}{p}} \\
& \leq C [u]_{W^{s,p}(B_{R+h_0})}^p + \left(\frac{R}{R-r} \right)^p \frac{C}{R^{sp}} \frac{1}{1-s} \int_{B_R} \left| \frac{\delta_h u}{|h|^{\frac{\gamma+t}{2}}} \right|^p dx,
\end{aligned}$$

for some $C = C(N, p, \Lambda) > 0$. Thus, from (3.7) by observing that $|h| < h_0 < 1$ and that $(\gamma + t)/2 \leq \gamma$, we get

$$\begin{aligned}
(3.8) \quad \mathcal{I}_1 & \geq \frac{1}{C} \left[\frac{\delta_h u}{|h|^{\frac{\gamma+t}{p}}} \eta \right]_{W^{s,p}(B_R)}^p \\
& - C [u]_{W^{s,p}(B_{R+h_0})}^p - \left(\frac{R}{R-r} \right)^p \frac{C}{(R-r)^{sp}} \frac{1}{s(1-s)} \int_{B_R} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx \\
& - C \int_{B_R} \int_{B_R} \left(\frac{|\delta_h u(x)|^p}{|h|^{\gamma+t}} + \frac{|\delta_h u(y)|^p}{|h|^{\gamma+t}} \right) \frac{|\eta(x) - \eta(y)|^p}{|x-y|^{N+sp}} dx dy,
\end{aligned}$$

where we also used that $R^{sp} \geq (R-r)^{sp}$ and that $s(1-s) \leq (1-s)$. By the Lipschitz character of η , the last integral is estimated by

$$\begin{aligned}
(3.9) \quad & \int_{B_R} \int_{B_R} \left(\frac{|\delta_h u(x)|^p}{|h|^{\gamma+t}} + \frac{|\delta_h u(y)|^p}{|h|^{\gamma+t}} \right) \frac{|\eta(x) - \eta(y)|^p}{|x-y|^{N+sp}} dx dy \\
& \leq \frac{C}{(R-r)^{sp}} \left(\frac{R}{R-r} \right)^p \frac{1}{s(1-s)} \int_{B_R} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx,
\end{aligned}$$

for some $C = C(N, p) > 0$. Observe that we again used the trivial estimates $R^{sp} \geq (R-r)^{sp}$ and $s(s-1) \leq (s-1)$, together with $h_0 < 1$ and $(\gamma + t)/p \leq \gamma$.

It is only left to observe that from the discrete Leibniz rule (2.1), we get

$$\begin{aligned}
\left[\frac{\delta_h(u\eta)}{|h|^{\frac{\gamma+t}{p}}} \right]_{W^{s,p}(B_R)}^p & \leq C \left[\frac{\delta_h u}{|h|^{\frac{\gamma+t}{p}}} \eta \right]_{W^{s,p}(B_R)}^p + C \left[\frac{\delta_h \eta}{|h|^{\frac{\gamma+t}{p}}} u_h \right]_{W^{s,p}(B_R)}^p \\
& \leq C \left[\frac{\delta_h u}{|h|^{\frac{\gamma+t}{p}}} \eta \right]_{W^{s,p}(B_R)}^p \\
& + C \left(\frac{R}{R-r} \right)^p \frac{h_0^{p-\gamma-t}}{(R-r)^p} \left([u]_{W^{s,p}(B_{R+h_0})}^p + \frac{1}{(1-s)R^{sp}} \|u\|_{L^p(B_{R+h_0})}^p \right).
\end{aligned}$$

Observe that by the hypothesis $h_0/(R-r) < 1$ and $h_0 < 1$. To get the last estimate, we used the Lipschitz character³ of $\nabla\eta$ (recall that $\eta \in C_0^2$).

By combining this and (3.9), from (3.8) we finally get

$$(3.10) \quad \begin{aligned} \mathcal{I}_1 &\geq \frac{1}{C} \left[\frac{\delta_h(u\eta)}{|h|^{\frac{\gamma+t}{p}}} \right]_{W^{s,p}(B_R)}^p \\ &\quad - C \left(\frac{R}{R-r} \right)^p h_0^{-\gamma-t} \left([u]_{W^{s,p}(B_{R+h_0})}^p + \frac{1}{(1-s)R^{sp}} \|u\|_{L^p(B_{R+h_0})}^p \right) \\ &\quad - \frac{C}{(R-r)^{sp}} \left(\frac{R}{R-r} \right)^p \frac{1}{s(1-s)} \int_{B_R} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx. \end{aligned}$$

Estimate of \mathcal{I}_2 . By recalling that η is supported on $B_{(R+r)/2}$, we have

$$\mathcal{I}_2 \geq - \int_{B_{(R+r)/2}} \int_{\mathbb{R}^N \setminus B_R} \left| J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right| \left| \frac{\delta_h u(x)}{|h|^{\gamma+t}} \right| \eta(x)^p d\mu(x, y).$$

Then we observe that by basic calculus

$$\begin{aligned} \left| J_p(u_h(x) - u_h(y)) - J_p(u(x) - u(y)) \right| &\leq (p-1) (|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}) \\ &\quad \times |(u_h(x) - u_h(y)) - (u(x) - u(y))| \\ &\leq (p-1) (|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}) \\ &\quad \times (|\delta_h u(x)| + |\delta_h u(y)|). \end{aligned}$$

Using once again the assumption (1.5) on the kernel K , we get

$$(3.11) \quad \begin{aligned} \mathcal{I}_2 &\geq -C \int_{B_{(R+r)/2}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} dy \right) \left| \frac{\delta_h u(x)}{|h|^{\frac{\gamma+t}{2}}} \right|^2 \eta(x)^p dx \\ &\quad - C \int_{B_{(R+r)/2}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} \left| \frac{\delta_h u(y)}{|h|^t} \right| dy \right) \left| \frac{\delta_h u(x)}{|h|^\gamma} \right| \eta(x)^p dx, \end{aligned}$$

where $C = C(p, \Lambda)$. We now estimate each term on the right-hand side separately: for the first one, we have

$$\begin{aligned} &\int_{B_{(R+r)/2}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} dy \right) \left| \frac{\delta_h u(x)}{|h|^{\frac{\gamma+t}{2}}} \right|^2 \eta(x)^p dx \\ &\leq \left(\int_{B_{(R+r)/2}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} dy \right)^{\frac{p}{p-2}} dx \right)^{\frac{p-2}{p}} \\ &\quad \times \left(\int_{B_{(R+r)/2}} \left| \frac{\delta_h u(x)}{|h|^{\frac{\gamma+t}{2}}} \right|^p dx \right)^{\frac{2}{p}}. \end{aligned}$$

³We used that

$$|\delta_h \eta(x) - \delta_h \eta(y)| \leq |x-y| \int_0^1 \left| \nabla \eta(x+t(y-x)+h) - \nabla \eta(x+t(y-x)) \right| dt \leq |x-y| |h| \|D^2 \eta\|_{L^\infty}.$$

Then by Jensen's inequality⁴ and with some simple manipulations, we get

$$\begin{aligned} & \int_{B_{\frac{R+r}{2}}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} dy \right)^{\frac{p}{p-2}} dx \\ & \leq C \left(\frac{1}{sR^{sp}} \right)^{\frac{2}{p-2}} \int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left(|(\mathcal{V}_p)_h|^2 + |\mathcal{V}_p|^2 \right) \frac{1}{|x-y|^{N+sp}} dy dx, \end{aligned}$$

for some $C = C(N, p) > 0$, where we recall the definition of \mathcal{V}_p , given in (3.2). For the second term in the right-hand side of (3.11), we have

$$\begin{aligned} & \int_{B_{\frac{R+r}{2}}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} \left| \frac{\delta_h u(y)}{|h|^t} \right| dy \right) \left| \frac{\delta_h u(x)}{|h|^\gamma} \right| \eta(x)^p dx \\ & \leq \left(\int_{B_{\frac{R+r}{2}}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} \left| \frac{\delta_h u(y)}{|h|^t} \right| dy \right)^{p'} dx \right)^{\frac{1}{p'}} \\ & \quad \times \left(\int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u(x)}{|h|^\gamma} \right|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

By proceeding similarly as before, i.e. by using Jensen's inequality we also have

$$\begin{aligned} & \int_{B_{\frac{R+r}{2}}} \left(\int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2}}{|x-y|^{N+sp}} \left| \frac{\delta_h u(y)}{|h|^t} \right| dy \right)^{p'} dx \\ & \leq C \left(\frac{1}{sR^{sp}} \right)^{\frac{1}{p-1}} \int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{\left(|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2} \right)^{p'}}{|x-y|^{N+sp}} \left| \frac{\delta_h u(y)}{|h|^t} \right|^{p'} dy dx. \end{aligned}$$

Thus from (3.11) we get the following lower-bound

(3.12)

$$\begin{aligned} \mathcal{I}_2 & \geq -C \left(\frac{1}{sR^{sp}} \right)^{\frac{2}{p}} \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left(|(\mathcal{V}_p)_h|^2 + |\mathcal{V}_p|^2 \right) \frac{dy dx}{|x-y|^{N+sp}} \right)^{\frac{p-2}{p}} \left(\int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^{\frac{\gamma+t}{2}}} \right|^p dx \right)^{\frac{2}{p}} \\ & \quad - C \left(\frac{1}{sR^{sp}} \right)^{\frac{1}{p}} \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{\left(|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2} \right)^{p'}}{|x-y|^{N+sp}} \left| \frac{\delta_h u(y)}{|h|^t} \right|^{p'} dy dx \right)^{\frac{1}{p'}} \\ & \quad \times \left(\int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

By a further application of Hölder's inequality with exponents

$$\frac{p}{p'} = p - 1 \quad \text{and} \quad \frac{p}{p - p'} = \frac{p - 1}{p - 2},$$

⁴With respect to the measure $|x-y|^{-N-sp} dy$ which is finite on $\mathbb{R}^N \setminus B_R$, for every $x \in B_{\frac{R+r}{2}}$.

the second term in the right-hand side of (3.12) is estimated by

$$\begin{aligned} & \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left(|u_h(x) - u_h(y)|^{p-2} + |u(x) - u(y)|^{p-2} \right)^{p'} \left| \frac{\delta_h u(y)}{|h|^t} \right|^{p'} \frac{dy dx}{|x-y|^{N+sp}} \right)^{\frac{1}{p'}} \\ & \leq C \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left(|(\mathcal{V}_p)_h|^2 + |\mathcal{V}_p|^2 \right) \frac{dy dx}{|x-y|^{N+sp}} \right)^{\frac{p-2}{p}} \\ & \quad \times \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left| \frac{\delta_h u(y)}{|h|^t} \right|^p \frac{dy dx}{|x-y|^{N+sp}} \right)^{\frac{1}{p}}. \end{aligned}$$

By using this estimate, we obtain for \mathcal{I}_2 the following lower bound

$$\begin{aligned} (3.13) \quad \mathcal{I}_2 & \geq -C \left(\frac{1}{s R^{sp}} \right)^{\frac{2}{p}} \left(\int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^{\frac{\gamma+t}{2}}} \right|^p dx \right)^{\frac{2}{p}} \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{|(\mathcal{V}_p)_h|^2 + |\mathcal{V}_p|^2}{|x-y|^{N+sp}} dy dx \right)^{\frac{p-2}{p}} \\ & \quad - C \left(\frac{1}{s R^{sp}} \right)^{\frac{1}{p}} \left(\int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx \right)^{\frac{1}{p}} \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{|(\mathcal{V}_p)_h|^2 + |\mathcal{V}_p|^2}{|x-y|^{N+sp}} dy dx \right)^{\frac{p-2}{p}} \\ & \quad \times \left(\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left| \frac{\delta_h u(y)}{|h|^t} \right|^p \frac{dy dx}{|x-y|^{N+sp}} \right)^{\frac{1}{p}}. \end{aligned}$$

Observe that the last term is the integral of a nonlocal quantity containing a difference quotient u . By recalling Definition 1.2 and using that $0 < |h| < h_0$, we get⁵

$$\begin{aligned} (3.14) \quad \int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left| \frac{\delta_h u(y)}{|h|^t} \right|^p \frac{1}{|x-y|^{N+sp}} dy dx & = \frac{C}{R^{sp}} \int_{B_{\frac{R+r}{2}}} \text{Snail} \left(\frac{\delta_h u}{|h|^t}; x, B_R \right)^p dx \\ & \leq \frac{C}{R^{sp}} \langle u \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{R+r}{2}}; B_R)}^p. \end{aligned}$$

⁵Observe that

$$\frac{1}{2} \text{dist} \left(B_{\frac{R+r}{2}}, \mathbb{R}^N \setminus B_R \right) = \frac{R-r}{4} > h_0,$$

thus we have

$$\sup_{0 < |h| < h_0} \int_{B_{\frac{R+r}{2}}} \text{Snail} \left(\frac{\delta_h u}{|h|^t}; x, B_R \right)^p dx \leq \langle u \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{R+r}{2}}; B_R)}^p,$$

by the very definition of the latter.

Finally, for the common \mathcal{V}_p term in (3.13), we have

$$\begin{aligned}
\int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \left(|(\mathcal{V}_p)_h|^2 + |\mathcal{V}_p|^2 \right) \frac{dy dx}{|x-y|^{N+sp}} &\leq C \int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{|u_h(x)|^p + |u_h(y)|^p}{|x-y|^{N+sp}} dy dx \\
&+ C \int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{|u(x)|^p + |u(y)|^p}{|x-y|^{N+sp}} dy dx \\
&\leq \frac{C}{s} \left(\frac{1}{R-r} \right)^{sp} \int_{B_{\frac{R+r}{2}+h_0}} |u|^p dx \\
&+ C \left(\frac{1}{R} \right)^{sp} \int_{B_{\frac{R+r}{2}}} \text{Snail}(u; x; B_R)^p dx \\
&+ C \left(\frac{1}{R} \right)^{sp} \int_{B_{\frac{R+r}{2}}} \text{Snail}(u_h; x; B_R)^p dx,
\end{aligned}$$

for some $C = C(N, p) > 0$. Moreover, by the monotonicity properties of Snails encoded in Lemma 2.9, we have

$$\int_{B_{\frac{R+r}{2}}} \text{Snail}(u; x; B_R)^p dx \leq C \left(\frac{R+r}{R-r} \right)^N \left(\frac{R+h_0}{R-r} \right)^{sp} \langle u \rangle_{\mathcal{X}_s^p(B_{\frac{R+r}{2}+h_0}; B_{R+h_0})}^p.$$

With a simple change of variables and by observing that

$$B_{(R+r)/2-h} \subset B_{(R+r)/2+h_0} \quad B_R - h \subset B_{R+h_0},$$

still by Lemma 2.9 we get again

$$\begin{aligned}
\int_{B_{\frac{R+r}{2}}} \text{Snail}(u_h; x; B_R)^p dx &= C R^{sp} \int_{B_{\frac{R+r}{2}-h}} \int_{\mathbb{R}^N \setminus (B_R-h)} \frac{|u(y)|^p}{|x-y|^{N+sp}} dy dx \\
&\leq C \left(\frac{R+r}{R-r} \right)^N \left(\frac{R+h_0}{R-r} \right)^{sp} \langle u \rangle_{\mathcal{X}_s^p(B_{\frac{R+r}{2}+h_0}; B_{R+h_0})}^p.
\end{aligned}$$

By keeping everything together and recalling Definition 1.2, we get

$$(3.15) \quad \int_{B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{|(\mathcal{V}_p)_h|^2 + |\mathcal{V}_p|^2}{|x-y|^{N+sp}} dy dx \leq \left(\frac{R+r}{R-r} \right)^N \left(\frac{R+h_0}{R-r} \right)^{sp} \frac{C}{s(R-s)^{sp}} \langle u \rangle_{\mathcal{X}_s^p(B_{\frac{R+r}{2}+h_0}; B_{R+h_0})}^p,$$

still for $C = C(N, p) > 0$. By using (3.14) and (3.15) in (3.13) in conjunction with Young's inequality, we finally end up with

$$\begin{aligned}
\mathcal{I}_2 &\geq -\frac{C}{(R-r)^{sp}} \frac{1}{s} \int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^{\frac{\gamma+t}{2}}} \right|^p dx - \frac{C}{(R-r)^{sp}} \frac{1}{s} \int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx \\
&- \left(\frac{R+r}{R-r} \right)^N \left(\frac{R+h_0}{R-r} \right)^{sp} \frac{C}{s(R-r)^{sp}} \langle u \rangle_{\mathcal{X}_s^p(B_{\frac{R+r}{2}+h_0}; B_{R+h_0})}^p - \frac{C}{R^{sp}} \langle u \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{R+r}{2}}; B_R)}^p.
\end{aligned}
\tag{3.16}$$

Estimate of \mathcal{I}_3 . This is estimated exactly in the same manner as \mathcal{I}_2 . We thus get

$$\begin{aligned}
\mathcal{I}_3 &\geq -\frac{C}{(R-r)^{sp}} \frac{1}{s} \int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^{\frac{\gamma+t}{2}}} \right|^p dx - \frac{C}{(R-r)^{sp}} \frac{1}{s} \int_{B_{\frac{R+r}{2}}} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx \\
&- \left(\frac{R+r}{R-r} \right)^N \left(\frac{R+h_0}{R-r} \right)^{sp} \frac{C}{s(R-r)^{sp}} \langle u \rangle_{\mathcal{X}_s^p(B_{\frac{R+r}{2}+h_0}; B_{R+h_0})}^p - \frac{C}{R^{sp}} \langle u \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{R+r}{2}}; B_R)}^p.
\end{aligned}
\tag{3.17}$$

Conclusion. From (3.4) we have

$$\begin{aligned} \mathcal{I}_1 &\leq |\mathcal{I}_2| + |\mathcal{I}_3| + \int_{\Omega} |\delta_h f| \left| \frac{\delta_h u}{|h|^{\gamma+t}} \right| \eta^p dx \\ &\leq |\mathcal{I}_2| + |\mathcal{I}_3| + (1-s)^{\frac{1}{p-1}} R^{sp'} \int_{B_R} \left| \frac{\delta_h f}{|h|^s} \right|^{p'} dx + \frac{1}{(1-s)R^{sp}} \int_{B_R} \left| \frac{\delta_h u}{|h|^{\gamma+t-s}} \right|^p dx. \end{aligned}$$

Thus by using (3.10), (3.16) and (3.17) we get the conclusion, by recalling that $|h| < h_0 < 1$ and that

$$\gamma + t - s \leq \gamma,$$

which follows from the hypothesis $t \leq s \leq \gamma$. \square

Remark 3.2 (Correction factor). Observe that the nonlocal terms $\langle u \rangle_{\mathcal{X}_s^p}$ and $\langle u \rangle_{\mathcal{Y}_s^{t,p}}$ in the right-hand side (3.3) do not contain the correction factor $(1-s)^{-1}$, as it is natural. Indeed, if we multiply (3.3) by $(1-s)$ these terms have to disappear in the limit $s \nearrow 1$, which corresponds to the equation becoming local.

3.2. Improving Lemma. The proof of Theorem 1.4 is based on a combination of Proposition 3.1 and of the following result, which is valid for general functions. This simple result is useful in order to handle the left-hand side of (3.3). Here second order differences and Besov spaces come into play.

Lemma 3.3 (Besov-Nikol'skii improvement). *Let $p \geq 2$, $0 < s < 1$ and $0 \leq t \leq s$. Let $B_r \Subset B_R \Subset \Omega$ be a couple of concentric balls. We take η a standard C^2 cut-off function such that*

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \quad \text{on } B_r, \quad \eta \equiv 0 \quad \text{on } \mathbb{R}^N \setminus B_{\frac{R+r}{2}}, \quad |\nabla \eta| \leq \frac{c_N}{R-r}, \quad |D^2 \eta| \leq \frac{c_N}{(R-r)^2}.$$

Let us assume that for some γ such that $s \leq \gamma \leq 1$ and some

$$0 < h_0 < \frac{1}{4} \min \left\{ \text{dist}(B_R; \partial\Omega), R-r, 1 \right\},$$

we have

$$\mathcal{M}_\gamma := \sup_{0 < |h| < h_0} \left[\frac{\delta_h(u\eta)}{|h|^{\frac{\gamma+t}{p}}} \right]_{W^{s,p}(B_R)}^p < +\infty.$$

Then, by setting for simplicity

$$(3.18) \quad \Gamma := \frac{\gamma + t + ps}{p},$$

we have the Besov-Nikol'skii estimate

$$(3.19) \quad [u\eta]_{\mathcal{B}_{\infty,p}^{\Gamma}(\mathbb{R}^N)}^p \leq C \left[(1-s) \mathcal{M}_\gamma + h_0^{-\Gamma p} \|u\|_{L^p(B_{R+h_0})}^p \right],$$

for some $C = C(N, p) > 0$. In particular, we have the following estimates, for a possibly different constant $C = C(N, p) > 0$:

- if $\boxed{\Gamma < 1}$

$$(3.20) \quad \sup_{0 < |h| < h_0} \left\| \frac{\delta_h u}{|h|^{\Gamma}} \right\|_{L^p(B_r)}^p \leq \frac{C}{(1-\Gamma)^p} \left[(1-s) \mathcal{M}_\gamma + h_0^{-\Gamma p} \|u\|_{L^p(B_{R+h_0})}^p \right];$$

- if $\boxed{\Gamma = 1}$, for every $0 < \tau < 1$

$$(3.21) \quad \sup_{0 < |h| < h_0} \left\| \frac{\delta_h u}{|h|^{\tau}} \right\|_{L^p(B_r)}^p \leq \frac{C}{(1-\tau)^p} \left[(1-s) \mathcal{M}_\gamma + h_0^{-p} \|u\|_{L^p(B_{R+h_0})}^p \right];$$

- if $\boxed{\Gamma > 1}$

$$(3.22) \quad \|\nabla u\|_{L^p(B_r)}^p \leq \|\nabla(u\eta)\|_{L^p(\mathbb{R}^N)}^p \leq \frac{C}{(\Gamma-1)^p} \left[(1-s)\mathcal{M}_\gamma + h_0^{-\Gamma p} \|u\|_{L^p(B_{R+h_0})}^p \right],$$

and for every $0 < \tau < \Gamma - 1$

$$(3.23) \quad [\nabla u]_{W^{\tau,p}(B_r)}^p \leq \frac{C}{(\Gamma-1-\tau)\tau} \left(\frac{h_0^{-1}}{(2-\Gamma)(\Gamma-1)} \right)^p \left[(1-s)\mathcal{M}_\gamma + h_0^{-\Gamma p} \|u\|_{L^p(B_{R+h_0})}^p \right].$$

Proof. By using the hypothesis and (2.9) with the choice

$$\psi = \frac{\delta_h(u\eta)}{|h|^{\frac{\gamma+t}{p}}},$$

for every $0 < |h| < h_0$ we get

$$\int_{\mathbb{R}^N} \left| \delta_\xi \left(\frac{\delta_h(u\eta)}{|h|^{\frac{\gamma+t}{p}}} \right) \right|^p \frac{1}{|\xi|^{sp}} dx \leq C(1-s)\mathcal{M}_\gamma, \quad \text{for every } \xi \in \mathbb{R}^N \setminus \{0\}.$$

If we now choose $\xi = h$, recall (2.2) and take the supremum over $0 < |h| < h_0$, we obtain

$$\sup_{0 < |h| < h_0} \int_{\mathbb{R}^N} \left| \left(\frac{\delta_h^2(u\eta)}{|h|^{\frac{\gamma+t+s}{p}}} \right) \right|^p dx \leq C(1-s)\mathcal{M}_\gamma$$

By joining (2.3) and the previous estimate we have

$$[u\eta]_{\mathcal{B}_\infty^{\Gamma,p}(\mathbb{R}^N)}^p \leq C \left[(1-s)\mathcal{M}_\gamma + h_0^{-\Gamma p} \|u\|_{L^p(B_{R+h_0})}^p \right],$$

where we used the expedient notation (3.18). This proves (3.19). We then treat each case separately.

Case $\boxed{\Gamma < 1}$. We now use Lemma 2.3 for $U = u\eta$, then from (3.19) we get

$$(3.24) \quad \sup_{0 < |h| < h_0} \left\| \frac{\delta_h(u\eta)}{|h|^\Gamma} \right\|_{L^p(\mathbb{R}^N)}^p \leq \frac{C}{(1-\Gamma)^p} \left[(1-s)\mathcal{M}_\gamma + \left(1 + h_0^{-\Gamma p}\right) \|u\|_{L^p(B_{R+h_0})}^p \right].$$

By using the discrete Leibniz rule (2.1), the triangle inequality and the Lipschitz character of η , we have

$$(3.25) \quad \left\| \frac{\delta_h u}{|h|^\Gamma} \right\|_{L^p(B_r)}^p \leq C \left\| \frac{\delta_h(u\eta)}{|h|^\Gamma} \right\|_{L^p(\mathbb{R}^N)}^p + \frac{C}{(R-r)^p} h_0^{p-\Gamma p} \|u\|_{L^p(B_{R+h_0})}^p,$$

for $C = C(p) > 0$. Then (3.20) follows by using the previous estimate in (3.24) and observing that $h_0 \leq (R-r)$ and that $h_0 < 1$.

Case $\boxed{\Gamma = 1}$. Let $\tau < 1$, we begin with the following remark

$$\sup_{0 < |h| < h_0} \left\| \frac{\delta_h^2(u\eta)}{|h|^\tau} \right\|_{L^p(\mathbb{R}^N)}^p \leq h_0^{(1-\tau)p} [u\eta]_{\mathcal{B}_\infty^{1,p}(\mathbb{R}^N)}^p.$$

By using (2.5) on the left and (3.19) on the right, we get

$$\sup_{0 < |h| < h_0} \left\| \frac{\delta_h(u\eta)}{|h|^\tau} \right\|_{L^p(\mathbb{R}^N)}^p \leq \frac{C}{(1-\tau)^p} \left[h_0^{(1-\tau)p} (1-s)\mathcal{M}_\gamma + (h_0^{-\tau p} + h_0^{-p} + 1) \|u\|_{L^p(B_{R+h_0})}^p \right],$$

possibly with a different constant $C = C(N, p) > 0$. Finally, we use again (3.25) to remove the dependence on η and the fact that $h_0 < 1$.

Case $\boxed{\Gamma > 1}$. We first observe that due to the restrictions on the parameters, we always have $\Gamma < 2$. Then we use Proposition 2.4 with $\psi = u\eta$ and from (3.19) we get

$$\|\nabla(u\eta)\|_{L^p(\mathbb{R}^N)}^p \leq C \left[\frac{(1-s)\mathcal{M}_\gamma}{(\Gamma-1)^p} + (1+h_0^{-\Gamma p}) \|u\|_{L^p(B_{R+h_0})}^p \right].$$

By recalling that $\eta \equiv 1$ on B_r and observing that $(\Gamma-1) < 1$, we get (3.22). As for (3.23), we observe that still by Proposition 2.4 we also have

$$\sup_{0 < |h| < h_0} \left\| \frac{\delta_h \nabla(u\eta)}{|h|^{\Gamma-1}} \right\|_{L^p(\mathbb{R}^N)}^p \leq \frac{C}{(2-\Gamma)^p (\Gamma-1)^p} \left[(1-s)\mathcal{M}_\gamma + h_0^{-\Gamma p} \|u\|_{L^p(B_{R+h_0})}^p \right].$$

If we now apply Proposition 2.7 to the compactly supported function $\psi = \nabla(u\eta)$ and the exponent $\beta = \Gamma-1$ we get

$$\begin{aligned} [\nabla u]_{W^{\tau,p}(B_r)}^p &\leq [\nabla(u\eta)]_{W^{\tau,p}(\mathbb{R}^N)}^p \\ &\leq C \left(\frac{h_0^{(\Gamma-1-\tau)p}}{\Gamma-1-\tau} \sup_{0 < |h| < h_0} \left\| \frac{\delta_h \nabla(u\eta)}{|h|^{\Gamma-1}} \right\|_{L^p(\mathbb{R}^N)}^p + \frac{h_0^{-\tau p}}{\tau} \|\nabla(u\eta)\|_{L^p(\mathbb{R}^N)}^p \right), \end{aligned}$$

for every $0 < \tau < \Gamma-1$. The right-hand side is now estimated by appealing to the previous two estimates, thus we conclude the proof with standard manipulations. \square

4. PROOF OF THEOREM 1.4

Let $R > 0$ and $B_R \Subset \Omega$, we want to prove the estimates (1.9) and (1.10)-(1.11) on the ball $B_{R/2}$. Without loss of generality, we can assume that B_R is centered at the origin. Observe that if we consider the rescaled functions

$$u_R(x) = u(Rx) \quad \text{and} \quad f_R(x) = R^{sp} f(Rx), \quad x \in R^{-1}\Omega,$$

then $u_R \in W_{loc}^{s,p}(R^{-1}\Omega) \cap \mathcal{Y}_s^{t,p}(R^{-1}\Omega)$ is a local weak solution in the rescaled set $R^{-1}\Omega$, with right-hand side f_R . Thus we just need to estimate

$$[u_R]_{W^{\tau,p}(B_{1/2})}, \quad \text{for every } \tau < \frac{t+sp}{p-1},$$

or

$$\|\nabla u_R\|_{L^p(B_{1/2})} + [\nabla u_R]_{W^{\tau,p}(B_{1/2})}, \quad \text{for every } \tau < \frac{t+sp}{p} - \frac{p-1}{p}.$$

The desired results will be then obtained by scaling back.

4.1. The general scheme. As explained in the Introduction, the desired estimates are proved by means on an iterative scheme. First of all, we define the sequence

$$(4.1) \quad \gamma_0 = s, \quad \gamma_{i+1} = \frac{\gamma_i + t + sp}{p}.$$

We observe γ_i is strictly increasing and

$$(4.2) \quad \lim_{i \rightarrow \infty} \gamma_i = \frac{t+sp}{p-1}.$$

We take any index $i_0 \in \mathbb{N} \setminus \{0\}$ such that

$$\gamma_{i_0-1} < 1,$$

the precise choice of i_0 will be done below. We define the decreasing sequence of radii

$$r_i := \frac{3}{4} - \frac{i}{i_0} \frac{1}{4}, \quad i \in \{0, \dots, i_0\}.$$

Accordingly, we consider the concentric balls B_{r_i} and observe that

$$B_{r_0} = B_{3/4} \quad \text{and} \quad B_{r_{i_0}} = B_{1/2}.$$

We point out that by construction, we have

$$(4.3) \quad r_i - r_{i+1} = \frac{1}{4i_0}, \quad \frac{1}{2} \leq r_i < 1, \quad \text{and} \quad \frac{r_i}{r_i - r_{i+1}} < 4i_0.$$

Then we define

$$(4.4) \quad h_0 = \frac{1}{100i_0},$$

thus with such a choice we have

$$h_0 < \frac{1}{4} \min \left\{ \text{dist} \left(B_{r_i}, \partial(R^{-1}\Omega) \right), r_i - r_{i+1}, 1 \right\}, \quad i = 0, \dots, i_0 - 1.$$

Finally, for every $i \in \{0, \dots, i_0 - 1\}$ we choose a standard cut-off function $\eta_i \in C_0^2(B_{r_i})$ such that

$$\begin{aligned} 0 \leq \eta_i \leq 1, \quad \eta_i \equiv 1 \quad \text{on } B_{r_{i+1}}, \quad \eta_i \equiv 0 \quad \text{on } \mathbb{R}^N \setminus B_{\frac{r_i+r_{i+1}}{2}}, \\ |\nabla \eta_i| \leq \frac{c_N}{r_i - r_{i+1}} = 4c_N i_0 \quad \text{and} \quad |D^2 \eta_i| \leq \frac{c_N}{(r_i - r_{i+1})^2} = 16c_N i_0^2. \end{aligned}$$

By taking into account (4.3) and (4.4), for every $0 < |h| < h_0$ by Proposition 3.1 with simple manipulations we get (recall the definition of h_0 and that $i_0 \geq 1$)

$$(4.5) \quad \begin{aligned} \left[\frac{\delta_h(u_R \eta_i)}{|h|^{\frac{\gamma_i+t}{p}}} \right]_{W^{s,p}(B_{r_i})}^p &\leq \frac{C}{(1-s)s} i_0^{2p} \left\| \frac{\delta_h u_R}{|h|^{\gamma_i}} \right\|_{L^p(B_{r_i})}^p \\ &\quad + C i_0^{p-\gamma_i-t} \left([u_R]_{W^{s,p}(B_{r_i+h_0})}^p + \frac{1}{s(1-s)} \|u_R\|_{L^p(B_{r_i+h_0})}^p \right) \\ &\quad + C (1-s)^{\frac{1}{p-1}} \left\| \frac{\delta_h f_R}{|h|^s} \right\|_{L^{p'}(B_{r_i})}^{p'} \\ &\quad + \frac{C}{s} i_0^{N+2p} \langle u_R \rangle_{\mathcal{X}_s^p(B_{\frac{r_i+r_{i+1}}{2}+h_0}, B_{r_i+h_0})}^p \\ &\quad + C \langle u_R \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{r_i+r_{i+1}}{2}}; B_{r_i})}^p, \quad i = 0, \dots, i_0 - 1. \end{aligned}$$

for some $C = C(N, p) > 0$. Before going on, we try to simplify the previous estimate.

By construction $B_{r_i+h_0} \subset B_1$ for every $i = 0, \dots, i_0$, then by Proposition 2.6 (*local case*) we get

$$(4.6) \quad (1-s)^{\frac{1}{p-1}} \sup_{0 < |h| < h_0} \left\| \frac{\delta_h f_R}{|h|^s} \right\|_{L^{p'}(B_{r_i})}^{p'} \leq \frac{C}{h_0^{p'}} (1-s^{p'}) \left[[f_R]_{W^{s,p'}(B_1)}^{p'} + \frac{1}{s(1-s)} \|f_R\|_{L^{p'}(B_1)}^{p'} \right],$$

where we used again that $h_0 < 1$. Also, by the monotonicity properties of Lemma 2.9

$$(4.7) \quad \langle u_R \rangle_{\mathcal{X}_s^p(B_{\frac{r_i+r_{i+1}}{2}+h_0}, B_{r_i+h_0})}^p \leq C i_0^{N+p} \langle u_R \rangle_{\mathcal{X}_s^p(B_{\frac{3}{4}}; B_1)}^p.$$

Finally, by observing that

$$\frac{1}{2} \text{dist} \left(B_{\frac{r_i+r_{i+1}}{2}}, \mathbb{R}^N \setminus B_{r_i} \right) = \frac{r_i - r_{i+1}}{4} = \frac{1}{16i_0} \leq \frac{1}{16} = \frac{1}{2} \text{dist} \left(B_{\frac{3}{4}}, \mathbb{R}^N \setminus B_{\frac{7}{8}} \right),$$

by (2.17) with the choices

$$K_1 = B_{\frac{r_i+r_{i+1}}{2}}, \quad E_1 = B_{r_i}, \quad K_2 = B_{\frac{3}{4}}, \quad E_2 = B_{\frac{7}{8}},$$

we get

$$(4.8) \quad \begin{aligned} \langle u_R \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{r_i+r_{i+1}}{2}}; B_{r_i})}^p &= \sup_{0 < |h| < \frac{r_i - r_{i+1}}{4}} \int_{B_{\frac{r_i+r_{i+1}}{2}}} \text{Snail} \left(\frac{\delta_h u_R}{|h|^t}; x, B_{r_i} \right)^p dx \\ &\leq \langle u_R \rangle_{\mathcal{Y}_s^{t,p}(B_{\frac{3}{4}}; B_{\frac{7}{8}})}^p + C i_0^{N+p} \sup_{0 < |h| < \frac{1}{16}} \left\| \frac{\delta_h u_R}{|h|^t} \right\|_{L^p(B_{\frac{7}{8}})}^p, \end{aligned}$$

for some $C = C(N, p) > 0$. The last local term can be further estimated by Proposition 2.6 (*local case*) as follows (recall that $t \leq s$)

$$(4.9) \quad \sup_{0 < |h| < \frac{1}{16}} \left\| \frac{\delta_h u_R}{|h|^t} \right\|_{L^p(B_R)}^p \leq C(1-s) \left[[u_R]_{W^{s,p}(B_1)}^p + \frac{1}{s(1-s)} \|u_R\|_{L^p(B_1)}^p \right].$$

By using (4.6), (4.7), (4.8) and (4.9) in (4.5) and observing that

$$h_0^{-p'} \leq C i_0^{2(N+p)}, \quad i_0^{p-\gamma_i-t} \leq C i_0^{2(N+p)}, \quad i_0^{N+p} \leq i_0^{3(N+p)},$$

for every $0 < |h| < h_0$ we obtain

$$(4.10) \quad \left[\frac{\delta_h(u_R \eta_i)}{|h|^{\frac{\gamma_i+t}{p}}} \right]_{W^{s,p}(B_{r_i})}^p \leq \frac{C}{(1-s)} i_0^{2p} \left\| \frac{\delta_h u_R}{|h|^{\gamma_i}} \right\|_{L^p(B_{r_i})}^p + C i_0^{3(N+p)} \mathcal{A}_1(u_R, f_R), \quad i = 0, \dots, i_0 - 1,$$

where \mathcal{A}_1 is the quantity defined in (1.8). In what follows, for simplicity we just write \mathcal{A}_1 in place of $\mathcal{A}_1(u_R, f_R)$. Observe that $\mathcal{A}_1 < +\infty$, thanks to the assumptions on u and f .

We now set

$$\mathcal{M}_{\gamma_i} := \sup_{0 < |h| < h_0} \left[\frac{\delta_h(u_R \eta_i)}{|h|^{\frac{\gamma_i+t}{p}}} \right]_{W^{s,p}(B_{r_i})}^p, \quad i = 0, \dots, i_0 - 1,$$

and claim that

$$(4.11) \quad \mathcal{M}_{\gamma_i} < +\infty, \quad \text{for every } i = 0, \dots, i_0 - 1.$$

This is true by a finite induction: for $i = 0$, by combining (4.10) and Proposition 2.6 (*local case*) we get

$$\mathcal{M}_{\gamma_0} \leq \frac{C}{s} \frac{i_0^{2p}}{h_0^p} \left[[u_R]_{W^{s,p}(B_1)}^p + \frac{1}{s(1-s)} \|u_R\|_{L^p(B_1)}^p \right] + C i_0^{3(N+p)} \mathcal{A}_1,$$

where we used again that $B_{r_0+h_0} \subset B_1$. Thus the claim is true for $i = 0$. Also, by using the definitions (1.8) and (4.4), we can infer

$$(4.12) \quad \mathcal{M}_{\gamma_0} \leq \frac{C_0}{s} i_0^{3(N+p)} \mathcal{A}_1,$$

where as usual $C_0 = C_0(N, p) > 0$.

Let us now assume that $\mathcal{M}_{\gamma_i} < +\infty$ for an index⁶ $i \in \{0, \dots, i_0 - 2\}$, then we can use Lemma 3.3. Namely, by combining (3.20) and (4.10) we get

$$\mathcal{M}_{\gamma_{i+1}} \leq \frac{C i_0^{2p}}{s(1-\gamma_{i+1})^p} \mathcal{M}_{\gamma_i} + \frac{C i_0^p}{(1-\gamma_{i+1})^p} h_0^{-\gamma_{i+1}p} \frac{i_0^p}{s(1-s)} \|u_R\|_{L^p(B_1)}^p + C i_0^{3(N+p)} \mathcal{A}_1,$$

where C is a possibly different constant still $C = C(N, p) > 0$ and we used the relation between γ_i and γ_{i+1} . This in turn shows that $\mathcal{M}_{\gamma_{i+1}} < +\infty$ and thus the validity of (4.11).

As before, at first we try to simplify the previous estimate. Observe that

$$h_0^{-\gamma_{i+1}p} \frac{i_0^p}{s(1-s)} \|u\|_{L^p(B_1)}^p \leq C i_0^{2p} \mathcal{A}_1, \quad i = 0, \dots, i_0 - 2,$$

⁶Of course, if $i_0 = 1$ there is nothing to prove.

where we used the definition (4.4) of h_0 and the fact that $\gamma_{i+1} \leq \gamma_{i_0-1} < 1$. From the previous discussion and (4.12) we thus obtain the iterative scheme

$$(4.13) \quad \begin{cases} \mathcal{M}_{\gamma_0} \leq \frac{C_0}{s} i_0^{3(N+p)} \mathcal{A}_1, \\ \mathcal{M}_{\gamma_{i+1}} \leq \frac{C_1 i_0^{2p}}{s(1-\gamma_{i+1})^p} \mathcal{M}_{\gamma_i} + \frac{C_2 i_0^{3(N+p)}}{(1-\gamma_{i+1})^p} \mathcal{A}_1, \quad \text{for } i = 0, \dots, i_0 - 2, \end{cases}$$

where $C_1 = C_1(N, p) > 0$ and $C_2 = C_2(N, p) > 0$. It is intended that the second estimate in (4.13) is void when $i_0 = 1$.

4.2. Case $t + sp \leq (p - 1)$. We fix a differentiability exponent

$$s \leq \tau < \frac{t + sp}{p - 1},$$

as in (1.9), then the index $i_0 \in \mathbb{N} \setminus \{0\}$ above is chosen so that

$$\tau < \gamma_{i_0} < 1.$$

This is possible thanks to (4.2). We observe that $\gamma_{i+1} \leq \gamma_{i_0} < 1$ for every $i = 0, \dots, i_0 - 1$. By using this observation in (4.13) and iterating, we get

$$(4.14) \quad \begin{aligned} \mathcal{M}_{\gamma_{i_0-1}} &\leq \left(\frac{C_1 i_0^{2p}}{s(1-\gamma_{i_0})^p} \right)^{i_0-1} \frac{C_0}{s} i_0^{3(N+p)} \mathcal{A}_1 \\ &+ \left[\sum_{i=0}^{i_0-2} \left(\frac{C_1 i_0^{2p}}{s(1-\gamma_{i_0})^p} \right)^i \right] \left[\frac{C_2 i_0^{3(N+p)}}{(1-\gamma_{i_0})^p} \mathcal{A}_1 \right] \leq \left(\frac{C_3}{s} \right)^{i_0} \left(\frac{i_0^{3(N+p)}}{(1-\gamma_{i_0})^p} \right)^{i_0} \mathcal{A}_1, \end{aligned}$$

where $C_3 = \max\{C_0, C_1, C_2, 1\}$. We are ready to perform the final step. We use again Lemma 3.3, then (3.19) and (4.14) yield the Besov-Nikol'skii estimate

$$[u_R \eta_{i_0-1}]_{\mathcal{B}^{\gamma_{i_0}, p}(\mathbb{R}^N)}^p \leq C \left[\left(\frac{C_3}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0})^p} \right)^{i_0} (1-s) \mathcal{A}_1 + h_0^{-\gamma_{i_0} p} \|u_R\|_{L^p(B_1)}^p \right].$$

The left-hand side is estimated from below thanks to (2.4), thus we get

$$\sup_{0 < |h| < h_0} \left\| \frac{\delta_h(u_R \eta_{i_0-1})}{|h|^{\gamma_{i_0}}} \right\|_{L^p(\mathbb{R}^N)}^p \leq \frac{C}{(1-\gamma_{i_0})^p} \left[\left(\frac{C_3}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0})^p} \right)^{i_0} (1-s) \mathcal{A}_1 + h_0^{-\gamma_{i_0} p} \|u_R\|_{L^p(B_1)}^p \right].$$

By recalling that i_0 has been chosen so that $\gamma_{i_0} > \tau$, by applying Proposition 2.7 we get

$$[u_R \eta_{i_0-1}]_{W^{\tau, p}(\mathbb{R}^N)}^p \leq \frac{h_0^{(\gamma_{i_0} - \tau)p}}{\gamma_{i_0} - \tau} \frac{C}{(1-\gamma_{i_0})^p} \left[\left(\frac{C_3}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0})^p} \right)^{i_0} (1-s) \mathcal{A}_1 + \frac{h_0^{-\gamma_{i_0} p}}{\tau} \|u_R\|_{L^p(B_1)}^p \right].$$

On the other hand $\eta_{i_0-1} \equiv 1$ on $B_{r_{i_0}} = B_{1/2}$ and by definition of h_0

$$\frac{h_0^{-\gamma_{i_0} p}}{\tau} \|u_R\|_{L^p(B_1)}^p \leq C \frac{(1-s)}{\tau} i_0^p \mathcal{A}_1.$$

Thus we conclude with the estimate (we use that $\tau \geq s$ and again $h_0 < 1$)

$$(4.15) \quad [u_R]_{W^{\tau, p}(B_{1/2})}^p \leq \frac{1}{(\gamma_{i_0} - \tau)} \left(\frac{C_4}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0})^p} \right)^{i_0+1} (1-s) \mathcal{A}_1,$$

where $C_4 > 0$ as usual depends on N and p only. We now scale back in order to catch the desired estimate for u in $B_{R/2}$. By recalling the definition (1.8) of \mathcal{A}_1 , from (4.15) with a simple change of variables we exactly get (1.9). The constant \mathcal{C}_1 appearing in (1.9) is given by

$$\mathcal{C}_1 := \frac{1}{(\gamma_{i_0} - \tau)} \left(\frac{C_4}{s} \frac{i_0^{3(N+p)}}{(1 - \gamma_{i_0})^p} \right)^{i_0+1}.$$

4.3. **Case $t + sp > (p - 1)$.** We still consider the sequence $\{\gamma_i\}_{i \in \mathbb{N}}$ defined by (4.1). Observe that in this case

$$\lim_{i \rightarrow \infty} \gamma_i = \frac{t + sp}{p - 1} > 1.$$

Then this time the index $i_0 \in \mathbb{N} \setminus \{0\}$ is chosen so that

$$\gamma_{i_0-1} < 1 \quad \text{and} \quad \gamma_{i_0} = \frac{\gamma_{i_0-1} + t + sp}{p} \geq 1,$$

which is feasible. From the scheme (4.13), by using that $\gamma_{i+1} < \gamma_{i_0-1} < 1$ for $i = 0, \dots, i_0 - 2$, we get

$$(4.16) \quad \mathcal{M}_{\gamma_{i_0-1}} \leq \left(\frac{C_3}{s} \right)^{i_0} \left(\frac{i_0^{3(N+p)}}{(1 - \gamma_{i_0-1})^p} \right)^{i_0} \mathcal{A}_1,$$

exactly as in (4.14) and the constant C_3 is the same. We need to make a distinction between two possible cases

$$\gamma_{i_0} > 1 \quad \text{or} \quad \gamma_{i_0} = 1.$$

Case $\boxed{\gamma_{i_0} > 1}$. Since $\gamma_{i_0} > 1$, we can apply (3.22) of Lemma 3.3 and get (recall that $h_0 < 1$)

$$\begin{aligned} \|\nabla u_R\|_{L^p(B_{1/2})}^p &\leq \|\nabla(u_R \eta_{i_0-1})\|_{L^p(\mathbb{R}^N)}^p \\ &\leq \frac{C}{(\gamma_{i_0} - 1)^p} \left[(1 - s) \mathcal{M}_{\gamma_{i_0-1}} + h_0^{-p\gamma_{i_0}} \|u_R\|_{L^p(B_1)}^p \right], \end{aligned}$$

which shows that $\|\nabla u_R\|_{L^p(B_{1/2})} < +\infty$. By using (4.16) in the previous estimate and the definitions of h_0 and of \mathcal{A}_1 , we end up with

$$(4.17) \quad \begin{aligned} \|\nabla u_R\|_{L^p(B_{1/2})}^p &\leq \|\nabla(u_R \eta_{i_0-1})\|_{L^p(\mathbb{R}^N)}^p \\ &\leq \frac{C}{(\gamma_{i_0} - 1)^p} \left(\frac{C_5}{s} \frac{i_0^{3(N+p)}}{(1 - \gamma_{i_0-1})^p} \right)^{i_0} (1 - s) \mathcal{A}_1, \end{aligned}$$

where $C_5 = C_5(N, p) > 1$. By going back to the original solution u with a scaling, we get (1.10) with the constant \mathcal{C}_2 given by

$$\mathcal{C}_2 := \frac{1}{(\gamma_{i_0} - 1)^p} \left(\frac{C_6}{s} \frac{i_0^{3(N+p)}}{(1 - \gamma_{i_0-1})^p} \right)^{i_0+1},$$

and $C_6 = C_6(N, p) > 0$ as usual. We still need to prove the fractional differentiability of the gradient. Observe that if we directly apply estimate (3.23) of Lemma 3.3 with γ_{i_0-1} , we would get the weaker result

$$[\nabla u_R]_{W^{\tau,p}(B_{1/2})} < +\infty, \quad \text{for every } \tau < \gamma_{i_0} - 1.$$

Thus, we have to proceed differently. We start by observing that for the compactly supported function $u \eta_{i_0-1}$ we have

$$\left\| \frac{\delta_h(u_R \eta_{i_0-1})}{|h|} \right\|_{L^p(\mathbb{R}^N)}^p \leq \|\nabla(u_R \eta_{i_0-1})\|_{L^p(\mathbb{R}^N)}^p.$$

By using again (3.25) we also obtain

$$(4.18) \quad \sup_{0 < |h| < h_0} \left\| \frac{\delta_h u_R}{|h|} \right\|_{L^p(B_{r_{i_0-1}})}^p \leq C \|\nabla(u_R \eta_{i_0-1})\|_{L^p(\mathbb{R}^N)}^p + C h_0^{-p} \|u_R\|_{L^p(B_1)}^p.$$

We can now use Proposition 3.1 in the limit case $\gamma = 1$, this gives

$$\begin{aligned} \mathcal{M}_1 &:= \sup_{0 < |h| < h_0} \left[\frac{\delta_h(u_R \eta_{i_0-1})}{|h|^{\frac{1+t}{p}}} \right]_{W^{s,p}(B_{r_{i_0-1}})}^p \\ &\leq \frac{C}{(1-s)s} i_0^{2p} \sup_{0 < |h| < h_0} \left\| \frac{\delta_h u_R}{|h|} \right\|_{L^p(B_{r_{i_0-1}})}^p + C i_0^{2(N+p)} \mathcal{A}_1. \end{aligned}$$

By combining (4.18) and (4.17), \mathcal{M}_1 can be further estimated by

$$\mathcal{M}_1 \leq \frac{C}{s} i_0^{2p} \left(\frac{C_5}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0-1})^p} \right)^{i_0+1} \frac{1}{(\gamma_{i_0}-1)^p} \mathcal{A}_1.$$

Then by using estimate (3.23) of Lemma 3.3 for $\gamma = 1$ and the previous inequality for \mathcal{M}_1 , we get

$$\begin{aligned} [\nabla u_R]_{W^{\tau,p}(B_{1/2})}^p &\leq \frac{h_0^{-\tau p}}{(\Gamma-1-\tau)\tau} \frac{C}{(2-\Gamma)^p(\Gamma-1)^p} \left[\frac{C}{s} i_0^{2p} \left(\frac{C_5}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0-1})^p} \right)^{i_0+1} \frac{1-s}{(\gamma_{i_0}-1)^p} \mathcal{A}_1 \right. \\ &\quad \left. + h_0^{-\Gamma p} \|u_R\|_{L^p(B_1)}^p \right], \end{aligned}$$

where $\Gamma = (1+t+sp)/p$. The usual elementary manipulations used so far then give

$$[\nabla u_R]_{W^{\tau,p}(B_{1/2})}^p \leq \frac{(2-\Gamma)^{-p}(\Gamma-1)^{-p}}{(\Gamma-1-\tau)\tau} \left(\frac{C_7}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0-1})^p} \right)^{i_0+2} \frac{1-s}{(\gamma_{i_0}-1)^p} \mathcal{A}_1.$$

By scaling we get (1.11) as desired, with the constant \mathcal{C}_3 given by

$$\mathcal{C}_3 = \left(\frac{C_7}{s} \frac{i_0^{3(N+p)}}{(1-\gamma_{i_0-1})^p} \right)^{i_0+2} \frac{1}{(\gamma_{i_0}-1)^p},$$

and $C_7 > 0$ depending on N and p only. This concludes the proof in the subcase $\gamma_{i_0} > 1$.

Case $\boxed{\gamma_{i_0} = 1}$. This case is subtle, due to the fact that $\mathcal{B}_\infty^{1,p} \not\subset W^{1,p}$. Rather than jumping directly from the ball $B_{r_{i_0-1}}$ to the final one $B_{r_{i_0}}$ as before, we need to slightly “rectify” the scheme.

First of all, we introduce the intermediate new ball

$$B_{r_{i_0}} \Subset \tilde{B} := B_{\frac{r_{i_0-1}+3r_{i_0}}{4}} \Subset B_{\frac{r_{i_0}+r_{i_0-1}}{2}} \Subset B_{r_{i_0-1}}.$$

Then we replace the cut-off function $\eta_{i_0-1} \in C_0^2(B_{r_{i_0}})$ with the new one $\tilde{\eta}$ such that

$$\begin{aligned} 0 \leq \tilde{\eta} \leq 1, \quad \tilde{\eta} \equiv 1 \quad \text{on } \tilde{B}, \quad \tilde{\eta} \equiv 0 \quad \text{on } \mathbb{R}^N \setminus B_{\frac{r_{i_0}+r_{i_0-1}}{2}}, \\ |\nabla \tilde{\eta}| \leq \frac{c'_N}{r_{i_0-1} - r_{i_0}} \quad \text{and} \quad |D^2 \tilde{\eta}| \leq \frac{c'_N}{(r_{i_0-1} - r_{i_0})^2}. \end{aligned}$$

Finally, we set

$$\tilde{\mathcal{M}}_{\gamma_{i_0-1}} := \sup_{0 < |h| < h_0} \left[\frac{\delta_h(u_R \tilde{\eta})}{|h|^{\frac{\gamma_{i_0-1}+t}{p}}} \right]_{W^{s,p}(\tilde{B})}^p.$$

We now proceed by iteration as in the proof (4.16), but in the last step we replace (4.13) with

$$\widetilde{\mathcal{M}}_{\gamma_{i_0-1}} \leq \frac{C_1 i_0^{2p}}{s(1-\gamma_{i_0-1})^p} \mathcal{M}_{\gamma_{i_0-2}} + \frac{C_2 i_0^{3(N+p)}}{(1-\gamma_{i_0-1})^p} \mathcal{A}_1.$$

The latter can be proved as before by combining (3.20) and (4.10). Thus this time we get

$$\widetilde{\mathcal{M}}_{\gamma_{i_0-1}} \leq \left(\frac{C_3}{s}\right)^{i_0} \left(\frac{i_0^{3(N+p)}}{(1-\gamma_{i_0-1})^p}\right)^{i_0} \mathcal{A}_1.$$

An application of estimate (3.21) of Lemma 3.3 gives⁷

$$\sup_{0 < |h| < h_0} \left\| \frac{\delta_h u_R}{|h|^\beta} \right\|_{L^p(\widetilde{B})}^p \leq \frac{C}{(1-\beta)^p} \left[(1-s) \widetilde{\mathcal{M}}_{\gamma_{i_0-1}} + h_0^{-p} \|u_R\|_{L^p(B_1)}^p \right],$$

for an arbitrary $0 < \beta < 1$. We now apply Proposition 3.1 with balls $B_{r_{i_0}} \Subset \widetilde{B}$, this would give as at the beginning of the proof

$$\left[\frac{\delta_h(u_R \eta)}{|h|^{\frac{\beta+t}{p}}} \right]_{W^{s,p}(\widetilde{B})}^p \leq \frac{C}{(1-s)s} i_0^{2p} \left\| \frac{\delta_h u_R}{|h|^{\gamma_i}} \right\|_{L^p(\widetilde{B})}^p + C i_0^{2(N+p)} \mathcal{A}_1, \quad 0 < |h| < h_0,$$

where η is as usual a C^2 cut-off function, such that $\eta \equiv 1$ on $B_{r_{i_0}} = B_{1/2}$. By choosing $\beta < 1$ such that

$$\frac{\beta + t + sp}{p} > 1,$$

we are then reduced to the previous subcase $\gamma_{i_0} > 1$. The proof can then be concluded accordingly. We leave the technical details to the reader.

Remark 4.1 (The number of iterations i_0). All the estimates above crucially depends on the number of iterations $i_0 \in \mathbb{N} \setminus \{0\}$. In particular, all the constants blow-up as i_0 goes to ∞ . It is thus useful to recall that if we set

$$\kappa = \kappa(t, s, p) := \frac{t + sp}{p - 1},$$

the sequence $\{\gamma_i\}_{i \in \mathbb{N}}$ has the following explicit expression

$$\gamma_i = \frac{1}{p^i} s + \frac{t + sp}{p} \sum_{j=0}^{i-1} \left(\frac{1}{p^j}\right) = \frac{1}{p^i} s + \kappa \left(1 - \frac{1}{p^i}\right), \quad i \in \mathbb{N}.$$

Then in the case $t + sp \leq (p - 1)$, the exponent i_0 is given by (recall that $s \leq \tau < \kappa$)

$$i_0 = \min \left\{ i \in \mathbb{N} : i > \frac{\ln(\kappa - s) - \ln(\kappa - \tau)}{\ln p} \right\},$$

while in the case $t + sp > (p - 1)$, this is given by

$$i_0 = \min \left\{ i \in \mathbb{N} : i \geq \frac{\ln(\kappa - s) - \ln(\kappa - 1)}{\ln p} \right\}.$$

and we have

$$\gamma_{i_0} = 1 \quad \iff \quad \frac{\ln(\kappa - s) - \ln(\kappa - 1)}{\ln p} \in \mathbb{N} \setminus \{0\}.$$

⁷Observe that by construction the difference of the radii of the two balls \widetilde{B} and $B_{r_{i_0-1}}$ is such that

$$r_{i_0-1} - \frac{r_{i_0-1} + 3r_{i_0}}{4} = \frac{3}{4} (r_{i_0-1} - r_{i_0}) = \frac{3}{16i_0} > 4h_0.$$

4.4. Robust estimate for $s \nearrow 1$. We now reprove (1.10) and (1.11), this time for s sufficiently close to 1 and with an exact control on the constants. In other words, we want to prove the estimates (1.12) and (1.13) claimed in Remark 1.5. We still denote by u_R the scaled solution. Let us thus fix $\ell_0 > p$ and consider $0 \leq t \leq s < 1$ such that

$$t + s(p + 1) \geq \ell_0.$$

Observe that sequence $\{\gamma_i\}_{i \in \mathbb{N}}$ defined in (4.1) is such that

$$(4.19) \quad \gamma_1 = \frac{t + s(p + 1)}{p} \geq \frac{\ell_0}{p} > 1,$$

thus $i_0 = 1$ and we can conclude in one step, i.e. there is no need to iterate the estimate, exactly like in the case of the local p -Laplacian. By using estimate (3.22) of Lemma 3.3 and (4.19), we immediately get

$$\|\nabla u_R\|_{L^p(B_{r_1})} \leq \|\nabla(u_R \eta_0)\|_{L^p(\mathbb{R}^N)} \leq \frac{C}{(\ell_0 - p)^p} \left[(1 - s) \mathcal{M}_{\gamma_0} + h_0^{-p \gamma_1} \|u_R\|_{L^p(B_1)}^p \right].$$

This leads directly to (recall (4.12) for \mathcal{M}_0)

$$\|\nabla u_R\|_{L^p(B_{r_1})} \leq \frac{C}{(\ell_0 - p)^p} (1 - s) \mathcal{A}_1.$$

By scaling back we get (1.12). As for the fractional differentiability of ∇u , we can reproduce the final step of the case $t + sp > (p - 1)$ above. That is, we use (4.18), i.e.

$$\sup_{0 < |h| < h_0} \left\| \frac{\delta_h u_R}{|h|} \right\|_{L^p(B_{r_{i_0-1}})}^p \leq C \|\nabla(u_R \eta_{i_0-1})\|_{L^p(\mathbb{R}^N)}^p + C h_0^{-p} \|u_R\|_{L^p(B_1)}^p,$$

then Proposition 3.1 in the limit case $\gamma = 1$ and once more estimate (3.23) of Lemma 3.3. We omit the details.

4.5. A note on more general lower order terms. We spend some words on the case of the more general equation

$$(4.20) \quad (-\Delta_{p,K})^s u = f + \Phi(u),$$

where $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ is a locally Lipschitz function. This in particular embraces the case of eigenfunctions of $(-\Delta_p)^s$, corresponding to $f = 0$, $K(z) = |z|^{N+sp}$ and $\Phi(t) = \lambda |t|^{p-2} t$ for some $\lambda > 0$. This nonlinear and nonlocal eigenvalue problem has been first introduced in [21]. For more general nonlinearities Φ , we address the reader to [12] for the existence theory.

It is not difficult to see that Theorem 1.4 still holds for local weak solutions $u \in W_{loc}^{s,p}(\Omega) \cap \mathcal{Y}_s^{t,p}(\Omega)$ (4.20) such that

$$u \in L_{loc}^\infty(\Omega).$$

Indeed, the only difference with the proof of Theorem 1.4 is the presence of the additional term in the right-hand side of (3.3)

$$\int |\Phi(u_h) - \Phi(u)| \left| \frac{\delta_h u}{|h|^{\gamma+t}} \right| \eta^2 dx.$$

This is of course a lower-order term, indeed it can be estimated as follows for $0 < |h| < h_0 < 1$

$$(4.21) \quad \int |\Phi(u_h) - \Phi(u)| \left| \frac{\delta_h u}{|h|^{\gamma+t}} \right| \eta^2 dx \leq L \int_{B_{R+\frac{x}{2}}} \left| \frac{\delta_h u}{|h|^{\frac{\gamma+t}{2}}} \right|^2 dx \leq C L R^N + C \int_{B_R} \left| \frac{\delta_h u}{|h|^\gamma} \right|^p dx,$$

where

$$L = \sup_{\xi \in [-M, M]} |\Phi'(\xi)| \quad \text{and} \quad M = \|u\|_{L^\infty(B_R)}.$$

The last term in (4.21) already appeared in the right-hand side of (3.3). Thus, the proof of Theorem 1.4 can be reproduced verbatim. Accordingly, estimates (1.9), (1.10) and (1.11) still hold for bounded local weak solutions of (4.20), with the term $\mathcal{A}_R(u, f)$ defined in (1.8) replaced by

$$\mathcal{A}'_R(u, f) := \mathcal{A}_R(u, f) + L R^N,$$

and L is as above. Remark 1.5 about the quality of the relevant constants still applies.

APPENDIX A. PROOF OF PROPOSITION 2.4

The proof is essentially the same as [27, Chapter 5, Section V, Propositions 8' & 9']. The only difference is the use of the heat kernel, in place of the Poisson's one⁸.

Proof. We introduce the heat kernel

$$\mathcal{K}_t(x) = \frac{1}{(4\pi t)^{\frac{N}{2}}} \exp\left(-\frac{|x|^2}{4t}\right),$$

then we set

$$\psi_t(x) = \mathcal{K}_t * \psi(x) = \frac{1}{(4\pi t)^{\frac{N}{2}}} \int_{\mathbb{R}^N} \exp\left(-\frac{|x-y|^2}{4t}\right) \psi(y) dy.$$

Observe that by the semigroup property of the heat kernel we have

$$\mathcal{K}_{t+s} = \mathcal{K}_t * \mathcal{K}_s,$$

thus we get

$$(A.1) \quad \frac{\partial}{\partial t} \nabla \psi_t = (\nabla \mathcal{K}_{t/2}) * \left(\frac{\partial}{\partial t} \psi_{t/2} \right),$$

where ∇ denotes the gradient with respect to the x variable. In order to estimate the right-hand side of (A.1) for $t > 0$, we observe that⁹

$$\begin{aligned} \nabla \mathcal{K}_{t/2}(x) &= -\frac{x}{t} \mathcal{K}_{t/2}(x), & \|\nabla \mathcal{K}_t\|_{L^1(\mathbb{R}^N)} &\leq \frac{C}{\sqrt{t}}, & \int_{\mathbb{R}^N} \frac{\partial}{\partial t} \mathcal{K}_t(y) dy &= 0 \\ \frac{\partial}{\partial t} \mathcal{K}_{t/2}(x) &= \frac{\partial}{\partial t} \mathcal{K}_{t/2}(-x), & \left| \frac{\partial}{\partial t} \mathcal{K}_{t/2}(x) \right| &\leq \frac{1}{2t} \mathcal{K}_{t/2}(x) \left| \frac{|x|^2}{t} - N \right|. \end{aligned}$$

Thus we get

$$\frac{\partial}{\partial t} \psi_{t/2}(x) = \frac{1}{2} \int_{\mathbb{R}^N} \frac{\partial}{\partial t} \mathcal{K}_{t/2}(y) \left[\psi(x+y) + \psi(x-y) - 2\psi(x) \right] dy.$$

From this, by Minkowski inequality we obtain

$$\begin{aligned} \left\| \frac{\partial}{\partial t} \psi_{t/2} \right\|_{L^p(\mathbb{R}^N)} &\leq \frac{1}{2} \int_{\mathbb{R}^N} \left| \frac{\partial}{\partial t} \mathcal{K}_{t/2}(y) \right| \left\| \psi_{-y} + \psi_y - 2\psi \right\|_{L^p(\mathbb{R}^N)} dy \\ &\leq \frac{1}{2} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} \int_{\mathbb{R}^N} \left| \frac{\partial}{\partial t} \mathcal{K}_{t/2}(y) \right| |y|^\alpha dy \\ &\leq \frac{1}{4t} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} \int_{\mathbb{R}^N} \mathcal{K}_{t/2}(y) \left| \frac{|y|^2}{t} - N \right| |y|^\alpha dy. \end{aligned}$$

⁸In [27] the space $\mathcal{B}_{\infty}^{\alpha,p}$ is noted by $\Lambda_{\alpha}^{p,\infty}$.

⁹We have

$$\frac{\partial}{\partial t} \mathcal{K}_t(x) = \mathcal{K}_t(x) \left[\frac{|x|^2}{4t^2} - \frac{N}{2t} \right].$$

With a simple change of variables, this gives

$$\left\| \frac{\partial}{\partial t} \psi_{t/2} \right\|_{L^p(\mathbb{R}^N)} \leq C [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} \left(\int_{\mathbb{R}^N} \mathcal{K}_1(z) \left| |z|^2 - N \right| |z|^\alpha dz \right) t^{\frac{\alpha}{2}-1}.$$

Observe that for $1 < \alpha < 2$

$$\int_{\mathbb{R}^N} \mathcal{K}_1(z) \left| |z|^2 - N \right| |z|^\alpha dz \leq \int_{\{|z| \leq 1\}} \mathcal{K}_1(z) \left| |z|^2 - N \right| |z| dz + \int_{\{|z| > 1\}} \mathcal{K}_1(z) \left| |z|^2 - N \right| |z|^2 dz,$$

and the last two terms depend only on N , so that in conclusion

$$(A.2) \quad \left\| \frac{\partial}{\partial t} \psi_{t/2} \right\|_{L^p(\mathbb{R}^N)} \leq C [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} t^{\frac{\alpha}{2}-1},$$

for some $C = C(N) > 0$. Then from (A.1) and (A.2) we obtain for every $t > 0$

$$(A.3) \quad \left\| \frac{\partial}{\partial t} \nabla \psi_t \right\|_{L^p(\mathbb{R}^N)} \leq \|\nabla \mathcal{K}_{t/2}\|_{L^1(\mathbb{R}^N)} \left\| \frac{\partial}{\partial t} \psi_{t/2} \right\|_{L^p(\mathbb{R}^N)} \leq C [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} t^{\frac{\alpha-3}{2}}.$$

We now integrate the previous inequality on the interval (s, τ) , by Minkowski inequality again we get

$$(A.4) \quad \begin{aligned} \|\nabla \psi_\tau - \nabla \psi_s\|_{L^p(\mathbb{R}^N)} &= \left\| \int_s^\tau \frac{\partial}{\partial t} \nabla \psi_t dt \right\|_{L^p(\mathbb{R}^N)} \leq \int_s^\tau \left\| \frac{\partial}{\partial t} \nabla \psi_t \right\|_{L^p(\mathbb{R}^N)} dt \\ &\leq \frac{2C}{\alpha-1} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} \left(\tau^{\frac{\alpha-1}{2}} - s^{\frac{\alpha-1}{2}} \right). \end{aligned}$$

Since $\alpha > 1$ by assumption, this shows that $\{\nabla \psi_t\}_{0 < t < 1}$ is a Cauchy net in the complete space $L^p(\mathbb{R}^N)$. Thus there exists a sequence $\{t_k\}_{k \in \mathbb{N}} \subset (0, 1)$ converging to 0 as k goes to ∞ , such that $\{\nabla \psi_{t_k}\}_{k \in \mathbb{N}}$ converges strongly in L^p . The limit function is the distributional gradient of ψ . Finally, this shows that $\nabla \psi \in L^p(\mathbb{R}^N)$. Moreover, by taking the limit in (A.4), we get the estimate

$$\|\nabla \psi\|_{L^p(\mathbb{R}^N)} \leq \|\nabla \psi_1\|_{L^p(\mathbb{R}^N)} + \frac{2C}{\alpha-1} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} \leq C \|\psi\|_{L^p(\mathbb{R}^N)} + \frac{2C}{\alpha-1} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)},$$

which is (2.7).

Once the existence of $\nabla \psi$ in L^p is established, we can now prove (2.8). We first need a decay estimate on the hessian $D^2 \psi_t$. For this, we observe that

$$|D^2 \mathcal{K}_t(x)| \leq \frac{\mathcal{K}_t(x)}{4t} \left[\frac{x \otimes x}{4t} - \text{Id}_N \right] \quad \text{and} \quad \|D^2 \mathcal{K}_t\|_{L^1(\mathbb{R}^N)} \leq \frac{C}{t}.$$

Then of course we have

$$(A.5) \quad \|D^2 \psi_t\|_{L^p(\mathbb{R}^N)} \leq \frac{C}{t} \|\psi\|_{L^p(\mathbb{R}^N)}.$$

Similarly as before, we can write

$$\frac{\partial}{\partial t} D^2 \psi_t = (D^2 \mathcal{K}_{t/2}) * \left(\frac{\partial}{\partial t} \psi_{t/2} \right),$$

then for every $t > 0$ we get

$$\left\| \frac{\partial}{\partial t} D^2 \psi_t \right\|_{L^p(\mathbb{R}^N)} \leq \|D^2 \mathcal{K}_{t/2}\|_{L^1(\mathbb{R}^N)} \left\| \frac{\partial}{\partial t} \psi_{t/2} \right\|_{L^p(\mathbb{R}^N)} \leq C [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} t^{\frac{\alpha-4}{2}}.$$

By integrating this estimate between s and $T \gg s$, as above we get

$$\|D^2 \psi_s\|_{L^p(\mathbb{R}^N)} \leq \|D^2 \psi_T\|_{L^p(\mathbb{R}^N)} + \frac{C}{2-\alpha} [\psi]_{\mathcal{B}_{\infty}^{\alpha,p}(\mathbb{R}^N)} \left(s^{\frac{\alpha-2}{2}} - T^{\frac{\alpha-2}{2}} \right).$$

By recalling that $\alpha < 2$, using (A.5) and taking the limit as T goes to ∞ , we get the desired decay estimate

$$(A.6) \quad \|D^2\psi_s\|_{L^p(\mathbb{R}^N)} \leq \frac{C}{2-\alpha} [\psi]_{\mathcal{B}_{\infty^p}^{\alpha,p}(\mathbb{R}^N)} s^{\frac{\alpha-2}{2}}.$$

Let $h \in \mathbb{R}^N \setminus \{0\}$, by using that ψ_t converges to ψ as t goes to 0, we have

$$(A.7) \quad \delta_h \nabla \psi = \delta_h \nabla \psi_t - \int_0^t \frac{\partial}{\partial s} (\delta_h \nabla \psi_s) ds.$$

By using the smoothness of ψ_t , we can write

$$\delta_h \nabla \psi_t = \int_0^{|h|} \frac{d}{d\tau} \nabla \psi_t \left(x + \frac{h}{|h|} \tau \right) d\tau = \int_0^{|h|} D^2 \psi_t \left(x + \frac{h}{|h|} \tau \right) \frac{h}{|h|} d\tau,$$

which implies

$$\|\delta_h \nabla \psi_t\|_{L^p(\mathbb{R}^N)} \leq \int_0^{|h|} \|D^2 \psi_t\|_{L^p(\mathbb{R}^N)} d\tau \leq \frac{C}{2-\alpha} [\psi]_{\mathcal{B}_{\infty^p}^{\alpha,p}(\mathbb{R}^N)} |h| t^{\frac{\alpha-2}{2}},$$

thanks to (A.6). On the other hand, by triangle inequality and invariance of the L^p norm by translations, we have

$$\left\| \frac{\partial}{\partial s} (\delta_h \nabla \psi_s) \right\|_{L^p(\mathbb{R}^N)} \leq 2 \left\| \frac{\partial}{\partial s} \nabla \psi_s \right\|_{L^p(\mathbb{R}^N)} \leq C [\psi]_{\mathcal{B}_{\infty^p}^{\alpha,p}(\mathbb{R}^N)} s^{\frac{\alpha-3}{2}},$$

where we also used (A.3). We can now use the two previous estimates in conjunction with (A.7), so to get

$$\begin{aligned} \|\delta_h \nabla \psi\|_{L^p(\mathbb{R}^N)} &\leq \|\delta_h \nabla \psi_t\|_{L^p(\mathbb{R}^N)} + \left\| \int_0^t \frac{\partial}{\partial s} (\delta_h \nabla \psi_s) ds \right\|_{L^p(\mathbb{R}^N)} \\ &\leq C [\psi]_{\mathcal{B}_{\infty^p}^{\alpha,p}(\mathbb{R}^N)} \left[\frac{|h|}{2-\alpha} t^{\frac{\alpha-2}{2}} + \int_0^t s^{\frac{\alpha-3}{2}} ds \right] = C [\psi]_{\mathcal{B}_{\infty^p}^{\alpha,p}(\mathbb{R}^N)} \left[\frac{|h|}{2-\alpha} t^{\frac{\alpha-2}{2}} + \frac{2}{\alpha-1} t^{\frac{\alpha-1}{2}} \right]. \end{aligned}$$

for some $C = C(N) > 0$. The previous estimate holds for every $t > 0$ and the right-hand side is minimal for $t = |h|^2/4$. With such a choice we thus get

$$\left\| \frac{\delta_h \nabla \psi}{|h|^{\alpha-1}} \right\|_{L^p(\mathbb{R}^N)} \leq \frac{C}{(2-\alpha)(\alpha-1)} [\psi]_{\mathcal{B}_{\infty^p}^{\alpha,p}(\mathbb{R}^N)},$$

as desired. \square

APPENDIX B. POINTWISE INEQUALITIES

For $p \geq 2$ we define the functions $J_p : \mathbb{R} \rightarrow \mathbb{R}$ and $V_p : \mathbb{R} \rightarrow \mathbb{R}$ by

$$J_p(t) = |t|^{p-2} t, \quad \text{and} \quad V_p(t) = |t|^{\frac{p-2}{2}} t.$$

Lemma B.1. *Let $p \geq 2$, for every $a, b \in \mathbb{R}$ we have*

$$(B.1) \quad \left(J_p(a) - J_p(b) \right) (a - b) \geq (p-1) \left(\frac{2}{p} \right)^2 |V_p(a) - V_p(b)|^2.$$

Proof. Since $J_p(a) - J_p(b)$ and $a - b$ share the same sign, we can assume without loss of generality that $a \geq b$. If $a = b$ there is nothing to prove. Let us assume that $a > b$, then we have

$$\begin{aligned} \left(J_p(a) - J_p(b) \right) (a - b) &= (p-1) \left(\int_b^a |t|^{p-2} dt \right) (a - b)^{p-1} \\ &\geq (p-1) \left(\int_a^b |t|^{\frac{p-2}{2}} dt \right)^2 = (p-1) \left(\frac{2}{p} \right)^2 |V_p(a) - V_p(b)|^2, \end{aligned}$$

which concludes the proof. \square

Lemma B.2. *Let $p \geq 2$, for every $a, b \in \mathbb{R}$ we have*

$$(B.2) \quad \left| J_p(a) - J_p(b) \right| \leq 2 \frac{p-1}{p} \left(|a|^{\frac{p-2}{2}} + |b|^{\frac{p-2}{2}} \right) |V_p(a) - V_p(b)|.$$

Proof. For $a = b$ there is nothing to prove. Let us consider the case $a \neq b$, without loss of generality, we can suppose that $a > b$. We set

$$G(t) = |t|^{\frac{p-2}{p}} t, \quad t \in \mathbb{R},$$

by basic calculus we have

$$G\left(|a|^{\frac{p-2}{2}} a\right) - G\left(|b|^{\frac{p-2}{2}} b\right) \leq \max\left\{G'\left(|a|^{\frac{p-2}{2}} a\right), G'\left(|b|^{\frac{p-2}{2}} b\right)\right\} \left(V_p(a) - V_p(b)\right).$$

By observing that

$$G\left(|t|^{\frac{p-2}{2}} t\right) = |t|^{p-2} t,$$

we get the conclusion. □

Lemma B.3. *Let $p \geq 2$, for every $a, b \in \mathbb{R}$ we have*

$$(B.3) \quad |V_p(a) - V_p(b)|^2 \geq |a - b|^p.$$

In particular, we also get

$$(B.4) \quad \left(J_p(a) - J_p(b)\right) (a - b) \geq (p-1) \left(\frac{2}{p}\right)^2 |a - b|^p.$$

Proof. Observe that if $a = 0$ or $b = 0$, the result trivially holds. Thus let us suppose that $a, b \neq 0$ and observe that the function $F : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$H(t) = |t|^{\frac{2-p}{p}} t,$$

is $2/p$ -Hölder continuous. More precisely, we have

$$|H(t) - H(s)| \leq |t - s|^{\frac{2}{p}}, \quad t, s \in \mathbb{R}.$$

By applying the previous with

$$t = V_p(a) \quad \text{and} \quad s = V_p(b),$$

we get (B.3). The last inequality (B.4) follows by combining (B.1) and (B.3). □

REFERENCES

- [1] R. A. Adams, Sobolev spaces. Pure and Applied Mathematics, **65**. Academic Press, New York-London, 1975. [5](#)
- [2] F. Andreu, J. M. Mazón, J. D. Rossi, J. Toledo, A nonlocal p -Laplacian evolution equation with nonhomogeneous Dirichlet boundary conditions, *SIAM J. Math. Anal.*, **40** (2009), 1815–1851. [2](#)
- [3] C. Bjorland, L. Caffarelli, A. Figalli, Non-Local Gradient Dependent Operators, *Adv. Math.*, **230** (2012), 1859–1894. [2](#)
- [4] J. Bourgain, H. Brezis, P. Mironescu, Limiting embedding theorems for $W^{s,p}$ when $s \rightarrow 1$ and applications, *J. Anal. Math.*, **87** (2002), 77–101. [7](#)
- [5] J. Bourgain, H. Brezis and P. Mironescu, Another look at Sobolev spaces, in *Optimal Control and Partial Differential Equations. A Volume in Honor of Professor Alain Bensoussan's 60th Birthday* (eds. J. L. Menaldi, E. Rofman and A. Sulem), IOS Press, Amsterdam, 2001, 439–455. [7](#)
- [6] L. Brasco, E. Lindgren, E. Parini, The fractional Cheeger problem, *Interfaces Free Bound.*, **16** (2014), 419–458. [5](#), [10](#)
- [7] L. Brasco, E. Parini, M. Squassina, Stability of variational eigenvalues for the fractional p -Laplacian, to appear on *Discrete Contin. Dyn. Syst.* (2015), available at <http://cvgmt.sns.it/paper/2649/> [7](#)
- [8] E. Chasseigne, E. Jakobsen, On nonlocal quasilinear equations and their local limits, preprint (2015), available at <http://arxiv.org/abs/1503.06939> [2](#)
- [9] M. Cozzi, Interior regularity of solutions of non-local equations in Sobolev and Nikol'skii spaces, preprint (2015), available at http://www.ma.utexas.edu/mp_arc/c/15/15-50.pdf [3](#), [6](#), [7](#)
- [10] A. Di Castro, T. Kuusi, G. Palatucci, Local behavior of fractional p -minimizers, to appear on *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2015), doi:10.1016/j.anihpc.2015.04.003 [2](#), [3](#)
- [11] A. Di Castro, T. Kuusi, G. Palatucci, Nonlocal Harnack inequalities, *J. Funct. Anal.*, **267** (2014), 1807–1836. [2](#)

- [12] A. Iannizzotto, S. Liu, K. Perera, M. Squassina, Existence results for fractional p -Laplacian problems via Morse theory, to appear on Adv. Calc. Var. (2015), available at <http://arxiv.org/abs/1403.5388> [30](#)
- [13] A. Iannizzotto, S. Mosconi, M. Squassina, A note on global regularity for the weak solutions of fractional p -Laplacian equations, preprint (2015), available at <http://arxiv.org/abs/1504.01006> [2](#)
- [14] A. Iannizzotto, S. Mosconi, M. Squassina, Global Hölder regularity for the fractional p -Laplacian, preprint (2014), available at <http://arxiv.org/abs/1411.2956> [2, 3](#)
- [15] H. Ishii, G. Nakamura, A class of integral equations and approximation of p -Laplace equations, Calc. Var. Partial Differential Equations, **37** (2010), 485–522. [2](#)
- [16] M. Kassmann, A priori estimates for integro-differential operators with measurable kernels, Calc. Var. Partial Differential Equations, **34** (2009), 1–21. [2](#)
- [17] T. Kuusi, G. Mingione, Y. Sire, Nonlocal self-improving properties, to appear on Anal. PDE, **8** (2015), 57–114. [3, 6](#)
- [18] T. Kuusi, G. Mingione, Y. Sire, Nonlocal equations with measure data, Comm. Math. Phys., **337** (2015), 1317–1368. [2](#)
- [19] T. Kuusi, G. Mingione, Y. Sire, A fractional Gehring lemma, with applications to nonlocal equations, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl., **24** (2014), 345–358. [3, 6](#)
- [20] E. Lindgren, Hölder estimates for viscosity solutions of equations of fractional p -Laplace type, preprint (2014), available at <http://arxiv.org/abs/1405.6612> [5](#)
- [21] E. Lindgren, P. Lindqvist, Fractional eigenvalues, Calc. Var. Partial Differential Equations, **49** (2014), 795–826. [30](#)
- [22] G. Mingione, The Calderón-Zygmund theory for elliptic problems with measure data, Ann. Sc. Norm. Super. Pisa Cl. Sci., **6** (2007), 195–261. [6](#)
- [23] G. Mingione, The singular set of solutions of non-differentiable elliptic systems, Arch. Rational. Mech. Anal., **166** (2003), 287–301. [2](#)
- [24] A. Ponce, A new approach to Sobolev spaces and connections to Γ -convergence, Calc. Var. Partial Differential Equations, **19** (2004), 229–255. [7](#)
- [25] A. Schikorra, Nonlinear commutators for the fractional p -Laplacian and applications, preprint (2015), available at <http://arxiv.org/abs/1506.02380> [3, 6, 7](#)
- [26] Y. Sire, E. Valdinoci, Rigidity results for some boundary quasilinear phase transitions, Commun. Partial Differ. Equations, **34** (2009), 765–784. [2](#)
- [27] E. Stein, Singular integrals and differentiability properties of functions. Princeton Mathematical Series, **30**. Princeton University Press, Princeton, N.J. 1970. [9, 31](#)
- [28] H. Triebel, Theory of function spaces. II. Monographs in Mathematics, **84**. Birkhäuser Verlag, Basel, 1992. [9](#)
- [29] K. Uhlenbeck, Regularity for a class of non-linear elliptic systems, Acta Math. **138** (1977), 219–240. [2](#)

(L. Brasco) AIX-MARSEILLE UNIVERSITÉ, CNRS
 CENTRALE MARSEILLE, I2M, UMR 7373, 39 RUE FRÉDÉRIC JOLIOT CURIE
 13453 MARSEILLE, FRANCE
E-mail address: lorenzo.brasco@univ-amu.fr

(E. Lindgren) DEPARTMENT OF MATHEMATICS, ROYAL INSTITUTE OF TECHNOLOGY
 10044 STOCKHOLM, SWEDEN
E-mail address: eriklin@kth.se