



**HAL**  
open science

# Profile decomposition and phase control for circle-valued maps in one dimension

Petru Mironescu

► **To cite this version:**

Petru Mironescu. Profile decomposition and phase control for circle-valued maps in one dimension. 2015. hal-01202156v1

**HAL Id: hal-01202156**

**<https://hal.science/hal-01202156v1>**

Preprint submitted on 22 Sep 2015 (v1), last revised 3 Jan 2016 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Profile decomposition and phase control for circle-valued maps in one dimension

Petru Mironescu\*

September 18, 2015

**Abstract.** When  $1 < p < \infty$ , maps  $f$  in  $W^{1/p,p}((0,1); \mathbb{S}^1)$  have  $W^{1/p,p}$  phases  $\varphi$ , but the  $W^{1/p,p}$ -seminorm of  $\varphi$  is not controlled by the one of  $f$ . Lack of control is illustrated by “the kink”:  $f = e^{i\varphi}$ , where the phase  $\varphi$  moves quickly from 0 to  $2\pi$ . A similar situation occurs for maps  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ , with Moebius maps playing the role of kinks. We prove that this is the only loss of control mechanism: each map  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  satisfying  $|f|_{W^{1/p,p}}^p \leq M$  can be written as

$f = e^{i\psi} \prod_{j=1}^K (M_{a_j})^{\pm 1}$ , where  $M_{a_j}$  is a Moebius map vanishing at  $a_j \in \mathbb{D}$ , while the integer  $K = K(f)$

and the phase  $\psi$  are controlled by  $M$ . In particular, we have  $K \leq c_p M$  for some  $c_p$ . When  $p = 2$ , we obtain the sharp value of  $c_2$ , which is  $c_2 = 1/(4\pi^2)$ . As an application, we obtain the existence of minimal maps of degree one in  $W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  with  $p \in (2 - \varepsilon, 2)$ .

**Résumé. Décomposition en profils et contrôle des phases des applications unimodulaires en dimension un.** Si  $1 < p < \infty$ , les applications  $f$  appartenant à  $W^{1/p,p}((0,1); \mathbb{S}^1)$  ont des phases  $\varphi$  dans  $W^{1/p,p}$ , mais la seminorme  $W^{1/p,p}$  de  $\varphi$  n’est pas contrôlée par celle de  $f$ . L’absence de contrôle est illustrée par “le pli”:  $f = e^{i\varphi}$ , où la phase  $\varphi$  augmente rapidement de 0 à  $2\pi$ . Pour des applications  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ , le même phénomène apparaît, avec les transformations de Moebius jouant le rôle des plis. Nous prouvons que cet exemple est essentiellement le seul : toute application  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  telle que  $|f|_{W^{1/p,p}}^p \leq M$  s’écrit  $f = e^{i\psi} \prod_{j=1}^K (M_{a_j})^{\pm 1}$ , où  $M_{a_j}$  est une transformation de Moebius s’annulant en  $a_j \in \mathbb{D}$ , tandis que l’entier  $K = K(f)$  et la phase  $\psi$  sont contrôlés par  $M$ . En particulier, nous avons  $K \leq c_p M$  pour une constante  $c_p$ . Pour  $p = 2$ , nous obtenons la valeur optimale de  $c_2$ , qui est  $c_2 = 1/(4\pi^2)$ . Comme application, nous obtenons l’existence d’une application minimale de degré un dans  $W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  avec  $p \in ]2 - \varepsilon, 2[$ .

## 1 Introduction

Let  $0 < s < 1$ ,  $1 \leq p < \infty$  and let  $f : (0,1) \rightarrow \mathbb{S}^1$  belong to the space  $W^{s,p}$ . Then  $f$  can be written as  $f = e^{i\varphi}$ , where  $\varphi \in W^{s,p}$  [4]. Once the existence of  $\varphi$  is known, a natural question is whether we can control  $|\varphi|_{W^{s,p}}$  in terms of  $|f|_{W^{s,p}}$ . For most of  $s, p$  the answer is positive. The exceptional cases are provided precisely by the spaces  $W^{1/p,p}((0,1); \mathbb{S}^1)$ , with  $1 < p < \infty$  [4]. In these spaces, lack of control is established via the following explicit example. For  $n \geq 1$ , we define  $\varphi_n$  as

---

\*Université de Lyon, Université Lyon 1, CNRS UMR 5208, Institut Camille Jordan, 43 blvd. du 11 novembre 1918, F-69622 Villeurbanne cedex, France. Email address: mironescu@math.univ-lyon1.fr

Keywords. Profile decomposition, circle-valued map, phase, Sobolev spaces

2002 AMS classification subject. 46E35

follows:

$$\varphi_n(x) := \begin{cases} 0, & \text{for } 0 < x < 1/2 \\ 2\pi n(x - 1/2), & \text{for } 1/2 < x < 1/2 + 1/n. \\ 2\pi, & \text{for } 1/2 + 1/n < x < 1 \end{cases}$$

Then  $|\varphi_n|_{W^{1/p,p}} \rightarrow \infty$  (since  $\varphi_n \rightarrow \varphi = 2\pi \chi_{(1/2,1)}$  a.e., and  $\varphi$  does not belong to  $W^{1/p,p}$ ). On the other hand, if we extend  $u_n := e^{i\varphi_n}$  with the value 1 outside  $(0,1)$  and still denote the extension  $u_n$  then, by scaling,

$$|u_n|_{W^{1/p,p}((0,1))} \leq |u_n|_{W^{1/p,p}(\mathbb{R})} = |u_1|_{W^{1/p,p}(\mathbb{R})} < \infty.$$

Thus  $|u_n|_{W^{1/p,p}((0,1))} \lesssim 1$  and  $|\varphi_n|_{W^{1/p,p}((0,1))} \rightarrow \infty$ . Finally, we invoke the fact that  $W^{1/p,p}$  phases are unique mod  $2\pi$  [4].

If one considers instead maps  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ , always in the critical case  $f \in W^{1/p,p}$ ,  $1 < p < \infty$ , then a new phenomenon occurs:  $f$  has a degree  $\deg f$ , and does not have a  $W^{1/p,p}$  phase at all when  $\deg f \neq 0$  [11, Remark 10]. However, even if  $\deg f = 0$  (and thus  $f$  has a  $W^{1/p,p}$  phase  $\varphi$ ), we have a loss of control phenomenon similar to the one on  $(0,1)$ . Indeed, let  $M_a(z) := \frac{a-z}{1-\bar{a}z}$ ,  $a \in \mathbb{D}$ ,  $z \in \bar{\mathbb{D}}$ , be a Moebius transform (that we identify with its restriction to  $\mathbb{S}^1$ ,  $M_a : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ ). Let  $f_a(z) := \bar{z}M_a(z)$ , so that  $f_a$  is smooth and  $\deg f_a = 0$ . One may prove (see below) that  $|M_a|_{W^{1/p,p}} = |\text{Id}|_{W^{1/p,p}}$ , and thus  $f_a$  is bounded in  $W^{1/p,p}$ . However, if  $a \rightarrow \alpha = e^{i\xi} \in \mathbb{S}^1$ , then the smooth phase  $\varphi_a$  of  $f_a$  converges a.e. to  $\varphi(e^{i\theta}) := \begin{cases} \xi - \theta, & \text{if } \xi - \pi < \theta < \xi \\ 2\pi + \xi - \theta, & \text{if } \xi < \theta < \xi + \pi \end{cases}$ , which does not

belong to  $W^{1/p,p}$ . [Here, uniqueness of the phases and convergence hold mod  $2\pi$ .] Thus  $\varphi_a$  is not bounded as  $a \rightarrow \alpha \in \mathbb{S}^1$ . On the other hand, the plot of  $\varphi_a$  shows that  $\varphi_a$  has a ‘‘kink shape’’, and thus we have here the analog of the example on  $(0,1)$ .

There are evidences that this loss of control mechanism is the only possible one. For example, the phase of the kink is not bounded in  $W^{1/p,p}$ , but clearly is in  $W^{1,1}$  (same for  $f_a$ ). Bourgain and Brezis [3] proved that for every  $f \in W^{1/2,2}((0,1); \mathbb{S}^1)$ , we may split  $f = e^{i\psi} v$ , with  $\psi$  and  $v = e^{i\eta}$  satisfying

$$|\psi|_{W^{1/2,2}} \lesssim |f|_{W^{1/2,2}} \text{ and } |\eta|_{W^{1,1}} = |v|_{W^{1,1}} \lesssim |f|_{W^{1/2,2}}^2. \quad (1)$$

Intuitively, one should think at  $v$  as at ‘‘the kink part of  $f$ ’’. The above result was extended by Nguyen [18] to  $1 < p < \infty$ : for every  $1 < p < \infty$  and every  $f \in W^{1/p,p}((0,1); \mathbb{S}^1)$ , we may split  $f = e^{i\psi} v$ , with  $\psi$  and  $v = e^{i\eta}$  satisfying

$$|\psi|_{W^{1/p,p}} \leq C_p |f|_{W^{1/p,p}} \text{ and } |\eta|_{W^{1,1}} = |v|_{W^{1,1}} \leq C_p |f|_{W^{1/p,p}}^p. \quad (2)$$

Here we present another result in this direction, written for simplicity on the unit circle.

**Theorem 1.** *Let  $1 < p < \infty$  and  $M > 0$ . Then there exist constants  $c_p$  and  $F(M)$  such that: every map  $f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  satisfying  $|f|_{W^{1/p,p}}^p \leq M$  can be written as  $f = e^{i\psi} \prod_{j=1}^K (M_{a_j})^{\varepsilon_j}$ , with  $\varepsilon_j \in \{-1, 1\}$ ,*

$$K \leq c_p M, \quad (3)$$

and

$$|\psi|_{W^{1/p,p}}^p \leq F(M). \quad (4)$$

When  $p = 2$ , we may take  $c_2 = 1/(4\pi^2)$ , and this constant is optimal.

**Corollary 1.** *Let  $1 < p < \infty$  and let  $f_n, f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  be such that  $f_n \rightharpoonup f$  in  $W^{1/p,p}$ . Then, up to a subsequence, there exist  $K \in \mathbb{N}$ ,  $\varepsilon_j \in \{-1, 1\}$ ,  $a_{j_n} \in \mathbb{D}$ ,  $\alpha_j \in \mathbb{S}^1$ ,  $j = 1, \dots, K$ ,  $\psi_n \in W^{1/p,p}(\mathbb{S}^1; \mathbb{R})$ , and a constant  $C$ , such that:*

- i)  $f_n = e^{i\psi_n} \prod_{j=1}^K (M_{a_{j_n}})^{\varepsilon_j} f$ ;
- ii)  $a_{j_n} \rightarrow \alpha_j$  as  $n \rightarrow \infty$ ;
- iii)  $\psi_n \rightharpoonup C$  in  $W^{1/p,p}$  as  $n \rightarrow \infty$ .

The theorem and the corollary are reminiscent of profile decompositions obtained in different, often geometrical, contexts. We mention e.g. the work of Sacks and Uhlenbeck [19] on minimal 2-spheres, the analysis of Brezis and Coron [6, 7, 8] of constant mean curvature surfaces, or the one of Struwe [20] of equations involving the critical Sobolev exponent. There are also abstract approaches to bubbling as in the work of Lions [16] about concentration-compactness or the characterization of lack of compactness of critical embeddings in Gérard [12], Jaffard [15] or Bahouri, Cohen and Koch [1].

Let us comment on the connection between (2) and our theorem. First, (2) has the following version for maps on  $\mathbb{S}^1$ : we may split  $f = e^{i\psi} v$ , with  $|\psi|_{W^{1/p,p}} \leq C_p |f|_{W^{1/p,p}}$  and  $|v|_{W^{1,1}} \leq C_p |f|_{W^{1/p,p}}$ . Next, a Moebius maps satisfies  $|M_\alpha|_{W^{1,1}} = 2\pi$ , and thus

$$\left| \prod_{j=1}^K (M_{a_j})^{\varepsilon_j} \right|_{W^{1,1}} \leq 2\pi K \leq 2\pi c_p M. \quad (5)$$

Estimate (5) shows that (3) is a refinement of the second part of (2). On the other hand, (4) is weaker than the first part of (2), since  $F(M)$  need not have a linear growth (and actually we do not have any control on  $F$ ). This suggests the following

**Conjecture.** Let  $1 < p < \infty$ . Then there exist constants  $c_p, d_p$  such that every  $f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  satisfying  $|f|_{W^{1/p,p}}^p \leq M$  can be decomposed as  $f = e^{i\psi} \prod_{j=1}^K (M_{a_j})^{\varepsilon_j}$ , with  $\varepsilon_j \in \{-1, 1\}$ ,

$$K \leq c_p M, \quad (6)$$

and

$$|\psi|_{W^{1/p,p}}^p \leq d_p M. \quad (7)$$

In addition, when  $p = 2$ , we may take  $c_2 = 1/(4\pi^2)$ .

## 2 Proofs

We start by recalling or establishing few auxiliary results. Given  $1 \leq p < \infty$ ,  $f, f_n$  will denote maps in  $W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$ . When  $1 < p < \infty$ , “ $\rightharpoonup$ ” refers to weak convergence in  $W^{1/p,p}$ .

**1.** Recall that, up to a multiplicative factor  $\alpha \in \mathbb{S}^1$ , the Moebius transforms give all the conformal representations  $u : \mathbb{D} \rightarrow \mathbb{D}$ . In particular,  $M_\alpha : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  is a smooth orientation preserving diffeomorphism, and thus  $\deg M_\alpha = 1$ . Consequence: if  $g : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  is continuous, then  $\deg[g \circ M_\alpha] = \deg g$ .

**2.** If  $1 \leq p < \infty$  and  $a \in \mathbb{D}$ , then  $|f \circ M_a|_{W^{1/p,p}} = |f|_{W^{1/p,p}}$ . [Here, we let  $|f|_{W^{1,1}} := \int_{\mathbb{S}^1} |\dot{f}| = \int_0^{2\pi} |d[f(e^{i\theta})]/d\theta| d\theta$  and, for  $1 < p < \infty$ ,  $|f|_{W^{1/p,p}}^p := \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} |f(x) - f(y)|^p / |x - y|^2 dx dy$ .] In order to prove the desired equality when  $p = 1$ , we write  $M_\alpha(e^{i\theta}) = e^{i\varphi(\theta)}$ ,  $0 \leq \theta \leq 2\pi$ , with  $\varphi$  smooth and increasing. Then

$$|f \circ M_\alpha|_{W^{1,1}} = \int_0^{2\pi} \left| \frac{d}{d\theta} [f(e^{i\varphi(\theta)})] \right| d\theta = \int_{\varphi^{-1}(0)}^{\varphi^{-1}(2\pi)} \left| \frac{d}{d\theta} [f(e^{i\theta})] \right| d\theta = \int_0^{2\pi} \left| \frac{d}{d\theta} [f(e^{i\theta})] \right| d\theta = |f|_{W^{1,1}}.$$

When  $1 < p < \infty$ , we rely on the following identity, valid for measurable functions  $F : \mathbb{S}^1 \times \mathbb{S}^1 \rightarrow [0, \infty]$ :

$$\int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{F(M_a(x), M_a(y))}{|x-y|^2} dx dy = \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \frac{F(x, y)}{|x-y|^2} dx dy. \quad (8)$$

Proof of (8): We have  $[M_a]^{-1} = M_a$  and thus, after change of variables, (8) amounts to

$$|x-y|^2 |\dot{M}_a(x)| |\dot{M}_a(y)| = |M_a(x) - M_a(y)|^2, \quad \forall x, y \in \mathbb{S}^1. \quad (9)$$

In turn, (9) follows immediately from the straightforward equality  $|\dot{M}_a(x)| = \frac{1-|a|^2}{|1-\bar{a}x|^2}$ .

**3.** If  $1 \leq p < \infty$  and  $a \in \mathbb{D}$ , then  $\deg[f \circ M_a] = \deg f$ . Indeed, to start with, such  $f$  has a degree, since  $W^{1/p,p} \hookrightarrow \text{VMO}$  and VMO maps gave a degree stable with respect to BMO convergence [11]. By item **1**, the desired equality holds true for smooth  $f$ . The general case follows by density of  $C^\infty(\mathbb{S}^1; \mathbb{S}^1)$  into  $W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$  [11, Lemmas A.11 and A.12] and by stability of the VMO degree.

**4.** If  $1 \leq p < \infty$  and the degree of  $f$  is  $d$ , then we may write  $f(z) = e^{i\psi(z)} z^d$ , with  $\psi \in W^{1/p,p}(\mathbb{S}^1; \mathbb{R})$ . This follows easily from the fact that maps  $f \in W^{1/p,p}((0, 1); \mathbb{S}^1)$  lift within  $W^{1/p,p}$  [4].

**5.** Let  $1 < p < \infty$ . For  $f \in W^{1/p,p}(\mathbb{S}^1; \mathbb{S}^1)$ , let  $u = u(f)$  be its harmonic extension. Set  $c'_p := \inf\{|f|_{W^{1/p,p}}^p; u(0) = 0\}$ . Clearly,  $c'_p$  is achieved, and therefore  $c'_p > 0$ .

**6.** When  $p = 2$ , we have the following straightforward calculations: if  $f = \sum_{n \in \mathbb{Z}} a_n e^{in\theta}$ , then  $|f|_{W^{1/2,2}}^2 = 4\pi^2 \sum_{n \in \mathbb{Z}} |n| |a_n|^2$  [10, Chapter 13], and  $\deg f = \sum_{n \in \mathbb{Z}} n |a_n|^2$  [11, eq (25)]. This leads to  $4\pi^2 |\deg f| \leq |f|_{W^{1/2,2}}^2$ , with equality e.g. when  $f(z) := z^d$ . On the other hand, if  $u(f)(0) = 0$ , then  $a_0 = 0$  and thus

$$|f|_{W^{1/2,2}}^2 = 4\pi^2 \sum_{n \neq 0} |n| |a_n|^2 \geq 4\pi^2 \sum_{n \neq 0} |a_n|^2 = 4\pi^2 \sum_{n \in \mathbb{Z}} |a_n|^2 = 2\pi \|f\|_{L^2}^2 = 4\pi^2.$$

Thus  $c'_2 \geq 4\pi^2$ , and the example  $f(z) := z$  shows that  $c'_2 = 4\pi^2$ .

**7.** For  $1 < p < \infty$ , there exists some constant  $c''_p$  such that  $c''_p |\deg f| \leq |f|_{W^{1/p,p}}^p$ ,  $\forall f \in W^{1/p,p}(\mathbb{S}^1, \mathbb{S}^1)$  [5, Corollary 0.5]. We let  $c''_p$  be the best constant such that this estimate holds, and set  $c_p^* := \min\{c'_p, c''_p\}$ . We also set  $c_p := 1/c_p^*$ . By item **6**, for  $p = 2$  we have  $c''_2 = c'_2 = c_2^* = 4\pi^2$ , and  $c_2 = 1/(4\pi^2)$ .

**8.** Let  $1 < p < \infty$ . Let  $\delta > 0$  and assume that  $|u(f)| \geq \delta$  in  $\mathbb{D}$ . Then there exists some  $C = C(\delta, p)$  such that

$$f = e^{i\psi}, \quad \text{with } \psi \in W^{1/p,p}(\mathbb{S}^1; \mathbb{R}) \text{ and } |\psi|_{W^{1/p,p}} \leq C |f|_{W^{1/p,p}}. \quad (10)$$

Indeed, set  $v := u/|u|$ , and write  $v = e^{i\varphi}$ , with smooth  $\varphi$ . By standard properties of the functional calculus and of trace theory, and by the lifting estimates in [4], we have  $\varphi \in W^{2/p,p}(\mathbb{D}; \mathbb{R})$ , and then  $\psi := \text{tr } \varphi \in W^{1/p,p}(\mathbb{S}^1; \mathbb{R})$  satisfies

$$|\psi|_{W^{1/p,p}} \leq C(p) |\varphi|_{W^{2/p,p}} \leq C(p) |v|_{W^{2/p,p}} \leq C(\delta, p) |u|_{W^{2/p,p}} \leq C(\delta, p) |f|_{W^{1/p,p}}.$$

**9.** Let  $1 < p < \infty$  and  $c < c'_p$ . If  $|f|_{W^{1/p,p}}^p \leq c$ , then there exists some  $\delta > 0$  such that  $|u(f)| \geq \delta$  in  $\mathbb{D}$ . Proof by contradiction: assume that  $|f_n|_{W^{1/p,p}}^p \leq c$ ,  $f_n \rightarrow g$  and  $|u(f_n)(a_n)| \leq 1/n$ . Since  $u(g \circ M_a) = [u(g)] \circ M_a$ , we may assume (by item **2**) that  $a_n = 0$ . We find that  $u(f)(0) = 0$  and  $|f|_{W^{1/p,p}}^p < c'_p$ , which is impossible.

**10.** Let  $1 < p < \infty$ . Assume that  $f_n \rightarrow f$  and  $f_n \rightarrow f$  a.e. Then  $|f_n|_{W^{1/p,p}}^p = |f|_{W^{1/p,p}}^p + |f_n \bar{f}|_{W^{1/p,p}}^p + o(1)$ . Indeed, if we set  $g_n := f_n \bar{f}$ , then this follows from the Brezis-Lieb lemma [9] and the identity

$$\overline{g_n}(x) [f_n(x) - f_n(y)] = f(x) - f(y) + \overline{g_n}(x) f(y) [g_n(x) - g_n(y)].$$

*Proof of the Theorem 1.* The proof is by complete induction on the integer part  $L := I(c_p M) = I(M/c_p^*)$  of  $c_p M$ . The case where  $L = 0$  follows from items **8** and **9**. Let  $L > 0$  and let  $M$  be such that  $I(M/c_p^*) = L$ . Assume, by contradiction, that the theorem does not hold for  $M$ . We may thus find a sequence  $(f_n)$  with the following properties:

(a)  $|f_n|_{W^{1/p,p}}^p \leq M$ .

(b) For any  $K \leq L$  and any choice of  $a_1, \dots, a_K \in \mathbb{D}$  and of signs  $\varepsilon_j = \pm 1$  such that  $\sum_{j=1}^K \varepsilon_j = \deg f_n$ , if we write  $f_n = e^{i\psi_n} \prod_{j=1}^K (M_{a_j})^{\varepsilon_j}$ , then we have  $|\psi_n|_{W^{1/p,p}} \rightarrow \infty$ . [It is always possible to take  $K$ ,  $a_j$ ,  $\varepsilon_j$  and  $\psi_n$  as above: it suffices to let  $K := |\deg f| \leq I(M/c_p'') \leq I(M/c_p^*) = L$ ,  $\varepsilon_j := \text{sgn } \deg f$ , and  $a_j = 0$ .]

By item **8** and property (b), there exist points  $a_n \in \mathbb{D}$  such that  $u(f_n)(a_n) \rightarrow 0$ . By item **2**, we may assume in addition that  $a_n = 0$ . Thus, in addition to (a) and (b), we may assume

(c)  $f_n \rightarrow f$  and  $\underline{f}_n \rightarrow f$  a.e., for some  $f$  with  $u(f)(0) = 0$ .

Set  $g_n := f_n \bar{f}$ . By item **10** and the definition of  $c_p'$ , we have  $|f|_{W^{1/p,p}}^p \geq c_p' \geq c_p^*$ , and  $|g_n|_{W^{1/p,p}}^p = M - |f|_{W^{1/p,p}}^p + o(1)$ . Let  $N > M - |f|_{W^{1/p,p}}^p$  be such that  $I(N/c_p^*) = I((M - |f|_{W^{1/p,p}}^p)/c_p^*) \leq L - 1$ . For large  $n$ , we have  $|g_n|_{W^{1/p,p}}^p \leq N$ . By the induction hypothesis, we may write (possibly up to a subsequence)  $g_n = e^{i\eta_n} \prod_{j=1}^R (M_{b_{j_n}})^{\varepsilon_j}$ , with  $|\eta_n|_{W^{1/p,p}}^p \leq F(N)$  and  $R \leq N/c_p^*$ . On the other hand, if  $d := \deg f$ ,  $b_{j_n} := 0$  and  $\varepsilon_j := \text{sgn } d$ , then we may write  $f = e^{i\eta} \prod_{j=R+1}^{R+|d|} (M_{b_{j_n}})^{\varepsilon_j}$ , with  $\eta \in W^{1/p,p}$  (item **4**). In addition, we have  $|d| \leq |f|_{W^{1/p,p}}^p / c_p''$  (item **7**). Finally, with  $\psi_n := \eta_n + \eta$  and  $K := R + |d| \leq M/c_p^*$ , we have  $f_n = e^{i\psi_n} \prod_{j=1}^K (M_{b_{j_n}})^{\varepsilon_j}$ , and  $(\psi_n)$  is bounded in  $W^{1/p,p}$ . This contradiction completes the proof of the first part of the theorem.

Optimality of (3) when  $p = 2$  follows from the fact that, by item **6**,  $f(z) := z^d$ ,  $d > 0$ , satisfies  $|f|_{W^{1/2,2}}^2 = c_2 d$  and requires at least  $d$  Moebius maps in its decomposition.  $\square$

*Proof of Corollary 1.* By replacing  $f_n$  with  $f_n \bar{f}$ , we may assume that  $f_n \rightarrow 1$ . Up to a subsequence, we may write  $f_n = e^{i\eta_n} \prod_{j=1}^P (M_{a_{j_n}})^{\varepsilon_j}$ , with  $a_{j_n} \rightarrow \alpha_j \in \overline{\mathbb{D}}$ ,  $j = 1, \dots, P$ , and  $\eta_n \rightarrow \eta$ . With no loss of generality, we assume that  $\alpha_1, \dots, \alpha_K \in \mathbb{S}^1$  and  $\alpha_{K+1}, \dots, \alpha_P \in \mathbb{D}$ . Since (clearly)  $M_{a_{j_n}} \rightarrow \alpha_j$ ,  $j = 1, \dots, K$ , we find that  $1 = e^{i(\eta_n - C)} \prod_{j=K+1}^P (M_{a_{j_n}})^{\varepsilon_j}$  for some appropriate  $C$ . Thus, with  $\zeta_n := \eta_n - \eta$ , we have

$$f_n = e^{i(\zeta_n + C)} \prod_{j=1}^K (M_{a_{j_n}})^{\varepsilon_j} \prod_{j=K+1}^P (M_{a_{j_n}} M_{\alpha_j}^{-1})^{\varepsilon_j} = e^{i\psi_n} \prod_{j=1}^K (M_{a_{j_n}})^{\varepsilon_j},$$

for some  $\psi_n$  such that  $\psi_n - \zeta_n \rightarrow C$  in  $W^{1/p,p}$ , and thus  $\psi_n \rightarrow C$ .  $\square$

*Remark.* The corollary implies the ‘‘bubbling-off of circles along a sequence of graphs’’. More specifically, the behavior of weakly converging sequences of manifold-valued maps can be investigated within the theory of Cartesian currents of Giaquinta, Modica and Souček [13]; see also [14, 17] for the specific case of  $W^{1/2,2}(\mathbb{S}^1; \mathbb{S}^1)$ . When  $p = 2$ , it is possible to define (as a current) the graph  $\mathcal{G}_f$  of  $f \in W^{1/2,2}(\mathbb{S}^1; \mathbb{S}^1)$ . With the notation in the corollary, if  $p = 2$  and  $f_n \rightarrow f$ , ‘‘bubbling-off’’ reads

$$\mathcal{G}_{f_n} \rightarrow \mathcal{G}_f + \sum_{j=1}^K \varepsilon_j \delta_{\alpha_j} \times [\mathbb{S}^1] \text{ in } \mathcal{D}_1(\mathbb{S}^1 \times \mathbb{S}^1). \quad (11)$$

This can be obtained directly from (1) [17, Proposition 3.1], but also as an immediate consequence of the corollary. Details are left to the reader.

### 3 Applications

We start with an immediate consequence of Theorem 1.

**Corollary 2.** *Let  $d$  be a non negative integer and  $\delta > 0$ . Then there exist a constant  $F(d, \delta)$  such that: every map  $f \in W^{1/2,2}(\mathbb{S}^1; \mathbb{S}^1)$  satisfying  $\deg f = d$  and  $|f|_{W^{1/2,2}}^2 \leq 4\pi^2(d+1) - \delta$  can be written as  $f = e^{i\psi} \prod_{j=1}^d M_{\alpha_j}$ , with  $|\psi|_{W^{1/2,2}}^2 \leq F(d, \delta)$ .*

Corollary 2 with  $d = 1$ , as well as a weak version of the corollary when  $d \geq 2$  were obtained in [2, Theorem 4.4, Theorem 4.8]. As an application of Corollary 2, we obtain

**Theorem 2.** *There exists some  $\varepsilon > 0$  such that, for  $p \in (2 - \varepsilon, 2]$ ,*

$$m_p := \min\{|f|_{W^{1/p,p}}^p; \deg f = 1\}$$

is achieved.

*Proof.* When  $p = 2$ , it follows from item **6** that  $m_2$  is achieved by multiples of Moebius maps.

When  $1 < p < 2$ , consider a minimizing sequence for  $m_p$ . Since  $m_p \leq |\text{Id}|_{W^{1/p,p}}^p := I_p$ , we may assume that

$$|f_n|_{W^{1/p,p}}^p \leq I_p \rightarrow I_2 = 4\pi^2 \text{ as } p \rightarrow 2. \quad (12)$$

On the other hand, when  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  we have  $|f|_{H^{1/2}}^2 \leq 2^{2-p} |f|_{W^{1/p,p}}^p$ . Thus

$$|f_n|_{H^{1/2}}^2 \leq J_p := 2^{2-p} I_p \rightarrow 4\pi^2 \text{ as } p \rightarrow 2. \quad (13)$$

For  $p$  sufficiently close to 2 and fixed  $\delta > 0$ , we have  $J_p \leq 8\pi^2 - \delta$ . We next apply Corollary 2 to  $f_n$  and write  $f_n = e^{i\psi_n} M_{\alpha_n}$ , with  $|\psi_n|_{W^{1/2,2}} \leq F(1, \delta)$ . Set  $g_n := f_n \circ M_{\alpha_n}$ . By item **2**,  $(g_n)$  is a minimizing sequence for  $m_p$ . On the other hand, we have  $g_n = e^{i\varphi_n} \text{Id}$ , with  $\varphi_n := \psi_n \circ M_{\alpha_n}$  bounded in  $W^{1/2,2}(\mathbb{S}^1; \mathbb{R})$  (by (8)). Therefore, up to a subsequence  $\varphi_n \rightharpoonup \varphi$  in  $W^{1/2,2}$ , and thus  $g_n \rightharpoonup g := e^{i\varphi} \text{Id}$  in  $W^{1/2,2}$ . We find that  $\deg g = 1$ . Since  $(g_n)$  is bounded in  $W^{1/p,p}$ , we obtain that  $g_n \rightharpoonup g$  in  $W^{1/p,p}$ . By a standard argument,  $g$  achieves  $m_p$ .  $\square$

## Acknowledgements

The author was supported by the ANR project ‘‘Harmonic Analysis at its Boundaries’’, ANR-12-BS01-0013-03, and by the LABEX MILYON (ANR-10-LABX-0070) of Universit e de Lyon, within the program ‘‘Investissements d’Avenir’’ (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR).

## References

- [1] H. Bahouri, A. Cohen, and G. Koch. A general wavelet-based profile decomposition in the critical embedding of function spaces. *Confluentes Math.*, 3(3):387–411, 2011.
- [2] L.V. Berlyand, P. Mironescu, V. Rybalko, and E. Sandier. Minimax critical points in Ginzburg-Landau problems with semi-stiff boundary conditions: existence and bubbling. *Comm. Partial Differential Equations*, 39(5):946–1005, 2014.
- [3] J. Bourgain and H. Brezis. On the equation  $\text{div} Y = f$  and application to control of phases. *J. Amer. Math. Soc.*, 16(2):393–426 (electronic), 2003.
- [4] J. Bourgain, H. Brezis, and P. Mironescu. Lifting in Sobolev spaces. *J. Anal. Math.*, 80:37–86, 2000.
- [5] J. Bourgain, H. Brezis, and P. Mironescu. Lifting, degree, and distributional Jacobian revisited. *Comm. Pure Appl. Math.*, 58(4):529–551, 2005.

- [6] H. Brezis. Problèmes de convergence dans certaines EDP non linéaires et applications géométriques. In *Goulaouic-Meyer-Schwartz seminar, 1983–1984*, pages Exp. No. 14, 11. École Polytech., Palaiseau, 1984.
- [7] H. Brezis and J.-M. Coron. Convergence de solutions de  $H$ -systèmes et application aux surfaces à courbure moyenne constante. *C. R. Acad. Sci. Paris Sér. I Math.*, 298(16):389–392, 1984.
- [8] H. Brezis and J.-M. Coron. Convergence of solutions of  $H$ -systems or how to blow bubbles. *Arch. Rational Mech. Anal.*, 89(1):21–56, 1985.
- [9] H. Brezis and E.H. Lieb. A relation between pointwise convergence of functions and convergence of functionals. *Proc. Amer. Math. Soc.*, 88(3):486–490, 1983.
- [10] H. Brezis and P. Mironescu. Sobolev Maps with Values into the Circle, Birkhäuser. In preparation.
- [11] H. Brezis and L. Nirenberg. Degree theory and BMO. I. Compact manifolds without boundaries. *Selecta Math. (N.S.)*, 1(2):197–263, 1995.
- [12] P. Gérard. Description du défaut de compacité de l’injection de Sobolev. *ESAIM Control Optim. Calc. Var.*, 3:213–233 (electronic), 1998.
- [13] M. Giaquinta, G. Modica, and J. Souček. *Cartesian currents in the calculus of variations. I*, volume 37 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1998. Cartesian currents.
- [14] M. Giaquinta, G. Modica, and J. Souček. On sequences of maps into  $S^1$  with equibounded  $W^{1/2}$  energies. *Selecta Math. (N.S.)*, 10(3):359–375, 2004.
- [15] S. Jaffard. Analysis of the lack of compactness in the critical Sobolev embeddings. *J. Funct. Anal.*, 161(2):384–396, 1999.
- [16] P.-L. Lions. The concentration-compactness principle in the calculus of variations. The limit case. II. *Rev. Mat. Iberoamericana*, 1(2):45–121, 1985.
- [17] P. Mironescu and A. Pisante. A variational problem with lack of compactness for  $H^{1/2}(S^1; S^1)$  maps of prescribed degree. *J. Funct. Anal.*, 217(2):249–279, 2004.
- [18] H.-M. Nguyen. Inequalities related to liftings and applications. *C. R. Math. Acad. Sci. Paris*, 346:957–962, 2008.
- [19] J. Sacks and K. Uhlenbeck. The existence of minimal immersions of 2-spheres. *Ann. of Math. (2)*, 113(1):1–24, 1981.
- [20] M. Struwe. A global compactness result for elliptic boundary value problems involving limiting nonlinearities. *Math. Z.*, 187(4):511–517, 1984.