

Deciphering genetic diversity and inheritance of tomato fruit weight and composition through a systems biology approach

Laura Pascual-Banuls, Jiaxin Xu, Benoit Biais, Mickael Maucourt, Patricia Ballias, Stéphane Bernillon, Catherine Deborde, Daniel Jacob, Aurore Desgroux, Mireille Faurobert, et al.

▶ To cite this version:

Laura Pascual-Banuls, Jiaxin Xu, Benoit Biais, Mickael Maucourt, Patricia Ballias, et al.. Deciphering genetic diversity and inheritance of tomato fruit weight and composition through a systems biology approach. Journal of Experimental Botany, 2013, 64 (18), pp.1-16. 10.1093/jxb/ert349. hal-01208665

HAL Id: hal-01208665 https://hal.science/hal-01208665v1

Submitted on 29 May 2020 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 Deciphering genetic diversity and inheritance of tomato fruit weight and composition

2 through a systems biology approach

Laura Pascual* (1), Jiaxin Xu* (1, 2), Benoît Biais (3, 4), Mickaël Maucourt (4, 5), Patricia
Ballias (3-5), Stéphane Bernillon (3, 5), Catherine Deborde (3), Daniel Jacob (3, 5), Aurore

6 Desgroux (1), Mireille Faurobert (1), Jean-Paul Bouchet (1), Yves Gibon (3, 5), Annick

7 Moing (3), Mathilde Causse (1)

9 (1) INRA, UR1052, Unité de Génétique et Amélioration des Fruits et Légumes, F-84143
10 Avignon, France

11 (2) Northwest A&F University, College of Horticulture, Yang Ling, Shaanxin, 712100, P.R.

12 China

13 (3) INRA-UMR 1332 Biologie du Fruit et Pathologie, Centre INRA de Bordeaux, F-33140

- 14 Villenave d'Ornon, France
- (4) Université de Bordeaux, UMR1332 Biologie du Fruit et Pathologie, Centre INRA de
 Bordeaux, F-33140 Villenave d'Ornon, France
- 17 (5) Metabolome Facility of Bordeaux Functional Genomics Center, IBVM, Centre INRA de
- 18 Bordeaux, F–33140 Villenave d'Ornon, France
- 19 * Equal contributors

Mail of the authors: Laura Pascual <Laura.pascual@avignon.inra.fr>, Jiaxin Xu

22 <jiaxin.xu@avignon.inra.fr>, Benoit Biais <benoit.biais@bordeaux.inra.fr>, Mickael Maucourt

<mickael.maucourt@bordeaux.inra.fr>, Patricia Ballias <patricia.ballias@bordeaux.inra.fr>, Stephane Bernillon

 $\label{eq:catherine.deborde} \texttt{24} \qquad < stephane.bernillon@bordeaux.inra.fr>, Catherine Deborde < catherine.deborde@bordeaux.inra.fr>, Daniel$

Jacob <daniel.jacob@bordeaux.inra.fr>, Aurore Desgroux <aurore.desgroux@rennes.inra.fr>, Mireille Faurobert

<mireille.faurobert@avignon.inra.fr>, Jean-Paul Bouchet <bouchet@avignon.inra.fr>, Yves Gibon

<yves.gibon@bordeaux.inra.fr>, Annick Moing <annick.moing@bordeaux.inra.fr>, Mathilde Causse

<mathilde.causse@avignon.inra.fr>

Corresponding author : Mathilde Causse (Email: <u>Mathilde.Causse@avignon.inra.fr</u>), Ph +33 4 32 72 26 60, Fax +33 4 32 72 27 02

- 32 Date of submission:
- **Total word counts: 6724** (From abstract to references)
- **Included** 7 colored online-only Figures; 3 Tables; 20 Supplementary data.

3

8

20

21

23

25

26

27

28

1 ABSTRACT

Integrative systems biology proposes new approaches to decipher the variation of phenotypic 2 traits. In an effort to link the genetic variation and the physiological and molecular bases of 3 fruit composition, we characterized the proteome (424 protein spots), metabolome (26 4 compounds), enzymatic profile (26 enzymes) and phenotypes of eight tomato accessions, 5 covering the genetic diversity of the species, and four of their F1 hybrids, at two fruit 6 7 developmental stages (cell expansion and orange-red). The contents in metabolites varied among the genetic backgrounds, while enzyme profiles were less variable, particularly at cell 8 9 expansion stage. Frequent genotype by stage interactions suggested that the trends observed for one accession at a physiological level may change in another one. In agreement with this, 10 the inheritance modes varied between crosses and stages. Although additivity was 11 predominant, 40% of the traits were non additively inherited. Relationships among traits 12 13 revealed associations between different levels of expression and provided information on several key proteins. Notably the role of frucktokinase, invertase and cysteine synthase in in 14 15 the variation of metabolites was underlined. Several stress related proteins also appeared related to fruit weight differences. These key proteins might be targets for improving 16 17 metabolite contents of the fruit. This systems biology approach provides better understanding of networks controlling the genetic variation of tomato fruit composition. Besides, the wide 18 data sets generated provide an ideal framework to develop innovative integrated hypothesis 19 20 and will be highly valuable for the research community.

21 Key words: Systems biology, tomato, fruit, metabolome, proteome.

1 INTRODUCTION

Identifying the genes controlling the variation of complex traits is a key goal of evolutionary 2 genetics and plant biology. Attempts to identify genetic variants underlying quantitative traits 3 have been achieved by traditional linkage mapping and genome wide association studies 4 using molecular markers. However, quantitative trait loci (QTL) resolution is limited and the 5 identification of the polymorphisms responsible for the variation not straightforward. 6 Furthermore, as several intermediate levels interact between the genotypes and the 7 phenotypes, DNA sequence variation (single nucleotide polymorphisms SNPs or Indel) may 8 9 not directly affect the traits. Intermediate molecular phenotypes such as gene expression, protein abundance and metabolite concentration also vary in populations and are themselves 10 quantitatively inherited (Rockman and Kruglyak 2006). Nowadays, rapid technological 11 advances in high-density experiments such as next-generation sequencing (NGS), RNA 12 13 expression analysis through microarray or RNAseq, mass spectrometry (MS) coupled to gas chromatography (GCMS) or to liquid chromatography (LCMS) and nuclear magnetic 14 resonance (NMR) metabolic profiling enable scientists to obtain large exhaustive datasets and 15 analyze biological systems as a whole. Integration of the genome expression products at 16 different levels should help dissecting the genetic variation of a given quantitative trait. 17

Systems biology proposes to relate the variation analyzed at different expression levels, from 18 phenotype to metabolome and proteome, studying the behavior of all the elements in a 19 biological system (Gutierez et al., 2008; Saito and Matsuda 2010). A bottom-up systems 20 biology approach consists in integrating 'omic' resources (genomic, transcriptomic, 21 proteomic, and metabolomic) and large physiological datasets, together with statistical 22 network analysis in order to identify candidate genes underlying phenotypes and construct 23 complex regulation networks (Kliebenstein 2010). This approach was first applied to yeast by 24 combining DNA microarrays and quantitative proteomics to describe the galactose pathway 25 (Ideker et al., 2001). It was then applied to gene expression analysis in E. coli (Rosenfeld et 26 al., 2002), and to Arabidopsis by Hirai et al. (2005) who elucidated gene to gene and 27 metabolite to gene networks by integrating metabolomic and transcriptomic data. Systems 28 biology has also been used to study the natural genetic variation at different levels, such as 29 metabolomics (Keurentjes 2009; Kliebenstein 2009a), proteomics (Stylianou et al., 2008) and 30 transcriptomics (Keurentjes et al., 2008; Kliebenstein 2009b). 31

Tomato (*Solanum lycopersicum*) is the model species for the study of fleshy fruit development and composition (Giovannoni et al., 2004). It is a self-pollinated species and derived from its closest wild ancestor *Solanum pimpinellifolium* (Nesbitt and Tanksley 2002). Cherry tomato

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

accessions (Solanum lycopersicum var. cerasiforme) have an intermediate position between 1 these two species, as their genome is a mosaic of those from S. lycopersicum and S. 2 pimpinellifolium (Ranc et al., 2008). During tomato domestication the diversification of fruit 3 aspect, as well as the adaptation to a wide range of environmental conditions was 4 simultaneous to a strong reduction of molecular diversity (Miller and Tanksley 1990; Blanca 5 et al., 2012). This lack of genetic variation in cultivated species led geneticists to study trait 6 variation mostly in distant crosses involving wild species and thus limited the exploitation of 7 intra-specific variation. Today the availability of the tomato genome sequence (Tomato 8 9 Genome Consortium 2012) and of a large number of SNP markers (Sim et al., 2012) allows a re-examination of the variation and inheritance of agronomical and fruit traits at the intra-10 specific level. 11

Systems biology approaches have been used in tomato to study fruit development. Carrari et 12 13 al. (2006) and Mounet et al. (2009) analyzed transcriptome and metabolome variation along fruit development. Garcia et al. (2009) combined phenotype, metabolome, transcriptome and 14 proteome profiles to study genes related to ascorbate metabolism in three transgenic lines. 15 Wang et al. (2009) compared transcriptome and metabolome to uncover the molecular events 16 underlying fruit set, while Osorio et al. (2011) compared enzyme activity, metabolite and 17 transcript profiles to analyze the connectivity between these groups of traits in fruit ripening 18 mutants. However, these studies were only focused on a few mutants or on the effect of 19 introgression in S. lycopersicum of wild species alleles. Little is known about the genetic 20 variation in metabolic, enzymatic and proteomic profiles contributing to phenotypic trait 21 variation inside the species. 22

In the present study, we aimed at deciphering the complex relationships between several 23 successive levels of omic profiles to characterize the genetic variation and physiological bases 24 of quantitative traits in tomato fruit. For this purpose, we first compared the variation of eight 25 genotypes representing a large range of phenotypic and genotypic diversity (four S. 26 lycopersicum and four S. l. var cerasiforme) and four of their corresponding hybrids at two 27 28 stages of fruit development (cell expansion and orange-red). We characterized their metabolic, enzymatic and proteome profiles. Genetic variability was analyzed for all traits, and 29 inheritance patterns of traits that were significantly different among genotypes were assessed. 30 Relationships among traits were analyzed within and between each group of traits at each 31 32 stage and networks were constructed using sparse partial least square regression. This systems biology approach combining proteome, metabolome and phenotypic analysis gave insights 33 34 into the diversity and relationships of quantitative traits at different levels.

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

1 MATERIALS AND METHODS

3 **Plant materials**

Eight tomato lines including four S. lycopersicum accessions (Levovil, Stupicke Polni Rane, 4 LA0147 and Ferum) and four S. l. lycopersicum var. cerasiforme accessions (Cervil, Criollo, 5 Plovdiv24A and LA1420) and four of their corresponding F1 hybrids (Levovil x Cevil, 6 Stupicke Polni Rane \times Criollo, LA0147 \times Plovdiv24A and Ferum \times LA1420) were used in 7 this study (details of the accessions are shown on Supplementary Table S1 and Supplementary 8 Figure S1). Lines were selected, based on a previous molecular characterization of 360 9 tomato accessions (Ranc et al., 2008), to include the maximum genetic diversity of the 10 species. Xu et al. (2013a) genotyped these lines and the line sequenced to obtain the reference 11 genome (Heinz1706, Tomato genome consortium 2012). From the 139 single nucleotide 12 13 polymorphism markers (SNPs) characterized, 133 were polymorphic (96%) showing the large range of molecular diversity represented by the eight lines. The range of polymorphism 14 between the lines and the reference genome (Heinz1706) ranged from 27% to 82% 15 (Supplementary Table S2) The genetic distances among the parents of F1 hybrids were 16 variable. According to Xu et al. (2013a) data, Levovil and Cervil were the two most distant 17 accessions (82% SNP polymorphic), followed by LA0147 x Plovdiv 24A (40%), Stupicke 18 Polni Rane x Criollo (34%) and Ferum x LA1420 (27%). 19

Plants were grown during 2010 spring under greenhouse conditions (16/20°C) in Avignon
(South of France). Plants were separated in two blocks, five plants per genotype were
included in each block.

For proteome, metabolome and enzymatic measurement two stages of development, cell 23 expansion (CE) and orange-red (OR) stage were selected. CE stage was chosen a 24 representative stage of the growing tomato fruit, OR stage was chosen because it is 25 unequivocally determined and is the key step where enzyme and protein concentrations are 26 changing and will determine the final characteristics of the fruit. The number of days after 27 28 anthesis to reach cell expansion varied among genotypes depending on their fruit size. Thus CE sampling was done at 14, 20 or 25 days after anthesis for small [Cervil], medium [Criollo, 29 30 Plovdiv 24A, Stupicke Polni Rane and the four F1 hybrids] or large [LA0147, Levovil and Ferum] fruited accessions respectively. OR sampling was done based on fruit color change. 31 32 Three biological replicates by stage were analyzed. Each replicate included 7 to 20 fruits from both greenhouse blocks to buffer environmental variations. Fruit pericarps were collected, 33 34 immediately frozen, ground in liquid nitrogen and stored at -80 °C until analysis. For fruit

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

phenotypic trait measurements, five fruits were harvested from the ten plants of each
genotype at the following six stages: (1) cell expansion stage, (2) cell expansion +7 days, (3)
+14, (4) +21, (5) orange-red stage and (6) red ripe. Fruits were evaluated for fresh weight
(FW), fruit diameter (FD, measured using a caliper) and dry matter content (DMC). Dry
matter content (expressed in g / 100 g FW) was assessed after 5 days in a ventilated oven at
80 °C.

7 8

9 Metabolome and enzyme activity analysis

Metabolome analyses were performed at the Metabolome Facility of Bordeaux, using 10 quantitative proton NMR (¹H-NMR) profiling of polar extracts and liquid chromatography 11 quadrupole time-of-flight tandem mass spectrometry (LC-QTOF-MS) profiling of semi polar 12 extracts. ¹H-NMR profiling was performed as described in Deborde et al. (2009) with minor 13 modifications. Briefly, polar metabolites were extracted on lyophilized powder (50 mg DW 14 per biological replicate) with an ethanol-water series at 80°C. The lyophilized extracts were 15 titrated to pH 6 and lyophilized again. Each dried titrated extract was solubilized in 0.5 mL 16 D₂O with (trimethylsilyl) propionic-2,2,3,3-d₄ acid (TSP) sodium salt (0.01% final 17 concentration) for chemical shift calibration and ethylene diamine tetraacetic acid (EDTA) 18 disodium salt (5 mM final concentration for CE and 2 mM for OR stage). ¹H-NMR spectra 19 were recorded at 500.162 MHz on a Bruker Avance III spectrometer (Bruker, Karlsruhe, 20 Germany) using an ATMA inverse 5 mm probe flushed with nitrogen gas and an electronic 21 reference for quantification (ERETIC2). Sixty-four scans of 32 K data points each were 22 acquired with a 90° pulse angle, a 6000 Hz spectral width, a 2.73 s acquisition time and a 25 s 23 recycle delay. Two technological replicates were used per biological replicate. Preliminary 24 data processing was conducted with TOPSPIN 3.0 software (Bruker Biospin, Wissembourg, 25 France). The assignments of metabolites in the ¹H-NMR spectra were made by comparing the 26 proton chemical shifts with values of the MeRy-B metabolomic database (Ferry-Dumazet et 27 al., 2011), by comparison with spectra of authentic compounds recorded under the same 28 solvent conditions and/or by spiking the samples. The metabolite concentrations were 29 calculated using AMIX (version 3.9.7, Bruker, Karlsruhe, Germany) software. The 144 ¹H-30 NMR spectra of the data set were converted into JCAMP-DX format and deposited with 31 associated metadata into Metabolomics Repository of Bordeaux MeRy-B (Ferry-Dumazet et 32 al., 2011, http://www.cbib.u-bordeaux2.fr/MERYB/view/project/34). 33

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

LC-QTOF-MS profiling of aqueous-methanol-0.1% formic acid extracts was performed from 1 lyophilized powder (20 mg in 1 ml). For each biological replicate, two extractions were 2 performed and two injections per extract were used. An Ultimate 3000 HPLC (Dionex, 3 Sunnyvale, CA, USA) was used to separate metabolites on a reversed phase C18 column (150 4 x 2.0 mm, 3 µm; Phenomenex, Torrance, CA, USA) using a 30 min linear gradient from 3 to 5 95% acetonitrile in water acidified with 0.1% formic acid. Metabolites were detected using a 6 quadrupole time-of-flight (QTOF) mass spectrometer (Bruker, Bremen, Germany). 7 Electrospray ionization in positive mode was used to ionize the compounds. Scan rate for ions 8 9 at m/z range 100-1500 was fixed at 2 spectra per second. Methyl vanillate was spiked in the extraction solvent and used as an internal standard. One sample was used as a QC sample and 10 injected each ten injections. Raw data were processed in a targeted manner using 11 QuantAnalysis 2.0 software (Bruker, Bremen, Germany). This resulted in eight compounds 12 13 identified based on accurate mass measurement and comparison with data from Gomez-Romero et al. (2010). 14

Ascorbic acid was measured using a spectrofluorometric method and values expressed as total ascorbate (ascorbic acid + dehydroascorbate) as previously described by Stevens et al. (2007). The maximum activity (Vmax) of 26 enzymes of the primary metabolism was assayed using a robotized platform as described in Gibon et al. (2004) and in Steinhauser et al. (2010). **Supplementary Tables S3 and S4** present the lists of the primary and secondary metabolites and the enzyme activities analyzed.

22 **Proteome analysis**

Methods for protein extraction, two-dimensional gel electrophoresis (2-DE), protein 23 identification and classification were as detailed in Xu et al. (2013b). Briefly, proteins were 24 extracted using the phenol extraction method developed by Faurobert et al. (2007). Later, 25 proteins were separated by 2-DE. After Coomassie colloidal staining, image analysis was 26 performed with Samespot software (version 4.1, city, country) and the normalized spot 27 volumes were obtained. Protein identification of 424 variable spots was performed at the 28 proteome platform of Le Moulon (Gif-sur-Yvette) using nano-LC-MS/MS method following 29 the procedure described in Xu et al. (2013b). The database search was run against the 30 International Tomato Annotation Group (ITAG) Release 2.3 of predicted proteins (SL2.40) 31 database (http://solgenomics.net/) with X!Tandem software (http://www.thegpm.org/TANDEM/, 32 version 2010.12.01.1). Fasta sequence of the identified proteins was employed to re-annotate 33 34 the proteins using the Blast2GO package (Conesa et al. 2005). Sequences were compared

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

1 against the NCBI-nr (version April 9, 2012) database of non-redundant protein sequence

2 using BLASTX with the default settings.

3

4 Statistical analysis and inheritance analysis

Metabolite contents and enzyme activities were expressed on a dry weight basis to be
comparable. All the analyses were performed using R (R Development Core Team 2012,
http://www.R-project.org/).

8 Data were submitted to a two-way ANOVA (P<0.05) with genotype, stage and interaction 9 effect and then to one-way ANOVA with genotype effect at each stage. Besides, to assess the 10 mode of inheritance of the traits, one-way ANOVA was also performed with genotype effect 11 for each cross (two parental lines and their hybrid) and stage.

Means and standard deviations were calculated for each trait (phenotypic traits, metabolite 12 13 contents, enzyme activities, protein spot volumes) in each genotype and stage. Significantly different (P < 0.05) traits in each cross were selected at each stage to estimate additive (A) and 14 15 dominance (D) components of genetic variation. A is equivalent to half of the difference between two parental lines. The S. lycopersicum line was systematically the first parent in a 16 17 cross. D is the difference between the hybrid value and the parental mean. The inheritance pattern of each trait was then assessed by the dominance/additivity (D/A) ratio and classified 18 as over-recessive (OR; D/A <-1.2), recessive (R; $-1.2 \le D/A \le -0.8$), additive (A; -0.8 < D/A19 < 0.8), dominant (D; $0.8 \le D/A \le 1.2$), over-dominant (OD; D/A>1.2). 20

Means of metabolite contents, enzyme activities and protein spot volumes were centered and 21 scaled to variance unit and used for the rest of the analysis. Principal component analyses 22 (PCA) were performed for metabolites, enzymes and phenotypic traits, as well as for protein 23 spot volumes for both development stages and at each stage, with the "pcaMethods" package 24 (Stacklies et al., 2007). Pearson correlations and p-values were calculated between 25 significantly variable traits at each stage. Correlations were considered to be significant when 26 $|\mathbf{r}| > 0.7$ (p-value <0.01). Significant correlations were plotted using R "corrplot" package (Wei 27 2012, http://CRAN.R-project.org/package=corrplot). To analyze the relationships among 28 protein spot volumes and other traits, networks were reconstructed and visualized using 29 sparse partial least squares correlation regression (sPLS) analysis with the "mixOmics" 30 package (Lé Cao et al., 2009). An arbitrary threshold of 0.7 was employed for network 31 reconstruction. Nodes represent the different traits and edges represent the relations between 32 variables belonging to different levels. 33

1 **RESULTS**

To represent a large range of the genetic diversity eight tomato accessions were chosen 2 according to previous studies (Ranc et. al., 2008; Xu et. al., 2013a). The eight accessions and 3 their four hybrids were characterized at phenotypic, metabolic and proteomic levels. The final 4 fruit weight of the eight parental lines and the four hybrids ranged from 5.3g to 134.4 g. Fruit 5 weights of the four hybrids were intermediate between the values of their parental lines all 6 along fruit development (Supplementary Figure S2). Fruit diameter (Fig. 1) was highly 7 correlated to fruit weight, as fruits were round. Dry matter content also showed a wide range 8 9 of variation (Fig. 1 and Supplementary Table S5).

11 Metabolome, enzyme and proteome profiles strongly differ among accessions

Metabolome profiling by ¹H-NMR and LC-QTOF-MS allowed the quantification of eighteen 12 13 metabolites from the central carbon metabolism and eight secondary metabolites (Supplementary Table S3). Besides, 26 enzyme activities were assessed by robotized assays 14 (Supplementary Table S4). These analyses provided a detailed characterization of sugars, 15 organic acids and amino acids metabolism pathways, as well as glycoalkaloids and phenolic 16 compounds (Fig. 2). Proteins were isolated from 2-D PAGE. A total of 1230 protein spots 17 were detected. A subset of 424 spots whose abundance was significantly different between 18 genotypes or stages, were sequenced by LC-MS/MS. Four hundred and twenty two spots 19 were identified (Xu et al., 2013b). Supplementary Table S5 lists the mean and standard 20 deviation of every trait for each genotype and stage. 21

The 12 accessions differed for most of the metabolites and phenotypic traits according to the 22 ANOVAs (Table 1). The means of most of the traits (27/29) were significantly different 23 across stages. The content of glucose, fructose, citrate, asparagine, aspartate and 24 phenylalanine increased from CE to OR while the other amino acids decreased. The 25 interactions between stage and genotype were significant for 93% of 29 metabolite and 26 phenotypic traits and 50% of these traits showed different trend according to the genotype. We 27 thus analyzed the data stage by stage (Table 1). A large range of variability was observed 28 among the 12 genotypes at each stage as all the trait means were significantly different except 29 for the crypto-chlorogenic acid. The fold-change difference between genotypes reached 30 values as high as 5.6 for threonine content at CE or 7.9 for malate at OR. 31

The activity of 26 enzymes from central carbon metabolism, including enzymes of the Calvin cycle, glycolysis, sucrose metabolism, TCA cycle and amino acid metabolism was quantified and expressed relative to dry weight to be comparable with the metabolome and proteome.

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

Enzyme activities exhibited a lower range of variation than metabolites (**Table 2**). The greatest differences were found between stages, where all the enzyme activities difered except alanine aminotransferase, fumarase and glyceraldehyde-3-phosphate dehydrogenase (NADP). The activity of 15 enzymes was greater at CE. Two and thirteen enzyme activities were significantly different among accessions at CE and OR, respectively.

The volume of the 424 protein spots was compared among the 12 genotypes. The genes 6 corresponding to most of these spots are identified (Xu et al., 2013b; Supplementary Table 7 S6). They include 133 protein spots related to primary metabolism. Several multi-spot 8 proteins (one gene corresponding to several spots) were detected, such as acid invertase (7 9 spots), phosphoglucomutase and enolase (5 spots). These multispots may be caused by post-10 transcriptional and post-traductional modifications or by allelic variations (Xu et al., 2013b). 11 A large range of variability was observed among genotypes and between stages for all the 12 13 protein spot amounts (Supplementary Table S6). As for metabolites and enzymes, the main differences were observed between stages (84% significantly variable spots; **Supplementary** 14 Table S6). with 46% in lower amount and 38% in higher amount at OR. When we analyzed 15 the data stage by stage, 256/424 spot amounts were significantly different among genotypes at 16 CE and 274/424 at OR. 17

The variation among the 12 accessions at the different levels was illustrated by PCA analysis. 18 When we analyzed the phenotypic traits, metabolite and enzyme profiles at both stages, two 19 main groups corresponding to each stage of development were detected (Supplementary 20 Figure S3A). Similar results were obtained for the protein spot volumes (Supplementary 21 Figure S3B). PCA were thus computed stage by stage (Fig. 3). In every case Cervil (the 22 accession with smaller fruits) was separated from the other genotypes, and the large fruited 23 accessions (Levovil, LA0147 and Ferum) were grouped together. Hybrids were usually 24 located in between their parental lines. 25

26

27 Inheritance of traits is predominantly additive

The four F1 hybrids derived from crosses among the eight lines and corresponded to different distances among parental lines. We assessed the mode of inheritance of the traits that were significantly different for each cross separately (**Supplementary Table S7; Fig. 4**).

The phenotypic traits, fruit diameter and fruit weight, were additive in the four crosses at each stage. The dry matter content was additive or over-recessive or not significant according to the stage and cross (**Fig. 1**). Most of metabolic contents were significantly variable at both stages. A large number of additive traits was found in the cross between the most distant lines

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

1 (Levovil x Cervil) (**Fig. 4A**). Traits could exhibit different inheritance modes at the two stages

2 for the same cross or in different crosses, as illustrated for citrate content on Fig. 5.

Most of the enzyme activities were not significantly variable within one cross, so the inheritance mode of only a few enzyme activities was assessed. At CE, Ferum x LA1420 was the most variable cross with four significantly variable enzymes, while for the other crosses, only one or non enzyme were variable. At OR, the predominant inheritance mode was additivity for Levovil x Cervil and Stupicke Polni Rane x Criollo but not for the two other crosses.

9 The number of protein spots significantly variable varied among crosses at CE in relation to 10 the genetic distance between the parental lines (**Fig. 4B**). As for the metabolites, proteins 11 showed different inheritance patterns at the two stages in the same cross or in different 12 crosses. On average, 40% of the variable traits showed a non additive mode of inheritance 13 without bias against recessivity or dominance.

14

15 Dissection of relationships among traits

Relationships among traits were only assessed among the traits significantly different between 16 genotypes at each stage (Supplementary Tables S8, S9, S10). The significant correlations 17 (summarized in Supplementary Table S8) were more frequent than expected by chance, with 18 an excess of positive correlations. Correlations among metabolites, phenotypic traits and 19 enzymes activities are illustrated on Figure 6). At CE, sugars (glucose and fructose) were 20 highly correlated together and negatively with most amino acids, tomatine and dry matter 21 content. Amino acid contents were highly correlated together (Fig. 6A). Very few correlations 22 were detected between metabolites and phenotypic traits at OR (Fig. 6B). For enzyme 23 activities correlations were significant between the glycolysis and TCA cycle enzymes at OR 24 (Fig. 6B). Very few correlations were significant between enzyme activities and metabolite 25 contents.Protein spot volumes were more frequently correlated with metabolite contents at CE 26 and with enzyme activities at OR, where a large number of positive correlations with spots 27 28 annotated as primary metabolism and stress response was detected. Correlations are summarized 1n Supplementary table S8 and provided on Supplementary Tables S9 and S10. 29 30 Correlations between protein spots corresponding to enzymes and their enzyme activities were analyzed at OR. In total 28 spots corresponding to 8 enzymes were analyzed 31 (Supplementary Table S11). Significant correlations, ranging from r=0.877 to 0.715, were 32 detected between enolase activity and the four spots annotated as two enolase genes 33 34 (Solyc09g009020 and Solyc10g085550), aldolase activity and one aldolase gene

Manuscrit d'auteur / Author manuscript

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

(Solyc09g009260) and acid invertase and two spots corresponding to the Solyc03g083910 acid invertase gene. For these enzymes, even when correlations were not significant (P<0.01) they were often positive with P-value<0.05 (Supplementary Figure S4). The other enzymes analyzed, pyruvate kinase, glyceraldehyde-3-phosphate dehydrogenase (NAD), isocitrate dehydrogenase, malic enzyme (NADP) and malate dehydrogenase were not significantly correlated with their corresponding spot volumes.

Table 3 lists the protein spots whose volume was strongly correlated with fruit weight or dry matter content. The numbers of spots were equivalent at both stages, with seven of the 15 spots related to stress response (heat shock proteins, NifU like protein, chaperonin). A larger proportion (13/32) of correlations was detected between dry matter content and spots related to primary metabolism (fructokinase, malate dehydrogenase, acid invertase, enolase). Most of the correlations with dry matter content were positive, on the contrary to those with fruit weight.

14

15 Reconstruction of networks integrating metabolic and protein profiles

Due to de large number of traits and correlations, we used sparse partial least square 16 regression (sPLS) for integrating protein expression data and metabolites, enzymes and 17 18 phenotypes. sPLS is a bidirectional multivariate regression method that allows separate modeling of covariance between two data sets. The main advantage of sparse methods over 19 20 non-sparse methods is that it sets the contribution of noise variables to zero to improve the prediction or classification performance (Filzmoser et al., 2012). sPLS networks relating 21 protein spot volumes with phenotypes, metabolites and enzymes were constructed at each 22 stage. We grouped the three levels (phenotypes, metabolites and enzymes) considering they 23 represent a global metabolic-related level to be related to the proteome level. 24

At CE, a network was constructed between, metabolites, phenotypic traits and enzyme activities (variable among genotypes) on one hand, and 77 variable protein spots related to primary and secondary metabolism and vitamin synthesis on the other hand. The network reconstructed connected 8 traits and 26 proteins by more than 50 edges (**Supplementary Figure S5 and Supplementary Table S12**), among which two fructokinase spots were connected to glucose, fructose and dry matter content.

A network connecting variable metabolites, phenotypic traits and enzymes activities with 87 protein spots (related with primary and secondary metabolism and vitamin synthesis) was also constructed for OR stage. Two main networks were obtained, with more connections than

for CE stage (Supplementary Figure S6 and Supplementary Table S13). Sucrose and dry 1 matter content played a pivotal role. They were linked to 11 and 17 proteins, including spots 2 corresponding to acid invertase, enolase, malate dehydrogenase and malic enzyme 3

As the variation of proteins expressed during CE may influence the metabolome and activome 4 at a later stage, we analyzed the connections between protein variations at CE and fruit 5 composition and enzyme activities at OR (Fig. 7, Supplementary Table S14). We detected a 6 relation between sucrose content at OR and the volumes of two spots corresponding to 7 fructokinase at CE. Besides those spots were also related with the fructose content at CE 8 (Supplementary Figure S5). An isocitrate dehydrogenase spot was related to isocitrate 9 activity at OR, and phosphoglycerate kinase from the Calvin cycle to shikimate 10 dehydrogenase activity, an enzyme downstream the erythrose-4P produced in that cycle. The 11 amounts of cysteine systehase and fructokinase 3 proteins had a pivotal role, each being 12 13 connected to several traits.

Networks were also constructed between phenotypes, metabolites and enzyme activities and 14 the proteins corresponding to other functions (data not shown). The most interesting 15 relationship involved a Chaperonin (JX383) that played an important role at CE, as it was 16 related to six enzymes activities, and to glucose and fructose content at OR. 17

DISCUSSION 19

The variation of tomato fruit composition has been widely studied, due to its role in sensory 20 and nutritional value. However, until now the variation of metabolic compounds has been 21 studied in tomato either along fruit development or according to environmental perturbations, 22 mainly in one accession or in lines resulting from the introgression of genome fragments from 23 a unique wild species (Schauer et al., 2006; Steinhauser et al., 2010). Results are subsequently 24 supposed to represent the variation of the species. In the present study, we aimed to analyze 25 the actual variation of the species, by comparing eight accessions selected to represent a large 26 part of the phenotypic and molecular diversity of S. lycopersicum (Ranc et al., 2008). We 27 28 described the variation, the inheritance and the relationships among metabolic, enzymatic and proteomic traits assessed at two developmental stages. 29

A large range of genetic variation is detected at all levels

32 A large range of variation was observed for most of the phenotypic and metabolic traits at least at one stage. Usually, the ratio of maximum to minimum values among genotypes varied in the range of two to three, showing the wide variation present in the species. The secondary

18

30

31

33

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

metabolites showed a higher range of variation, some of them being present in one line and 1 2 almost absent in another. This may be due to the inclusion in the study of S. lycopersicum var cerasiforme accessions which are not fully domesticated, as domestication caused great 3 alterations on those compounds (reviewed by Meyer et al., 2012). Among enzyme activities, 4 we detected significant differences between stages, and the genetic variation was less 5 significant at CE. Steinhauser et al. (2011) observed the same tendency, with enzyme 6 activities having a lower heritability than metabolites, suggesting that metabolites have a tight 7 8 regulation, while enzyme activities can be compensated by coordinated changes in other enzymes. 9

Until now proteome variation in tomato fruit and protein amount inheritance were poorly 10 documented (reviewed by Faurobert et al., 2013). Faurobert et al. (2007) described the 11 proteome variation of tomato pericarp in one line along fruit development. They identified 12 13 could identify the function of 90 spots. We studied 424 protein spots that were variable among stages or genotypes (Xu et al., 2013b). Thanks to the release of the tomato genome sequence, 14 we identified the function of almost every spot and detected 307 unique proteins 15 corresponding to 424 spots. Most of the spots variable at both stages showed the same 16 tendency (increase or decrease along fruit development) in all the genotypes. Nevertheless 17 57% of the spots revealed significant Genotype by Stage interactions, indicating that the trend 18 observed in one genotype at a given physiological level (stage) may change in another 19 20 genotype.

The observed variation may be related to the genetic distance among accessions. In the PCA 21 analysis, all the large-fruited lines closely related at the molecular level were grouped 22 together, while the small cherry tomatoes, genetically more diverse, were more spread. 23 Besides, Cervil, the most distant line from all others at the molecular level, presented a very 24 specific profile for every trait, leading to most of the extreme values (lowest fruit weight, 25 highest dry matter, sugar and acid contents). It was also specific in terms of secondary 26 metabolites, with high content in chlorogenic acid, dehydrotomatin and rutin. The large 27 28 variation detected and the differences between genotypes along fruit development showed the important effect of genetic diversity in fruit composition and enhances the value of the 29 30 presented dataset.

32 Diversity in the modes of inheritance among crosses and traits

Hybrids are widely used in modern agriculture, either for heterosis (the advantage of a hybrid compared to both parents) or for the combination of dominant traits. Agronomical traits often

31

33

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

show heterosis in the F1 when distant cultivated accessions or cultivated and wild species are
 crossed (Springer and Stupar 2007; Li et al., 2008). The molecular origin of heterosis has been

3 studied for years and is usually related to a combination of dominance or over-dominance

4 effects and to epistatic interactions (Stuber, 2010).

In tomato, fewer traits than in maize, a highly heterotic crop, show a systematic heterosis 5 trend (Lipman and Zamir, 2007). Steinhauser et al. (2011) studied the enzyme activities in 6 introgression lines derived from the wild species S. pennellii and found approximately 7 equivalent ratio of QTL showing additive, recessive and dominant modes of inheritance, with 8 only 5% showing overdominance. In the present study, the number of traits significantly 9 variable within each cross (one hybrid and its two parents) differed from one cross to the 10 other and was related to the genetic distance at the proteomic level. Regarding to phenotypic 11 traits, in accordance with the absence of heterosis, fruit weight and diameter were additive in 12 13 the four crosses. Around 60% of the other traits showed an additive inheritance, with a number of traits exhibiting an over-dominant or over-recessive mode of inheritance, but no 14 specific trend towards one of them. The higher rate of additivity in this study compared to 15 previous studies involving S. pennellii introgression lines (Schauer et al., 2008; Steinhauser et 16 al., 2011) may result from the lower distance between the parental lines, which are all from 17 the same species. 18

The inheritance mode in one cross was not systematically the same in another cross and appeared relatively independent from one stage to the other, as a consequence of the complex genetic control of the traits studied. Enzyme activities for example are suggested to be controlled by a network of trans-acting genes (Steinhauser et al., 2011), thus dealing with different genetic backgrounds that carry different combinations of haplotypes, may lead to different inheritance modes.

26 Systems approach revealed complex connectivity among the different levels analyzed

We dissected the genetic variation at several levels, from phenotype to metabolite and 27 28 proteome profiles, in eight unrelated tomato accessions and four F1 hybrids at two developmental stages. In such experimental design, a significant correlation may reveal the 29 30 effect of a polymorphic gene acting on two related traits, but also fortuitous association, as a correlation between two traits may not be due to a causal relationship but to linkage 31 32 disequilibrium between genes controlling the variation of both traits. Nevertheless, Osorio et al. (2011) in a similar approach described co-varying genes or proteins as "guilty by 33 34 association", as closer the functions implicated more meaningful the relationships. Besides,

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

the two stages studied in the present study correspond to very distinct physiological processes 1

(Gillaspy et al., 1993; Giovannoni 2004), increasing the complexity but also the impact of the 2

3 study.

4

5

8

At every expression level, we detected more significant correlations than expected by chance. The bias towards positive correlations among enzyme activities and metabolites suggested a coordinated regulation of these traits. Some of the detected relationships were already 6 established in previous studies, as for instance the coordinated variation between several 7 amino acids and sugars (fructose and glucose) at CE (Prudent et al., 2010).

9 Different studies have tried to uncover the relationships between different levels of traits (phenotypes, metabolites, enzymes and transcripts) in tomato (Schauer et al., 2006; Carrari et 10 al., 2006). As the enzyme activities assessed correspond to Vmax, and thus mainly reflect the 11 corresponding protein amount, one might hypothesize that proteins and enzyme activities 12 13 should be correlated. We only found correlations between three out of eight enzymes and the protein spot amount corresponding to the same function at OR, all of them showing a positive 14 15 correlation, as expected from a causal relationship. This lack of relationship between enzyme activities and their protein amounts is consistent with the results obtained when comparing 16 enzyme activities and their corresponding gene expression (Gibon et al., 2006, Morcuende et 17 al., 2007; Steinhauser et al., 2010). This may be due to the fact that enzyme activities result 18 from a combination of several proteins (subunits) or that most of the primary metabolism 19 enzymes belong to multigene families. Besides, protein spots may also be the product of 20 complex post-translational modifications, where only one of the forms will be the functional 21 one (Faurobert et al., 2007). 22

One problem when dealing with omic data is that the number of traits is much larger than the 23 number of samples. Sparse methods were developed for dealing with high-dimensional data. 24 Such method sets the contribution of noise variables to zero and thus improve the prediction 25 of correlations or classification performance (Filzmoser et al., 2012). The networks 26 reconstructed with SPLs methods showed complex patterns of connectivity, relating several 27 nodes together and different pathways or metabolisms. In each network, a few hubs could be 28 identified relating many different compounds or proteins. 29

At CE, several correlations with dry matter content and metabolite contents involved two of 30 the protein spots coding for fructokinase, an enzyme participating in the sugar 31 phosphorylation. Fructokinase plays a role in sugar import and in starch biosynthesis (Dai et 32 al., 2002). Several isoforms were detected, being correlated with the variation of sugars. 33

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

1 The network constructed at OR revealed the key role of invertase in sucrose breakdown, as 2 already documented (Faurobert et al., 2007). Two spots corresponding to this function were 3 strongly related to the content in sucrose and dry matter content. Besides, they were also 4 correlated with several enzyme activities and with fruit weight. Schauer et al., (2006) have 5 also detected an association between phenotypic traits and metabolic compounds in tomato.

Previous studies on Arabidopsis suggested that the relations between transcript modifications 6 and enzyme activities showed greater agreement at long term (Osuna et al., 2007). We thus 7 analyzed the relationships between protein amounts at CE and the other traits at OR. We then 8 9 detected relationships between isocitrate dehydrogenase enzyme activity and the protein spots corresponding to this protein. Besides, the interconnections suggested a possible role of 10 regulator for cysteine synthase whose amount at CE is correlated with several enzymes, 11 glucose and fruit diameter at OR. This enzyme has been suggested to play a key role in highly 12 13 metabolically active cells (Wang et al., 2003). Those facts should be taken into account when trying to modify gene expression or protein contents in order to alter metabolite contents. 14 15 According to our finding this approach will just work for a small subset of metabolites, so researchers should focus on proteins like fructokinase, or cysteine synthase that affect several 16 17 metabolites instead of just the enzymes that regulate the direct synthesis of a target compound. 18

Our analysis provided a detailed characterization of fruit metabolism, at several levels in a set 19 of accessions representing a wide range of genetic variation and led to interesting conclusions. 20 First, the contents in primary and secondary metabolites are quite variable depending on the 21 genetic background, while enzyme activities seem to be less variable, particularly at CE. 22 Besides, significant genotype by stage interactions showed that the trends observed in one 23 genotype at a physiological level may change in another genotype. In agreement with this, the 24 inheritance modes varied between crosses and stages, showing the multigenic nature of the 25 traits studied, although additivity was predominant. The network reconstruction revealed 26 associations between different levels of expression and provided information on several key 27 proteins that might be targets for improving metabolite contents. This study is the starting 28 point of a broad experiment including the development of a multi-allelic population derived 29 30 from the eight parental lines. QTLs for fruit composition will be mapped in the population 31 and will be related to the variations observed at various levels in the parental lines.

Version définitive du manuscrit publiée dans / Final version of the manuscript published in : Journal of experimental botany (2013), DOI: 10.1093/jxb/ert349 Journal homepage: http://jxb.oxfordjournals.org/open_access.html

SUPPLEMENTARY DATA 1

- 2 Supplementary Table S1. Accessions origins.
- Supplementary Table S2. Polymorphism rate between the eight parental lines and 3
- Heinz1706 (the tomato reference genome) detected with 139 SNP (from Xu et al., 2013a) 4
- Supplementary Table S3. List of the 19 primary metabolites and 8 secondary metabolites* 5
- analyzed. 6
- Supplementary Table S4. List of the enzyme activities analyzed. 7
- 8 Supplementary Table S5. Means and standard deviations for all the traits studied
- 9 Supplementary Table S6. Annotation and analysis of variation of protein spots
- Supplementary Table S7. Analysis of variation cross by cross at each stage 10
- Supplementary Table S8. Overview of the number of significant correlations within and 11
- among different levels of analysis 12
- 13 Supplementary Table S9. Correlations at cell expansion stage.
- Supplementary Table S10. Correlations at orange-red stage. 14
- **Supplementary Table S11.** Correlations between enzyme activities and the spots coding for 15
- those enzymes at orange-red stage. 16
- Supplementary Table S12. Annotation of spots included in the cell expansion stage network 17
- (Supplementary Figure S5) 18
- Supplementary Table S13. Annotation of spots included in the orange-red stage network 19 (Supplementary Figure S6) 20
- Supplementary Table S14. Annotation of spots included in the cell expansion and orange-red 21 network (Figure 7) 22
- **Supplementary Figure S1.** Fruits of the eight tomato lines and four F1 hybrids studied. 23
- Supplementary Figure S2. Fruit fresh weight variation during fruit development in the eight 24 lines and four F1 hybrids. 25
- Supplementary Figure S3. First plan of the principal component analysis showing the 26 variation of 12 genotypes at two stages. 27
- 28 **Supplementary Figure S4.** Correlations between the volumes of protein spots corresponding to acid invertase and the enzyme activity. 29
- 30 Supplementary Figure S5. Network reconstruction based on sPLS between protein spot 31 volumes and metabolites, phenotypes and enzyme activities at cell expansion.
- 32 Supplementary Figure S6. Network reconstruction based on sPLS between protein spot volumes and metabolites, phenotypes and enzyme activities at orange-red. 33

1

2 ACKNOWLEDGMENT

- 3 We thank Esther Pelpoir for her help in fruit sampling and Caroline Callot and Karine Leyre
- 4 for technical help. We also thank the experimental team and Yolande Carretero for taking care
- 5 of the plants in the greenhouse. This work was supported by ANR Genomic project
- 6 MAGICTomSNP project. Jiaxin Xu was supported by China Scholarship Council.

REFERENCES

- Blanca J, Canizares J, Cordero L, Pascual L, Diez MJ, Nuez F. 2012. Variation Revealed by SNP Genotyping and Morphology Provides Insight into the Origin of the Tomato. PloS One 7(10), e48198.
- Carrari F, Baxter C, Usadel B, et al. 2006. Integrated analysis of metabolite and transcript levels reveals the metabolic shifts that underlie tomato fruit development and highlight regulatory aspects of metabolic network behavior. Plant Physiology 142(4), 1380-1396.
- Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. Bioinformatics 21(18), 3674-3676.
- **Deborde C, Maucourt M, Baldet P, et al.** 2009. Proton NMR quantitative profiling for quality assessment of greenhouse-grown tomato fruit. Metabolomics **52**, 183-198.
- Dai N, German MA, Matsevitz T, Hanael R, Swartzberg D, Yeselson Y, Petreikov M, Schaffer AA, Granot D. 2002. LeFRK2, the gene encoding the major fructokinase in tomato fruits, is not required for starch biosynthesis in developing fruits. Plant Science 162, 423-430.
- Faurobert M, Mihr C, Bertin N, Pawlowski T, Negroni L, Sommerer N, Causse M. 2007. Major proteome variations associated with cherry tomato pericarp development and ripening. Plant Physiology 143(3), 1327-1346.
- Faurobert M, Iijima Y, Aoki K. 2013. Proteomics and Metabolomics. In: Liedl B, Labate JA, Stommel JR, Slade A, Kole C, eds. *Genetics, genomics and breeding of tomato*. USA: CRC Press Taylor & Francis Group, 403-421.
- Ferry-Dumazet H, Gil L, Deborde C, Moing A, Bernillon S, Rolin D, Nikolski M, de Daruvar A, Jacob D. 2011. MeRy-B: a web knowledgebase for the storage, visualization, analysis and annotation of plant NMR metabolomic profiles. BMC Plant Biology 11, 104.
- Filzmoser P, Gschwandtner M, Todorov V. 2012. Review of sparse methods in regression and classification with application to chemometrics. Journal of Chemometrics 263(4), 42-51.
- Garcia V, Stevens R, Gil L, et al. 2009. An integrative genomics approach for deciphering the complex interactions between ascorbate metabolism and fruit growth and composition in tomato. Comptes Rendus Biologies 332(11), 1007-1021.

- Gibon Y, Blaesing OE, Hannemann J, Carillo P, Hohne M, Hendriks JHM, Palacios N, Cross J, Selbig J, Stitt M. 2004. A robot-based platform to measure multiple enzyme activities in Arabidopsis using a set of cycling assays: Comparison of changes of enzyme activities and transcript levels during diurnal cycles and in prolonged darkness. Plant Cell 16(12), 3304-3325.
- Gibon Y, Usadel B, Blaesing OE, Kamlage B, Hoehne M, Trethewey R, Stitt M. 2006. Integration of metabolite with transcript and enzyme activity profiling during diurnal cycles in Arabidopsis rosettes. Genome Biology 7(8), R76.
- Gillaspy G, Bendavid H, Gruissem W. 1993. Fruits A Developmental Perspective. Plant Cell 5(10), 1439-1451.
- Giovannoni JJ. 2004. Genetic regulation of fruit development and ripening. Plant Cell 16, S170-S180.
- Gomez-Romero M, Segura-Carretero A, Fernandez-Gutierrez A. 2010. Metabolite profiling and quantification of phenolic compounds in methanol extracts of tomato fruit. Phytochemistry **71(16)**, 1848-1864.
- Gutierez RA, Stokes TL, Thum K, *et al.* 2008. Systems approach identifies an organic nitrogen-responsive gene network that is regulated by the master clock control gene CCA1. Proceedings of the National Academy of Sciences of the United States of America 105(12), 4939-4944.
- Hirai MY, Klein M, Fujikawa Y, Yano M, Goodenowe DB, Yamazaki Y, Kanaya S, Nakamura Y, Kitayama M, Suzuki H. 2005. Elucidation of gene-to-gene and metabolite-to-gene networks in Arabidopsis by integration of metabolomics and transcriptomics. Journal of Biological Chemistry 280(27), 25590-25595.
- Ideker T, Galitski T, Hood L. 2001. A new approach to decoding life: Systems biology. Annual Review of Genomics and Human Genetics 2, 343-372.
- Keurentjes, JJB, Sulpice R, Gibon Y, Steinhauser MC, Fu J, Koornneef M, Stitt M, Vreugdenhil D. 2008. Integrative analyses of genetic variation in enzyme activities of primary carbohydrate metabolism reveal distinct modes of regulation in Arabidopsis thaliana. Genome Biology 9(8), R129.
- **Keurentjes JJB.** 2009. Genetical metabolomics: closing in on phenotypes. Current Opinion in Plant Biology **12(2)**, 223-230.

- Kliebenstein DJ. 2009a. Advancing Genetic Theory and Application by Metabolic Quantitative Trait Loci Analysis. Plant Cell **21(6)**, 1637-1646.
- Kliebenstein DJ. 2009b. Quantification of variation in expression networks. Methods in Molecular Biology 553, 227-245.
- Kliebenstein DJ. 2010. Systems Biology Uncovers the Foundation of Natural Genetic Diversity. Plant Physiology 152(2), 480-486.
- Le Cao KA, Gonzalez I, Dejean S. 2009. integrOmics: an R package to unravel relationships between two omics datasets. Bioinformatics 25(21), 2855-2856.
- Li L, Lu K, Chen Z, Mu T, Hu Z, Li X. 2008. Dominance, Overdominance and Epistasis Condition the Heterosis in Two Heterotic Rice Hybrids. Genetics **180(3)**, 1725-1742.
- Lippman ZB, Zamir D. 2007. Heterosis: revisiting the magic. Trends in Genetics 23, 60-66.
- Meyer RS, DuVal AE, Jensen HR. 2012. Patterns and processes in crop domestication: an historical review and quantitative analysis of 203 global food crops. New Phytologist 196(1), 29-48.
- Miller JC, Tanksley SD. 1990. RFLP analysis of phylogenetic-relationships and geneticvariation in the genus *Lycopersicon*. Theoretical and Applied Genetics **80(4)**, 437-448.
- Morcuende R, Bari R, Gibon Y, *et al.* 2007. Genome-wide reprogramming of metabolism and regulatory networks of Arabidopsis in response to phosphorus. Plant Cell and Environment **30(1)**, 85-112.
- Mounet F, Moing A, Garcia V, *et al.* 2009. Gene and Metabolite Regulatory Network Analysis of Early Developing Fruit Tissues Highlights New Candidate Genes for the Control of Tomato Fruit Composition and Development. Plant Physiology **149(3)**, 1505-1528.
- Nesbitt, TC, Tanksley SD. 2002. Comparative sequencing in the genus *Lycopersicon*: Implications for the evolution of fruit size in the domestication of cultivated tomatoes. Genetics 162(1), 365-379.
- **OsorioS, Alba R, Damasceno CMB, et al. 2011.** Systems Biology of Tomato Fruit Development: Combined Transcript, Protein, and Metabolite Analysis of Tomato Transcription Factor nor, rin and Ethylene Receptor Nr Mutants Reveals Novel Regulatory Interactions. Plant Physiology **157**(1), 405-425.

- **Osuna D, Usadel B, Morcuende R**, *et al*, 2007. Temporal responses of transcripts, enzyme activities and metabolites after adding sucrose to carbon-deprived Arabidopsis seedlings. Plant Journal 49, 463–491
- Prudent M, Bertin N, Genard M, Munos S, Rolland S, Garcia V, Petit J, Baldet P, Rothan C, Causse M. 2010. Genotype-dependent response to carbon availability in growing tomato fruit. Plant Cell and Environment 33(7), 1186-1204.
- Ranc N, Munos S, Santoni S, Causse M. 2008. A clarified position for Solanum lycopersicum var. cerasiforme in the evolutionary history of tomatoes (solanaceae). BMC Plant Biology 8, 130.
- Rockman MV, Kruglyak L. 2006. Genetics of global gene expression. Nature Reviews Genetics 711, 862-872.
- Rosenfeld N, Elowitz MB, Alon U. 2002. Negative autoregulation speeds the response times of transcription networks. Journal of Molecular Biology **323(5)**, 785-793.
- Saito K, Matsuda F. 2010 Metabolomics for Functional Genomics, Systems Biology, and Biotechnology. Annual Review of Plant Biology. 61, 463-489.
- Schauer N, Semel Y, Balbo I, Steinfath M, Repsilber D, Selbig J, Pleban T, Zamir D, Fernie AR. 2008. Mode of inheritance of primary metabolic traits in tomato. Plant Cell 20(3), 509-523.
- Schauer N, Semel Y, Roessner U, et al. 2006. Comprehensive metabolic profiling and phenotyping of interspecific introgression lines for tomato improvement. Nature Biotechnology 24(4), 447-454.
- Sim SC, Durstewitz G, Plieske J, S *et al.* 2012. Development of a Large SNP Genotyping Array and Generation of High-Density Genetic Maps in Tomato. PloS One 7(7), e40563.
- Springer NM, Stupar RM. 2007. Allele-specific expression patterns reveal biases and embryo-specific parent-of-origin effects in hybrid maize. Plant Cell **19(8)**, 2391-2402.
- Stacklies W, Redestig H, Scholz M, Walther D, Selbig J. 2007. pcaMethods a bioconductor package providing PCA methods for incomplete data. Bioinformatics 23(9), 1164-1167.
- Steinhauser MC, Steinhauser D, Gibon Y, Bolger M, Arrivault S, Usadel B, Zamir D, Fernie AR, Stitt M. 2011. Identification of Enzyme Activity Quantitative Trait Loci in

a Solanum lycopersicum X Solanum pennellii Introgression Line Population. Plant Physiology **157(3)**, 998-1014.

- Steinhauser MC, Steinhauser D, Koehl K, Carrari F, Gibon Y, Fernie AR, Stitt M. 2010. Enzyme Activity Profiles during Fruit Development in Tomato Cultivars and *Solanum pennellii*. Plant Physiology 153(1), 80-98.
- Stevens R, Buret M, Duffe P, Garchery C, Baldet P, Rothan C, Causse M. 2007. Candidate genes and quantitative trait loci affecting fruit ascorbic acid content in three tomato populations. Plant Physiology 143(4), 1943-1953.
- Stylianou IM, Affourtit JP, Shockley KR, Wilpan RY, Abdi FA, Bhardwaj S, Rollins J, Churchill GA, Paigen B. 2008. Applying gene expression, proteomics and singlenucleotide polymorphism analysis for complex trait gene identification. Genetics 178(3), 1795-1805.
- Stuber CW. 2010. Heterosis in plant breeding. Plant Breeding Reviews 12. Oxford, UK: John Wiley & Sons, Inc.
- **Tomato Genome Consortium.** 2012. The tomato genome sequence provides insights into fleshy fruit evolution. Nature **485(7400)**, 635-641.
- Wang Z, Potter RH, Jones MG. 2003. Differential display analysis of gene expression in the cytoplasm of giant cells induced in tomato roots by *Meloidogyne javanica*. Molecular Plant Pathology. 4(5), 361-71.
- Wang H, Schauer N, Usadel B, Frasse P, Zouine M, Hernould M, Latche A, Pech JC, Fernie AR, Bouzayen M. 2009. Regulatory Features Underlying Pollination-Dependent and -Independent Tomato Fruit Set Revealed by Transcript and Primary Metabolite Profiling. Plant Cell 21(5), 1428-1452.
- Xu J, Ranc N, Munos S, Rolland S, Bouchet JP, Desplat N, Le Paslier MC, Liang Y,
 Brunel D, Causse M. 2013a. Phenotypic diversity and association mapping for fruit quality traits in cultivated tomato and related species. Theoretical and Applied Genetics. 126(3), 567-581.
- Xu J, Pascual L, Aurant R, Bouchet JP, Valot B, Zivy M, Causse M, Faurobert M.
 2013b. An extensive proteome map of tomato (*Solanum lycopersicum*) fruit pericarp.
 Proteomics. Doi:10.1002/pmic201200438.

Table 1. Analysis of variation for phenotypic and metabolite contents in 8 tomato accessions and 4 F1 (see Supplementary Table S2 for metabolite

abbreviations).

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manuscript

| | G | lobal analy | sis | CE/ | /OR | | CE stage | | | | OR stage | | |
|--------|-----|-------------|------|-------|--------|-----|-----------|------------|---------|-----|-----------|-----------|---------|
| Traits | Fs | Fg | Fgxs | min | max | Fg | min | max | max/min | Fg | min | max | max/min |
| FW | *** | *** | *** | 0.10 | 0.41 | *** | 1.26 | 30.84 | 24.41 | *** | 5.33 | 134.36 | 25.22 |
| FD | *** | *** | *** | 0.46 | 0.74 | *** | 14.19 | 42.50 | 3.00 | *** | 22.73 | 69.95 | 3.08 |
| DMC | *** | *** | ns | 1.12 | 1.42 | *** | 6.16 | 10.59 | 1.72 | *** | 4.95 | 8.81 | 1.78 |
| Glc | *** | *** | *** | 0.42 | 0.90 | *** | 68996.30 | 193536.82 | 2.81 | *** | 158574.12 | 233208.35 | 1.47 |
| Suc | *** | *** | *** | 0.31 | 1.08 | *** | 6417.25 | 11856.59 | 1.85 | *** | 7596.82 | 28049.81 | 3.69 |
| Fru | *** | *** | *** | 0.38 | 0.85 | *** | 65853.14 | 198775.33 | 3.02 | *** | 173280.21 | 238532.61 | 1.38 |
| Ala | *** | *** | *** | 0.79 | 5.51 | *** | 288.06 | 1356.30 | 4.71 | *** | 171.01 | 492.55 | 2.88 |
| Asn | *** | *** | *** | 0.31 | 0.95 | *** | 752.84 | 2461.92 | 3.27 | *** | 1200.65 | 4593.94 | 3.83 |
| Asp | *** | *** | *** | 0.20 | 0.41 | *** | 579.58 | 1278.39 | 2.21 | *** | 1564.39 | 3598.28 | 2.30 |
| Abu | *** | *** | *** | 0.80 | 2.25 | *** | 4099.47 | 8485.63 | 2.07 | *** | 2130.82 | 5854.16 | 2.75 |
| Gln | *** | *** | *** | 0.48 | 1.45 | *** | 5976.22 | 26264.35 | 4.39 | *** | 6605.55 | 24155.80 | 3.66 |
| Ile | ** | *** | *** | 0.57 | 3.30 | *** | 168.70 | 865.48 | 5.13 | *** | 232.59 | 687.61 | 2.96 |
| Leu | *** | *** | *** | 0.40 | 1.26 | *** | 306.62 | 927.64 | 3.03 | *** | 401.47 | 1042.48 | 2.60 |
| Phe | *** | *** | *** | 0.35 | 0.94 | *** | 1209.44 | 4905.71 | 4.06 | *** | 1988.74 | 7429.73 | 3.74 |
| Tyr | *** | *** | *** | 0.50 | 2.20 | *** | 153.13 | 760.58 | 4.97 | *** | 185.86 | 656.84 | 3.53 |
| Val | *** | *** | *** | 1.15 | 10.95 | *** | 206.82 | 1004.13 | 4.86 | *** | 91.66 | 396.69 | 4.33 |
| Thr | *** | *** | *** | 0.38 | 2.12 | *** | 125.21 | 745.30 | 5.95 | *** | 221.38 | 768.59 | 3.47 |
| Asc | *** | *** | *** | 0.48 | 1.13 | *** | 1304.64 | 2236.94 | 1.71 | *** | 1509.86 | 3021.01 | 2.00 |
| Cit | *** | *** | *** | 0.37 | 0.57 | *** | 26295.26 | 62463.29 | 2.38 | *** | 49525.87 | 149942.47 | 3.03 |
| Mlt | ns | *** | *** | 0.59 | 3.78 | *** | 13562.91 | 25447.13 | 1.88 | *** | 3894.13 | 30910.20 | 7.94 |
| Fum | *** | *** | *** | 0.88 | NA | *** | 5.54 | 18.71 | 3.38 | *** | 0.00 | 12.75 | NA |
| Tom* | *** | *** | *** | 22.16 | 154.19 | *** | 983224.09 | 4556008.75 | 4.63 | *** | 16992.93 | 98910.75 | 5.82 |
| DHTom* | *** | *** | *** | 9.15 | 63.02 | *** | 196588.35 | 2644852.27 | 13.45 | *** | 7239.96 | 80840.49 | 11.17 |

| OH Lyc* | *** | *** | *** | 0.19 | 1.03 | *** | 4384.91 | 9953.68 | 2.27 | *** | 5766.39 | 46972.34 | 8.15 |
|---------|-----|-----|-----|------|------|-----|----------|-----------|------|-----|----------|-----------|--------|
| 5CQA* | *** | *** | *** | 0.91 | 1.62 | *** | 51444.47 | 111458.40 | 2.17 | ** | 43965.78 | 68607.80 | 1.56 |
| 3CQA* | ns | ns | ns | 0.45 | 2.33 | ns | 4404.24 | 19467.52 | 4.42 | ns | 5977.49 | 13005.07 | 2.18 |
| Nch* | *** | *** | *** | 0.00 | 0.30 | * | 408.94 | 2137.62 | 5.23 | *** | 2788.25 | 684629.20 | 245.54 |
| Rut* | *** | *** | *** | 0.63 | 4.26 | *** | 63505.59 | 243368.28 | 3.83 | *** | 28927.20 | 265669.06 | 9.18 |
| RutP* | *** | *** | *** | 0.40 | 1.09 | *** | 21497.91 | 59076.96 | 2.75 | *** | 25248.73 | 79202.85 | 3.14 |

Fs: significance level of the ANOVA for the stage factor

Fg: significance level of the ANOVA for genotype factor

Fgxs: significance level of the ANOVA for the interaction between genotype and stage

Min: minimum average values for each variable among the 12 genotypes

Max: maximum average values for each variable among the 12 genotypes

CE/ OR ratio value to cell expansion stages and orange-red (min and max) for each genotype

*: 0.01 <P <0.05. **: 0.001 <P <0.01. ***: P<0.001. ns: P>0.05

NA: non available

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manu

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manuscript

Table 2. Analysis of variation for enzyme activities in 8 tomato accessions and 4 F1 (see Supplementary Table S3 for enzyme abbreviations) 1

| | | Global | | | | | | | | | | | |
|--------------|-----|----------|------|------|-------|----------------------|-----------|-----------|---------|-----|----------|-------------|---------|
| | | analysis | | CE/ | OR | Cell expansion stage | | | | | Orang | e-red stage | |
| Enzymes | Fs | Fg | Fgxs | min | max | Fg | min | max | max/min | Fg | min | max | max/min |
| PEPC | *** | ns | ns | 1.63 | 5.01 | ns | 4743.07 | 10166.94 | 2.14 | ns | 1393.72 | 3988.74 | 2.86 |
| ALS | *** | * | ns | 1.11 | 6.96 | ns | 23342.95 | 39076.09 | 1.67 | * | 4955.98 | 27286.39 | 5.51 |
| G6PDH | *** | ns | ns | 1.41 | 3.22 | ns | 1948.75 | 2991.71 | 1.54 | ** | 736.48 | 1494.57 | 2.03 |
| PGM | *** | ns | ns | 1.56 | 3.58 | ns | 18767.93 | 31388.58 | 1.67 | ns | 6272.80 | 14070.21 | 2.24 |
| РК | *** | ns | ns | 0.98 | 3.04 | ns | 4214.45 | 7365.46 | 1.75 | * | 1736.57 | 4838.01 | 2.79 |
| Ppi-PFK | *** | ns | ns | 1.42 | 5.97 | ns | 7380.06 | 13238.75 | 1.79 | *** | 1778.74 | 6372.06 | 3.58 |
| ACO | *** | ns | ns | 0.85 | 11.98 | ns | 1115.56 | 3838.66 | 3.44 | ** | 125.70 | 1940.54 | 15.44 |
| ATP-PFK | *** | ns | ns | 0.71 | 2.05 | ns | 922.01 | 1640.44 | 1.78 | ns | 709.66 | 1295.85 | 1.83 |
| FRK | *** | * | ** | 1.29 | 8.99 | * | 1027.06 | 5020.16 | 4.89 | ns | 309.56 | 1373.80 | 4.44 |
| InvN | *** | *** | ** | 0.12 | 1.60 | ns | 881.42 | 4517.10 | 5.12 | *** | 1798.65 | 13163.36 | 7.32 |
| InvA | *** | ns | ns | 0.12 | 1.03 | ns | 1769.30 | 5805.68 | 3.28 | * | 3542.76 | 22718.43 | 6.41 |
| NAD-MDH | *** | ns | ns | 1.57 | 4.69 | ns | 212499.39 | 372186.20 | 1.75 | ** | 65345.21 | 186895.32 | 2.86 |
| AlaAT | ns | ns | ns | 0.30 | 2.83 | ns | 8355.30 | 27841.06 | 3.33 | ns | 3908.25 | 60730.09 | 15.54 |
| FH | ns | ns | ns | 0.59 | 10.56 | ns | 2227.13 | 7340.03 | 3.30 | ns | 266.42 | 5426.77 | 20.37 |
| AspAT | *** | ns | ns | 0.63 | 2.75 | ns | 31619.55 | 88013.71 | 2.78 | ns | 23952.50 | 55452.74 | 2.32 |
| NAD-ME | ** | ns | ns | 0.89 | 3.06 | ns | 4699.19 | 10772.77 | 2.29 | ns | 3248.43 | 9464.11 | 2.91 |
| NADP-ME | *** | *** | ** | 0.59 | 4.00 | ** | 1585.95 | 5869.58 | 3.70 | ** | 960.60 | 3947.79 | 4.11 |
| GAPDH (NAD) | *** | ns | ns | 1.33 | 5.27 | ns | 22716.25 | 37476.27 | 1.65 | ns | 7059.38 | 24964.31 | 3.54 |
| GAPDH (NADP) | ns | ns | ns | 0.43 | 7.19 | ns | 4232.25 | 11154.22 | 2.64 | * | 966.42 | 22461.89 | 23.24 |
| GK | *** | ns | ns | 2.57 | 7.59 | ns | 979.49 | 1914.96 | 1.96 | ns | 166.27 | 443.60 | 2.67 |
| IcDH | *** | ** | *** | 0.39 | 1.54 | ns | 1210.69 | 3118.82 | 2.58 | *** | 1482.13 | 4029.86 | 2.72 |
| ENO | *** | ns | * | 1.79 | 7.08 | ns | 3420.96 | 6643.23 | 1.94 | ** | 773.15 | 2156.06 | 2.79 |

| TPI | *** | ns | ns | 1.64 | 6.26 | ns | 627079.82 | 1212604.25 | 1.93 | ns | 187195.55 | 505691.43 | 2.70 |
|-------|-----|----|----|------|------|----|-----------|------------|------|-----|-----------|-----------|------|
| PGK | *** | ns | ns | 0.72 | 7.58 | ns | 52497.52 | 105333.98 | 2.01 | ns | 13903.37 | 73080.01 | 5.26 |
| SuSy | ** | ns | ns | 0.66 | 4.78 | ns | 1857.82 | 11726.22 | 6.31 | ns | 1520.16 | 7106.26 | 4.67 |
| ShKDH | *** | ** | ns | 1.07 | 2.57 | ns | 1568.76 | 2787.32 | 1.78 | *** | 725.68 | 2212.66 | 3.05 |

Fg: significance level of the ANOVA for genotype factor

Fs: significance level of the ANOVA for the stage factor

Fgxs: significance level of the ANOVA for the interaction between genotype and stage

Min: minimum average values for each variable among the 12 genotypes

Max: maximum average values for each variable among the 12 genotypes

CE/ OR ratio value to cell expansion stages and orange-red (min and max) for each genotype

*: 0.01 <P <0.05. **: 0.001 <P <0.01. ***: P<0.001. ns: P>0.05

1

Author manus

Manuscrit d'auteur / 1

- 1 Table 3. Spots highly correlated with fruit weight (a) and dry matter content (b) in 8 tomato accessions and 4 F1 hybrids. Pearson correlations are indicated
- 2 with spot volumes assessed at cell expression (rCE) or orange-red (rOR) stage. Range of variation of the spots at the same stage (min and max) in the 12
- 3 genotypes

Manuscrit d'auteur / Author manu

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manuscript

4 Only highly significantly correlated spots (p-value<0.001 and |r|>0.78) are indicated

| | spot | gene | annotation | Process | | | |
|-------------|-----------|--------------------|--|--------------------|-------|--------|--------|
| a. Spots co | orrelated | with Fruit weight | | | | | |
| Cell expan | nsion | | | | rCE | CE.min | CE.max |
| | JX031 | Solyc08g076970.2.1 | Acetylornithine deacetylase | macromolecule | -0.82 | 0.31 | 0.55 |
| | JX074 | Solyc08g082820.2.1 | Heat shock protein | stress response | -0.88 | 0.75 | 1.21 |
| | JX141 | Solyc03g082920.2.1 | Heat shock protein | stress response | -0.83 | 0.46 | 1.00 |
| | JX152 | Solyc06g075010.2.1 | chaperonin | macromolecule | -0.81 | 0.57 | 0.87 |
| | JX289 | Solyc01g079220.2.1 | NifU like protein | stress response | 0.89 | 4.82 | 14.57 |
| | JX390 | Solyc05g053470.2.1 | chaperonin | macromolecule | -0.81 | 1.56 | 2.16 |
| | | | | | | | |
| Orange-re | d | | | | rOR | OR.min | OR.max |
| | JX051 | Solyc03g083910.2.1 | Acid beta-fructofuranosidase | primary metabolism | -0.84 | 0.24 | 0.71 |
| | JX059 | Solyc06g083790.2.1 | Succinyl-CoA ligase | primary metabolism | 0.84 | 1.66 | 4.11 |
| | JX135 | Solyc01g057000.2.1 | Universal stress protein family protein | stress response | -0.88 | 2.00 | 3.85 |
| | JX149 | Solyc08g082820.2.1 | Heat shock protein | stress response | -0.83 | 0.26 | 0.56 |
| | JX157 | Solyc08g082430.2.1 | Nucleoside diphosphate kinase | primary metabolism | -0.80 | 0.58 | 1.18 |
| | JX164 | Solyc01g106430.2.1 | Inorganic pyrophosphatase family protein | primary metabolism | -0.80 | 1.12 | 1.69 |
| | JX188 | Solyc05g008460.2.1 | ATP synthase subunit beta | regulation | -0.87 | 2.43 | 3.41 |
| | JX317 | Solyc01g104170.2.1 | Ankyrin repeat domain-containing protein 2 | regulation | 0.95 | 0.41 | 0.76 |
| | JX367 | Solyc08g076220.2.1 | Phosphoribulokinase/uridine kinase | primary metabolism | 0.84 | 0.59 | 1.15 |
| | | | | | | | |

b. Spots correlated with dry matter content

| Cell expansion | | | rCE | CE.min | CE.max |
|--------------------------|-----------------------------|---------------|------|--------|--------|
| JX012 Solyc06g005940.2.1 | Protein disulfide isomerase | macromolecule | 0.86 | 0.25 | 0.53 |

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manuscript

Manuscrit d'auteur / Author manuscript

| JX028 | Solyc02g080420.2.1 | RNA Binding Protein 45 | regulation | -0.83 | 0.49 | 0.96 |
|-------|--------------------|---|--------------------|-------|------|-------|
| JX040 | Solyc09g089580.2.1 | 1-aminocyclopropane-1-carboxylate oxidase | maturation | 0.83 | 0.35 | 1.32 |
| JX056 | Solyc01g099190.2.1 | Lipoxygenase | primary metabolism | 0.87 | 0.31 | 0.73 |
| JX120 | Solyc06g005940.2.1 | Protein disulfide isomerase | macromolecule | 0.83 | 0.95 | 1.66 |
| JX137 | Solyc06g005940.2.1 | Protein disulfide isomerase | macromolecule | 0.92 | 0.18 | 0.40 |
| JX154 | Solyc03g115990.1.1 | Malate dehydrogenase | primary metabolism | 0.84 | 1.41 | 2.36 |
| JX218 | Solyc06g073190.2.1 | Fructokinase-like | primary metabolism | 0.81 | 3.46 | 5.84 |
| JX219 | Solyc02g082800.2.1 | Ubiquilin-1 | macromolecule | 0.79 | 0.76 | 1.12 |
| JX311 | Solyc12g044740.1.1 | Ubiquitin carboxyl-terminal hydrolase | primary metabolism | 0.81 | 0.40 | 1.02 |
| JX325 | Solyc01g011000.2.1 | Eukaryotic translation initiation factor 5A | translation | 0.86 | 6.29 | 10.83 |
| JX359 | Solyc04g011400.2.1 | UDP-glucose 4-epimerase | cell wall | 0.79 | 1.72 | 2.44 |
| JX363 | Solyc02g091490.2.1 | Fructokinase 3 | primary metabolism | -0.84 | 0.44 | 1.13 |
| JX377 | Solyc09g090980.2.1 | Major allergen Mal d 1 | stress response | 0.82 | 3.37 | 6.87 |
| JX395 | Solyc02g086730.1.1 | 50S ribosomal protein L12-C | translation | 0.81 | 2.70 | 4.32 |
| | | | | | | |

| Orange-red stage | | | | rOR | OR.min | OR.max |
|------------------|--------------------|--|---------------------|-------|--------|--------|
| JX032 | Solyc01g094200.2.1 | NAD-dependent malic enzyme 2 | primary metabolism | 0.89 | 0.62 | 0.97 |
| JX035 | Solyc02g078540.2.1 | Unknown Protein | Unknown | 0.91 | 0.41 | 1.43 |
| JX051 | Solyc03g083910.2.1 | Acid beta-fructofuranosidase | primary metabolism | 0.81 | 0.24 | 0.71 |
| JX085 | Solyc09g015000.2.1 | class I heat shock protein | stress response | 0.89 | 0.25 | 4.38 |
| JX091 | Solyc05g050120.2.1 | Malic enzyme | primary metabolism | 0.93 | 0.10 | 0.70 |
| JX100 | Solyc10g083650.1.1 | Peroxiredoxin ahpC/TSA family | oxidation-reduction | 0.85 | 2.00 | 4.16 |
| JX103 | Solyc08g075210.1.1 | Acyltransferase-like protein | regulation | 0.83 | 0.21 | 0.55 |
| JX112 | Solyc06g009020.2.1 | Glutathione S-transferase | stress response | 0.79 | 0.95 | 1.89 |
| JX127 | Solyc10g084050.1.1 | 26S protease regulatory subunit 6B homolog | macromolecule | 0.81 | 0.85 | 1.29 |
| JX139 | Solyc05g056230.2.1 | Calreticulin 2 calcium-binding protein | macromolecule | 0.79 | 0.29 | 0.70 |
| JX160 | Solyc12g010040.1.1 | Leucyl aminopeptidase | macromolecule | 0.79 | 0.32 | 0.68 |
| JX214 | Solyc09g009020.2.1 | Enolase | primary metabolism | 0.82 | 2.24 | 3.83 |
| JX223 | Solyc01g106320.2.1 | Octicosapeptide/Phox/Bem1p domain protein | Unknown | -0.91 | 0.19 | 0.49 |
| JX239 | Solyc03g083910.2.1 | Acid beta-fructofuranosidase | primary metabolism | 0.87 | 1.32 | 10.71 |

| JX262 Solyc06g068860.2.1 | Alpha-mannosidase | primary metabolism | -0.84 | 0.33 | 0.95 |
|--------------------------|--|--------------------|-------|------|------|
| JX311 Solyc12g044740.1.1 | Ubiquitin carboxyl-terminal hydrolase | primary metabolism | 0.85 | 0.37 | 0.88 |
| JX320 Solyc12g009060.1.1 | Charged multivesicular body protein 2a | localization | 0.82 | 0.33 | 0.65 |

FIGURE LEGENDS

Figure 1. Fruit size (FD) and dry matter content (DMC) of the 8 tomato accessions studied and their four hybrids (measured at Orange-Red stage)

Figure 2. Assignment of the metabolites and enzymes studied to pathways. A total of 26 metabolites are indicated in continuous line squares, 26 enzymes are highlighted in dot squares.

Figure 3. First plans of the principal component analysis showing the variation of 12 genotypes based on (A) metabolite contents, enzyme activities and phenotypic traits at cell expansion stage; (B) proteins at cell expansion; (C) metabolite contents, enzyme activities and phenotypic traits at OR and (D) proteins at orange-red stage. Values along the axes indicate the percentage of total variation accounted for each component. Genotypes are indicated with different symbols, *S. lycopersicum* squares, *S. lycopersicum var cerasiforme* circles and F1 triangles. Levovil x Cervil (1), Stupicke Polni Rane x Criollo (2), LA0147 x Plovdiv 24A (3), Ferum x LA1420 (4).

Figure 4. Inheritance mode of the two groups of traits in the four crosses: (A) metabolite contents, enzyme activities and phenotypic traits; (B) protein spot volumes. From top to bottom over-recessive, recessive, additive, dominant and over-dominant. Left panels: cell expansion stage, right panels: orange-red stage.

Figure 5. Inheritance of citrate content in tomato fruit at (A) cell expansion stage, (B) orange-red stage. Genotypes are indicated with different symbols, *S. lycopersicum* squares, *S. lycopersicum* var *cerasiforme* circle and F1 triangle. From left to right Levovil x Cervil, Stupicke Polni Rane x Criollo, LA0147 x Plovdiv 24A, Ferum x LA1420.

Figure 6. Correlations among significantly variable phenotypic and metabolic traits, and enzyme activities at (A) cell expansion stage and (B) orange-red stage. Only correlations where $|\mathbf{r}|$ >0.7 (p-value<0.01) are shown. Red or dark gray indicates positive correlations, green or light gray negative correlations.

Figure 7. Network reconstruction based on sPLS between protein spot volumes at cell expansion stage (circular nodes) and metabolite contents, phenotypes and enzyme activities at orange-red stage (square nodes). Positive and negative relations are shown in continuous and dot lines. Annotation of spots is detailed in Supplemental Table S14.

Figure 1.



ompos

Figure 2.



Figure 3.



Figure 4.



Manuscrit d'auteur / Author manuscript

D cor





Figure 6.



Figure 7.



mposit