



**HAL**  
open science

## Identification and validation of reference genes for normalization of transcripts levels in roots of potatoes infected by nematodes

Patricio Castro Quezada, Jawad Aarouf, Michel Claverie, Eric Grenier, Marie-Claire Kerlan, Bruno Favery, Véronique Lefebvre, Bernard Caromel

### ► To cite this version:

Patricio Castro Quezada, Jawad Aarouf, Michel Claverie, Eric Grenier, Marie-Claire Kerlan, et al.. Identification and validation of reference genes for normalization of transcripts levels in roots of potatoes infected by nematodes. 5. Congrès de la Société Française de Phytopathologie, Jun 2012, Paris, France. 2012. hal-01208737

**HAL Id: hal-01208737**

**<https://hal.science/hal-01208737>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## SFP2012/163

### Identification and validation of reference genes for normalization of transcripts levels in roots of potatoes infected by nematodes.

P. Castro-Quezada<sup>a</sup>, J. Aarrouf<sup>b</sup>, M. Claverie<sup>a</sup>, E. Grenier<sup>c</sup>, M.-C. Kerlan<sup>c</sup>, B. Favery<sup>d</sup>, V. Lefebvre<sup>a</sup> et B. Caromel<sup>a</sup>

<sup>a</sup>INRA-UR1052, GAFL, BP 94, Allée des Chênes, 84143 Montfavet, France

<sup>b</sup>UAPV Avignon, Labo EA 4279, Pôle Agrosociétés, BP 21239, 301 Rue Baruch de Spinoza, 84916 Avignon, France

<sup>c</sup>INRA UMR IGEPP, Domaine de la motte, 35653 Le Rheu, France

<sup>d</sup>INRA, Institut Sophia-Agrobiotech, 400 route des Chappes, 06903 Sophia Antipolis, France

patricio.castro@avignon.inra.fr

Potato cyst nematodes induce changes in gene expression after infecting the roots. For studying gene expression, reverse transcription quantitative real-time PCR (RT-qPCR) is now a standard method. However, for accurate quantification of transcripts, a set of internal control genes, stable under different experimental conditions, is necessary for transcript normalization. Very few experimental data on suitable reference genes are available for plant-nematode interactions. In this study, we tested eight potential candidate reference genes identified through a microarray study performed on the pathosystem *Arabidopsis/Meloigodygne* for normalizing levels of potato gene transcripts, after invasion by nematodes. The transcription profiles of these genes were assessed by RT-qPCR. Samples used in these experiments were RNAs, issued from roots of infected and uninfected potato plants, carrying resistant or susceptible alleles against *Globodera pallida*, following a kinetic of four days. Analyses were performed using the geNorm and NormFinder softwares. Our results showed that the eight candidate reference genes were stably expressed in our conditions. For accurate normalization in RT-qPCR studies, RPN7 (26S proteasome regulatory subunit, involved in protein synthesis) and MST2 (mercaptopyruvate sulfurtransferase) were the top ranked genes and, therefore, are reliable reference genes in compatible and incompatible potato-nematode interactions.

Nombre de mots du résumé: 193

Mots-clé: gène de référence - nématode - résistance - pomme de terre

Thème: Réponse de la plante aux bioagresseurs et aux organismes phytobénéfiques

Session spéciale: Non spécifié(e)

Présentation: Présentation poster de préférence

Équipement particulier: Pas d'équipement particulier

Comment citer ce document :

Castro Quezada, P., Aarrouf, J., Claverie, M., Grenier, E., Kerlan, M.-C., Favery, B., Lefebvre, V., Caromel, B. (2012). Identification and validation of reference genes for normalization of transcripts levels in roots of potatoes infected by nematodes. Presented at 5. Congrès de la Société Française de Phytopathologie, Paris, FRA (2012-06-05 - 2012-06-08).