



## **Spatial prediction of potentiel wetlands at the French national scale based on hydroecoregions stratification and inference modelling**

Lionel Berthier, Lubomira Guzmova, Bertrand Laroche, Sébastien Lehmann, Hervé Squidadant, Manuel Martin, Jean-Philippe Chenu, Emmanuel Thiry, Blandine Lemerancier, Marion Bardy, et al.

### **► To cite this version:**

Lionel Berthier, Lubomira Guzmova, Bertrand Laroche, Sébastien Lehmann, Hervé Squidadant, et al.. Spatial prediction of potentiel wetlands at the French national scale based on hydroecoregions stratification and inference modelling. EGU General Assembly 2014, Apr 2014, Vienne, Austria. 2014, Geophysical Research Abstracts, vol.16. hal-01209219

**HAL Id: hal-01209219**

**<https://hal.science/hal-01209219>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## **Spatial prediction of potential wetlands at the French national scale based on hydroecoregions stratification and inference modelling.**

Lionel Berthier (1), Lubomira Guzmova (2), Bertrand Laroche (2), Sébastien Lehmann (2), Hervé Squvident (1), Manuel Martin (2), Jean-Philippe Chenu (2), Emmanuel Thiry (4), Blandine Lemerrier (1), Marion Bardy (2), Philippe Mérot (3), and Christian Walter (1)

(1) AGROCAMPUS OUEST, UMR 1069, Sol Agro et hydrosystème Spatialisation, F-35000 Rennes, France, (2) INRA, US1106, InfoSol, F-45075 Orléans, France, (3) INRA, UMR 1069, Sol Agro et hydrosystème Spatialisation, F-35000 Rennes, France, (4) MEDDE, Direction de l'eau et de la biodiversité, sous-direction des espaces naturels.

The prediction of wetlands at the national scale is crucial to design and implement consistent national policies aiming at preserving these fragile but essential ecosystems involved in the production of many fundamental ecosystem services. Depending on the topographic situation, the occurrence of potential wetlands is mostly driven by geomorphology and rainfall and the vertical distance to the channel network. This study aims to test for France the nationwide implementation of a method based on a topo-climatic index in combination with the vertical distance to the channel network to predict the spatial distribution of potential wetlands.

Following the PEEW approach used in this study (based on potential, existing, and efficient wetland functional assessment), potential wetlands include all geographical situations where geomorphological and/or climatic criteria may entail high probability of wetland occurrence. Potential wetlands include therefore former wetlands areas destructured for instance by artificial drainage, urbanization or river straightening. Topo-climatic index and the vertical distance to the channel network were calculated nationwide. To determine the threshold of these indices, France was stratified into hydroecoregions, based on geology, climate, topography, vegetation and channel network density. Within each hydroecoregion, cumulative frequencies of the topo-climatic index were calculated and a threshold was determined according to pedological data (percentage of hydromorphic soils) available in the area and using a probabilistic approach, whereas the threshold of the vertical distance to the channel network was computed alongside the potential flood zone. Both were combined according to a decision tree based on permeability of the soil surface and topographic situations in order to get a national prediction of the location of potential wetlands. Results were validated against point-specific soil data and detailed soil maps.

Combination of climato-topographic index and the vertical distance to the channel network was better at predicting the nationwide spatial distribution of wetlands with a highly improved accuracy by comparison with monofactorial modelling. The first validation test using point-specific soil data were satisfactory, the overall accuracy of validation equaled 67.8%, the accuracy for potential wetlands equaled 54.6% and the Kappa index amounted to 0.17. The accuracy seems to be poor for areas where few soil data were available to determine the threshold, thus stressing the need for further acquisition of soil data. The potential of this strategy to predict wetlands over large areas was therefore confirmed.