



HAL
open science

Amino acid incorporation into pig feeds reduces the environmental impacts of pig production

Florence Garcia-Launay, Hayo van Der Werf, Thi Tuyet Hanh Nguyen, Loïc Le Tutour, Jean-Yves Dourmad

► **To cite this version:**

Florence Garcia-Launay, Hayo van Der Werf, Thi Tuyet Hanh Nguyen, Loïc Le Tutour, Jean-Yves Dourmad. Amino acid incorporation into pig feeds reduces the environmental impacts of pig production. Orffa Symposium Danemark, Jan 2015, Billund, Denmark. hal-01209279

HAL Id: hal-01209279

<https://hal.science/hal-01209279>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Amino acid incorporation into pig feeds reduces the environmental impacts of pig production




AJINOMOTO.

AGRO
CAMPUS
OUEST


AJINOMOTO ANIMAL NUTRITION
AJINOMOTO EUROLYSINE S.A.S.

*Florence GARCIA-LAUNAY (1,2), Hayo VAN DER WERF (2,3),
Thi Tuyet Hanh NGUYEN (2,3), Loïc LE TUTOUR (4)
and Jean-Yves DOURMAD (1,2)*

- (1) INRA, UMR PEGASE, 35590 Saint-Gilles, France*
- (2) Agrocampus Ouest, F-35000 Rennes, France*
- (3) INRA, UMR1069 SAS, 35000 Rennes, France*
- (4) Ajinomoto Eurolysine SAS, 75817 Paris, France*





Nutrients flows on a pig farm



Potential environmental impacts

- Negative impacts
 - On atmosphere
 - Green House Gases (N_2O , CH_4) => Global warming
 - Ammonia (NH_3) => Acidification – Fine particles
 - On water
 - Nitrates => Eutrophication
 - Phosphates => Eutrophication
 - on soil
 - Accumulation of trace elements => Terrestrial ecotoxicity
 - Utilisation of ressources => Energy, phosphates, surface...
 - ...
- Positive impacts
 - Recycling of nutrients - valorisation of co-products
 - Fertility of soils- Organic matter
 - Biodiversity – landscapes ?

↘ protein content of feeds reduces nitrogen excretion and the associated emissions ...



...without lowering the performances

if the contents in essential amino acids are maintained

Reducing feed protein content

Feed with reduced protein content
(with same Net Energy and Essential AA contents)

Soybean meal ↘

Mosnier et al. 2011
Bourdon et al. 1995
Portejoie et al. 2004
Quiniou et al. 2011

Intake \approx

ADG \approx

FCR \approx

Bourdon et al. 1995
Portejoie et al. 2004
Quiniou et al. 2011

N excreted ↘

N₂O, NH₃ ↘

Bourdon et al. 1995
Portejoie et al. 2004
Osada et al. 2011

Objective : evaluate the potential for mitigation of environmental impacts when incorporating feed-use amino acids into pig feeds


Objectives and Layout

- Study 1:
Evaluate the environmental impacts of the pig production for different strategies of incorporation of feed-use amino acids
- Study 2:
Evaluate the potential for mitigation of GHG emissions from 2010 to 2030 in France for promising strategies



Evaluation of the environmental implications of the incorporation of feed-use amino acids in pig production using Life Cycle Assessment

F. Garcia-Launay^{a,c,*}, H.M.G. van der Werf^{b,c}, T.T.H. Nguyen^{b,c},
L. Le Tutour^d, J.Y. Doumad^{a,c}



QUELLE CONTRIBUTION DE L'AGRICULTURE FRANÇAISE À LA RÉDUCTION DES ÉMISSIONS DE GAZ À EFFET DE SERRE ?

POTENTIAL D'ATTÉNUATION ET CÔÛT DE DIX ACTIONS TECHNIQUES

Document de travail de l'Institut National de la Recherche Agronomique (INRA)

<http://institut.inra.fr/en/Missions/Inform-public-decision-making/Advanced-Studies/All-the-news/Study-on-reduction-of-GHG-in-agriculture>

Livestock Science 161 (2014) 158–175



Contents lists available at ScienceDirect

Livestock Science

journal homepage: www.elsevier.com/locate/livsci




Evaluation of the environmental implications of the incorporation of feed-use amino acids in pig production using Life Cycle Assessment



F. Garcia-Launay^{a,c,*}, H.M.G. van der Werf^{b,c}, T.T.H. Nguyen^{b,c},
L. Le Tutour^d, J.Y. Dourmad^{a,c}

The stages of Life Cycle Analysis




Iso 14002 Normalization


According to Geier, 1999

Aggregation of emissions per category of impacts


(from Owens, 1999 and Guinée et al. 2002)



Description of the system




Flow diagram for the production of pig



**Functional unit:
Kg of live pig at the
farm gate**

Description of formulation scenarios


- Farrow-to-finish farm / average performance for Brittany
- Sows and weaners : two feed per stage
- Fattening : single feed (1P) vs. two-phase (2P) vs. Multiphase (MP)



Two types of manure management : slurry (SL) et straw bedding (LI)
 Protein sources : soybean (SOJ) or soybean-rapeseed-peas (SOJRP)

Formulation of diets

- Digestible amino-acid requirements , and others nutrients (InraPorc, 2006, Ajinomoto Eurolysine, 2011)
- Fixed Net Energy contents
- **Formulation to achieve the same performance**
- Optimisation of diets using « OpenSolver for Excel » (Mason et Dunning, 2010)



Minimum nutritional contents for feed formulation

	Sow		Weaned Piglet		Fattening pig				
	Gestating	Lactating	Pre-starter	Starter	1P	Growing 2P	Finishing 2P	Growing MP	Finishing MP
Dry matter (g/kg)	860	860	860	860	860	860	860	860	860
Energy (MJ/kg)	9.0	9.5	10.0	10.0	9.6	9.6	9.6	9.6	9.6
SID AA									
Lysine (g/kg)	4.95	8.30	11.90	10.00	8.65	8.65	7.21	8.80	5.60
Threonine (% Lys)	72%	66%	65%	65%	67%	67%	68%	67%	68%
Methionine (% Lys)	28%	30%	30%	30%	30%	30%	30%	30%	30%
Met + Cystine (% Lys)	65%	60%	60%	60%	60%	60%	60%	60%	60%
Tryptophan (% Lys)	20%	20%	22%	22%	20%	20%	19%	20%	19%
Isoleucine (% Lys)	65%	60%	53%	53%	53%	53%	53%	53%	53%
Valine (% Lys)	75%	85%	70%	70%	65%	65%	65%	65%	65%
Leucine (% Lys)	100%	115%	100%	100%	100%	100%	100%	100%	100%
Phenylalanine (% Lys)	60%	60%	50%	50%	50%	50%	50%	50%	50%
Phe + Tyrosine (% Lys)	100%	115%	95%	95%	95%	95%	95%	95%	95%
Histidine (% Lys)	30%	42%	32%	32%	32%	32%	32%	32%	32%
Arginine (% Lys)	-	67%	42%	42%	42%	42%	42%	41%	42%
Minerals (g/kg)									
Digestible P	2.50	3.30	5.00	3.80	2.50	2.50	2.00	2.50	2.00
Ca ¹	9.00	10.50	13.50	11.00	7.25	7.25	7.25	7.25	7.25


Main inputs used, yield and nitrate-N emitted for the main crops used as feed ingredients

		Maize Brittany	Wheat France	Barley France	Pea France	Rapeseed France	Soybean Brazil C.West	Soybean Brazil South
N mineral fertiliser	kg/ha	32	165	130	0	165	9	0
N manure	kg/ha	210	10	10	0	16	0	5
P ₂ O ₅	kg/ha	29	26	37	44	50	90	58
K ₂ O	kg/ha	0	24	34	69	50	90	58
CaO	kg/ha	167	167	167	167	167	450	250
Seed	kg/ha	20	140	125	240	3	55	50
Pesticide	kg/ha	1.05	1.97	8.96	5.67	1.13	2.50	2.10
Diesel	kg/ha	85	83	84	89	92	80	67
Agricultural machinery	kg/ha	21.7	18.6	18.7	22.0	20.4	19.5	15.1
Irrigation water	m ³ /ha	0	0	0	88	0	0	0
Moisture at harvest	%	32	15	15	15	8	14	14
Reference moisture	%	15	15	15	15	8	13	13
Yield (dry matter)	kg/ha	6860	6010	5520	3550	3040	2400	2180
Nitrate-N emitted	kg/ha	40	40	40	70	40	16	21

Potential environmental impacts due to the production and delivery of feed ingredients


Feed ingredients	Climate change	Eutrophication	Acidification	Terrestrial ecotoxicity	Cumulative energy	Land occupation
	(kg CO ₂ -eq.)	(g PO ₄ -eq.)	(g SO ₂ -eq.)	(g 1.4-DCB-eq.)	(MJ)	(m ² year)
Wheat France	0.54	4.0	4.4	1.6	3.6	1.44
Maize France Brittany	0.43	4.0	5.2	7.4	3.1	1.26
Barley France	0.50	4.2	4.1	1.6	3.6	1.58
Wheat bran France	0.26	1.7	2.0	1.1	2.5	0.60
Pea France	0.37	9.0	1.7	1.7	4.0	2.56
Rapeseed meal France	0.52	4.0	4.4	2.3	4.3	1.39
Rapeseed oil France	1.95	15.7	16.9	8.8	15.4	5.57
Soybean meal Brazil						
Centre West	1.87	5.8	6.9	4.1	12.7	1.52
Average mix	1.47	5.8	6.3	3.6	11.2	1.56
Extruded soybean Brazil						
Centre West	2.15	7.0	7.7	5.2	13.7	1.85
Average mix	1.67	7.0	7.0	4.7	12.0	1.91
L-lysine.HCl / L-threonine	4.30	9.6	13.8	23.3	119.2	2.37
L-tryptophan / L-valine	8.54	19.2	27.4	46.4	237.5	4.74
DL-methionine	3.05	1.6	7.2	3.0	90.9	0.01
Whey powder France	1.32	0.3	1.2	0.6	23.1	0.00
Phytase	1.90	2.2	4.8	0.0	26.0	0.15
Monocalcium phosphate	1.17	16.1	30.7	7.6	17.7	0.31
Salt	0.19	0.7	1.0	1.3	3.5	0.02
Calcium carbonate	0.04	0.0	0.2	0.1	0.8	0.00
Premix	0.04	0.0	0.2	0.1	0.8	0.00

Average CP content of diets according to the feeding scenarios




Least-cost formulation without minimal CP constraints

Average incorporation of feed-use AA according to the feeding scenario




Least-cost formulation without minimal CP constraints

Average incorporation of soybean meal according to the feeding scenario









Least-cost formulation without minimal CP constraints

Average feeding cost according to the feeding scenario



**Least-cost formulation without
minimal CP constraints**

Comparison to literature results



Reduction of impacts per kg of live pig - for two-phase feeding using soybean/rapeseed/peas -

slurry



Straw bedding

Climate Change

Energy

Climate Change

Energy



Reduction of impacts per kg of live pig - for two-phase feeding using soybean/rapeseed/peas -

slurry

Straw bedding

Acidification

Eutrophication

Acidification



Eutrophication

kg SO₂-eq

kg PO₄-eq


kg SO₂-eq

kg PO₄-eq




Impacts for the different scenarios


- Slurry -




Impacts of the different scenarios - straw bedding -



Synergy between AA supp. & phase feeding



Synergy between AA supp. & phase feeding



Intermediate Conclusion

- Substitution of soybean meal by cereals and feed-use AA reduces the environmental impacts per kg pig produced especially for acidification, eutrophication and climate change
- Least cost formulation with incorporation of feed-use AA reduces both environmental impacts and feed cost whilst maintaining performance
 - **interest of suppressing minimal protein constraints**
- Synergy between «incorporation of feed-use AA» and «multiphase feeding»
 - **Maximal reduction of impacts**
 - **Potential of precision feeding to be evaluated**



Mitigating GHG emissions of pig farms by reducing the protein content of feeds: strategies and perspectives

F. Garcia-Launay, J-Y. Dourmad, P. Faverdin, S. Espagnol

INRA UMR1348 PEGASE, Saint-Gilles
IFIP, Le Rheu



QUELLE CONTRIBUTION DE L'AGRICULTURE FRANÇAISE
À LA RÉDUCTION DES ÉMISSIONS DE GAZ À EFFET DE SERRE ?

POTENTIEL D'ATTÉNUATION ET COÛT DE DIX ACTIONSTÉCHNIQUES

Résumé du rapport de l'étude réalisée par l'INRA
pour le compte de l'ADEME, du MAAF et du MEDDE - Juillet 2013






Strategies investigated | France

Reference 2010	2PMin€	MPMin€
<p>Protein content of pig feeds in accordance with recommendations Corpen 2003 (LowCP)</p> <p>80% of pig farms with two-phase feeding for growing-finishing</p>	<p>Increased utilization of feed-use amino-acids and 100% of farms with two-phase feeding for growing-finishing</p>	<p>Increased utilization of feed-use amino-acids and 100% of farms with multiphase feeding for growing-finishing</p>
<p>Least-cost formulation with minimum amino-acids contents to cover the requirements</p>		



Inventory of emissions



Impacts of feed ingredients :

Nguyen et al.
Mosnier et al.
Garcia-Launay et al.
 Dia'Terre



N intake

N excreted

Inventory of direct and indirect emissions


EMEP/EEA air pollutant emission inventory
 guidebook 2009

European Environment Agency



Functional unit and Hypothesis of strategy adoption

Least-cost formulation
Free protein content



/sow and production / year


28.2 weaned piglets /sow / year
26.6 pigs/sow/year




**Calculation of the mitigation potential
per sow and per year**

X

Sows in France 2010




Mitigation of GHG emissions



MPMin€ → Improved reduction of emissions through the reduction of protein supply



Abatement potential until 2030




Cumulated mitigation 2PMin€ > MPMIn€ because of the adoption hypothesis for MPMIn€




- Least cost formulation with incorporation of feed-use AA
 → **win-win strategies :**
 - ↘ **various impacts**
 - ↘ **both induced and on-farm emissions (at least for GHG)**
 - ↘ **Feed cost**
- Abatement of GHG emissions at negative cost :
 - 2PMin€: -96€/tCO2-eq**
 - MPPMin €: -72€/tCO2-eq**
- Decision support tools and training | and to demonstrate the improvements



Thank you for your attention




Coût annuel de la tonne de CO₂e évité (€/tCO₂e)






Win-win strategies



■ MPAA+ ■ 2PAA+

↘ **eutrophication, acidification...**
Garcia-Launay et al. 2014



■ MPAA+ ■ 2PAA+ ■ 2010

↘ **coûts d'alimentation**
Garcia-Launay et al. 2014

Some results on broilers feed
*Evaluation of the environmental implications of the
incorporation of feed-use amino acids in the
manufacturing of pig and broiler feeds using LCA*


***E. Mosnier, H. van der Werf, J. Boissy,
Jean-Yves Dourmad***

INRA Agrocampus Ouest
UMR Pegase, UMR SAS
35590 Saint-Gilles, France

Ref : Animal (2011), 5:12, pp 1972–1983



Description of the system



Simulation of broilers feeds

Six feeding strategies


1. a starter feed with 3 AA, a growing and a finishing feed with incorporation of FU-Methionine (Min€-M),
2. a starter feed with the 3 AA, a growing and a finishing feed with incorporation of FU-Methionine and L-Lysine.HCl (Min€-ML)
3. a starter, a growing, and a finishing feed with incorporation of the 3 AA (Min€-MLT).
4. maximize the AA incorporation (MaxAA)
5. minimize the CP content (MinCP) with AA incorporation
6. minimize the emission of greenhouse gas (1F-noAA),

Four contexts for feed ingredients


- 2 types of soybean meal (associated or not with deforestation)
- 2 origins for Maize (west or south of France)
- 2 protein sources (SBM only or SBM + RSM + Peas)

=> 48 diets formulated (6 x 2 x 2 x 2)


Global warming impact of different broilers feeding strategies based on cereals and soybean meal.
- for 1 kg feed -




Acidification impact of different broilers feeding strategies based on cereals and soybean meal - for 1 kg feed -



Eutrophication impact of different broilers feeding strategies based on cereals and soybean meal - for 1 kg feed -



Energy use impact of different broilers feeding strategies based on cereals and soybean meal
- for 1 kg feed -



Conclusion

- Similar results as in pigs but effects are generally less pronounced
 - not possible to formulate diets without FU-AA
 - constraints on minimal protein content
- More research needed to explain the effect of protein content on performance
- The effect on the whole system of production remains to be evaluated