

[i]Ludwigia grandiflora subsp. hexapetala[/i](Onagraceae) becomes an invasive weed on flood-prone meadows in Brière marsh (Western France)

Jacques J. Haury, Gaëtan Pottier, Michel M. Bozec, Julie Coudreuse, Jean-Patrice Damien

▶ To cite this version:

Jacques J. Haury, Gaëtan Pottier, Michel M. Bozec, Julie Coudreuse, Jean-Patrice Damien. [i]Ludwigia grandiflora subsp. hexapetala[/i] (Onagraceae) becomes an invasive weed on flood-prone meadows in Brière marsh (Western France). 4. International Symposium – Environmental weeds and invasive plants, May 2014, Montpellier, France. 2014, Environmental weeds and invasive plants. hal-01210275

HAL Id: hal-01210275

https://hal.science/hal-01210275

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



4th International Symposium



Environmental Weeds and Invasive Plants

May 19th to 23rd, 2014 Montpellier SupAgro, Montpellier, France



Ludwigia grandiflora subsp. hexapetala (Onagraceae) becomes an invasive weed on flood-prone meadows in Brière marsh (Western France)

Haury J., Pottier G., Bozec M., Coudreuse J. & J.-P. Damien

Agrocampus Ouest & INRA, UMR INRA AO ESE, France

E-mail (corresponding author): jacques.haury@agrocampus-ouest.fr

Large Waterprimrose, *Ludwigia grandiflora* subsp. *hexapetala* (Hook. & Arn.) G.L.Nesom & Kartesz (Onagraceae), is expanding on flood-prone meadows in many places, especially in Brière marsh. Thus can we explain such expansion and what are its effects on the biodiversity and the cattle breeding?

Our hypotheses are:

- (i) that tropical plant becomes invasive due to its large ecological range, but some patterns of distribution can appear, leading both to local risk analysis and areas to survey;
- (ii) adaptations to terrestrial conditions lead to changes in biomass and plant morphology,
- (iii) it has an effect on the meadow biodiversity.

Considering field research since 1998 on its distribution and local plant diversity, phenology, biomass and distribution mapping, adaptations are summarized and consequences for managers are underlined.

Keywords: Wetlands; Biological invasion; Ludwigia; biomass; Phenoplasticity