

## Biology, ecology and fertility of Water Primroses [i](Ludwigia grandiflorasubsp. hexapetala[/i] and [i]Ludwigia peploides subsp. montevidensis)[/i]: variability and consequences for management in Western France

Jacques J. Haury, Julie Coudreuse, Michel M. Bozec, Dominique D. Barloy

### ▶ To cite this version:

Jacques J. Haury, Julie Coudreuse, Michel M. Bozec, Dominique D. Barloy. Biology, ecology and fertility of Water Primroses [i](Ludwigia grandiflorasubsp. hexapetala[/i] and [i]Ludwigia peploides subsp. montevidensis)[/i]: variability and consequences for management in Western France. 4. International Symposium – Environmental weeds and invasive plants, May 2014, Montpellier, France. 2014, Environmental weeds and invasive plants. hal-01210277

## HAL Id: hal-01210277 https://hal.science/hal-01210277

Submitted on 2 Jun2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



4<sup>th</sup> International Symposium

# Environmental Weeds and Invasive Plants

May 19<sup>th</sup> to 23<sup>rd</sup>, 2014 Montpellier SupAgro, Montpellier, France

**Program - Abstracts** List of participants

> Guillaume Fried (coord.) et al. <u>Publishers:</u> EWRS, Anses, Montpellier SupAgro, MAAF-DGAI, USDA ARS-EBCL, CSIRO, Cirad, FCBN

### Biology, ecology and fertility of Water Primroses (Ludwigia grandiflora subsp. hexapetala and Ludwigia peploides subsp. montevidensis): variability and consequences for management in Western France

#### Haury J., Coudreuse J., Bozec M. & D. Barloy

Agrocampus Ouest & INRA, UMR INRA AO ESE, France

E-mail (corresponding author): jacques.haury@agrocampus-ouest.fr

Waterprimroses (*Ludwigia grandiflora* subsp. *hexapetala* (Hook. & Arn.) G.L.Nesom & Kartesz and *Ludwigia peploides* subsp. *Montevidensis* (Spreng.) P.H.Raven) are expanding in Western France and become fertile, depending on species, populations and sites. A review on existing data in literature on biology and ecology of both species in Western France is presented in a first part.

Results on fertility obtained on more than twenty populations/years are presented. Differences between species exist: in many cases, *L. peploides* has higher germination rates than *L. grandiflora*. Vernalisation increases much germination rates in any case, up to more than 80%. Aquatic and terrestrial populations present small differences in the same site, but autumn collection followed by laboratory cold exposure are less efficient than spring collection after winter cold. Field seed production is important with more than 40,000 seeds produced per m<sup>2</sup>.

The possibility of seed dispersal when flooding exists, and we observed *in situ* seedlings on some meadows. Thus fertility modifies cattle-breeder practices and all wetland management in heavily colonised sites, with special attention paid to early stages of establishment.

Keywords: Ludwigia; Population; Fertility; Adaptative strategy