



Linkage analysis of the XVIth QTLMAS simulated dataset using QTLMAP

Olivier Demeure, Olivier Filangi, Hélène Gilbert, Carole Moreno-Romieux,
Andres Legarra, Jean Michel Elsen, Pascale Le Roy

► To cite this version:

Olivier Demeure, Olivier Filangi, Hélène Gilbert, Carole Moreno-Romieux, Andres Legarra, et al.. Linkage analysis of the XVIth QTLMAS simulated dataset using QTLMAP. 16th QTLMAP, AGRIS. Algero, ITA. Università degli Studi di Sassari (UniSS), ITA., May 2012, Alghero, Italy. hal-01210359

HAL Id: hal-01210359

<https://hal.science/hal-01210359>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Linkage analysis of the XVIth QTLMAS simulated dataset using QTLMMap

O. Demeure^{1,2,*}, O. Filangi^{1,2,*}, H.Gilbert³, C. Moreno⁴, A. Legarra⁴, J.M. Elsen⁴ and P. Le Roy^{1,2}

¹ INRA, UMR1348 PEGASE, F-35590 Saint-Gilles, France

² Agrocampus Ouest, UMR1348 PEGASE, F-35590 Saint-Gilles, France

³ INRA, UMR444 Génétique Cellulaire, 31326 Castanet-Tolosan, France

⁴ INRA, UR0631 SAGA, 31326 Castanet-Tolosan, France

*: these authors contributed equally to this work




Aim

More and more association analyses, often without any use of linkage

- Does LA analyses still add some information ?
- Is LA better than association in false positive discovery ?
- It could be an interesting comparison point for the organizers

Dataset



5 chromosomes / 10000 SNP (1/0.05Mb) / 3 traits

QTLMap

A software for trait dissection in mixture of full and half-sib families using genomic markers

Single QTL	X	LA
Multi QTL	X	LD
		LDLA

Multi Trait	X	LA
-------------	---	----



ALIMENTATION
AGRICULTURE
ENVIRONNEMENT

NRA

QTLMap

A software for trait dissection in mixture of full and half-sib families using genomic markers

Single QTL
Multi QTL

X

LA
LD
LDLA

Multi Trait

X

LA



ALIMENTATION
AGRICULTURE
ENVIRONNEMENT

NRA

Linkage Analysis

- Full Mixture model, a likelihood approach

$$Likelihood = \prod_i \sum_{hs_i} p(hs_i / M) \prod_j \sum_{hd_j} p(hd_j / hs_i, M)$$

Parental phase probability


$$\prod_k \sum_{t_{ijk}} p(t_{ijk} / hd_j, hs_i, M). f(y_{ijk} / t_{ijk}, \sigma^2)$$

Transmission probability at position x

Penetrance function

Linkage Analysis

- linearized model, the regression approach



- The only most probable sire and dam phases are considered
 - The penetrance is a normal with as a mean the weighted QTL effect

$$\longrightarrow y \approx \mathcal{N}(\mu, V=I\sigma^2)$$


Linkage Analysis

- mixed model (including an animal effect)

$$\longrightarrow \mathbf{y} \approx \mathcal{N}(\mu, \mathbf{V} = \mathbf{A}\sigma_a^2 + \mathbf{I}\sigma_e^2)$$

- The user gives the pedigree and heritability
- QTLMap computes the relationship matrix A
- The effects are estimated by a BLUP

LA vs. LA Animal



LA vs. LA Animal

- No difference in QTL detection was observed in this dataset
 - It would be interesting to test this method in a small family structured design
- Focus on LA analysis
Use of all the design (G0-G1-G2-G3)
→ 60 independant families (3000 phenotypes)

Single QTL

Detection of 9 QTL using 3000 animals:

- Two were not observed using only the G3 animals
- Surprisingly QTL affecting trait 2 on chromosome 5 is less significant
- A few QTL have a location shift

→ There are more than one QTL location on Chr 1 and 5
Chr1: 19 (trait 2), 52 (trait 1) and 80cM (trait 3)
Chr5: 71 (trait 2) and 79cM (trait 1)

Single QTL

Detection of 9 QTL using 3000 animals:

- Two were not observed using only the G3 animals
- Surprisingly QTL affecting trait 2 on chromosome 5 is less significant
- A few QTL have a location shift

- There are more than one QTL location on Chr 1 and 5
- Some of the QTL affecting different traits share the same location (pleiotropy ?)
Chr4: 23 and 26cM for traits 1 and 2, respectively

Multi-QTL

0 vs 2 QTL

-No significant QTL

1 vs 2 QTL

- Chr1 / Trait 3: 60 and 84cM
 - Chr5 / Trait 2: 25 and 68cM
- Chr1 seems to carry 3 QTL, around 20, 60 and 84 cM
Chr5 seems to carry a QTL at 25cM in addition of the QTL detected around 70 and 80cM

Multi-Trait

Chromosome 1:

Single- and Multi-QTL analyses:

Chr1 seems to carry 3 QTL, around 20, 60 and 84 cM

Multi-trait analyses:

Single-QTL (three traits): 1 QTL at 60cM

Single-QTL (traits 1 / 2): 1 QTL at 19cM

Single-QTL (traits 1 / 3): 1 QTL at 78cM

3QTL (traits 1/ 2 / 3) : QTL at 19, 70 and 80cM

- QTL at 60cM is probably a false positive
There are 2 pleiotropic QTL at 19 and 78cM

Multi-Trait

Chromosome 4:

Multi-trait using traits 1 and 2: 1 QTL at 23.5cM

→ The QTL at 23.5cM is pleiotropic

The off topic slide

With the Linkage Disequilibrium (LD) analysis, we do not find the QTL on Chromosome 2 while the QTL on Chr 3 is shifted

	Chr2	Chr3
LA	0.79 ***	0.36***
LD	0.80 NS	0.49***

- Using the LA seems to be interesting for Chr2. Hard to conclude for Chr3

Conclusion

Chromosome	Trait	Location (M)	Effect (2a)	Effect (trait std)
1	1	0.19	137.32	0.87
1	2	0.19	5.93	0.87
1	1	0.78	125.35	0.79
1	3	0.78	0.02	0.96
2	3	0.79	0.02	0.7
3	3	0.365	0.02	0.875
4	1	0.235	92.03	0.58
4	2	0.235	5.74	0.67
5	1	0.785	116.42	0.79
5	2	0.25	6.71	0.775
5	2	0.68	7.38	0.852

Multi-QTL +
Multi-trait
Single-QTL
Single-QTL
Multi-Trait
Single-QTL
Multi-QTL

- 8 QTL were detected, three of them being pleiotropic
- Using different models allows the identification of additional QTL but also to precise locations or avoid false positive
- LA sometimes is more interesting than LD



We will see tomorrow how wrong we are !

Thank you for your attention



Copyright (c) 2000 Les Editions Albert René / Goscinny-Uderzo