



HAL
open science

Manure management for greenhouse gas mitigation

S O Petersen, Mélanie Blanchard, D Chadwick, A del Prado, Nadège
Edouard, J Mosquera, S G Sommer

► **To cite this version:**

S O Petersen, Mélanie Blanchard, D Chadwick, A del Prado, Nadège Edouard, et al. Manure management for greenhouse gas mitigation. *Animal*, 2013, 7 (Suppl. 2), pp.266-282. 10.1017/S1751731113000736 . hal-01210633

HAL Id: hal-01210633

<https://hal.science/hal-01210633>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Manure management for greenhouse gas mitigation

S O Petersen¹*, M. Blanchard², D. Chadwick³, A. Del Prado⁴, N. Edouard⁵, J Mosquera⁶ and S G Sommer⁷

¹Department Agroecology, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark; ²Crad, UMR Sismet, Montpellier, F-34398, France; ³Crad, UMR Sismet, RSA, Antsirabe, Madagascar; ⁴School of Environment, Natural Resources and Geography, Bangor University, Deiniol Road, Bangor LL57 2UW, Wales; ⁵Basque Centre for Climate Change, Alameda Urquijo 4, 48008 Bilbao, Spain; ⁶INRA Agrocampus Ouest, UMR1348 PEGASE, F-35590 Saint Gilles, France; ⁷Wageningen UR Livestock Research, PO Box 135, 6700 AC Wageningen, The Netherlands; *E-mail: soren.o.petersen@agrsci.dk

(Received 21 November 2012; Accepted 25 March 2013)

Ongoing intensification and specialisation of livestock production lead to increasing volumes of manure to be managed, which are a source of the greenhouse gases (GHGs) methane (CH₄) and nitrous oxide (N₂O). Net emissions of CH₄ and N₂O result from a multitude of microbial activities in the manure environment. Their relative importance depends not only on manure composition and local management practices with respect to treatment, storage and field application, but also on ambient climatic conditions. The diversity of livestock production systems, and their associated manure management, is discussed on the basis of four regional cases (Sub-Saharan Africa, Southeast Asia, China and Europe) with increasing levels of intensification and priorities with respect to nutrient management and environmental regulation. GHG mitigation options for production systems based on solid and liquid manure management are then presented, and potentials for positive and negative interactions between pollutants, and between management practices, are discussed. The diversity of manure properties and environmental conditions necessitate a modelling approach for improving estimates of GHG emissions, and for predicting effects of management changes for GHG mitigation, and requirements for such a model are discussed. Finally, we briefly discuss drivers for, and barriers against, introduction of GHG mitigation measures for livestock production. There is no conflict between efforts to improve food and feed production, and efforts to reduce GHG emissions from manure management. Growth in livestock populations are projected to occur mainly in intensive production systems where, for this and other reasons, the largest potentials for GHG mitigation may be found.

Keywords: methane, nitrous oxide, storage, treatment, farm model

Implications

Livestock manure is a source of greenhouse gas (GHG) emissions, mainly as methane and nitrous oxide. GHG emissions are biogenic and regulated by manure characteristics, and therefore emissions can be manipulated via handling, treatment and storage conditions. Globally, livestock production systems vary widely, and this is also true for GHG mitigation potentials, but generally efforts to conserve nutrients in manure for crop production will also reduce GHG emissions. Future growth in livestock production is projected to occur mainly in confined animal feeding operations, which also appear to have the greatest potential for GHG mitigation.

Introduction

Since the mid 20th century, there has been a growing pressure on land resources for production of food and feed

for livestock and, increasingly, crops for energy production (Hoogwijk *et al.*, 2005). To fulfil the demand for meat, milk and eggs, livestock production in developing countries is expanding, especially in peri-urban areas (Gerber *et al.*, 2005), and worldwide becomes more specialised (Steinfeld *et al.*, 2006). In consequence of these trends, increasing volumes of livestock manure are produced, which are a source of greenhouse gases (GHGs) contributing to radiative forcing (Forster *et al.*, 2007). Using a life cycle approach, the relative contribution of global livestock production to anthropogenic GHG emissions was estimated to be 18% (Steinfeld *et al.*, 2006), whereas a similar analysis for the European Union arrived at 12.8%, or 9.1% without land use and land use change-related emissions (Leip *et al.*, 2011).

GHG emissions from agriculture are biogenic, and the GHG balance of manure management reflects a multitude of microbial activities, that is: emissions of methane (CH₄) are the net result of methanogenesis and CH₄ oxidation; nitrous oxide (N₂O) is a product of several processes, but may also

* E-mail: soren.o.petersen@agrsci.dk

be consumed via denitrification before escaping to the atmosphere; and the carbon dioxide (CO₂) balance is influenced by manures via (net) soil carbon stock changes upon field deposition and any production of bioenergy.

Within the lifecycle of livestock products, most variability occurs at the level of farming system (Oenema *et al.*, 2003), and globally livestock manure differs widely in volume, composition and storage conditions. This is discussed with reference to four specific cases of livestock production that differ in production intensity and regulation. The range of manure characteristics is extended by treatments such as composting, forced aeration, solid-liquid separation, anaerobic digestion or use of additives. Moreover, feeding is part of the manure management chain by determining the quality and nutrient composition of excreta. The paper includes an overview of management practices with a potential for GHG mitigation. Management at one stage affects manure composition and emissions at subsequent stages, and therefore the entire continuum of manure management and treatment must be taken into account when evaluating individual or combinations of GHG mitigation measures (Chadwick *et al.*, 2011), which calls for the use of whole-farm models (Del Prado *et al.*, 2013). The paper finally discusses opportunities for, and barriers against, GHG mitigation. Although excretal returns during grazing should also be considered in the context of manure management, this review will mainly focus on the management of manure collected during confinement.

CH₄ and N₂O emissions from manure: sources and sinks

Understanding the microbial ecology of manure environments is critically important for proper estimation of GHG emissions from manure management, and for efforts to predict effects of management changes and develop GHG mitigation strategies. Here, we briefly review the response of microorganisms in manure environments to key environmental controls.

Methanogenesis

CH₄ emissions may occur from all manure environments, but is mainly associated with liquid or compacted manure (Osada *et al.*, 2000; Chadwick *et al.*, 2011). Methanogenesis occurs only under strictly anaerobic conditions where it is coupled to other processes involved in the breakdown of manure organic matter (Valentine, 2007). Little is known about the microbiological basis of methanogenesis in fresh manure, but the methanogenic potential in fresh manure is low, probably because of inhibitory concentrations of ammonia (NH₃) derived from urine (Chen *et al.*, 2008). There are, however, slow-growing methanogens capable of adapting to as much as 7 g total ammoniacal N/l (i.e., *Methanosarcina* spp.), which are known to develop in manure and anaerobic digesters (Rastogi *et al.*, 2008; de Vrieze *et al.*, 2012).

Owing to a marginal energy yield (Valentine, 2007), methanogenesis is sensitive to low temperature, and therefore cooling is a potential CH₄ mitigation option (Sommer *et al.*, 2004; Umetsu *et al.*, 2005). For cool temperate climates, frequent removal of manure from the housing to an

outside store has been proposed as a low-cost strategy to reduce CH₄ emissions (Sommer *et al.*, 2009). However, the predicted effect of frequent removal assumes that there is no difference in the potential for CH₄ production between slurry pits and the outside storage facility, which may not be the case if adapted organisms have not yet developed in the slurry pit, for example, because of inhibitory concentrations of NH₃, or cleaning after emptying (Haeussermann *et al.*, 2006).

Presumably, during the gradual filling of an outside storage tank or lagoon, there will be a fast inoculation of newly added slurry, such that in the final storage facility CH₄ production and emissions can be predicted by the physical and chemical properties of the slurry, that is, substrate availability for methanogens (Wood *et al.*, 2012). This is also the assumption of the 2006 methodology of the Intergovernmental Panel on Climate Change (IPCC), which links CH₄ emissions from liquid manure to mean storage temperature (IPCC, 2006).

CH₄ oxidation

CH₄ is oxidised mainly by aerobic bacteria (Hanson and Hanson, 1996). Both methanogens and CH₄-oxidising bacteria (MOB) are present in solid manure (Sharma *et al.*, 2011). There are few studies on the potential for CH₄ oxidation, but MOB population dynamics during composting and maturation phases have been described for other organic residues (Wilshusen *et al.*, 2004; Halet *et al.*, 2006), and thus MOB activity may play a role in mitigating CH₄ emissions from solid manure, as proposed by Szanto *et al.* (2007).

An organic surface crust is often present on livestock slurry stores, which may contain a significant potential for CH₄ oxidation (Ambus and Petersen, 2005; Petersen *et al.*, 2005; Hansen *et al.*, 2009). However, Nielsen *et al.* (2013) found that potential methane oxidation (PMO) in two natural crusts from piggeries remained low until late autumn, indicating that MOB activity was low during summer and early autumn where most of the annual emissions occur (Husted, 1994). The reason for this delay is unknown; however, a crust overlying liquid manure will be rich in NH₃, and also nitrification and denitrification activity could potentially interfere with CH₄ oxidation activity. Duan *et al.* (2013) found PMO in a surface crust to be 50 to 100 times more sensitive to nitrite (NO₂²⁻) than to NH₄¹⁺ or nitrate (NO₃²⁻), and, 1 mM NO₂²⁻ significantly inhibited CH₄ oxidation activity.

N₂O emissions via nitrification and denitrification

The conversion of oxidised N to gaseous forms during manure handling, storage and after field application can represent a significant loss of plant-available N (Rotz, 2004). Both N₂O and, indirectly, NH₃ volatilisation and NO₃²⁻ leaching contribute importantly to the GHG balance of manure management. Whereas CH₄ production and oxidation processes are associated with anoxic and oxic conditions, respectively, emissions of N₂O are stimulated under O₂-limited conditions.

Nitrification is a two-step conversion of NH₃ via NO₂²⁻ to NO₃²⁻, primarily carried out by ammonia-oxidising (AOB) and nitrite-oxidising bacteria (NOB). Ammonia-oxidising archaea

Table 1 Regional differences in manure management practices for dairy production

System ¹	North America	CSA	Western Europe	Eastern Europe	Russian Federation	NENA	SSA	South Asia	East Asia	Oceania
Lagoon	12	0	0	0	0	0	0	0	0	4
Liquid/slurry	32	0	38	22	0	2	0	4	4	0
Solid storage	31	29	36	61	78	29	32	26	26	0
Drylot	0	19	0	0	0	20	21	18	18	0
Pasture/range	16	52	22	14	22	48	47	48	48	94
Daily spread	9	0	4	3	0	0	0	0	0	2

CSA:5 Central and South America; NENA:5 Near East and North Africa; SSA:5 Sub-Saharan Africa.

The table shows the relative importance (in %) of liquid and soil manure management practices, including systems dominated by grazing.

¹Lagoons are typically earthen sedimentation basins for dilute waste from housed animals; sludge is degraded anaerobically, while the liquid evaporates or is pumped to spray fields. Liquid manure/slurry typically has a higher dry matter content and may include bedding material and is stored in tanks and may form a crust during storage. Solid storage is used for manure from housed animals on deep litter, or with separate collection of faeces, bedding material and urine, respectively. Drylots are large confinements where a compacted pad forms on the ground, which is infrequently scraped and may be stacked for composting. In pasture/range-dominated systems, most manure will be deposited during grazing.

Source: Food and Agriculture Organization of the United Nations (FAO) (2010).

have also been found in composting manure and soil, but are not responsive to N inputs, and hence their role is uncertain (Ja and Conrad, 2009; Yamamoto *et al.*, 2010). N₂O is produced by AOB, either as a side-product in the conversion of NH₃ or by a subsequent conversion of NO₂² via a process called *nitrifier denitrification* (Goreau *et al.*, 1980; Kool *et al.*, 2011). Nitrifier denitrification may be a mechanism to avoid toxic levels of NO₂², which can occur when there is an imbalance between AOB and NOB activity, for example, because of competition for O₂ (Laanbroek and Gerards, 1991).

Denitrification is carried out by a phylogenetically diverse group of heterotrophic bacteria, although the process has also been found among fungi and archaea (Wallenstein *et al.*, 2006). Most denitrifiers are facultative anaerobes, that is, they prefer O₂ as electron acceptor, but can use NO₃² or NO₂² in the absence of O₂. The expression of genes for denitrifying enzymes is stimulated at low O₂ levels, but N₂O reductase is unstable even with traces of O₂ (Thomson *et al.*, 2012). Therefore, N₂O can accumulate around oxic-anoxic interfaces in the manure.

Manure environments – during storage and after field application – are characterised by steep gradients in O₂ because of intense decomposer activity, which enables potentials for nitrification and denitrification to develop at close proximity around oxic-anoxic interfaces (Petersen *et al.*, 1992 and 1996). Chemical gradients can also develop that may influence N₂O emissions. The pH of fresh manure is neutral to alkaline, and dominated by NH₄¹ 1 NH₃ and carbonates derived from decomposer activity (Husted *et al.*, 1991). Around air-liquid interfaces, CO₂ and NH₃ will be lost to the gas phase, thereby removing buffering capacity and alkalinity. As NH₃ oxidation is an acidifying process, there is thus a potential for significant reduction in pH because of nitrification. For example, Petersen *et al.* (1992) observed a drop in pH from 6.5 to 4.5 around manure hot spots in a sandy loam soil that coincided with a more than 10-fold increase in potential NH₃ oxidation. If pH declines, rates of nitrification and denitrification may also decline (Cuhel *et al.*, 2010; Cytryn *et al.*, 2012), but the N₂O-to-N₂ ratio of

denitrification will increase (Baggs *et al.*, 2010). It can be speculated that this environment is also conducive to N₂O emissions via nitrifier denitrification if NO₂² accumulates because of selective inhibition of NOB (de Boer and Kowalchuk, 2001; Kool *et al.*, 2011). Indeed, Fukumoto (2006) showed that addition of NOB to a compost prevented NO₂² accumulation and N₂O emission.

Manure management practices – a global view

The relative importance of microbial processes leading to CH₄ and N₂O emissions will depend on the manure environment as determined by local management practices and climate, factors that differ greatly between regions, as exemplified for dairy production in Table 1. For dairy production, grassland systems predominate in Central and South America, Africa and Asia/Oceania; confined animals with solid manure management in Eastern Europe and the Russian Federation; and some regions have a high proportion of liquid manure management for confined animals.

In grassland-based systems, faeces and urine from grazing animals is deposited on pastures or rangeland and not handled (Oenema *et al.*, 2007). Dry areas of the tropics, and continental climates of Asia and North America, are dominated by extensive grazing by ruminants (cattle, sheep, goats, camels), often in a rotational way (Steinfeld *et al.*, 2006). Intensive grazing, mainly by cattle, is found in medium to high population density areas, mostly in temperate climate zones of Europe, North and South America and, increasingly, in the humid tropics. In both extensive and intensive grazing systems, substantial N losses may occur through leaching and volatilisation from point sources of urine and faeces where N concentrations may exceed the equivalent of 200 and 2000 kg N/ha, respectively (Lantinga *et al.*, 1987).

If livestock production is partly or completely based on confined animal feeding, manure is normally collected and must be managed from the time of excretion, during storage, possibly by treatment, and finally during spreading to land (Chadwick *et al.*, 2011), although in some regions a

proportion is also directly discharged to the environment. On-farm manure management systems can vary widely depending on animal category, housing design and manure collection system. Burton and Turner (2003) distinguished three categories of manure management: (a) systems collecting liquid manure (slurry) from animals kept on slatted or solid floors regularly swept clear of any excreta, sometimes with some dilution from washing water; (b) systems producing solid manure (farmyard manure) from animals kept on bedding material, which is collected together with all excreta; and (c) systems producing mixed manure from animals kept on bedding material, but with drainage and separate collection of liquids. In dry climates, animals may also be kept in unpaved feedlots where manure is periodically removed (IPCC, 2006). Before storage and field application, manure can be treated by different methods for improved handling, nutrient use or energy generation (see next section). As basis for further discussion of GHG mitigation potentials via changes in manure management, we present four specific livestock production situations. It must be stressed that these examples are not intended to represent the true diversity, but rather four different levels of intensification.

Sub-Saharan Africa – subsistence farming

Sub-Saharan Africa is characterised by extensive subsistence farming. In West Africa, agricultural land covers 30% to 60% of the area, and population density is 45 to 80 per km². Farming systems are dominated by cereal (corn, sorghum and millet) and cotton production, and the area available for grazing is limited. Livestock consists mainly of cattle at 0.08 to 4.8 tropical livestock units (TLU)/ha (Anon., 2007), where 1 TLU = 250 kg live weight (Hoffmann *et al.*, 2001). During the dry season, animals are confined and fed crop residues. When the agricultural season begins (rainy season), shepherds lead their livestock to graze on pastures either near the farm or over long distances (transhumance). There are three main categories of farms. Farms dominated by crop production (cropped area 5 to 11 ha, 1 to 6 TLU) constitute 68% of all. Agro-pastoralist farms with more emphasis on livestock (11 to 24 ha, 11 to 30 TLU) represent 30% of all farms. The last 2% are specialist breeders with cattle herds up to 15 TLU and small areas (on average 0.6 ha) with cereals for their own consumption (Vall *et al.*, 2006).

There is considerable diversity in management of animal manure between farms (Manlay *et al.*, 2002; Blanchard, 2010); the main priority is recycling of nutrients for crop production. Garbage piles with domestic waste, daily sweepings and faeces from small ruminants, and some soil, may be produced in the homestead area. Confining animals helps produce organic fertiliser in significant quantities, by facilitating manure collection. Some farmers add bedding material and feed leftovers to the pen or animal shed (Landais and Guérin, 1992; Landais and Lhoste, 1993; Ganry *et al.*, 2001), which further increases the quantity and nutrient content of manure, as nutrients in urine are trapped by the litter. Household compost may be produced in pits near the homestead area on the basis

of animal faeces, feed and crop residues, and domestic waste (Ganry *et al.*, 2001). Farmers may irrigate the pit, turn the compost and use a cover to limit N losses and promote decomposition.

Nutrient cycling and losses associated with the management of manure have been estimated for farms with 10 to 75 TLU in South Mali (Blanchard, 2010). Between 38% and 50% of animal faeces (6 to 40 t/year) is deposited during grazing on common pastures. Deposition of faeces during transhumance represents 0 to 25 t/year. The monitoring study indicates that, for farms of West Africa, 46% of the N in crop residues and manure is returned to the soil of common pastures or areas of transhumance, whereas 13% is lost in gaseous form at the time of excretion. Organic manure produced on the farm represents 24% of the N, and 17% is lost through leaching or in gaseous form during handling and storage of manure and compost (Blanchard, 2010). The N cycling efficiencies were close to those reported by Rufino *et al.* (2007) of 13% to 28%.

With the rising price of mineral fertilisers, reduction in fertiliser subsidies and programmes promoting organic manure quality, there is an increasing focus on efficient use of nutrients in livestock manure (Blanchard and Vall, 2010). To increase nutrient conservation, recommendations are to compost under roofs and on floors (Rufino and Rowe, 2006), and to limit storage time (Titttonell *et al.*, 2010). Farmers aim to keep animals longer in confinement by improving forage availability (Landais and Lhoste, 1993). Bio-digesters on farms to provide energy for light and cooking are still new to this region, but will produce a new type of manure to be managed.

Vietnam – smallholder confined animal feeding operations (CAFOs)

In Vietnam, total livestock production has increased significantly since the late 20th century, but with little adaptation of existing manure management practices. Traditionally, solid manure is removed manually from pig and cattle houses on a daily basis and floors cleaned with water (Vu *et al.*, 2007). As pigs are also cooled with water, there may be a 10-fold dilution of manure (Vu *et al.*, 2012). Asian farming systems typically include crop production, fish farming and livestock. A survey in two North Vietnamese provinces found that 5% and 35% of the manure, respectively, was used for cash crops (Vu *et al.*, 2007). Addition of untreated animal manure to rice paddies may increase N₂O and CH₄ emissions significantly, but emissions can be reduced by water management; it has been shown that intermittent irrigation and/or midseason drainage reduces CH₄ emissions from paddy fields compared with continuous flooding (Minamikawa *et al.*, 2006).

Both liquid and solid manure (fresh or composted) are applied to fish ponds. If dilute liquid manure is not drained to a fish pond, it is mostly discharged to rivers or ditches, as the nutrient content is low. This discharge can account for 7% to 15% of livestock feed N intake, 10% to 17% of P intake and 9% to 23% of K intake (Vu *et al.*, 2012).

Biogas production is used at the farm scale in most Asian countries, where it substitutes other energy sources while also reducing odour and improving nutrient use efficiency (Cu *et al.*, 2012; Wang and Zhang, 2012). In Vietnam, 110 000 digesters were in use in 2010, which may increase to 300 000 by 2018 (Cu *et al.*, 2012). For comparison, in China, 30 million biogas plants were in use in 2009, a number that is also expected to increase (Jang *et al.*, 2011). For a small farm with a total production of 12 pigs/year, the GHG balances of manure management without and with anaerobic digestion have been estimated at 2271 and 2 504 kg CO₂ Eq/year, respectively (S G. Sommer, unpublished data). However, even well-managed biogas plants have minor leakages of CH₄ to the environment (Sommer *et al.*, 2004), and small-scale digesters in Asian countries are often not well managed (Jang *et al.*, 2011; Cu *et al.*, 2012). A study from Thailand estimated the release to account for 15% of the gas produced (Prapasongsa, 2010), and recently Bruun *et al.* (2013) concluded that, for Vietnam, the loss of CH₄ from digesters could be as high as 40%, which would make biogas production a net source of GHG.

In summary, livestock production in Vietnam and other countries of Southeast Asia is characterised by significant growth, but with little progress in manure management, and therefore GHG emissions are increasing. On-farm biogas production is a potential GHG mitigation option, but management should be improved.

China – small- and large-scale CAFOs

Livestock numbers in China have increased markedly over the past two decades (MOA, 2010), with the associated manure generation reaching 3000 to 3500 Mt in 2007 (MEP, 2010). This manure is generated through a combination of backyard low-productivity household farms and highly productive intensive livestock enterprises, similar to those seen throughout the United States and Europe. However, as the demand for livestock products increases within China, so will livestock numbers and volumes of manure generated, with an estimated increase of around 1000 Mt between the years 2000 and 2030 (Chadwick *et al.*, 2012). A recent survey suggests that the proportion of livestock reared in intensive systems is increasing, especially in the peri-urban areas (MOA, 2010). For example, MOA data show that , 80% of broilers and laying hens, and 50% of pigs, are now reared in CAFOs in China.

Another recent survey (2007 data) has suggested that almost 20% of manure is 'wasted' and not applied to land, which has implications for water quality and indirect N₂O emissions via NO₃². Of the remaining 80%, ca. 26% is composted, 8% used for biogas production before spreading and the rest of the manure (66%) is spread directly to land. A range of subsidies will lead to more livestock manure being managed via the digestion and composting routes. Simple containment to ensure that manure is applied to land will improve both air and water quality.

More centralised livestock production presents opportunities for GHG emission control during the manure production

(housing) and handling phases (centralised biogas production), although careful planning is necessary to ensure that manure nutrients are used effectively on agricultural land (both proximate to the production unit, and at distance where, e.g., composted faecal matter is sold). Local-, household- and village-scale anaerobic digestion schemes have been implemented, but the lack of mechanisation for removal of solids from digesters and land spreading means that use of nutrients in the digestate is not optimised.

Manure and fertiliser N use in China is excessive, especially for high-value crops such as fruit and vegetables (Gao *et al.*, 2012). It can be as high as 1 to 2 t N/ha per year. If greater account was made of manure nutrients, especially N, for crop supply, then less fertiliser N would need to be applied, reducing the carbon footprint of production and use of fertilisers (Zhang *et al.*, 2013). A reduction in these excessive applications would reduce direct and indirect (via NO₃² leaching) emissions of N₂O from soil, especially if one considers the non-linear relationship between application rate and direct emissions (e.g., Cardenas *et al.*, 2010).

EU – highly regulated livestock production

For socio-economic reasons, European livestock production is increasingly intensive, with specialisation and mechanisation leading to larger farms (Burton and Turner, 2003). Intensive systems are dominated by three animal categories: cattle, pigs and poultry. They are often 'landless' in the sense that , 10% of animal feed is produced on the farm (Kruska *et al.*, 2003). The geographic uncoupling of feed production from animal production is a fundamental challenge because of the concentration of nutrients in livestock-intensive areas – here, as in other parts of the world.

Large proportions of nutrient intake are excreted, for example, 60% to 70% of ingested N for fattening pigs and laying hens, and 70% to 90% for cattle depending on physiological stage (Peyraud *et al.*, 2012). Manure is commonly used as a fertiliser on the farm, but transfer between farms is also seen in regions with high livestock densities (e.g., Netherlands, Western France). Regulations have been introduced in almost all countries to prevent discharge to rivers and streams, guidelines are available for storage and land application (timing, location, rate, method). The EU Nitrates Directive stipulates a maximum annual application of 170 kg/ha of manure N (European Commission (EC), 1991), although some derogations exist that allow higher rates for crops with a high N uptake potential. Nutrient recycling is a challenge for large livestock farms with little or no land (e.g., a farm with 20 000 fattening pig places requires 2 to 3000 ha of cropland for manure recycling; Menzi *et al.*, 2010). By assigning an N fertiliser value to the manure, regulations and recommendations can ensure that manure nutrients are considered as a primary source of macro-nutrients (Anon., 2012).

In Europe, 30% to 40% of livestock excreta are deposited during grazing and thus not handled. The remaining 60% to 70% is collected in housing systems, a percentage that tends to increase. Manure management systems producing solid manure represent 20% to 30% of excreta, whereas the

Table 2 Effects of different management options on CH₄, N₂O and combined CH₄ + N₂O emissions from housing

Management option	Animal category	N ₂ O	CH ₄	CH ₄ + N ₂ O	References
Solid v. liquid manure Straw bedding	Fatteners	1 106	2 2	1 29	Philippe <i>et al.</i> (2007)
	Gestating sows	1 383	2 9	1 131	Philippe <i>et al.</i> (2011)
	Weaned pigs	1 ¹	2 18	1 22	Cabaraux <i>et al.</i> (2009)
	Dairy cattle	1 85	1 33	1 48	Eclouard <i>et al.</i> (2012)
	Weaned pigs	1 286	2 51	1 195	Nicks <i>et al.</i> (2003)
Sawdust v. straw	Fatteners	1 6867	2 33	1 286	Nicks <i>et al.</i> (2004)
	Fatteners	1 7600	1 100	1 667	Kaiser (1999)
	Laying hens	1 259	1 319	1 275	Mennicken (1998)
Wood shavings v. straw	Figs		2 31		Sommer <i>et al.</i> (2004)
	Fatteners		2 43		Groenestein <i>et al.</i> (2012)
	Nursing sows		2 46		Groenestein <i>et al.</i> (2012)
	Gestating sows		2 33		Groenestein <i>et al.</i> (2012)
	Weaned pigs		2 30		Groenestein <i>et al.</i> (2012)
Cooling	Figs	2 39	2 56	2 51	Amon <i>et al.</i> (2007)
	Figs		2 40		Haeussermann <i>et al.</i> (2006)
	Weaned pigs	0	2 50	2 50	Groenestein <i>et al.</i> (2011)
	Fatteners	0	2 86	2 86	Groenestein <i>et al.</i> (2011)

¹ '1' represents higher emissions (%) and '–' lower emissions (%) compared with the reference (untreated) manure. The comparison of systems is based on CO₂ equivalents.

¹N₂O emissions from slurry were not measurable.

remainder is handled as slurry that is either stored in pits beneath animal confinements or in outside tanks (Oenema *et al.*, 2007). The proportion of manure in liquid form varies considerably between countries; it is generally higher (. 65%) in central and northern Europe, even reaching more than 95% in the Netherlands, and lower (. 50%) in the United Kingdom, France and some parts of Eastern Europe where housing systems are often associated with bedding materials (e.g., deep litter). For both liquid and solid manure, the appropriate storage capacity depends on the maximum length of time during which manure cannot be applied to land. In Europe, because of the winter break in vegetation growth, the required storage capacity ranges from 4 to 9 months (Menzi *et al.*, 2010).

There are several drivers leading farmers towards liquid manure management systems. In addition to easier handling, liquid manure has a higher mineral N-to-organic N ratio and thus a higher percentage of plant-available N, and there are several options for treatment with a potential to improve manure quality and reduce losses towards the environment. In addition, production of 25 t solid manure requires straw from approximately 1 ha of cereals (Schröder, 2005), and hence availability and price of straw is a constraint. Mechanical separation of liquid manure is practised to varying extent in European countries, for example, 90% of pig slurry in Greece, 10% of all slurry in Spain and in Italy 15% of cattle and 40% of pig slurry (Burton and Turner, 2003). Aeration of slurry is increasingly practised in France and the Netherlands. Biogas production is expanding in several countries, but the proportion of livestock slurry that is treated varies greatly (Anon., 2010).

GHG emissions, direct and indirect, occur at all stages of manure management. Although livestock production in EU is

already highly regulated, this does not include GHG emissions, and hence there is scope for further improvements in manure management. NH₃ and GHG emissions may be higher for either slurry or solid manure depending on management stage (housing, storage, spreading) and animal species (Gac *et al.*, 2007), and hence it is important to identify mitigation strategies for both categories.

Manure handling and treatment for GHG mitigation

In this section, we briefly discuss GHG mitigation potentials of the most relevant options for solid and liquid manure management. Liquid manure is collected in slurry channels below slatted floors or deposited on soiled slats and surfaces, whereas solid manure accumulates as litter on the floor. GHG emissions tend to be higher for solid manure-based systems (Table 2), particularly for N₂O and with fattening pigs. Manure removed from animal houses or confined areas may be treated or stored outside, and is mostly field-applied to recycle nutrients. At all stages, management has an impact on GHG emissions.

Housing – diet

Diet has not only a direct effect on CH₄ emissions from enteric fermentation, but also an indirect effect on CH₄ emissions during storage, by affecting manure composition (e.g., Hindrichsen *et al.*, 2005); see also section 'On-farm interactions' below. The effect of diet on denitrification and N₂O emission is related to the protein balance, as excess N is excreted, and a reduction in manure N concentration will also reduce N₂O emissions (Misselbrook *et al.*, 1998). Inclusion of some natural compounds (such as tannins via

Table 3 Effects of different management options on CH₄, N₂O and combined CH₄ + N₂O emissions from storage

Type of storage	Management option	N ₂ O	CH ₄	CH ₄ + N ₂ O	References	
Solid manure	Forced v. passive composting	2 35	2 90	2 78	Amon <i>et al.</i> (2001); cattle farmyard manure; summer measurements	
		2 41	1 32	2 7	Amon <i>et al.</i> (2001); cattle farmyard manure; winter measurements	
		1 44	2 81	2 34	Pattey <i>et al.</i> (2005)	
	Straw cover		2 28			Hao <i>et al.</i> (2001); farmyard manure
		2 42	2 45	2 42	Yamulki (2006); cattle farmyard manure; conventional farm	
		2 11	2 50	2 14	Yamulki (2006); cattle farmyard manure; organic farm	
	Plastic sheet cover	2 70	2 6	2 36	Chadwick (2005); cattle solid manure, period 1	
		1 2000	2 81	2 17	Chadwick (2005); cattle solid manure, period 2	
		2 54	1 120	1 111	Chadwick (2005); cattle solid manure, period 3	
		2 99	2 87	2 98	Hansen <i>et al.</i> (2006); solid fraction of digested pig manure	
		2 32			Thorman <i>et al.</i> (2006); poultry manure	
		1 304			Thorman <i>et al.</i> (2006); poultry manure	
	Liquid manure	Straw cover	1 57	2 25	2 23	VanderZaag <i>et al.</i> (2009); cattle slurry; straw layer: 15 cm
			1 100	2 27	2 24	VanderZaag <i>et al.</i> (2009); cattle slurry; straw layer: 30 cm
			1 37		Guarino <i>et al.</i> (2006); cattle slurry; straw layer: 7 cm	
			1 3		Guarino <i>et al.</i> (2006); cattle slurry; straw layer: 14 cm	
			1 7		Guarino <i>et al.</i> (2006); pig slurry; straw layer: 7 cm	
			2 28		Guarino <i>et al.</i> (2006); pig slurry; straw layer: 14 cm	
1 432			1 22	1 238	Berg <i>et al.</i> (2006); pig slurry; straw layer: 6 to 8 cm	
Solid cover		1 30	2 32	1 1	Amon <i>et al.</i> (2007); pig slurry; warm period (50 days)	
		2 4	2 70	2 52	Amon <i>et al.</i> (2007); pig slurry; warm period (200 days)	
		2 50	2 37	2 48	Amon <i>et al.</i> (2007); pig slurry; cold period (50 days)	
		2 13	2 14	2 13	Cemens <i>et al.</i> (2006); cattle slurry (winter period)	
		1 20	2 16	2 11	Cemens <i>et al.</i> (2006); cattle slurry (summer period)	
		1 2	2 29	2 4	Cemens <i>et al.</i> (2006); digested cattle slurry (winter period)	
		2 19	2 14	2 16	Cemens <i>et al.</i> (2006); digested cattle slurry (summer period)	

'1' represents higher emissions (%) and '-' lower emissions (%) compared with the reference (untreated) manure. The comparison of systems is based on CO₂ equivalents

birdsfoot trefoil) in the diet can increase the proportion of N excreted in organic compounds via faeces, rather than in the urine as urea, which in turn reduces the potential for NH₃ emissions and N₂O emissions (Misselbrook *et al.*, 2005), but potentially also plant availability.

Housing – manipulation of storage temperature

The temperature dependency of methanogenesis was already discussed. Active cooling of slurry channels may be a cost-effective CH₄ mitigation option if the exchanged heat can be used (Sommer *et al.*, 2004). Cooling of slurry below slatted floors to 10°C has been found to reduce CH₄ emissions by 30% to 46% compared with the situation without cooling (Table 2). Manure cooling can also mitigate NH₃ emissions from in-house manure storage (Groenestein *et al.*, 2011).

Frequent removal of manure to an outside store relies on a significant temperature difference between housing and outside store (Sommer *et al.*, 2009) and is therefore most relevant for cold and temperate climates. Efficacy will depend on the methanogenic potential of the slurry, as discussed above, but several studies did find significant (40% to 86%) reductions in GHG emissions from pig housing with frequent manure removal (Table 2).

Solid manure – composting

Composting is a process where microorganisms transform degradable organic matter into CO₂ and water under (predominantly) aerobic conditions. Manure can either be left

undisturbed during the composting process (passive composting), mechanically turned (extensive composting) or actively aerated (intensive composting). Aeration may reduce CH₄ emissions (Table 3), but increase N₂O and NH₃ emissions (Pattey *et al.*, 2005; Webb *et al.*, 2012). Combined CH₄ and N₂O emissions are generally lower after forced aeration and turning compared with passive composting (Table 3).

Solid manure – cover during storage

Covering solid manure with straw or a plastic sheet reduces in general both N₂O and CH₄ emissions, and therefore total GHG emissions, compared with the situation without cover (Table 3). However, Chadwick (2005) showed both a reduction (up to 36%) and an increase (by 11%) of total GHG emissions when covering compacted cattle solid manure with a plastic sheet. Thorman *et al.* (2006) reported both a reduction and an increase in N₂O emissions after covering poultry solid manure with a plastic sheet. NH₃ emissions may also be reduced by covering the heap (Chadwick, 2005; Webb *et al.*, 2012).

Liquid manure – cover during storage

Covers on liquid manure stores are mainly adopted to reduce NH₃ emissions. N₂O emissions from liquid manure are negligible during storage, unless a surface crust is present (e.g., VanderZaag *et al.*, 2009). With a crust, potentials for nitrification and denitrification can develop and lead to N₂O emissions, as the crust dries and oxygen enters the crust

Table 4 Effects of different management options on CH₄, N₂O and combined CH₄ + N₂O emissions from manure treatment

Management option	Type of manure	N ₂ O	CH ₄	CH ₄ + N ₂ O	Reference	
Manure separation	Fg slurry (58C)	0	2.8	2.8	Dinuccio <i>et al.</i> (2008)	
	Fg slurry (258C)	1 ¹	1.3	1.41	Dinuccio <i>et al.</i> (2008)	
	Cattle slurry (58C)	0	1.4	1.4	Dinuccio <i>et al.</i> (2008)	
	Cattle slurry (258C)	0	2.9	2.9	Dinuccio <i>et al.</i> (2008)	
	Cattle slurry	1.1133	2.34	2.23	Fangueiro <i>et al.</i> (2008)	
	Cattle slurry + wooden lid	1.10	2.42	2.39	Amon <i>et al.</i> (2006)	
	Fg slurry	1 ¹	2.93	2.29	Mosquera <i>et al.</i> (2011)	
	Cattle slurry	1 ¹	2.42	1.25	Mosquera <i>et al.</i> (2011)	
	Fg slurry		2.18		Martinez <i>et al.</i> (2003)	
	Cattle slurry		2.40		Martinez <i>et al.</i> (2003)	
Anaerobic digestion	Cattle slurry	2.9	2.32	2.14	Clemens <i>et al.</i> (2006)	
	Cattle slurry	1.49	2.68	2.48	Clemens <i>et al.</i> (2006)	
	Cattle slurry + wooden lid	1.41	2.67	2.59	Amon <i>et al.</i> (2006)	
Aeration	Cattle slurry	1.144	2.57	2.43	Amon <i>et al.</i> (2006)	
	Fg slurry (period 1)		2.99		Martinez <i>et al.</i> (2003)	
	Fg slurry (period 2)		2.70		Martinez <i>et al.</i> (2003)	
Dilution	Fg slurry		2.35		Martinez <i>et al.</i> (2003)	
	Cattle slurry		2.57		Martinez <i>et al.</i> (2003)	
Additives	NX ₂₃		2.47		Martinez <i>et al.</i> (2003)	
	Stalosan		2.54		Martinez <i>et al.</i> (2003)	
	Biosuper		2.64		Martinez <i>et al.</i> (2003)	
	Sulphuric acid (pH 6)	Cattle slurry (pH 5.5)		2.87		Petersen <i>et al.</i> (2012)
		Fg slurry (in-house, pH 5.6)		2.99		Petersen <i>et al.</i> (submitted)
	Fg slurry (in-store, pH 6.6)		2.94		Petersen <i>et al.</i> (submitted)	

¹ '1' represents higher emissions (%) and '-' lower emissions (%) compared with the reference (untreated) manure. The comparison of systems is based on CO₂ equivalents.

¹N₂O emissions from untreated slurry were not measurable.

(Sommer *et al.*, 2000; Petersen *et al.*, 2013). As explained above, surface crusts also develop a potential for CH₄ oxidation, although the importance of this process is not known at present. However, significant stimulation may be feasible as the half saturation constant for this process is three to four orders of magnitude above atmospheric concentrations (Petersen and Ambus, 2006; Duan *et al.*, 2013). Reported values (Table 3) show that covering slurry with either a solid cover, straw or a natural surface crust results in lower CH₄ emissions, higher N₂O emissions and, in general, a reduction of overall GHG emissions, when compared with uncovered slurry. Emissions of CH₄ and N₂O were higher when using straw instead of a solid cover (Amon *et al.*, 2006), but in a related study a straw crust in combination with a solid cover on the store gave the lowest emissions (Clemens *et al.*, 2006), possibly by increasing CH₄ availability to MOB as discussed by Petersen *et al.* (2013).

Treatment technologies – manure separation

Manure separation is a process whereby a fraction of slurry particles is isolated by one of several mechanical separation processes (Burton, 2007). Storage of the liquid fraction may result in lower N₂O emissions than untreated slurry (and higher potential NH₃ emissions) if crust formation is prevented. However, N₂O emissions from the solid fraction during storage can be high (Fangueiro *et al.*, 2008), and thus overall N₂O emissions

during storage may increase significantly after separation without additional measures (Table 4). Separate storage of the liquid and solid fractions after manure separation have in most cases, but not always, resulted in lower CH₄ emissions (Table 4). Similarly, combined CH₄ and N₂O emissions from storage of both separation products have usually, but not always, been lower than from untreated manure (cf. Dinuccio *et al.*, 2008; Mosquera *et al.*, 2011). This indicates that slurry separation requires additional measures to achieve GHG mitigation during subsequent storage. The efficiency of covering for both solid and liquid fractions was discussed above, but anaerobic digestion of the solid fraction is also an option (Sutaryo *et al.*, 2012).

Treatment technologies – anaerobic (co-)digestion

Anaerobic (co-)digestion is a treatment technology specifically designed to optimise methanogenesis from manure and other residues. During the process, easily degradable organic matter in manure and other organic substrates is transformed into biogas (mainly CO₂ and CH₄). Besides energy substituting fossil fuel, this treatment reduces the potential for CH₄ emissions during subsequent storage, but an enriched methanogenic microflora in digested slurry will continue to produce CH₄ at high rates during the cooling phase (Sommer *et al.*, 2000). It is important that CH₄ is collected during this phase, or a significant part of potential

GHG mitigation may be lost. Available data (e.g., Table 4) show a reduction in CH₄, and in combined CH₄ and N₂O emissions from storage of digested manure compared with untreated cattle slurry. Anaerobic digestion will reduce the amount of C recycled to soil, but a new study has found that the long-term stabilisation of manure C in the soil is the same at 12% to 14% irrespective of pre-treatment (Thomsen *et al.*, 2012).

Treatment technologies – aeration

Amon *et al.* (2006) reported a reduction in CH₄ emission (by 57%), an increase in N₂O emission (by 144%) and a decrease in total GHG emissions (by 43%) with aeration of cattle slurry. Martinez *et al.* (2003) reported reductions in CH₄ emissions of 70% to 99% after aeration of pig slurry. The overall potential for loss of N as NH₃ or denitrification products will be high, and N₂O emissions as high as 19% of total N in pig slurry have been reported (see Chadwick *et al.*, 2011, for references). Hence, measures to conserve N during aeration may be needed to ensure GHG mitigation via this treatment.

Treatment technologies – additives

Chemical additives have been evaluated that change the chemical environment of slurry and thereby prevent unwanted transformations. Martinez *et al.* (2003) reported reductions in CH₄ emission of 47% to 64% by different chemical additives in pig slurry. In Denmark, slurry acidification to a pH around 6 by sulphuric acid is increasingly used as an NH₃ mitigation strategy. In 2012, around 10% of the total slurry volume was acidified by one of several technologies, which adjust slurry pH either in slurry channels, in the store before spreading, or during spreading. Interestingly, acidification by sulphuric acid has also been found to reduce CH₄ emissions from cattle slurry by 67% to 87% (Petersen *et al.*, 2012), and from pig slurry by 94% to 99%, (S. Q. Petersen; unpublished results) during 3-month storage periods. Mechanisms of inhibition likely involve sulphur transformations, as significant CH₄ mitigation is also achieved with sulphate or reduced S amendment alone (Petersen *et al.*, 2012), and with even a moderate reduction of pH by sulphuric acid (Table 4). Mitigation of CH₄ emissions is of course achieved only if slurry is acidified before storage.

Land spreading – application method, rate and timing

Emissions of CH₄ after land spreading of manures are insignificant (Collins *et al.*, 2011) relative to the large losses from manure storage and enteric fermentation. Measures to reduce N₂O emissions after land spreading include choice of application method, optimising rate and timing of application to match crop requirements. Emissions of NH₃ can be minimised by proper selection of application method, but different studies (e.g., Vallejo *et al.*, 2005; Velthof *et al.*, 2010) have shown that this may instead stimulate direct N₂O emissions. A stimulation of N₂O is not always observed, possibly owing to a complex interaction with soil type and soil moisture (Thomsen *et al.*, 2010). According to the Tier 1 approach of the IPCC guidelines (IPCC, 2006), direct N₂O

emissions after land spreading are independent of crop and soil type, and of the type of manure applied, but this is clearly not the case (Van Groenigen *et al.*, 2004; Mosquera *et al.*, 2007; Velthof *et al.*, 2010). When both direct and indirect (because of NH₃ emissions and NO₃²⁻ leaching) N₂O emissions are considered, the choice of manure application technique appears to have little impact on overall N₂O emissions (Velthof *et al.*, 2010). On the other hand, a reduction of N losses will potentially increase crop yields and thereby reduce emissions per unit product. Apparently, N₂O emissions increase curvi-linearly when N application rates exceed crop N requirements (Van Groenigen *et al.*, 2004; Cardenas *et al.*, 2010). Similarly, proper timing of application has been shown to influence both direct and indirect N₂O emissions after land spreading of manures (Weslien *et al.*, 1998; Chambers *et al.*, 2000; Thorman *et al.*, 2007). These various observations suggest that, by optimising application method, rate and timing of manure application relative to the need of growing crops, N₂O emissions from land spreading may be kept to a minimum.

Use of nitrification inhibitors (NI)

Synthetic NI have been developed to promote plant N uptake by reducing losses via NO₃²⁻ leaching or denitrification. Effects of NI on crop yields are mostly moderate or absent, although better in situations with a high potential for loss of plant-available N (Nelson and Huber, 1992; Subbarao *et al.*, 2006). In recent years, research has refocussed to mainly consider effects of NI on both direct and indirect (via NO₃²⁻ leaching) N₂O emissions from N amendments to soil (e.g., Di and Cameron, 2012). In the main, NI appear successful at reducing N₂O emissions from urine (Zaman and Nguyen, 2012), fertilisers (Ding *et al.*, 2011) and livestock slurries (Dittert *et al.*, 2001; Vallejo *et al.*, 2006) in a range of climates and soil types throughout the world, although their efficacy varies between studies (Akiyama *et al.*, 2010). It is not entirely clear what controls this inconsistency. The fact that laboratory studies (e.g., Hatch *et al.*, 2005) tend to report greater inhibition of N₂O than field studies (Dittert *et al.*, 2001) suggests that soil conditions, such as variations in temperature or leaching/runoff after excessive rainfall, reduces the effect of NI. The most well-documented compounds, that is, nitrapyrin (N-serve), dicyandiamide (DCD) and 3,4-dimethylpyrazol phosphate (DMPP), have very different properties with respect to volatility and water solubility (Subbarao *et al.*, 2006). Soil temperature is important for the effect of NI, and the efficiency declines linearly above 10°C owing to the combined effect of higher nitrification rates and more rapid NI degradation (Subbarao *et al.*, 2006). Zaman and Nguyen (2012) measured greater efficacy of DCD to reduce N₂O emissions following urine deposition in autumn than in spring, which they attributed to the cooler autumn temperatures. Ongoing research should elucidate the controlling factors, but the additional challenge will be in developing cost-effective strategies for the use of NI, either via N-containing amendment themselves or directly to the soil.

Table 5 On the basis of experimental data and literature data, GHG balances for three basic diets, with and without a supplementary source of fat were calculated

		Early grass		Late grass		Maize		
		Control	High-fat	Control	High fat	Control	High fat	
A.	Enteric fermentation	L CH ₄ /kg DMI	28.9	27.4	31.9	31.0	26.5	25.1
	Net effect of high-fat diet	g CO ₂ Eq/kg DMI		2 25.1		2 15.1		2 23.5
B.	Anaerobic digestion	L CH ₄ /kg DMI	53.7	63.2	58.5	76.9	72.0	80.5
	Substitution, natural gas ¹	g CO ₂ Eq/kg DMI		2 14.4		2 27.9		2 12.9
	Post-digestion storage ³	L CH ₄ /kg DMI	3.8	4.4	4.1	5.4	5.0	5.6
		g CO ₂ Eq/kg DMI		11.1		21.6		9.9
C.	Liquid manure storage	L CH ₄ /kg DMI	5.4	6.3	5.9	7.7	7.2	8.0
	Effect of high-fat diet ²	g CO ₂ Eq/kg DMI		15.9		30.8		14.2
	Combined effect, A1 B	g CO ₂ Eq/kg DMI		2 28.4		2 21.4		2 26.4
	Combined effect, A1 C	g CO ₂ Eq/kg DMI		2 9.2		15.8		2 9.2

Effects of feeding, anaerobic digestion and storage on CH₄ emissions and energy substitution are expressed as CO₂ equivalents (bold). Negative numbers indicate a reduction in emissions. Source of experimental data: Brask *et al.* (submitted).

¹Substitution of natural gas, assuming 39.7 MJ/kg CH₄, 57 g CO₂/MJ natural gas and 80% conversion efficiency.

²A 30% reduction compared with untreated dairy slurry is assumed (Nielsen *et al.*, 2010).

³Estimated assuming a methane conversion factor, of 10% (Nielsen *et al.*, 2010).

On-farm interactions

GHG mitigation measures may influence several gases and may influence emissions at other stages of manure management. Negative interactions occur when efforts to mitigate an emission lead to higher emissions of other pollutants, or of the same pollutant at a different stage; this is referred to as pollution swapping (Monteny *et al.*, 2006). The trade-off between NH₃ and N₂O in connection with land spreading was already mentioned. However, interactions may also be positive. In a recent study, Brask *et al.* (submitted) fed dairy cattle one of three basic diets, with or without a supplement of rapeseed to increase the fat intake. Across a lactation period, they observed a reduction in enteric CH₄ emissions with all three diets (Table 5), but there was also an increase in the CH₄ production potential (b₀) of faeces from cattle on the high-fat diets. The calculations in Table 5 suggest that combining a high-fat diet for cattle with anaerobic digestion of the manure (A1 B) could maximise GHG mitigation, representing a positive interaction. On the other hand, the example also implies that, in the absence of biogas treatment, high-fat diets for cattle could lead to higher CH₄ emissions from the manure during storage (A1 C), resulting in a partial or complete loss of any mitigation achieved for CH₄ from enteric fermentation. According to Møller *et al.* (2012), this negative interaction will be compounded in warmer climates where the CH₄ conversion factor for liquid manure storage is much higher than the 10% used in the example of Table 5 (IPCC, 2006).

GHG mitigation at the farm level

It is clear from the previous sections that effects of GHG mitigation measures must be evaluated at the farm level to account for effects on C and/or N flows and associated GHG emissions in other farm components (e.g., Schils *et al.*, 2007).

Several examples of on-farm interactions were already presented above. Whole-farm modelling can reduce the need for costly experimental verification and may capture potential trade-offs in emissions, also relative to land use management and local conditions (climate and soil), and thus help predict overall environmental impact.

Model complexity – pros and cons

Farm models simulate flows and losses of N and C (including GHG emissions) associated with manure management using either emission factors, empirical equations, or process-oriented mechanisms, with principles and complexity depending on their origin and scope. In order to describe the various on-farm interactions, models need a mass balance approach where nutrient, water and matter is calculated at each step as the difference between inputs and outputs, and where processes at one stage depend on what happened at previous stages.

Farm models based on empirical or semi-empirical principles have their individual strengths and limitations. For example, MELODIE (Chardon *et al.*, 2012) translates general farm objectives and constraints into an activity plan for the year without detailed user-input data, and the model has been successfully used to simulate GHG emissions from manure handling activities in pig production systems (Rigolot *et al.*, 2010a and 2010b). Other farm models include a comprehensive calculation of GHG emissions at the manure management level (including biogenic C emissions), for example, DAIRYGHG (Rotz *et al.*, 2010) and FARMGHG (Clesen *et al.*, 2006), or they are developed to assess trade-offs with other ecosystem services, for example, SMS_{DAIRY} (Del Prado *et al.*, 2011), or economic performance, for example, DAIRYMSE (Schils *et al.*, 2006). To the best of our knowledge, there is no single farm model that represents all stages of manure management with a full and detailed process-based approach. MANURE-DNDC (Li *et al.*, 2012) is a new, mostly mechanistic approach based

on the DNDC biogeochemical model for soils (Li *et al.*, 1992) that was recently developed to simulate GHG emissions in the soil–plant–manure system.

While fully mechanistic approaches offer greater robustness in capturing effects and interactions, and flexibility in comparing systems, they are also sometimes difficult to parameterise, and data requirements may limit their use. On the other hand, models that are over-simplified may be imprecise if not accounting for, for example, management practices and climatic conditions. Hence, there is a need for balanced system-based models that can effectively account for farmer practices and local climate while complexity and parameterisation is still manageable. In general terms, whole-farm models should be evaluated for their ability to simulate temporal, spatial and genetic variability, coupling of water, nutrients and energy flows and transformations, farmer decision making and economics, indirect emissions, and uncertainty of results. For manure management, it is important that the nutrient feedback loops are simulated; manure N applied to soil influences the productivity where feed is grown, and is itself an output from the animal feed intake. Del Prado *et al.* (2013) discuss in more detail the requirements of whole-farm models for simulating GHG mitigation in ruminant systems.

Effects of climate

Within and between regions, there are large gradients in climatic conditions, and farm models have different output sensitivity to factors such as temperature and precipitation. To the best of our knowledge, models involved in GHG mitigation all ignore the direct effect of climate (e.g., heat stress) on animal performance and metabolism. Following excretion, models may consider volume and simulate the dilution caused by rainfall (e.g., MELDIE). Modelled C and N transformations in manure, and derived losses, are generally related to temperature using empirical equations, for example, CH₄ from manure storage may be based on an Arrhenius equation (FARMGHG), or use a more process-based approach where chemical thermodynamics and kinetics are controlled by a group of environmental factors including temperature, moisture and redox potential (MANURE-DNDC).

Model applications

Validation of whole-farm models is difficult, and generally partial validations are performed for individual farm components. Alternatively, models may be evaluated for specific scenarios to assess their usefulness. Several studies have looked at GHG mitigation measures affecting emissions from manure management. Li *et al.* (2012), using MANURE-DNDC, predicted a 17% decrease in N excretion, and a 13% decrease in total GHG emissions, after reducing the CP content of a dairy cow diet from 18% to 15%. This model does not predict a change in the urine N-to-dung N ratio; however, according to, for example, Panetta *et al.* (2006), an increase in animal N retention will reduce mainly urine N excretion and therefore total ammoniacal N flows. This in turn may be predicted to reduce overall GHG emissions per unit of product, mainly because of an increase in animal

N use efficiency and reductions in N₂O and NH₃ emissions at the different stages of manure management. Hence, the GHG mitigation potential of reducing CP in the diet could be even higher than predicted by MANURE-DNDC. If CP reduction is achieved by replacing grass for maize (or another arable crop), there is, however, concern that soil carbon may be lost as CO₂ owing to ploughing of existing grassland (Vellinga and Hoving, 2011). Some authors have proposed the use of mechanistic models of rumen function (Dijkstra *et al.*, 2011), but generally they are too complex for integration in a farm scale model (Del Prado *et al.*, 2013).

Whole-farm model simulations have shown that mitigation measures that do not involve changes in manure management practices may still have implications for manure-derived GHG emissions. For example, Schils *et al.* (2007), evaluating several models, predicted that a reduction in the length of the grazing period would give a modest decrease in GHG emissions because of a decrease in N₂O emissions from N deposited during grazing, and because of a decrease in CH₄ from enteric fermentation as a result of better feed quality, although there is also an increase in CH₄ emissions from manure stores and a reduction in potential soil C storage. Del Prado *et al.* (2013) compared three UK farms and concluded that, relative to confinement, half-year grazing had slightly lower, but extensive grazing considerably higher GHG emissions, both on area and product basis. It was further concluded that soil conditions and climate will greatly affect the potential for N₂O emissions from N deposition during grazing, hampering the development of general recommendations for mitigation.

An important application of whole-farm models is for the evaluation of GHG mitigation strategies that include measures adopted simultaneously for manure management and other farm components. This may be useful for identifying potential synergies or incompatible measures. Li *et al.* (2012) used the MANURE-DNDC model to simulate effects of three different measures, that is, reducing dietary CP (AL-1), covering the lagoon (AL-2), and replacing corn and soya with alfalfa (AL-3) and their combined effect (AL-4), see Figure 1. A reduction of GHG emissions (30% total, and 16% per unit of product) was estimated, although mainly because of an increase in soil C sequestration, and despite a decline in animal productivity and enhanced CH₄ emissions from the lagoon when covered.

Whole-farm models need to account for effects of manure treatment. In the example of Table 5, a high-fat diet for cattle in combination with anaerobic digestion of the manure resulted in a positive interaction with respect to GHG mitigation, whereas storage without treatment of manure from cattle on a high-fat diet could outbalance or even reverse GHG mitigation achieved for enteric fermentation. There is evidently a need to verify such basic calculations experimentally via feeding-storage experiments, and to evaluate the importance of these effects at the farm level via modelling. Del Prado *et al.* (2010) tested increased dietary fat intake as one of eight mitigation measures applied in different combinations for dairy systems using SIMS_{DAIRY}.




Figure 1 Contribution of different sources to annual GHG emissions from a dairy farm as determined with the whole-farm model Manure-DNDC. Emissions of CH₄ and N₂O were calculated for each source with a baseline and alternative management scenarios and expressed (a) as total emissions (t CO₂ Eq/year), and (b) relative to the farm productivity in meat and milk (kg CO₂ Eq/kg product). The alternative management scenarios were: Reduced dietary crude protein (AL-1), covering of lagoon (AL-2), replacing corn/soya with alfalfa (AL-3) and a combination of all these measures (AL-4). Adapted from Li *et al.* (2012).

The inclusion of fat supplements in the diet reduced overall GHG emissions by 8% to 14%, and the adoption of additional measures to improve plant N use efficiency, such as the use of NI, enabled maximum GHG mitigation of about 45%. However, SIMS_{DAIRY} did not account for changes in manure composition because of a high-fat diet, or for consequences of unsaturated fat supplementation on rumen function and DM intake. Some models simulate energy production by anaerobic digestion of manure, but not effects of this treatment on C and N transformations in the soil after field application (e.g., FARMGHG), or only account for the changed proportion of ammoniacal N in the digestate (SIMS_{DAIRY}). Thus, besides more experimental evidence for effects of mitigation measures, there is also a need for further model development to simulate the multiple effects of GHG mitigation measures and strategies.

How to achieve GHG mitigation?

The previous sections have highlighted the diversity of manure environments and management practices encountered across

regions. Considering also the complex regulation of microbial processes behind CH₄ and N₂O emissions, GHG mitigation for manure management is clearly a significant challenge. The economic and cultural context of livestock production varies greatly between regions, and this is also true for opportunities to improve manure management for GHG mitigation. While models suggest that part-time grazing could lead to reductions in GHG emissions compared with confinement under business-as-usual scenarios, the magnitude of on-farm emissions and emissions from excretal returns to pastures will depend on local conditions with respect to climate and soil type. In addition, land resources for agriculture are finite and under pressure from population growth and increasing demands for food and feed production; thus, in many cases, increasing access to grazing may not be an option. Instead, we believe that key to GHG mitigation is *containment* of nutrients by limiting leakage and atmospheric losses, as the dosing of nutrient cycles also serves to prevent direct and indirect GHG emissions. Integrated crop and livestock production (mixed farming) has been suggested to improve nutrient use efficiency and reduce environmental impacts under both North American (Russelle *et al.*, 2007), European (Ryschawy *et al.*, 2012) and tropical conditions (Ogburn and White, 2011). In this paper, via four selected cases, we have argued that in subsistence farming a main priority must be to improve nutrient use efficiency for increasing crop yields, for example via improved storage conditions and more targeted use of manure nutrients for crop production. Losses of C and N during traditional composting may be considerable, and options for anaerobic storage should be explored. Where farm effluents are to some extent 'wasted' by direct discharge into water courses, infrastructure is required to enable farmers to store livestock manures. Policy intervention via legislation, and by provision of subsidies, may be needed to ensure this. Containment is also an issue in large-scale intensive livestock production, where NH₃ emissions in particular represent a threat to natural environments and human health (Sutton *et al.*, 2011).

Also key to GHG mitigation is *constraining inputs* for food and feed production. The imbalance between nutrients in excretal returns from livestock and the land available for manure recycling can be a significant challenge, in developing countries, as well as in regions where livestock production is already highly intensified, as spreading of manure N in excess of crop requirements increases the potential for environmental losses, including emissions of NH₃, N₂O and other N compounds. For example, a reduction in average surplus N from 175 to 123 kg/ha was achieved by Danish agriculture between 1980 and 2004 by adoption of fertiliser plans that take the availability of N in manure into account, together with measures to reduce environmental losses during storage and field application (Kjellingsbæk and Hansen, 2007).

Projected changes in livestock numbers by 2050 were recently published in a study by Bouwman *et al.* (2012), see Figure 2. They include dramatic increases in South and Central America (cattle), Africa (cattle, sheep/goats) and South Asia (cattle, pigs, sheep/goats). Around 75% of the




Figure 2 Global animal stocks for 1900, 1950, 2000 and 2050 for cattle (a), pigs (b), and sheep and goats (c). Adapted from Bouwman *et al.* (2012).

increase in livestock production is foreseen to occur by intensification (IAASTD, 2009). As described in previous sections, intensification is characterised by a higher degree of animal confinement, and a shift towards liquid manure management with more opportunities for containment of nutrients, manure treatment and nutrient recycling. From an economic perspective, intensive livestock production is also characterised by a greater cash flow, and hence better opportunities to fund investments in facilities and treatment technologies for improved manure management. These all suggest that the greatest GHG mitigation potentials, also in many developing countries, will be associated with new or expanding CAFOs, and regulations may be easier to enforce in these livestock production systems.

GHG emission inventories are to a large extent based on annual emission factors, which do not capture effects of all management options discussed above. Another major difficulty is the large variability in manure characteristics behind average emission factors, which makes the effect of GHG mitigation efforts highly uncertain. Therefore, modelling is required for more precise estimates of GHG emissions and mitigation potentials. It is important to identify models with sufficient responsiveness towards factors controlling the microbial processes leading to CH₄ and N₂O emissions,

notably temperature and oxygen status, but at the same time with a (limited) data requirement that will allow wide adoption. Evidently, microbial activities in manure during storage will be affected by local climatic conditions, and currently the documentation of GHG emissions is strongly biased towards temperate climatic conditions, as reflected in the reference list to this paper. Future research should fill the gap of documenting GHG emissions from livestock manure as influenced by management in warmer climates, and efforts should continue to develop models that can support the evaluation of GHG mitigation strategies across climate zones.

Acknowledgements

Contributions of S O Petersen, M. Blanchard and N. Edouard were supported by the AnimalChange project funded by the European Community (FP7/2007–2013) under the grant agreement no. 266018. The contributions of S O Petersen and S G Sommer were supported by the Danish Strategic Research Council project no. 12-132631 (BioChain). A. del Prado would like to thank the Spanish National R1 DI i Plan (grant no. CGL2009-10176) and Basque Department of Education, Universities and Research (grant no. FC2010-33A) for funding part of this study.

This paper was published as part of a supplement to *animal*, publication of which was supported by the Greenhouse Gases & Animal Agriculture Conference 2013. The papers included in this supplement were invited by the Guest Editors and have undergone the standard journal formal review process. They may be cited. The Guest Editors appointed to this supplement are R. J. Dewhurst, D. R. Chadwick, E. Charmley, N. M. Holden, D. A. Kenny, G. Lanigan, D. Moran, C. J. Newbold, P. O’Kely, and T. Yan. The Guest Editors declare no conflict of interest.

References

Akiyama H, Yan X and Yagi K 2010. Evaluation of effectiveness of enhanced-efficiency fertilizers as mitigation options for N₂O and NO emissions from agricultural soils: a meta-analysis. *Global Change Biology* 16, 1837–1846.

Ambus P and Petersen SO 2005. Oxidation of ¹³C-labeled methane in surface crusts of pig- and cattle slurry. *Isotopes in Environmental and Health Studies* 41, 125–133.

Amon B, Amon T, Boxberger J and Alt C 2001. Emissions of NH₃, N₂O and CH₄ from dairy cows housed in a farmyard manure tying stall (housing, manure storage, manure spreading). *Nutrient Cycling in Agroecosystems* 60, 103–113.

Amon B, Kryvoruchko V, Amon T and Zechmeister-Boltenstern S 2006. Methane, nitrous oxide and ammonia emissions during storage and after application of dairy cattle slurry and influence of slurry treatment. *Agriculture, Ecosystems and Environment* 112, 153–162.

Amon B, Kryvoruchko V, Fröhlich M, Amon T, Pöllinger A, Mösenbacher I and Hausleitner A 2007. Ammonia and greenhouse gas emissions from a straw flow system for fattening pigs: housing and manure storage. *Livestock Science* 112, 199–207.

Anon. 2007. Valoriser les savoirs locaux sur l’intégration agriculture élevage pour une gestion durable des écosystèmes des savanes subhumides de l’Afrique. *Projet Agriculture-Élevage de Duras* (DCG 2-50). Bobo-Dioulasso (Burkina Faso), CIRDES Rapport Final: 99.

Anon. 2010. Biogas barometer – November 2010. *Le journal des énergies renouvelables* No. 200, pp. 104–119. Retrieved from <http://www.eurobserv-er.org/downloads.asp>

Anon. 2012. Fertiliser manual (RB209), 8th edition. TSO, Norwich. 252pp. Retrieved from <http://www.defra.gov.uk/publications/2011/03/25/fertiliser-manual-rb209/>

- Baggs EM, Smales CL and Bateman EJ 2010. Changing pH shifts the microbial source as well as the magnitude of N_2O emission from soil. *Biology and Fertility of Soils* 46, 793–805.
- Berg W, Brunsch R and Pазiczki I 2006. Greenhouse gas emissions from covered slurry compared with uncovered during storage. *Agriculture, Ecosystems and Environment* 112, 129–134.
- Blanchard M 2010. Gestion de la fertilité des sols et rôle du troupeau dans les systèmes coton-céréales-élevage au Mali-Sud, savoirs techniques locaux et pratiques d'intégration agriculture élevage (Soil fertility management and herd role in the cotton-cereals-livestock systems in south Mali, technical local knowledge and crop-livestock farming practices). Thèses, UPEC.
- Blanchard M and Vall E 2010. Production et utilisation de la fumure organique au Mali-Sud: savoirs des paysans face aux savoirs des agronomes, quels enseignements pour le développement? FA Thibaud Bénédicte. Paris, Karthala, pp. 59–76.
- Bouwman L, Goldewijk KK, Van Der Hoek KW, Beusen AHW, Van Vuuren DP, Willmers J, Rufino MC and Stehfest E 2012. Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900–2050 period. *Proceedings of the National Academy of Sciences*. DOI: 10.1073/pnas.1012878108.
- Brask M, Lund P, Hellwing ALF, Poulsen M and Weisbjerg MR. Enteric methane production, digestibility and rumen fermentation in dairy cows fed different forages with and without rapeseed fat supplementation. Submitted.
- Bruun S, Jensen LS, Vu Thi Khanh Van and Sommer SG 2013. Small-scale household biogas digesters: an option for global warming mitigation or a potential climate bomb? Policy Analysis, submitted.
- Burton C 2007. The potential contribution of separation technologies to the management of livestock manure. *Livestock Science* 112, 208–216.
- Burton CH and Turner C 2003. Manure management. Treatment strategies for sustainable agriculture, 2nd edition. Silsoe Research Institute, Bedford, UK.
- Cabaraux JF, Philippe FX, Laitat M, Canart B, Van den Heede M and Nicks B 2009. Gaseous emissions from weaned pigs raised on different floor systems. *Agriculture, Ecosystems and Environment* 130, 86–92.
- Cardenas LM, Thorman R, Ashlee N, Butler M, Chadwick DR, Chambers B, Quttie S, Donovan N, Kingston H, Lane S and Scholefield D 2010. Quantifying annual N_2O emission fluxes from grazed grassland under a range of inorganic fertiliser nitrogen inputs. *Agriculture, Ecosystems and Environment* 136, 218–226.
- Chadwick DR 2005. Emissions of ammonia, nitrous oxide and methane from cattle manure heaps: effect of compaction and covering. *Atmospheric Environment* 39, 787–799.
- Chadwick D, Qing C, Yan'an T, Guanghui Y and Qirong S 2012. Improving manure nutrient management towards sustainable intensification in China. A Policy Brief No. 6, UK-China Sustainable Agriculture Innovations Network (SAIN), 6pp.
- Chadwick D, Sommer SG, Thorman R, Fanguero D, Cardenas L, Amon B and Misselbrook T 2011. Manure management: implications for greenhouse gas emissions. *Animal Feed Science and Technology* 166–167, 514–531.
- Chambers BJ, Smith KS and Pain BF 2000. Strategies to encourage better use of nitrogen in animal manures. *Soil Use and Management* 16, 157–161.
- Chardon X, Rigolot C, Baratte C, Espagnol S, Raison C, Martin-Clovaire R, Rellier JP, Le Gall A, Dourmad JY, Flouquet B, Leterme P, Paillat JM, Delaby L, Garcia F, Peyraud JL, Poupau JC, Morvan T and Faverdin P 2012. MELODIE a whole-farm model to study the dynamics of nutrients in dairy and pig farms with crops. *Animal* 6, 1711–1721.
- Chen Y, Cheng JJ and Creamer KS 2008. Inhibition of anaerobic digestion process: a review. *Bioresource Technology* 99, 4044–4064.
- Clemens J, Trimbom M, Weiland P and Amon B 2006. Mitigation of greenhouse gas emissions by anaerobic digestion of cattle slurry. *Agriculture, Ecosystems and Environment* 112, 171–177.
- Collins HP, Alva AK, Streubel JD, Fransen SF, Frear C, Chen S, Kruger C and Granatstein D 2011. Greenhouse gas emissions from an irrigated silt loam soil amended with anaerobically digested dairy manure. *Soil Science Society of America Journal* 75, 2206–2216.
- Qu TTT, Pham HC, Le TH, Nguyen VC, Le XA, Nguyen XT and Sommer SG 2012. Manure management practices on biogas and non-biogas pig farms in developing countries using livestock farms in Vietnam as an example. *Journal of Cleaner Production* 27, 64–71.
- Quhel J, Smek M, Laughlin RJ, Bru D, Chêneby D, Watson CJ and Philippot L 2010. Insights into the effect of soil pH on N_2O and N_2 emissions and denitrifier community size and activity. *Applied and Environmental Microbiology* 76, 1870–1878.
- Cytryn E, Levkovitch I, Negreanu Y, Dowd S, Frenk Sand Silber A 2012. Impact of short-term acidification on nitrification and nitrifying bacterial community dynamics in soilless cultivation media. *Applied and Environmental Microbiology* 78, 6576–6582.
- De Boer W and Kowalchuk GA 2001. Nitrification in acid soil: microorganisms and mechanisms. *Soil Biology and Biochemistry* 33, 853–866.
- De Vrieze J, Hennebel T, Boon N and Verstraete W 2012. Methanosarcina: the rediscovered methanogen for heavy duty biomethanation. *Bioresource Technology* 112, 1–9.
- Del Prado A, Crosson P, Olesen JE and Rotz A 2013. Whole-farm models to quantify GHG emissions and their potential use for linking climate change mitigation and adaptation in ruminant systems. *Animal (suppl. s2)*, 373–385.
- Del Prado A, Chadwick D, Cardenas L, Misselbrook T, Scholefield D and Merino P 2010. Exploring systems responses to mitigation of GHG in UK dairy farms. *Agriculture, Ecosystems and Environment* 136, 318–332.
- Del Prado A, Misselbrook T, Chadwick D, Hopkins A, Dawhurst RJ, Davison P, Butler A, Schröder J and Scholefield D 2011. SIMSDAIRY: a modelling framework to identify sustainable dairy farms in the UK. Framework description and test for organic systems and N fertiliser optimisation. *Science of The Total Environment* 409, 3993–4009.
- Di HJ and Cameron KC 2012. How does the application of different nitrification inhibitors affect nitrous oxide emissions and nitrate leaching from cow urine in grazed pastures? *Soil Use and Management* 28, 54–61.
- Dijkstra J, Oenema O and Bannink A 2011. Dietary strategies to reducing N excretion from cattle: implications for methane emissions. *Current Opinion in Environmental Sustainability* 3, 414–422.
- Ding WX, Yu HY and Cai ZC 2011. Impact of urease and nitrification inhibitors on nitrous oxide emissions from fluvo-aquic soil in the North China Plain. *Biology and Fertility of Soils* 47, 91–99.
- Dinuccio E, Berg W and Balsari P 2008. Gaseous emissions from the storage of untreated slurries and the fractions obtained after mechanical separation. *Atmospheric Environment* 42, 2448–2459.
- Dittert K, Bol R, King R, Chadwick D and Hatch D 2001. Use of a novel nitrification inhibitor to reduce nitrous oxide emissions from ^{15}N -labelled slurry injected into soil. *Rapid Communications in Mass Spectrometry* 15, 1291–1296.
- Duan YF, Elsgaard L and Petersen SO 2013. Inhibition of methane oxidation in slurry surface crust by inorganic nitrogen: an incubation study. *Journal of Environmental Quality* 42, 507–515.
- European Commission (EC) 1991. Council Directive of 12 December 1991 concerning the protection of waters against pollution caused by nitrates from agricultural sources. Directive 91/676/EEC, Brussels.
- Edouard N, Charpiot A, Hassouna M, Faverdin P, Robin P and Dollé JB 2012. Ammonia and greenhouse gases emissions from dairy cattle buildings: slurry vs farm yard manure management systems. International Symposium on Emission of Gas and Dust from Livestock. INRA, Saint-Malo, France, p. 31.
- Fanguero D, Coutinho J, Chadwick D, Moreira N and Trindade H 2008. Cattle slurry treatment by screw-press separation and chemically enhanced settling: effect on greenhouse gases and ammonia emissions during storage. *Journal of Environmental Quality* 37, 2322–2331.
- Food and Agriculture Organization of the United Nations (FAO) 2010. Greenhouse gas emissions from the dairy sector. A life cycle assessment. Rome, Italy, 94pp. Retrieved from <http://www.fao.org/docrep/012/k7930e/k7930e00.pdf>.
- Forster P, Ramaswamy V, Artaxo P, Bernsten T, Betts R, Fahey DW, Haywood J, Lean J, Lowe DC, Myhre G, Nganga J, Prinn R, Raga G, Schulz M and Van Dorland R 2007. Changes in atmospheric constituents and in radiative forcing. In *Climate change 2007: The physical science basis. Contribution of working group I to the fourth assessment report of the Intergovernmental Panel on Climate Change* (ed. S Solomon, D Qin, M Manning, Z Chen, M Marquis, KB Averyt, M Tignor and HL Miller), pp. 129–234. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Fukumoto Y 2006. Reduction of nitrous oxide emission from pig manure composting by addition of nitrite-oxidizing bacteria. *Environmental Science and Technology* 40, 6787–6791.
- Gac A, Béline F, Bioteau T and Maguet K 2007. A French inventory of gaseous emissions (CH_4 , N_2O , NH_3) from livestock manure management using a mass-flow approach. *Livestock Science* 112, 252–260.
- Gany F, Feller C, Harmand JM and Guilbert H 2001. Management of soil organic matter in semiarid Africa for annual cropping systems. *Nutrient Cycling in Agroecosystems* 61, 105–118.

- Gao JJ, Bai XL, Zhuo B, Zhou JB and Chen ZJ 2012. Soil nutrient content and nutrient balances in newly-built solar greenhouses in northern China. *Nutrient Cycling and Agroecosystems* 94, 63–72.
- Gerber P, Chilonda P, Franceschini G and Menzi H 2005. Geographical determinants and environmental implications of livestock production intensification in Asia. *Bioresource Technology* 96, 263–276.
- Goreau TJ, Kaplan WA, Wofsky SC, McElroy MB, Valois FW and Watson SW 1980. Production of NO_2^- and N_2O by nitrifying bacteria at reduced concentrations of oxygen. *Applied and Environmental Microbiology* 40, 526–532.
- Groenestein CM, Smits MCJ, Huijsmans JFM and Oenema O 2011. Measures to reduce ammonia emissions from livestock manures; now, soon and later. Wageningen UR Livestock Research Report 488. Wageningen, 53 pp.
- Groenestein K, Mosquera J and Van der Suis S 2012. Emission factors for methane and nitrous oxide from manure management and mitigation options. *Journal of Integrative Environmental Sciences* 9 (suppl. 1), 139–146.
- Guarino M, Fabbri C, Brambilla M, Valli L and Navarotto P 2006. Evaluation of simplified covering systems to reduce gaseous emissions from livestock manure storage. *Transactions of the ASABE* 49, 737–747.
- Häussermann A, Hartung E, Callmann E and Jungbluth T 2006. Influence of season, ventilation strategy, and slurry removal on methane emissions from pig houses. *Agriculture, Ecosystems and Environment* 112, 115–121.
- Halet D, Boon N and Verstraete W 2006. Community dynamics of methanotrophic bacteria during composting of organic matter. *Journal of Bioscience and Bioengineering* 101, 297–302.
- Hansen MN, Henriksen K and Sommer SG 2006. Observations of production and emission of greenhouse gases and ammonia during storage of solids separated from pig slurry: effects of covering. *Atmospheric Environment* 40, 4172–4181.
- Hansen RR, Nielsen DA, Schramm A, Nielsen LP, Revsbech NP and Hansen MN 2009. Greenhouse gas microbiology in wet and dry straw crust covering pig slurry. *Journal of Environmental Quality* 38, 1311–1319.
- Hanson RS and Hanson TE 1996. Methanotrophic bacteria. *Microbiology Reviews* 60, 439–471.
- Hao X, Chang C, Lamey FJ and Travis GR 2001. Greenhouse gas emissions during cattle feedlot manure composting. *Journal of Environmental Quality* 30, 376–386.
- Hatch D, Trinidade H, Cardenas L, Carneiro J, Hawkins J, Scholfield D and Chadwick D 2005. Laboratory study of the effects of two nitrification inhibitors on greenhouse gas emissions from a slurry-treated arable soil: impact of diurnal temperature cycle. *Biology and Fertility of Soils* 41, 225–232.
- Hindrichsen IK, Wettstein HR, Madmüller A, Jörg B and Kreuzer M 2005. Effect of the carbohydrate composition of feed concentrates on methane emissions from dairy cows and their slurry. *Environmental Monitoring and Assessment* 107, 329–350.
- Hoffmann I, Gerling D, Kyogwom UB and Mané-Biefeldt A 2001. Farmers' management strategies to maintain soil fertility in a remote area in northwest Nigeria. *Agriculture, Ecosystems and Environment* 86, 263–275.
- Hoogwijk M, Faaij A, Eickhout B, de Vries B and Turkenburg W 2005. Potential of biomass energy out to 2100, for four IPCC SRES land-use scenarios. *Biomass and Bioenergy* 29, 225–257.
- Husted S 1994. Seasonal variation in methane emission from stored slurry and solid manures. *Journal of Environmental Quality* 23, 585–592.
- Husted S, Jensen LS and Jørgensen SS 1991. Reducing ammonia loss from cattle slurry by the use of acidifying additives: the role of the buffer system. *Journal of the Science of Food and Agriculture* 57, 335–349.
- IAASTD 2009. Agriculture at a crossroads Global report. In International assessment of agricultural knowledge, science and technology for development (ed. BD McIntyre, HR Herren, J Wakhungu and RT Watson), 590pp. IAASTD, Washington DC. Retrieved from <http://www.unep.org/dewa/Assessments/Ecosystems/IAASTD/tabid/105853/Default.aspx/>.
- IPCC 2006. Guidelines for national greenhouse gas inventories. Agriculture, forestry and other land use, vol. 4. Intergovernmental Panel on Climate Change. IGES, Hayama, Kanagawa, Japan.
- Ja Z and Conrad R 2009. Bacteria rather than archaea dominate microbial ammonia oxidation in an agricultural soil. *Environmental Microbiology* 11, 1658–1671.
- Jiang X, Sommer SG and Christensen KV 2011. A review of the biogas industry in China. *Energy Policy* 39, 6073–6081.
- Kaiser S 1999. Analyse und bewertung eines zweiraumkompoststalls für matschweine unter besonderer berücksichtigung der gasförmigen stoffströme. PhD Thesis, VDI-MEG Schrift 334, University of Göttingen, Germany.
- Kool DM, Dolfing J, Wrage N and Van Groenigen JW 2011. Nitrifier denitrification as a distinct and significant source of nitrous oxide from soil. *Soil Biology and Biochemistry* 43, 174–178.
- Kruska RL, Reid RS, Thornton FK, Henninger N and Kristjanson PM 2003. Mapping livestock-oriented agricultural production systems for the developing world. *Agricultural Systems* 77, 39–63.
- Kyllingsbæk A and Hansen JF 2007. Development in nutrient balances in Danish agriculture 1980–2004. *Nutrient Cycling in Agroecosystems* 79, 267–280.
- Laanbroek HJ and Gerards S 1991. Effects of organic manure on nitrification in arable soils. *Biology and Fertility of Soils* 12, 147–153.
- Landais E and Guérin H 1992. Systèmes d'élevage et transferts de fertilité dans la zone des savanes africaines. I. La production des matières fertilisantes. *Cahiers Agriculture* 1, 225–238.
- Landais E and Lhoste P 1993. Systèmes d'élevage et transferts de fertilité dans la zone des savanes africaines. II. Les systèmes de gestion de la fumure animale et leur insertion dans les relations entre l'élevage et l'agriculture. *Cahiers Agriculture* 2, 9–25.
- Lantinga EA, Keuning JA, Groenwold J and Deenan PJAG 1987. Distribution of excreted nitrogen by grazing cattle and its effects on sward quality, herbage production and utilization. In *Animal manure on grassland and fodder crops. Fertilizer or waste?* (ed. HG van der Meer, RJ Unwin, TA van Dijk and GC Ennik), pp. 103–107. Martinus Nijhoff, Dordrecht.
- Leip A, Weiss F, Wassenaar T, Perez I, Fellmann T, Loudjani P, Tubiello F, Grandgirard D, Monni S and Biala K 2011. Evaluation of the Livestock Sector's Contribution to the EU Greenhouse Gas Emissions (GGELS). European Commission Joint Research Centre. Web. 5 May 2011. Retrieved from http://ec.europa.eu/agriculture/analysis/external/livestock-gas/full_text_en.pdf
- Li C, Frolking S and Frolking TA 1992. A model of nitrous oxide evolution from soil driven by rainfall events: 1. Model structure and sensitivity. *Journal of Geophysical Research* 97, 9759–9776.
- Li C, Salas W, Zhang R, Krauter C, Rotz A and Mitloehner F 2012. Manure-DNDC: a biogeochemical process model for quantifying greenhouse gas and ammonia emissions from livestock manure systems. *Nutrient Cycling in Agroecosystems* 93, 163–200.
- Manlay RJ, Chotte JL, Masse D, Laurent JY and Feller C 2002. Carbon, nitrogen and phosphorus allocation in agro-ecosystems of a West African savanna: III. Plant and soil components under continuous cultivation. *Agriculture, Ecosystems and Environment* 88, 249–269.
- Martinez J, Guizou F, Feu P and Gueutier V 2003. Influence of treatment techniques for pig slurry on methane emissions during subsequent storage. *Biosystems Engineering* 85, 347–354.
- Mennicken L 1998. Biobett für Legehennen – ein Beitrag zum Umweltschutz? [Biobeds for laying hens – a contribution to environmental protection?] *DGS* 13, 12–20.
- Menzi H, Oenema O, Burton C, Shipin Q, Gerber P, Robinson T and Franceschini G 2010. Impacts of intensive livestock production and manure management on the environment. In *Livestock in a changing landscape, vol. 1: drivers, consequences, and responses* (ed. H Steinfeld, HA Mooney, F Schneider and LE Neville), pp. 139–163. Island Press, Washington, DC, USA.
- MEP 2010. Ministry of Environmental Protection, Bulletin of National Environmental Statistics 2009 MEP, Beijing (in Chinese).
- Minamikawa K and Sakai N 2006. The practical use of water management based on soil redox potential for decreasing methane emission from a paddy field in Japan. *Agriculture, Ecosystems and Environment* 116, 181–188.
- Misselbrook TH, Chadwick DR, Pain BF and Headon DM 1998. Dietary manipulation as a means of decreasing N losses and methane emissions and improving herbage N uptake following application of pig slurry to grassland. *The Journal of Agricultural Science* 130, 183–191.
- Misselbrook TH, Powell JM, Broderick GA and Grabber JH 2005. Dietary manipulation in dairy cattle: laboratory experiments to assess the influence on ammonia emissions. *Journal of Dairy Science* 88, 1765–1777.
- MCA 2010. The Chinese agricultural statistical data in 2009. China Agricultural Publishing House, Beijing.
- Monteny GJ, Bannink A and Chadwick D 2006. Greenhouse gas abatement strategies for animal husbandry. *Agriculture, Ecosystems and Environment* 112, 163–170.
- Mosquera J, Höl JMG, Rappoldt C and Dolfing J 2007. Precise soil management as a tool to reduce CH_4 and N_2O emissions from agricultural soils. *Animal Sciences Group Report* 28, 42 pp.

- Mosquera J, Schils R, Groenestein CM, Hoeksma P, Velthof G and Hummelink E 2011. Emissies van lachgas, methaan en ammoniak uit mest na scheiding. Wageningen UR Livestock Research Report 427.
- Møller HB, Sarker S, Hellwin ALF and Weisbjerg MR 2012. Quantification of methane production and emission from anaerobic digestion of cattle manure derived from different feeding. Paper presented at 3rd Int. Conf. CIGR, July 8–12 in Valencia, Spain. Retrieved from http://cigr.ageng2012.org/images/fotos/tabla_137_C0327.pdf
- Nelson DW and Huber D 1992. Nitrification inhibitors for corn production. National Corn Handbook, NHC-55.
- Nicks B, Laitat M, Van den Heede M, Désiron A, Verhaeghe C and Canart B 2003. Emissions of ammonia, nitrous oxide, methane, carbon dioxide and water vapor in the raising of weaned pigs on straw-based and sawdust-based deep litter. *Animal Research* 52, 299–308.
- Nicks B, Laitat M, Farnir F, Van den Heede M, Désiron A, Verhaeghe C and Canart B 2004. Gaseous emissions from deep-litter pens with straw or sawdust for fattening pigs. *Animal Science* 78, 99–107.
- Nielsen DA, Schramm A, Nielsen LP and Revsbech NP 2013. Seasonal methane oxidation potential in manure crusts. *Applied and Environmental Microbiology* 79, 407–410.
- Nielsen OK, Lyck E, Mikkelsen MH, Hoffmann L, Gydenkæne S, Winther M, Nielsen M, Fauser P, Thomsen M, Plejdrup MS, Albrektsen R, Hjelgaard K, Johansen VK, Vesterdal L, Rasmussen E, Arfaoui K and Baunbæk L 2010. Denmark's National Inventory Report 2010. Emission Inventories 1990–2008 – Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. National Environmental Research Institute, Aarhus University, Denmark. 1178pp. – NER Technical Report No 784. Retrieved from <http://www.dmu.dk/Pub/FR784.pdf>.
- Oenema Q, Kros H and de Vries W 2003. Approaches and uncertainties in nutrient budgets: implications for nutrient management and environmental policies. *European Journal of Agronomy* 20, 3–16.
- Oenema Q, Oudendag D and Velthof GL 2007. Nutrient losses from manure management in the European Union. *Livestock Science* 112, 261–272.
- Ogburn DM and White I 2011. Integrating livestock production with crops and saline fish ponds to reduce greenhouse gas emissions. *Journal of Integrative Environmental Analyses* 8, 39–52.
- Olesen JE, Schelde K, Weiske A, Weisbjerg MR, Asman WAH and Djurhuus J 2006. Modelling greenhouse gas emissions from European conventional and organic dairy farms. *Agriculture, Ecosystems and Environment* 112, 207–220.
- Osada T, Kuroda K and Yonaga M 2000. Determination of nitrous oxide, methane, and ammonia emissions from a swine waste composting process. *Journal of Material Cycles and Waste Management* 2, 51–56.
- Panetta DM, Powers WJ, Xin H, Kerr BJ and Stalder KJ 2006. Nitrogen excretion and ammonia emissions from pig fed modified diets. *Journal of Environmental Quality* 35, 1297–1308.
- Pattey E, Trzinski MK and Desjardins RL 2005. Quantifying the reduction of greenhouse gas emissions as a result of composting dairy and beef cattle manure. *Nutrient Cycling in Agroecosystems* 72, 173–187.
- Petersen SO and Ambus P 2006. Methane oxidation in pig and cattle slurry storages, and effects of surface crust moisture and methane availability. *Nutrient Cycling in Agroecosystems* 74, 1–11.
- Petersen SQ, Amon B and Gattinger A 2005. Methane oxidation in slurry storage surface crusts. *Journal of Environmental Quality* 34, 455–461.
- Petersen SQ, Andersen AJ and Eriksen J 2012. Effects of slurry acidification on ammonia and methane emission during storage. *Journal of Environmental Quality* 41, 88–94.
- Petersen SQ, Nielsen AL, Haarder K and Henriksen K 1992. Factors controlling nitrification and denitrification: a laboratory study with gel-stabilized liquid cattle manure. *Microbial Ecology* 23, 239–255.
- Petersen SQ, Nielsen TH, Frostegård Å and Olesen T 1996. O_2 uptake, C metabolism and denitrification associated with manure hot-spots. *Soil Biology and Biochemistry* 28, 341–349.
- Petersen SQ, Dorno N, Lindholm S, Feilberg A and Eriksen J 2013. Emissions of CH_4 , N_2O , NH_3 and odorants from pig slurry during winter and summer storage. *Nutrient Cycling in Agroecosystems* 95, 103–113.
- Petersen SQ, Højberg Q, Poulsen M, Schwab C and Eriksen J. Methane mitigation and methanogen community changes with acidification of pig slurry. Submitted.
- Peyraud JL, Cellier P, Donnars C and Réchauchère O 2012. Les flux d'azote liés aux élevages, réduire les pertes, rétablir les équilibres. Expertise scientifique collective, synthèse du rapport. INRA (France), pp. 1–68.
- Philippe FX, Laitat M, Canart B, Van den Heede M and Nicks B 2007. Gaseous emissions during the fattening of pigs kept either on fully slatted floors or on straw flow. *Animal* 1, 1515–1523.
- Philippe FX, Laitat M, Wavreille J, Bartiaux-Thill N, Nicks B and Cabaraux JF 2011. Ammonia and greenhouse gas emission from group-housed gestating sows depends on floor type. *Agriculture, Ecosystems and Environment* 140, 498–505.
- Prapaspongsa T 2010. Sustainable piggery waste management: A study based on examples and cases from Denmark and Thailand. PhD thesis, Aalborg University, Aalborg, Denmark, ISBN 978-87-90033-67-5.
- Rastogi G, Ranade DR, Yeole TY, Gupta AK, Patole MS and Shouche YS 2008. Molecular analyses of methanogen diversity associated with cattle dung. *World Journal of Microbiology and Biotechnology* 24, 2973–2979.
- Rigolot C, Espagnol S, Pomar C and Dourmad J-Y 2010a. Modelling of manure production by pigs and NH_3 , N_2O and CH_4 emissions. Part I: animal excretion and enteric CH_4 , effect of feeding and performance. *Animal* 4, 1401–1412.
- Rigolot C, Espagnol S, Robin P, Hassouna M, Béline F, Paillat JM and Dourmad J-Y 2010b. Modelling of manure production by pigs and NH_3 , N_2O and CH_4 emissions. Part II: effect of animal housing, manure storage and treatment practices. *Animal* 4, 1413–1424.
- Rotz CA 2004. Management to reduce nitrogen losses in animal production. *Journal of Animal Science* 82, E119–E137.
- Rotz CA, Montes F and Chianese DS 2010. The carbon footprint of dairy production systems through partial life cycle assessment. *Journal of Dairy Science* 93, 1266–1282.
- Rufino MC and Rowe EC 2006. Nitrogen cycling efficiencies through resource-poor African crop-livestock systems. *Agriculture, Ecosystems and Environment* 112, 261–282.
- Rufino MC, Tittone P, van Wijk MT, Castellanos-Navarrete A, Delve RJ, de Ridder N and Giller KE 2007. Manure as a key resource within smallholder farming systems: analysing farm-scale nutrient cycling efficiencies with the NUANCES framework. *Livestock Science* 112, 273–287.
- Russelle MP, Entz MH and Franzuebbers AJ 2007. Reconsidering integrated crop-livestock systems in North America. *Agronomy Journal* 99, 325–334.
- Ryszchawy J, Choisis N, Choisis JP, Jannon A and Gibon A 2012. Mixed crop-livestock systems: an economic and environmental-friendly way of farming? *Animal* 6, 1722–1730.
- Schils RLM, Olesen JE, del Prado A and Soussana JF 2007. A review of farm level modelling approaches for mitigating greenhouse gas emissions from ruminant livestock systems. *Livestock Science* 112, 240–251.
- Schils RLM, Verhagen A, Aarts HFM, Kuikman PJ and Šebek LBJ 2006. Effect of improved nitrogen management on greenhouse gas emissions from intensive dairy systems in the Netherlands. *Global Change Biology* 12, 382–391.
- Schröder J 2005. Revisiting the agronomic benefits of manure: a correct assessment and exploitation of its fertilizer value spares the environment. *Bioresource Technology* 96, 253–261.
- Sharma R, Ryan K, Hao X, Lamey FJ, McAllister TA and Topp E 2011. Real-time quantification of *mcrA*, *pmoA* for methanogen, methanotroph estimations during composting. *Journal of Environmental Quality* 40, 199–205.
- Sommer SG, Petersen SO and Sægaard HT 2000. Greenhouse gas emission from stored livestock slurry. *Journal of Environmental Quality* 29, 744–750.
- Sommer SG, Petersen SO and Møller HB 2004. Algorithms for calculating methane and nitrous oxide emissions from manure management. *Nutrient Cycling in Agroecosystems* 69, 143–154.
- Sommer SG, Olesen JE, Petersen SQ, Weisbjerg MR, Valli L, Rohde L and Béline F 2009. Region-specific assessment of greenhouse gas mitigation with different manure management strategies in four agroecological zones. *Global Change Biology* 15, 2825–2837.
- Steinfeld H, Gerber P, Wassenaar T, Castel V, Rosales M and de Haan C 2006. Livestock's long shadow: environmental issues and options. Food and Agriculture Organization, Rome, Italy.
- Subbarao GV, Ito Q, Sahrawat KL, Berry WL, Nakahara K, Ishikawa T, Watanabe T, Suenaga K, Rondon M and Rao IM 2006. Scope and strategies for regulation of nitrification in agricultural systems – challenges and opportunities. *Critical Reviews in Plant Sciences* 25, 303–335.

- Sutaryo S, Ward A and Møller HB 2012. Thermophilic anaerobic co-digestion of separated solids from acidified dairy cow manure. *Bioresource Technology* 114, 195–200.
- Sutton MA, Oenema O, Erisman JW, Leip A, van Grinsven H and Winiwarter W 2011. Too much of a good thing. *Nature* 472, 159–161.
- Szanto GL, Hamelers HMM, Rulkens WH and Veeken AHM 2007. NH_3 , N_2O and CH_4 emissions during passively aerated composting of straw-rich pig manure. *Bioresource Technology* 98, 2659–2670.
- Thomsen IK, Pedersen AR, Nyord T and Petersen SO 2010. Effects of slurry pre-treatment and application technique on short-term N_2O emissions as determined by a new non-linear approach. *Agriculture, Ecosystems and Environment* 136, 227–235.
- Thomsen IK, Olesen JE, Møller HB, Sørensen P and Christensen BT 2012. Carbon dynamics and retention in soil after anaerobic digestion of dairy cattle feed and faeces. *Soil Biology and Biochemistry* 58, 82–87.
- Thomson AJ, Giannopoulos G, Pretty J, Baggs EM and Richardson DJ 2012. Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philosophical Transactions of the Royal Society B* 367, 1157–1168.
- Thorman RE, Chadwick DR, Harrison R, Boyles LO and Matthews R 2007. The effect on N_2O emissions of storage conditions and rapid incorporation of pig and cattle farmyard manure into tillage land. *Biosystems Engineering* 97, 501–511.
- Thorman RE, Chadwick D, Boyles LQ, Matthews R, Sagoo E and Harrison R 2006. Nitrous oxide emissions during storage of solid manure and following application to arable land. In Proceedings of the 12th RAMIRAN Conference, Aarhus, Denmark, 11–13 September (ed. SO Petersen). Danish Institute of Agricultural Sciences, Tjele, Denmark.
- Tittonell P, Rufino MC, Janssen BH and Giller KE 2010. Carbon and nutrient losses during manure storage under traditional and improved practices in smallholder crop-livestock systems—evidence from Kenya. *Plant and Soil* 328, 253–269.
- Umetsu K, Kimura Y, Takahashi J, Kishimoto T, Kojima T and Young B 2005. Methane emission from stored dairy manure slurry and slurry after digestion by methane digester. *Animal Science Journal* 76, 73–79.
- Valentine DL 2007. Adaptations to energy stress dictate the ecology of the archaea. *Nature Reviews – Microbiology* 5, 316–323.
- Vall E, Dugué P and Blanchard M 2006. Le tissage des relations agriculture-élevage au fil du coton. *Cahiers Agricultures* 15, 72–79.
- Vallejo A, García-Torres L, Díez JA, Arce A and López-Fernández S 2006. Comparison of N losses (NO_3^- , N_2O , NO) from surface applied, injected or amended (DOD) pig slurry of an irrigated soil in a Mediterranean climate. *Plant and Soil* 272, 313–325.
- Vallejo A, Skiba UM, García-Torres L, Arce A, López-Fernández and Sánchez-Martín L 2006. Nitrogen oxides emission from soils bearing a potato crop as influenced by fertilization with treated pig slurries and composts. *Soil Biology and Biochemistry* 38, 2782–2793.
- Van Groenigen JW, Kasper GJ, Velthof GL, van den Pol-van Dassel A and Kuikman PJ 2004. Nitrous oxide emissions from silage maize fields under different mineral nitrogen fertilizer and manure applications. *Plant and Soil* 263, 101–111.
- VanderZaag AC, Gordon RJ, Jamieson RC, Burton DL and Stratton GW 2009. Gas emissions from straw covered liquid dairy manure during summer storage and autumn agitation. *Transactions of the ASABE* 52, 599–608.
- Vellinga TV and Hoving IE 2011. Maize silage for dairy cows: mitigation of methane emissions can be offset by land use change. *Nutrient Cycling in Agroecosystems* 89, 413–426.
- Velthof GJ, Mosquera J, Huis in't Veld JWH and Hummelink E 2010. Effect of manure application technique on nitrous oxide emission from agricultural soils. Alterra report 1992, Wageningen.
- Vu QD, Tran TM, Nguyen PD, Vu CC, Vu VTK and Jensen LS 2012. Effect of biogas technology on nutrient flows for small- and medium-scale pig farms in Vietnam. *Nutrient Cycling in Agroecosystems* 94, 1–13.
- Vu TKV, Tran MT and Dang TTS 2007. A survey of manure management on pig farms in Northern Vietnam. *Livestock Science* 112, 288–297.
- Wallenstein MD, Myrold DD, Firestone M and Voytek M 2006. Environmental controls of denitrifying communities and denitrification rates: insights from molecular methods. *Ecological Applications* 16, 2143–2152.
- Wang CB and Zhang LX 2012. Life cycle assessment of carbon emission from a household biogas digester: implications for policy. *Proceedings in Environmental Science* 13, 778–789.
- Webb J, Sommer SG, Kupper T, Groenestein K, Hutchings NJ, Eurich-Menden B, Rodhe L, Misselbrook TH and Amon B 2012. Emissions of ammonia, nitrous oxide and methane during the management of solid manures. *Sustainable Agriculture Reviews* 8, 67–107.
- Weslien P, Klemmedtsson L, Svensson L, Galle B, Kasimir-Klemmedtsson A and Gustafsson A 1998. Nitrogen losses following application of pig slurry to arable land. *Soil Use and Management* 14, 200–208.
- Wilshusen JH, Hettiaratchi JA, De Visscher A and Saint-Fort R 2004. Methane oxidation and formation of EPS in compost: effect of oxygen concentration. *Environmental Pollution* 129, 305–314.
- Wood JD, Gordon RJ, Wagner-Riddle C, Dunfield KE and Madani A 2012. Relationships between dairy slurry total solids, gas emissions, and surface crusts. *Journal of Environmental Quality* 41, 694–704.
- Yamamoto N, Otawa K and Nakai Y 2010. Diversity and abundance of ammonia-oxidizing bacteria and ammonia-oxidizing archaea during cattle manure composting. *Microbial Ecology* 60, 807–815.
- Yamulki S 2006. Effect of straw addition on nitrous oxide and methane emissions from stored farmyard manures. *Agriculture, Ecosystems and Environment* 112, 140–145.
- Zaman M and Nguyen ML 2012. How application timings of urease and nitrification inhibitors affect N losses from urine patches in pastoral system. *Agriculture, Ecosystems and Environment* 156, 37–48.
- Zhang W-F, Dou Z, He P, Ju X-T, Powlson D, Chadwick D, Norse D, Lu Y-L, Zhang Y, Wu L, Chen X-P, Cassman KG and Zhang F-S 2013. New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China. *Proceedings of the National Academy of Sciences* (in press).

