



Adult stem cells in adipose tissue and skeletal muscle of pigs differing in body composition

Marie-Hélène Perruchot, Louis Lefaucheur, Florence Gondret, Isabelle Louveau

► To cite this version:

Marie-Hélène Perruchot, Louis Lefaucheur, Florence Gondret, Isabelle Louveau. Adult stem cells in adipose tissue and skeletal muscle of pigs differing in body composition. 65. Annual Meeting of the European Federation of Animal Science (EAAP), Aug 2014, Copenhagen, Denmark. Wageningen Academic Publishers, Annual Meeting of the European Association for Animal Production, 20, 2014, Annual Meeting of the European Association for Animal Production. hal-01210672

HAL Id: hal-01210672

<https://hal.science/hal-01210672>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Book of Abstracts of the 65th Annual Meeting of the European Federation of Animal Science



**Book of abstracts No. 20 (2014)
Copenhagen, Denmark
25 - 29 August 2014**

Session 02

Poster 8

mRNA abundance of the lactate receptor HCA1 in two different adipose tissue depots of dairy cows

M. Weber¹, L. Locher², K. Huber³, J. Rehage², S. Dänicke⁴, H. Sauerwein¹ and M. Mielenz^{1,5}

¹University of Bonn, Institute of Animal Science, Katzenburgweg 7, 53115 Bonn, Germany, ²University of Veterinary Medicine, Clinic for Cattle, Bischofsholer Damm 15, 30173 Hannover, Germany, ³University of Veterinary Medicine, Institute of Physiology, Bischofsholer Damm 15, 30173 Hannover, Germany, ⁴Friedrich Loeffler Institute, Institute of Animal Nutrition, Bundesallee 50, 38116 Braunschweig, Germany, ⁵Present address: Leibniz Institute for Farm Animal Biology, Institute of Nutritional Physiology, Wilhelm-Stahl-Allee 2, 18196 Dummerstorf, Germany; sauerwein@uni-bonn.de

The G-protein coupled receptor HCA1 mediates antilipolytic effects via binding lactate, an intermediate of energy metabolism. HCA1 is mainly located on adipocytes and enables them to respond to metabolic changes through fine-tuning of lipolysis. Thus we hypothesized that different amounts of concentrate in the diet, resulting in differences in the energy balance, will affect the HCA1 mRNA abundance in bovine adipose tissue (AT). Furthermore, we aimed to determine whether the mRNA abundance between subcutaneous (SC) and retroperitoneal (RP) AT differs among late pregnancy and lactation. Biopsy samples of SCAT and RPAT from 20 pluriparous Holstein Friesian cows were collected at day (d) -42, 1, 21 and 100 relative to parturition. Until d -42 all cows were fed the same silage-based diet. Between d -42 and d -1, 10 animals each were assigned to either a high-concentrate (HC, 60:40 concentrate:roughage) or a low concentrate group (LC, 30:70). From d 1 to 24 the concentrate proportion was increased from 30 to 50% for all cows. HCA1 mRNA abundance was not affected by treatment, but showed a time-dependent increase in SCAT from d -42 to d 100 ($P < 0.05$). In RPAT, no time-dependent changes were observed. Comparing the fat depots, HCA1 mRNA abundance was higher in RPAT than in SCAT on days -42 and 21 ($P < 0.001$). The presented results indicate that HCA1 mRNA is regulated independently of the feeding regimen. Differences between SCAT and RPAT suggest differences in the lipolytic regulation.

Session 02

Poster 9

Adult stem cells in adipose tissue and skeletal muscle of pigs differing in body composition

M.H. Perruchot, L. Lefaucheur, F. Gondret and I. Louveau

INRA, UMR 1348 Pegase, Domaine de la Prise, 35590 Saint-Gilles, France;
marie-helene.perruchot@rennes.inra.fr

The control of body composition is of the utmost importance for animal production. It is now recognized that adult stem cells are essential for tissue homeostasis. The current study was undertaken to determine whether changes in the relative proportion of muscle and adipose tissue mass in pigs are controlled by the plasticity of adult stem cells. Between 10 and 20 weeks of age, 24 Large White barrows from two lines divergently selected for residual feed intake (RFI), an indicator of feed efficiency, were fed *ad libitum* either a high-starch (LF) diet or a high-fat/high-fiber (HF) diet ($n=6$ per line and diet). At 20 weeks of age, stroma-vascular cells were isolated by collagenase treatment from subcutaneous adipose tissue and longissimus muscle. Cell immuno-phenotypes were then determined by flow cytometry using cell surface markers (CD34, CD56 and CD90). Irrespective of diet, low RFI pigs were leaner than high RFI pigs. In the two lines, pigs fed the HF diet exhibited a reduced body fatness compared with pigs fed the LF diet. Whatever the experimental group considered, adipose and muscle cells were predominantly positive for CD56 (muscle precursor marker) and CD90 (mesenchymal stem cell marker) markers. Cells expressing the CD34 hematopoietic stem cell marker were identified in muscle but not in adipose tissue. The proportion of CD90+ and CD56+ cells in adipose tissue did not differ between the experimental groups. In muscle, the proportion of CD34+ cells was higher ($P < 0.005$), whereas that of CD90+ cells was lower ($P < 0.005$) in the low than in the high RFI pigs, whatever the diet. The current study indicates that divergent selection for RFI in growing pigs had strongly affected the relative proportions of adult stem cell populations in skeletal muscle but not in adipose tissue. Altogether, there is no simple relationship between the proportion of adult stem cells and body adiposity.