Synergies and trade-offs in sustainable dairy farming systems: a novel efficiency approach



A.D. Soteriades¹, P. Faverdin^{2, 3}, M. March⁴ and A.W. Stott¹

- ¹ SRUC, West Mains Road, Edinburgh, EH9 3JG, Scotland, Email: andreas.soteriades@sruc.ac.uk
- ² INRA, UMR1080, Production du Lait, F-35590 St-Gilles, France
- ³ Agrocampus Ouest, UMR1080, Production du Lait, F-35000 Rennes, France
- ⁴ Scotland's Rural College, Hestan House, The Crichton Business Park, Dumfries, DG1 4TA, UK






Introduction

World food production must increase to meet greater future demand without exacerbating climate change and despite dwindling resources. Appropriate measures of efficiency are therefore essential for farms to become- and remain- economically, socially and environmentally sustainable. This study used a slacks-based measure of efficiency (SBM) (Tone, 2001) within a Data Envelopment Analysis (DEA) framework to identify synergies and trade-offs between the technical (TE), environmental (EE) and economic (cost) efficiencies (CE) of dairy farms; and between four contrasting dairy systems; by using data from a long-term genetic line x feeding systems experiment (the "Langhill" experiment; Pollott and Coffey, 2008). The slacks represented farm resource/pollutant excesses and helped identify specific aspects in which each system could perform better.



Methods



Results

- ☐ Kendall's *tau* rank correlation coefficient:
- 1. TE & EE strongly correlated


 See conclusions (1st bullet)

 2. EE & CE modestly correlated


 Trade-offs between systems?
- 3. TE & CE weakly correlated
- ☐ Kruskal-Wallis test + Figure 1:
- 1. HFS **SB*** LFC & LFS for CE
- 2. HFC **SB** LFC & LFS for CE
- Unexpected!
 See Conclusions (2nd bullet)

Further inspection below

- 3. LFS & LFC **SB** HFC for Land Use & Fertilizer **SP****
- 4. HFS & LFS **SB** HFC for greenhouse gas (GHG) SP



Novel finding!
See Conclusions (4th bullet)






Figure 1. Box plots for efficiency scores (row 1) and input/undesirable output savings potentials (**SP**; row 2). SP = ratio of input/undesirable output slacks over their corresponding input/undesirable output.

Conclusions

- Synergy: Technically efficient farms also efficient in their production of pollution. Result in line with Shortall and Barnes (2013).
- Trade-offs for CE: This is dependent upon the specific cost structures and assumptions applied to the Langhill herd and so needs further investigation in commercial units.
- Significantly better performance of LF over HFC for Land Use and Fertilizer expected as LF systems housed all year round and their diet largely based on bought-in concentrates. Nevertheless, what would have been the result had land use and fertilizer applications been accounted-for regarding bought-in feeds?
- In line with Toma et al. (2013), we found that genetic merit could have differential effects on the systems' environmental performance. However the SBM indicated specific reasons for this difference (i.e. GHG emissions performance), through the disaggregation of efficiency into SP. Thus, SBMs superior to "traditional" DEA models.
- This study helps guide policy in sustainable dairy farming. Combining the SBM framework with an abundance of data available in SRUC's Langhill database, there exist numerous opportunities for further research.

References

Pollot, G.E. and Coffey, M.P. 2008. Journal of Dairy Science 91 (9), 3649-3660. Shortall, O.K. and Barnes, A.P. 2013. Ecological Indicators 29, 478-488. Toma, L., March, M., Stott, A.W. and Roberts, D.J. 2013. Journal of Dairy Science 96 (11), 7014-7031.

Tone, K. 2001. European Journal of Operational Research 130 (3), 498-509.

Acknowledgements

This project is jointly funded by Scotland's Rural College (SRUC) and the French National Institute for Agricultural Research (INRA). It is also supported by, and benefits from, Scottish Government's Strategic Research Programmes.