

Banana meal for feeding pigs: digestive utilization, growth performance and feeding behavior

David Renaudeau, Jérémy Brochain, Mario Giorgi, Bruno Bocage, M Hery, E

Crantor, Carine Marie-Magdeleine, Harry Archimède

▶ To cite this version:

David Renaudeau, Jérémy Brochain, Mario Giorgi, Bruno Bocage, M Hery, et al.. Banana meal for feeding pigs: digestive utilization, growth performance and feeding behavior. Animal, 2014, 8 (4), pp.565-571. 10.1017/S175173111400010X . hal-01210711

HAL Id: hal-01210711 https://hal.science/hal-01210711

Submitted on 28 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Banana meal for feeding pigs: digestive utilization, growth performance and feeding behavior

D. Renaudeau^{1†}, J. Brochain¹, M. Giorgi¹, B. Bocage¹, M. Hery², E. Crantor³, C. Marie-Magdeleine¹ and H. Archimède¹

¹INRA UR143, Unité de Recherches Zootechniques, F-97170 Petit Bourg, Guadeloupe, France; ²SICA Les Producteurs de Guadeloupe, F-97100 Basse Terre, Guadeloupe, France; ³Grands Moulins des Antilles, F-97122 Baie Mahault, Guadeloupe, France

(Received 14 January 2013; Accepted 19 December 2013; First published online 17 February 2014)

The main objective of the present work was to determine the nutritional value and the strategies of using green banana meal (BM) in growing pigs. Two trials involving a total of 96 growing pigs were designed to study the effect of the harvest stage on the nutritional and energy values of BM (trial 1) and to evaluate the consequence of feeding gradual levels of BM on growth performance and feeding behavior in growing pigs (trial 2). In trial 1, the digestive utilization of three diets including 40% BM were compared with a control (C) soybean meal-corn diet in two batches of 12 pigs. BM was obtained from fruits harvested at 750 degrees-days (DD; early harvesting stage), 900 DD (normal harvesting stage) and 1150 DD (late harvesting stage). In trial 2, 72 Large White pigs were grouped in pens of nine animals and were given ad libitum access to one of the four dietary treatments (two pens/diet) differing from the rate of inclusion of 900 DD BM (0%, 20%, 40%, 60%). The estimated energy apparent digestibility coefficients of BM increased with the harvest stage (75.5%, 80.7% and 83.2% for BM at 750, 900 and 1150 DD, respectively). Digestible energy and metabolizable energy values were higher for BM at 1150 DD (13.56 and 13.05 MJ/kg DM, respectively) than at 900 DD (13.11 and 12.75 MJ/kg DM, respectively) or at 750 DD (12.00 and 11.75 MJ/kg DM, respectively). In trial 2, average daily gain and feed conversion ratio were not affected (P > 0.05) by the rate of BM inclusion (822 g/day and 2.75 kg/kg on average, respectively). Feed intake and feeding behavior parameters were not significantly influenced by the dietary treatments except for the rate of feed ingestion with a lower value for the diet with 40% of BM (27.4 v. 32.2 g/min on average; P < 0.01) when compared with the other diets. Results of this study indicate that the energy value of BM increases with the harvest stage and that BM can be incorporated up to 60% in growing finishing pig diets.

Keywords: growing pigs, banana meal, energy, harvest stage

Implications

The present paper has confirmed that banana meal is a valuable energy source for feeding pigs and demonstrated that delayed harvest stage can improve the energy value of this raw material. In our experimental conditions, up to 60% of meal produced from banana harvested at 900 degrees-days can be fed to the growing-finishing pigs without any detrimental consequences on growth performance.

Introduction

Many tropical countries such as French West Indies (FWI) do not produce grain for livestock feeding and have to import it. As feed represents about 60% to 70% of total cost of production in the pig industry, the high cost of imported feed is a major constraint to the development of the local pig production especially when local meat must compete with cheap imported meat. Within tropical countries, if properly utilized, several local feed resources could greatly contribute to reduce the amount of imported raw material for livestock. Banana is an important agricultural export commodity for many tropical and subtropical countries. In Guadeloupe, bananas accounted for 50% of export earnings in 2011. In practice, bananas are harvested green, and the fruits whose guality does not meet the expected standards are rejected for export. About 15% to 20% of the banana production in the FWI is wasted (Archimède et al., 2011). These discarded bananas (i.e., from 45 000 to 60 000 T/year in the FWI) could constitute a very good source of energy for livestock. However, the high moisture content and the short conservation time of green banana make its transportation difficult, and limit the direct use by animal

[†] Present address: INRA UMR, 1348 PEGASE, F-35590 Rennes, France. E-mail: david.renaudeau@rennes.inra.fr

Renaudeau, Brochain, Giorgi, Bocage, Hery, Crantor, Marie-Magdeleine and Archimède

production units located far from the plantations. Alternatively, 'rejected' bananas could be dried and the meal could be used as a substitute of imported corn in pig feeds (Clavijo and Maner, 1975; Le Dividich et al., 1978). The main costs of the use of banana meal (BM) would be transportation, the processing and the storage since feed grade bananas can be obtained at almost no cost. There is limited information regarding the nutritional value of BM in pigs (Le Dividich and Canope, 1974; Clavijo and Maner, 1975) and this information needs to be reevaluated with actual banana cultivars and pig genotypes.

The purpose of the present work was to determine the nutritional value of green BM in growing pigs according to different harvesting stages and to quantify the effect of gradual substitution of cereals by BM on growth performance and feeding behavior in growing-finishing pigs.

Material and methods

Two trials were designed to study the effect of harvest stage on the nutritional value of BM (trial 1) and to evaluate the consequences of feeding gradual levels of BM on growth performance and feeding behavior in growing pigs (trial 2). Both trials were performed at the experimental facilities of the Institut National de la Recherche Agronomigue in Guadeloupe, FWI (latitude 16°N, longitude 61°W). Banana ('Grande Naine'; AAA genome, Cavendish subgroup) used in both trials was harvested in Guadeloupe. The time of harvesting for green banana, as to other crops, is based on the concept of thermal unit accumulation, expressed as degrees-days (DD). This method sums, from the flower emergence to harvest, the daily average temperature minus a base temperature (22°C for 'Grande Naine' cultivar). For export markets, the optimal harvesting time is 900 DD. For this study, banana, once harvested, was sliced with its peal, dried at 60°C for 3 days, using an industrial oven and ground through a 2 mm screen.

Experimental design and animal management

Trial 1 was conducted on 24 castrated Large White (LW) pigs in two batches of 12 pigs each. Within each batch, three blocks of four littermates were chosen where each littermate was randomly assigned to one of the four dietary treatments. At about 40 kg BW, pigs were moved to individual digestibility cages in an open front experimental room. Climatic parameters were not controlled and followed those of the outdoor conditions. Three diets including BM were compared with a control (C) soybean meal-corn diet. The 750 DD, 900 DD and the 1150 DD diets were obtained by substituting 40% (on a DM basis) of the C diet by BM obtained from fruits harvested at 750 DD (early harvesting stage), 900 DD (normal harvesting stage) and 1150 DD (late harvesting stage). The control diet was a commercial pig-grower feed. Experimental diets were prepared by mixing BM with the control diet, water (1:1.5, W/v), and mineral and vitamins mixture (4 g/kg) to maintain a constant level (i.e. 1%) of minerals and vitamins in the final feed. Chemical compositions of C diet and BMs are given in Table 1. The same BMs were used in the two batches. Within each replicate, pigs were adapted to

Table 1	Chemical composition of control diet ¹	and banana meals (BM)
(trial 1)		

			BM ²	
	Control	750 DD	900 DD	1150 DD
Chemical composition (% MS)				
Organic matter	94.1	94.5	95.4	95.3
Ash	5.9	5.4	4.6	4.7
CP (N×6.25)	17.2	5.0	4.5	4.0
Starch	45.9	63.5	67.3	70.0
Crude fat	8.0	1.4	1.0	1.0
Free sugars	8.4	3.6	3.9	3.3
NDF	14.3	16.6	11.8	9.3
ADF	3.5	8.4	5.8	5.1
ADL	0.4	4.4	2.7	2.2
Gross energy (MJ/kg DM)	18.18	16.90	16.89	17.01

¹Diet formulated with corn (55%), wheat middling (22%), and soybean meal

(17%). $^{2}750\ \text{DD},$ 900 DD and 1140DD, indicated as early, normal and late stages of harvesting of banana bunch.

the diet for 10 days before total collection of feces and urine for 8 days. During the adaptation and the collection periods, feed allowance was fixed at about 150 g DM/day per kg BW^{0.60}. At the beginning of the collection period, the daily DM allowance was calculated for each pig on the average BW basis, estimated form the initial BW and the average BW gain measured during the adaptation period assuming that BW gain was similar in the adaptation and the collection periods. Pigs had free access to water with drinkers designed to avoid spillage.

Trial 2 was conducted on a total 72 LW pigs (36 castrated males and 36 gilts). At 10 weeks of age, pigs were weighed and divided into eight groups of nine animals. Each group was constituted by five castrated males and four gilts or four castrated males and five gilts. While being weighed, each pig also had a transponder ear tag inserted to enable individual identification by 'Acema 64' (Acemo, Pontivy, France) feed dispenser system. Animals were moved in one of the eight pens $(5.7 \times 2.7 \text{ m})$ in an open-front fattening unit and allowed to one of the four dietary treatments (2 pens/diet) differing in the rate of inclusion of BM. These experimental diets were based on wheat, corn, soybean meal, wheat byproducts and BM (900 DD) and contained the same amount of digestible lysine and metabolizable energy (ME). We used a basal control diet without BM and diets D20, D40 and D60 containing 20%, 40% and 60% of BM, respectively at the expense of cereals. The experimental diets were processed in a commercial factory according to standard procedures. The feed was provided as pellets and was replenished every day between 0700 and 0800 h to ensure a continuous supply in the feeder. Details on the composition of the experimental diets are given in Table 2. Only four of the eight pens had feed intake recording, so pigs were switched between pens every 2 weeks after being weighed in the morning according to the procedure described by Young et al. (2011). Alternate pens were in the same room and had the same characteristics

Tab	ole 2	Composition	of	experimental	diets	(trial	2)
-----	-------	-------------	----	--------------	-------	--------	----

Items	Control	D20	D40	D60
Ingredients (g/kg)				
Corn	310.0	160.0	80.0	
Wheat	170.0	170.0	120.0	35.1
Soybean meal	240.0	260.0	285.0	311.5
Wheat middling	80.0	87.7	47.2	
Wheat bran	152.3	75.0	20.0	
BM		200.0	400.0	600.0
Soybean oil				5.0
DL-Methionine		0.2	0.6	1.0
L-Threonine			0.1	0.3
Sepiolite	10.0	10.0	10.0	10.0
Monocalcium phosphate	4.3	4.3	4.3	4.3
Calcium carbonate	21.7	21.7	21.7	21.7
Salt	1.1	1.1	1.1	1.1
Minerals and vitamins ¹	10.0	10.0	10.0	10.0
Analyzed composition (% DM)				
Ash	7.1	8.6	9.3	8.6
СР	21.5	21.0	19.5	18.2
Starch	36.0	37.7	38.9	38.6
Lipids	2.7	2.2	2.0	2.7
Free sugars				
NDF	16.2	14.1	10.0	12.1
ADF	5.0	5.0	4.9	5.1
ADL	0.7	0.9	1.3	1.4
Calculated composition (% DM) ²				
Lysine	1.08	1.08	1.06	1.06
Methionine + cystine	0.69	0.67	0.65	0.63
Threonine	0.78	0.75	0.73	0.72
Tryptophan	0.27	0.26	0.24	0.22
Calculated nutritional value ³				
SID lysine (g/kg DM)	9.4	9.4	9.4	9.4
ME (MJ/kg)	13.4	13.4	13.4	13.4

BM = banana meal; SID = standardized ileal digestible; ME = metabolizable energy.

¹Minerals and vitamins mixture supplied (g/kg of diet): 10 of Cu (as CuSo₄); 80 of Fe (as FeSO₄.7H₂O); 40 of Mn (as MnO); 100 of Zn (as ZnO); 0.6 of I (as Ca(IO₃)2); 0.10 of CO (as CoSO₄.7H₂O); 0.15 of Se (as Na₂SeO₃); 5,000 IU of vitamin A; 1000 IU of vitamin D₃; 15 IU of vitamin E; 2 mg of vitamin K₃; 2 mg of thiamin; 4 mg of riboflavin; 20 mg of nicotinic acid; 10 mg of D-panthothenic acid; 3 mg of pyroxidine; 0.02 mg of vitamin B₁₂; 1.0 mg of folic acid; and 0.2 mg of biotin.

²Estimated from INRA French tables (Sauvant *et al.*, 2002) considering that AA contents in BM is negligible.

³SID lysine and ME content were calculated from INRA Tables (Noblet et al. 2002).

(except for the feeder). Feeding behavior data from days when pigs were switched between pens were not taken into account. The experiment started after 10 days of adaptation and ended when pigs reached approximately 100 kg live BW.

Measurements

Trial 1: All the pigs were weighted at the beginning and the end of the adaptation period, and at the end of the collection period. Every morning, any feed refusal and spillage were collected and dried to measure the DM content (24 h at 103°C). One sample of each experimental diet was also taken daily and pooled at the end of the collection period for DM determination and further chemical analysis. During the 8 days collection period, feces and urine were daily and individually collected and stored at 4°C. Urine was collected with sulphuric acid (0.1 N; 10% v/v) in order to avoid ammonia losses during collection and storage. At the end of the collection period, feces and urine were weighed, homogenized and sub-sampled. One sample of feces was dried (48 h at 103°C) for DM determination and the second was freeze-dried for further analyses.

Trial 2: All the pigs were individually weighed at the beginning of the experiment, and thereafter every 2 weeks until the end of the experiment. An additional weighing was performed at the end of the experiment after a 24 h fasting period. The backfat thickness was measured ultrasonically (Agroscan; E.C.M., Angoulême, France) at the last rib at 45 mm from the midline at the end of the experiment. For pigs housed in pen without electronic feed dispenser, feed intake was measured by the difference between the feed allowed during each 2 weeks period and refusal collected the days when pigs were switched between pens. For the pens equipped with a feed dispenser, the equipment was calibrated at the start of the experiment using a 500 g test weight. Each feeding stalls allows access to only one pig at a time. After each visit to the feeder, the identity of the animal (via the ear-tag transponder), the feeder entry and exit times and the amount of feed consumed were registered and stored in a computer.

Chemical analyses

Diet samples were analyzed for DM, ash, organic matter (AOAC Method N°923.03), CP (N × 6.25; AOAC Method N° 992.15), and ether extract (AOAC Method N°945.16) (AOAC, 1990), for free sugars according to Tollier and Robin (1979) and for gross energy (GE) content using an adiabatic bomb calorimeter (C5000; IKA, Staufen, Germany). Starch content was determined using the UV method for the determination of starch in foods and other materials (Enzy Plus Starch kit Biocontrol). Dry fecal samples were analyzed for DM, GE, ash, ether extract and CP and fresh samples of urine for N. Cell wall components (NDF, ADF and ADL; AOAC Methods 200.04 and 973.18) in diet samples were determined according to the Van Soest and Wine method (1967).

Calculation and statistical analyses

From data produced in trial 1, apparent digestibility coefficients of nutrient and energy were calculated according to standard procedures. Digestible energy (DE) content was calculated as the difference between measured GE intake and energy loss in feces. ME content of diets was calculated as the difference between DE and energy loss in urine estimated from N loss (g/day) according to the equation of Le Goff and Noblet (2001; energy loss in urine (kJ/day) = 345 + $31.1 \times N$ loss in urine (g/day), $R^2 = 0.94$, r.s.d. = 110). Data were analyzed by using an ANOVA including the effects of diet, batch, block effect within each batch and their interactions each animals being considered the experimental unit. The digestive utilization and energy values of the different BMs were calculated according to the difference method assuming that the nutrient digestibility of the control diet was constant in the experimental diets.

Renaudeau, Brochain, Giorgi, Bocage, Hery, Crantor, Marie-Magdeleine and Archimède

	Control	750 DD	900 DD	1150 DD	r.s.d.	Statistics ²
Number of pigs	6	6	6	6		
Average BW (kg)	53.7	57.4	51.6	52.5	9.0	
DM intake (g/day)	1544	1652	1465	1489	168	
DM intake (g/day per kg BW ^{0.60})	142	146	138	138	10	
Average BW gain (g/day)	783 ^a	606 ^b	534 ^b	555 ^b	75	D*
Digestibility coefficients (%)						
Dry matter	88.4 ^a	84.5 ^b	86.7 ^c	87.5 ^c	1.4	D**
Organic matter	90.2 ^a	85.9 ^b	88.1 ^c	88.9 ^c	1.4	D**
Cprotein	88.4 ^a	68.3 ^b	71.4 ^{bc}	73.5 ^c	3.0	D**
Starch	99.7	99.0	99.7	99.7	0.5	
NDF	62.5ª	50.3 ^b	48.6 ^b	51.6 ^b	5.6	D**, R*
Energy	88.2 ^a	83.1 ^b	85.2 ^c	86.2 ^c	1.5	D**
Energy values (MJ/kg DM)						
Digestible energy	15.96ª	14.36 ^b	14.82 ^c	14.96 ^c	0.26	D**
Metabolizable energy	15.51ª	13.96 ^b	14.41 ^c	14.52 ^c	0.26	D**

 Table 3 Digestive utilization of experimental diets¹ (least square means)

r.s.d. = residual standard deviation.

¹750 DD, 900 DD and 1150 DD diets composing 40% banana meal and 60% of control diet.

²ANOVA, taking into account the effect of dietary treatment (D), Replicate (R), block within each replicate and their interaction. Least square means with different superscripts differed (P < 0.05) from dietary treatments. Each animals (n = 6/treatment) was the experimental unit.

P*<0.05, *P*<0.01.

To allow comparison with results from other studies, successive feeder visits were grouped into the same meal using a meal criterion defined as the maximum length of a withinmeal interval (i.e. short pauses belonging to the same meal) between visits. Visits separated by intervals shorter than the meal criterion were considered to be part of the same meal. A meal criterion was estimated using the log survivor curve technique as described by Labroue et al. (1994). The meal criterion of 2 min was established by the latter authors from LW pigs in experimental conditions similar to the present ones. According to this 2-min meal criterion, the components of feeding behavior were calculated for each pig as following: daily number of meals, daily feed intake (g), total ingestion time per day (i.e. total duration of all feeding bouts (min)), ingestion rate (i.e. total feed intake/total ingestion time (g/min)), the average feed intake per meal (total feed consumption/number of meals (g)), and the average ingestion time per meal (total ingestion time/number of meals (min)). Feeding behavior data were pooled by 2 weeks period and were submitted to an ANOVA including the effect of diet, sex, period of measurements, block and their interactions. The animal (n = 17 or 18/treatment) was considered the experimental unit. Mean growth performance (except feed conversion ratio (FCR)) were submitted to an ANOVA taking into account the effect of diet, sex, pen, block and interactions. The FCR was calculated on a pen basis for each period of measurement and computed with an ANOVA with diet, pen, and interactions as main effects. Each pen of pigs (2/treatment) was the experimental unit.

Results

Digestibility study (trial 1)

As planned, average BW and DM intake were not affected by diet composition (Table 3). During the collection period, the

ADG was significantly higher for control diet than for diets including BM (783 v. 565 g/day; P < 0.05). This can be partly explained by the reduced CP contents when compared with the control diet (12.3% on average v. 17.2%; data not shown). The inclusion of 40% of BM reduced the digestibility coefficients of nutrients and energy. This effect was related to the stage of harvesting. When compared with the control diet, the digestibility coefficients of organic matter and energy were reduced by 4.3% and 5.1%, respectively for the 750 DD diet (*P* < 0.05) and by 1.7% and 2.5% for the 900 DD and 1050 DD diets. Whatever the stage of harvesting, the digestibility coefficient of NDF was significantly lower in diets with 40% BM than in the control ones (50.2% v. 62.5%; P < 0.01). The effect of dietary treatment on DE and ME values followed the same trends as those observed for digestibility coefficients of energy. Higher DE and ME values were found for control diet (15.96 and 15.51 MJ/kg DM, respectively) than for diets with BM. Among the three other diets, DE and ME values were greater (P < 0.05) in diet 900 and 1150 DD than in diet 750 DD (Table 3).

The experimental diets were prepared to calculate digestibility coefficients of energy and energy values of BM according to different stages of harvesting. These values were calculated by difference. Data shown in Table 4 indicate that the digestibility coefficient of energy increased with the degree of maturity. It was increasing by 5.2% between 750 and 900 DD and by only 2.5% between 900 and 1150 DD. Similar results were obtained for DE and ME values (Table 4).

Growth trial (trial 2)

The chemical composition of experimental diets given in Table 2 was in agreement with the objectives of the experiment. At the beginning of the experiment, pigs were 83 days old and averaged 29.5 kg BW. Two pigs were removed at the

 Table 4 Apparent nutrient digestibility coefficients and energy values
 of banana meal according to stages of harvesting¹

	750 DD	900 DD	1150 DD
Digestibility coefficients (%)			
Organic matter	79.5	85.0	87.0
Cprotein	38.2	45.9	51.2
NDF	32.0	27.8	35.3
Energy	75.5	80.7	83.2
Energy values (MJ/kg DM)			
DE	12.00	13.11	13.56
ME	11.75	12.75	13.05
NE ²	9.22	10.10	10.40

DE = digestible energy; ME = metabolizable energy; NE = net energy.

¹Value calculated according to difference methods.

²The NE values were estimated by NEg8 equation, published by Noblet et al. (1994).

 Table 5 Effect of banana meal on performance of growing pigs (least
 square means)

Item	Control	D20	D40	D60	r.s.d.	Statistics ¹
Number of animals	17	18	17	17		
Initial BW (kg)	29.1	30.2	29.0	29.8	5.1	P**
Final BW (kg)	94.9	96.7	93.0	94.5	8.9	S*
ADG (g/day)	820	826	795	805	75	P**
FCR (kg/kg gain) ²	2.63	2.65	2.74	2.69	0.40	М*
Final backfat thickness (mm) ³	15.3	15.3	14.7	14.8	1.0	S**, P**

r.s.d. = residual standard deviation.

ANOVA taking into account the effect of dietary treatment (D), sex (S), pen (P) within dietary treatment and block. Each animal was the experimental unit. ²Adjusted for an average dry matter of 88.0% and calculated per pen for each 2 week measurement period. These data computed by ANOVA, involving the effect of dietary treatment (D), period of measurements (M), their interactions. Fach pen of pigs was the experimental unit.

³Mean values of ultra sound measurements performed on the back and shoulder. *P<0.05, **P<0.01.

beginning of the experiment because of leg problems (diets C diet and D20) and one pig died for unknown reasons after 27 days of experiment (diet D40). As shown in Table 5, the diet composition did not influence the ADG and FCR (812 g/day and 2.75 kg/kg on average, respectively; P > 0.05). The backfat thickness at the end of the experiment was similar (P = 0.19) for all diets (15.0 mm on average). The final BW and backfat thickness were found significantly higher in barrows than in gilts (97.1 v. 92.5 kg and 15.4 v. 14.7 mm, respectively). The ADG tended to be higher in barrows than in gilts (828 v. 794 q/day; P = 0.079).

During the growing period, pigs were measured three times for their individual feed intake and feeding behavior. The average BW and age for these measurements were 67.2 kg at 131 days, respectively and they did not differ among diets (Table 6). Logically, most of the feeding behavior parameters were influenced by the period of measurement in

 Table 6 Effect of banana meal on feeding behavior of growing pigs
 (least square means)

ltem	Control	D20	D40	D60	r.s.d.	Statistics ¹
Number of animals	17	18	17	17		
Mean BW (kg)	67.6	69.1	66.3	65.8	7.9	S**, M**
Mean age (day)	131	131	131	131	1	M**
No. of meals	7.4	8.1	8.3	7.6	2.7	S**, M**
Feed intake ²						
g/day	2023	2060	1930	2007	291	S**, M**
g/meal	333	333	294	328	119	M**
Feed ingestion time						
min/day	68.7	64.5	71.8	67.2	15.8	S**, M**
min/meal	10.5	10.6	10.4	10.3	5.8	
Rate of feed intake (g/min)	31.7 ^a	34.0 ^a	27.4 ^b	30.9 ^a	6.5	D**, M**

r.s.d. = residual standard deviation.

¹ANOVA taking into consideration the effect of dietary treatment (D), sex (S), period of measurement (M) and block within D. Least square means with a different superscripts differed (P < 0.05) from dietary treatment each individual pig was the experimental unit. *P < 0.05, **P < 0.01. ²Values adjusted for 88.0% dry matter.

connection with the increase of BW. The rate of feed intake was significantly lower in pigs fed D40 compared with the other diets (32.2 v. 27.4 g/min; P < 0.01). As the average daily feed intake was similar among the diets (2.0 kg/day; P = 0.150), the numerical increase in the feed ingestion time in D40 (71.8 v. 66.8 min/day; P = 0.128) was mainly related to the higher rate of feed intake. Meal characteristics were not influenced (P > 0.05) by the dietary treatment. Gilts compared with barrows made less frequent meals (7.0 v. 8.7 meals/day; P < 0.01), ate less (2132 v. 1878 g/day; P < 0.01) and spent less time eating (64.5 v. 71.6 min/day; P < 0.01).

Discussion

Whatever the stage of harvesting, the unpeeled green BM was lower in starch (67.0 v. 74.2 g/100 g DM), CP (5.0 v. 9.4 g/100 g DM), fat (1.1 v. 4.3 g/100 g DM), and subsequently lower in GE (16.9 v. 18.7 MJ/kg DM) compared to corn (Sauvant et al., 2002). The NDF content was rather similar in BM and corn (12.4 v. 10.4/100 g) but detergent fiber fractions (DF) of BM was more lignified compared with corn (25 v. 5 g lignin/100 g NDF). The low energy digestibility of BM compared with corn (75.5% to 83.2% v. 88% in corn; Noblet et al., 2002) can be mainly explained by both its reduced concentration in highly digestible nutrients (fat, starch) and the high proportions of non-fermentable fiber and lignin in the DF. According to the low energy digestibility coefficient but also to the low GE content in BM, DE and ME contents of BM was 18% to 27% lower than corn depending on the age of harvesting (Noblet et al., 2002). The DE content reported in the present study for BM (12.0MJ/kg to 13.6 MJ/kg DM with an average value of 12.8 MJ/kg DM) is rather similar to values reported in young pigs by Le Dividich and Canope (1974) (13.3 MJ/kg DM) and in growing pigs by

Renaudeau, Brochain, Giorgi, Bocage, Hery, Crantor, Marie-Magdeleine and Archimède

References		Items			Treatments		
Oliva (1970)	% of BM	0	12	24	36	48	
	ADG (g/day)	620	600	610	590	540	
	FCR (kg/kg)	4.24	4.33	4.36	4.48	5.23	
Celleri <i>et al</i> . (1971)	% of BM	0	25	50	75	_	
	ADG (g/day)	670	650	630	610	_	
	FCR (kg/kg)	3.66	3.88	4.04	4.19	_	
Restrepo and Gonzalo (1973)	% of BM	0	15	30	_	_	
	ADG (g/day)	660	720	710	_	_	
	FCR (kg/kg)	2.90	3.03	2.71	_	_	
Fomunyam (1991)	% of BM	0	10	20	30	_	
	ADG (g/day)	540	550	540	530	_	
	FCR (kg/kg)	4.20	4.21	4.33	4.39	_	
Present study	% of BM	0	20	40	60	_	
	ADG (g/day)	820	826	795	805	_	
	FCR (kg/kg)	2.63	2.65	2.74	2.69	_	

Table 7 Effect of various level of green BM on performance of growing finishing pigs: literature survey

BM = banana meal; FCR = feed conversion ratio.

Clavijo and Maner (1975) (13.4 MJ/kg DM) but it was lower that the value obtained from plantain meal by Restrepo and Gonzalo (1973) (15.1 MJ/kg DM). This higher DE value in plantains is mainly related to its greater content of starch compared with bananas (Ly, 2004; Heuzé *et al.*, 2011).

To our knowledge few papers have been published on the effect of the harvest stage on the nutritional values of BM in pigs. The present study has shown clearly that energy digestibility coefficient, DE and ME values of BM increase with the age of harvesting. Expressed as a percentage of ME measured at 750 DD, ME content increased by 8% and 10%, respectively at 900 and 1150 DD. This can be related to a change in chemical composition (more starch and less DF) with the development of the fruit and the increase in the pulp to peel ratio from 750 to 1150 DD. In particular, NDF content is an important component limiting the total tract digestibility of nutrient and energy due to the low digestibility of DF fraction and its possible interaction with other nutrients.

The first practical application of the study is to provide valuable information to help producers to make a good decision for the best harvesting stage of bananas for producing BM for feeding pigs. When compared with the standard harvest time of 900, 1150 DD delays the date of harvest by about 30 days. Practically, the yield of fruits (T DM/ha) is improved by about 15% during this time lag (Lassoudière et al., 1978) which potentially represents a gain of about 20% of energy (ME) for pigs per unit of surface. In other words, for banana plantations specifically dedicated for meal production for feeding pigs, a delayed harvesting stage at 1150 DD could be advised. However, in practice only nonmarketed fruits harvested at 900 DD are currently valued for feeding livestock. In consequence, we decided to use BM produced from bananas harvested at 900 DD for determining the optimal incorporation rate of BM for growing pigs.

In trial 2, BM was used as a substitute for cereal grain to supply 0%, 20%, 40% or 60% of the total diet for growing-finishing pigs. Based on the results obtained in trial 1, the

experimental diets were formulated in order to maintain constant ME and digestible lysine contents. Increasing the level of incorporation of BM did not have negative effect on ADG and FCR. This means that up to 60% of BM can be fed to growing/ finishing pigs without any adverse effects on growth performance. Similarly, growth performance remained constant in post-weaning pigs fed by diets with 15% or 30% of green BM (Restrepo and Gonzalo, 1973) (Table 7). However, some other works performed in Equator (Oliva, 1970), in Colombia (Celleri et al., 1971) and in Cameroon (Fomunyam, 1991) in growing finishing pigs showed a gradual decrease in both ADG and FCR when level of BM increase (Table 7). In most of the latter studies, corn has been substituted on an equal weight basis with BM leading to an overestimation of energy value of BM with negative effects on growth performance. In addition, in the present study, the amino acids supply by BM was likely negligible (Ly, 2004). Thus, the major part of digestible amino acids was provided by soybean meal and synthetic amino acids. Accordingly, it can be hypothesized that the linear depression in growth performance with increasing level of BM observed in the literature is also related to a dietary amino acid deficiency.

The inclusion of high-dietary level of BM adversely affected the pellets processing. Problems during pellets processing (low hardness and durability, resistance to flow of the mash feed through the die, etc.) have been detected by the manufacturer at the beginning of the D40 feed production. This problem concerned to only 5% on the total fabrication (200 kg on a total of 4 T). According to the limited availability of BM, this amount of feed was not discarded and it was mixed with 95% of 'normal' feed. The pelleting problem was solved by increasing the retention time of the mash feed into the steam conditioner before the pelleting processing. These changes were also applied for pelleting D60 diet. In practice, these results suggest that pelletability of diets would decrease with increased rate of BM without any changes during the processing. During feed pelleting, heat and water added in the conditioner will alter components like starch and protein in the feed mash in a way that binding properties come into effect (Abdollahi, 2011). As the gelatinization temperature of banana starch is higher than that of cereals (corn, wheat) (Ling *et al.*, 1982). more heat is required for the banana starch to be gelatinized. In other words, the pelleting problems encountered at the beginning of the D40 diet fabrication could result from an insufficient temperature in the conditioner. As a longer retention time in the conditioner indirectly increases the temperature (Stark, 2012), logically the change in this parameter in the present study has resulted to an improved pellet quality. However, the pelleting properties in diets with high level of BM have been decreased in relation to the substitution of raw materials with high binding properties (wheat products; Wood, 1987) by-products with low pelletability characteristics (BM and/or fat).

A striking result of the present study was the reduced rate of feed intake in pigs fed D40 diet. As the rate of feed intake and feed acceptance are generally positively related (Bigelow and Houpt, 1988), it could be suggested that the D40 diet was less palatable when compared with the three others diets. Solà-Oriol *et al.* (2009) suggested that feed characteristics such as texture, taste or odor may be responsible for the change in feed preference in pigs. Undesirable chemical reactions and/ or physical changes are supposed to have occurred during the process of D40 diet which affected its palatability. This hypothesis is purely speculative and remains to be confirmed by further works.

Acknowledgements

The authors would like to acknowledge the staff of the experimental station of PTEA for their technical assistance in the care of pigs, and T. Etienne, S. Calif, Y. Jaguelin for the laboratory analysis. This work was supported by the European Union (FEOGA FEDER), SICA Les Producteurs de Guadeloupe and les Grands Moulins des Antilles.

References

Abdollahi MR 2011. Influence of feed processing on the performance, nutrient utilisation, and gut development of poultry and feed quality. Thesis PhD, Massey University, New Zeland.

AOAC 1990. Official methods of analysis, 15th edition. Association of Official Analytical Chemists, Arlington, Virginia.

Archimède H, Gourdine JL, Fanchone A, Alexandre G, Marie Magdeleine C, Calif E, Fleury J, Anais C and Renaudeau D 2011. Le bananier et ses produits dans l'alimentation animale. Revue Innovations Agronomiques 16, 181–192.

Bigelow JA and Houpt TR 1988. Feeding and drinking patterns in young pigs. Physiology & Behavior 43, 99–109.

Celleri H, Oliva F and Maner JH 1971. Green banana meal for feeding growingfinishing pigs. Memoria, Associacion Latinoamericana de Produccion Animal 6, 148 (abstract).

Clavijo H and Maner J 1975. The use of waste bananas for swine feed. In Proceedings of the Conference on Animal Feeds of Tropical and Subtropical (ed. D Haliday), pp. 99–106. Tropical Products Institute, London, UK. Fomunyam RT 1991. Economic aspects of banana and plantain use in animal feeding: the Cameroon experience. In Roots, tubers and bananas in animal feeding (ed. D Machin and S Nyvold), pp. 277–289. CIAT, California, Colombia.

Heuzé V, Tran G, Archimède H, Lessire R and Renaudeau D 2011. Bananas fruits. In Feedipedia.org. A project by INRA, CIRAD, AFZ and FAO, Retrieved on 16 September 2012 from http://www.feedipedia.org/node/4670

Labroue F, Guéblez R, Sellier P and Meunier-Salaün MC 1994. Feeding behaviour of group-housed Large White and Landrace pigs in French central test stations. Livestock Production Science 40, 303–312.

Lassoudière A, Frossard P and Sarah JL 1978. Le bananier et sa culture en Côte d'Ivoire. Première partie: Connaissance de la plante. Intéraction avec le milieu écologique. Deuxième partie: Techniques culturales. Adaptation aux conditions écologiques de la bananeraie ivoirienne. GERDAT-IRFA, Abidjan, Cote d'Ivoire.

Le Dividich J and Canope I 1974. Valeur alimentaire de la farine de banane et de manioc dans le régime du porcelet sevré à 5 semaines: influence du taux de protéines de la ration. Annales de Zootechnie 23, 161–169.

Le Dividich J, Geoffroy F, Canope I and Chenost M 1978. Utilisation des déchets de banane pour l'alimentation du bétail. Revue Mondiale de Zootechnie 20, 22–30.

Le Goff G and Noblet J 2001. Comparative total tract digestibility of dietary energy and nutrients in growing pigs and adult sows. Journal of Animal Science 79, 2418–2427.

Ling LH, Osman EM, Fernandes JB and Reilly PJ 1982. Physical properties of starch from Cavendish banana fruit. Starch – Stärke 34, 184–188.

Ly J 2004. Bananas y platanos para alimentar cerdos: aspectos de la composicion quimica de las frutas y de su palatabilidad. Revista Computadorizada de Produccion Porcina 11, 5–24.

Noblet J, Seve B and Jondreville C 2002. Valeur nutritive pour le porc. In Tables de composition et de valeur nutritive des matières premières destinées aux animaux d'élevage (ed. D Sauvant, JM Perez and G Tran), pp. 25–36. INRA-AFZ, Paris.

Noblet J, Fortune H, Shi XS and Dubois S 1994. Prediction of net energy value of feeds for growing pigs. Journal of Animal Science 72, 344–354.

Oliva F 1970. Evaluacion de la harina de banano verde con cascara, en crecimiento y acabo de cerdos en confinamiento. Thesis PhD, Central University of Equator, Department of Agronomy and Veterinary Medecine, Quito, Equator.

Restrepo A and Gonzalo L 1973. Digestibilidad, valor nutritivo y energético del plátano, Musa paradisiaca L., en cerdos. Thesis PhD, Universidade National de Colombia, Bogota, Colombia.

Sauvant D, Perez JM and Tran G 2002. Tables de composition et de valeur nutritive des matières premières destinées aux animaux d'élevage. In Tables de composition et de valeur nutritive des matières premières destinées aux animaux d'élevage (ed. D Sauvant, JM Perez and G Tran), 301 pp. INRA-AFZ, Paris.

Solà-Oriol D, Roura E and Torrallardona D 2009. Feed preference in pigs: Relationship with feed particle size and texture. Journal of Animal Science 87, 571–582.

Stark CR 2012. Feed processing to maximize feed efficiency. In Feed efficiency in swine (ed. JF Patience), pp. 131–151. Wageningen Academic Publisher, Wageningen.

Tollier MT and Robin JP 1979. Adaptation de la méthode à l'orcinol sulfurique au dosage automatique des glucides neutres totaux: conditions d'application aux extraits d'origine végétale. Annales de Technologies Agricoles 28, 1–15.

Van Soest PJ and Wine RH 1967. Use of detergents in the analysis of fibrous feeds. IV. Determination of plant cell-wall constituents. Journal of the AOAC 50, 50–55.

Wood JF 1987. The functional properties of feed raw materials and their effect on the production and quality of feed pellets. Animal Feed Science and Technology 18, 1–17.

Young JM, Cai W and Dekkers JCM 2011. Effect of selection for residual feed intake on feeding behavior and daily feed intake patterns in Yorkshire swine. Journal of Animal Science 89, 639–647.