



**HAL**  
open science

## Long-lasting effects of thermal manipulations during embryogenesis in broiler chickens

Thomas Loyau, Sonia Metayer-Coustard, Cécile Berri, Sandrine Grasteau, Christelle Hennequet-Antier, Michel Jacques M.J. Duclos, Sophie Tesseraud, Christophe Praud, N. Everaert, Marco Moroldo, et al.

### ► To cite this version:

Thomas Loyau, Sonia Metayer-Coustard, Cécile Berri, Sandrine Grasteau, Christelle Hennequet-Antier, et al.. Long-lasting effects of thermal manipulations during embryogenesis in broiler chickens. 66. Annual Meeting of the European Federation of Animal Science (EAAP), Aug 2015, Varsovie, Poland. hal-01211070

**HAL Id: hal-01211070**

**<https://hal.science/hal-01211070v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Book of Abstracts of the 66<sup>th</sup> Annual Meeting of the European Federation of Animal Science



**Book of abstracts No. 21 (2015)**  
**Warsaw, Poland**  
**31 August - 4 September 2015**

# Book of Abstracts of the 66<sup>th</sup> Annual Meeting of the European Federation of Animal Science

Warsaw, Poland, 31 August – 4 September, 2015



## EAAP Scientific Committee:

J. Conington  
M. Klopčič  
C. Lauridsen  
G. Pollott  
A. Santos  
H. Sauerwein  
H. Simianer  
H. Spoolder  
M. Tichit  
G. Van Duinkerken  
O. Vangen (chair)

Proceedings publication and Abstract Submission System (OASES) by



*Wageningen Academic*  
*P u b l i s h e r s*

**Long-lasting effects of thermal manipulations during embryogenesis in broiler chickens**

*T. Loyau<sup>1</sup>, S. Métyer-Coustard<sup>1</sup>, C. Berri<sup>1</sup>, S. Mignon-Grasteau<sup>1</sup>, C. Hennequet-Antier<sup>1</sup>, M.J. Duclos<sup>1</sup>, S. Tesseraud<sup>1</sup>, C. Praud<sup>1</sup>, N. Everaert<sup>2</sup>, M. Moroldo<sup>3</sup>, J. Lecarbonnel<sup>3</sup>, P. Martin<sup>4</sup>, V. Coustham<sup>1</sup>, S. Lagarrigue<sup>5</sup>, S. Yahav<sup>6</sup> and A. Collin<sup>1</sup>*

<sup>1</sup>INRA, UR83 Recherches Avicoles, 37380 Nouzilly, France, <sup>2</sup>University of Liège, Gembloux Agro-Bio Tech, Animal Science Unit, Gembloux, Belgium, <sup>3</sup>INRA, CRB GADIE, Domaine de Vilvert, 78350 Jouy-en-Josas, France, <sup>4</sup>INRA, UMR1313, GABI, 78350 Jouy-en-Josas, France, <sup>5</sup>INRA, Agrocampus Ouest, UMR1348 Pegase, 35590 Saint Gilles, France, <sup>6</sup>ARO, The Volcani Center P.O. Box 6, 50250 Bet Dagan, Israel; [thomas.loyau@tours.inra.fr](mailto:thomas.loyau@tours.inra.fr)

Climate change may affect animal physiology, having deleterious consequences for poultry farms. Thermal manipulation (TM) during embryogenesis has been shown to improve heat acclimation in the long term in broiler chickens. TM consists of increasing cyclically incubation temperature from 37.8 °C to 39.5 °C from day (d) 7 to d16 of embryogenesis. Our study aimed to determine the effects of TM and of a subsequent heat challenge (HC) on chickens performance, physiology and metabolism, in order to better understand heat adaptation acclimation mechanisms. In our conditions, TM did not modify hatchability and performance until d28 post hatch, but slightly decreased body weight and composition at d34. TM decreased body temperature (bT) until d28 as well as plasma triiodothyronine concentrations. Our results suggest that TM changes respiratory physiology and decreases stress response during HC at d34. TM associated with low bT affected the expression of genes involved in muscle energy metabolism, or mitigated the effects of acute heat stress. The transcriptomic study performed on breast muscle also pointed out new pathways modified by TM in the long term regulating vascularization and epigenetic mechanisms. TM animals activated more genes in response to HC than controls, possibly explaining their better heat adaptation. This integrated study demonstrated that TM induced long term physiological and metabolic adaptations to changes in the thermal environment of chickens.

***In ovo* stimulation of the chicken embryo with synbiotics influences adult's molecular responses**

*M. Siwek, A. Slawinska, A. Plowiec, A. Dunislawaska and M. Bednarczyk*

*University of Technology and Sciences, Animal Biochemistry and Biotechnology, Mazowiecka 28, 84-085 Bydgoszcz, Poland; [siwek@utp.edu.pl](mailto:siwek@utp.edu.pl)*

Chicken embryo development takes place inside the egg and is influenced by the genetic background, composition of the egg yolk (e.g. nutrients and IgY antibodies deposited by the hen) and conditions during incubation (e.g. temperature and humidity). However, *in ovo* technology allows the supply of additional stimuli to the growing embryo to optimise conditions for embryonic and post-hatch growth and development. In commercial settings, the newly hatched chicks are kept in a sterile environment, separate from the adults. Therefore, the neonates are deprived of the stimuli needed for the development of a healthy intestinal microflora. With the properly selected combinations of pre- and probiotics (synbiotics) delivered *in ovo*, it is possible to modulate the optimal composition of the chicken intestinal microflora, fully developed at hatching. In this study we analyzed the immune-related gene expression regulation in adult broiler chickens (Cobb) that had been treated *in ovo* at 12<sup>th</sup> day of embryo development with different synbiotic combinations. Intestinal (jejunum, cecal tonsils) and splenic tissues were harvested at several time points post hatching (day: 1, 7, 14, 21, 35). Gene expression regulation was analyzed using RT-qPCR to determine the expression of the selected gene panel, including IL-4, IL-6, IL-8, IL-18, IL-12p40, cluster of differentiation (CD) 80, interferon-beta (IFN-β) and interferon-gamma (IFN-γ). Preliminary results indicate that *in ovo* modulation of the microbiome results in the response of the host at the molecular level. Results of this study will provide an insight into development of the host-microbiome relations throughout the chickens' lifespan in terms of molecular responses of the periphery of the immune system (spleen), intestine (jejunum) and the gut-associated lymphoid tissue (cecal tonsils) to *in ovo* delivered synbiotics. Acknowledgements: FP7-KBBE-2012-6-singlestage project from European Commission.