



HAL
open science

COMPLETENESS OF SUMS OF SUBSPACES OF BOUNDED FUNCTIONS AND APPLICATIONS

Joël Blot, Philippe Cieutat

► **To cite this version:**

Joël Blot, Philippe Cieutat. COMPLETENESS OF SUMS OF SUBSPACES OF BOUNDED FUNCTIONS AND APPLICATIONS. 2015. hal-01211149

HAL Id: hal-01211149

<https://hal.science/hal-01211149v1>

Preprint submitted on 5 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMPLETENESS OF SUMS OF SUBSPACES OF BOUNDED FUNCTIONS AND APPLICATIONS

JOËL BLOT¹ AND PHILIPPE CIEUTAT²

ABSTRACT. We give a new proof of a characterization of the closeness of the range of a continuous linear operator and of the closeness of the sum of two closed vector subspaces of a Banach space. Then we state sufficient conditions for the closeness of the sum of two closed subspaces of the Banach space of bounded functions and apply this result on various pseudo almost periodic spaces and pseudo almost automorphic spaces.

2010 Mathematic Subject Classification: 42A75, 42A99, 47A05.

Keywords: Closed range operator, closed sum of subspaces, completeness theorem, pseudo almost periodic function, pseudo almost automorphic function.

1. INTRODUCTION

The origin of this work is the important work of Zheng and Ding [34] on the completeness of the space of weighted pseudo almost automorphic functions. Here we identify two abstract general tools which we used to give a new proof and to generalize the result of Zheng and Ding. Then we apply these general tools to state the completeness of various spaces of pseudo almost periodic functions or pseudo almost automorphic functions.

The first general tool is a characterization of the closeness of the sum of two closed vector subspaces of a Banach space. Such a characterization is established from a characterization of the closeness of the image of a continuous linear operator. The second general tool is a radial retraction, a notion which is associated to the names of Dunkl and Williams [21].

We describe the contents of this work. In Section 2 we give a new proof of a theorem on the characterization of the closeness of the image of a continuous linear operator, we prove that this result of characterization is equivalent to Inverse Mapping Theorem of Banach, and we extend this characterization to unbounded closed linear operators. In Section 3 we provide a characterization of the closeness of the sum of two closed vector subspaces of a Banach space. In Section 4 we provide sufficient conditions to ensure the closeness of the sum of two closed vector subspaces of the Banach space of bounded functions under the supremum norm by using a result on a radial projection. In Section 5 we apply these general tools to state the completeness of the spaces of μ -pseudo almost periodic and μ -pseudo almost automorphic functions without restrictive hypothesis on the measure μ . In Section 6 we study the completeness of the spaces of square-mean μ -pseudo almost periodic and square-mean μ -pseudo almost automorphic processes which are used in stochastic evolution equations. We states similar results in Section 7 for the space of

¹ Laboratoire SAMM EA4543, Paris 1 Panthéon-Sorbonne, centre P.M.F., 90 rue de Tolbiac, 75634 Paris cedex 13, France. E-mail address: blot@univ-paris1.fr

² Laboratoire de Mathématiques de Versailles, Université Versailles-Saint-Quentin-en-Yvelines, 45 avenue des États-Unis, 78035 Versailles cedex, France. E-mail address: philippe.cieutat@uvsq.fr

μ -pseudo almost periodic functions defined on the half-line and in Section 8 for the spaces of weighted pseudo almost periodic sequences and weighted pseudo almost automorphic sequences.

2. CLOSED RANGES

We provide a characterization of the closeness of the range of a linear operators.

Theorem 2.1. *Let E and F be two Banach spaces and $L : E \rightarrow F$ be a continuous linear mapping. The two following assertions are equivalent.*

- i) $\text{Im } L$ is closed into F .
- ii) $\exists c > 0, \forall y \in \text{Im } L, \exists x \in E$ such that $L(x) = y$ and $\|x\| \leq c \|y\|$.

Remark 2.2. A proof of $ii) \implies i)$ is given in [5], Lemma 3.4, and the proof of $i) \implies ii)$ is given in [20], Lemma 1 in Section 6 of Chapter IV. Following our own knowledge, this result was first established by Ganesa Moorthy and Johnson in [25]. We give now another proof of Theorem 2.1. The proof given below is very different and shorter than the two previously cited.

Proof of Theorem 2.1. We consider the following equivalence relation on E :

$$x \sim y \iff L(x) = L(y) \iff x - y \in \text{Ker } L.$$

We consider the quotient space $E^\bullet := E / \sim$. An element x^\bullet of E^\bullet is $x^\bullet = x + \text{Ker } L$ when $x \in E$. We know that E^\bullet is a vector space. Since L is continuous, $\text{Ker } L$ is closed into E , that permits to $\|x^\bullet\|^\bullet := \inf\{\|u\| : u \in x + \text{Ker } L\}$ to be a norm on E^\bullet . Since E is complete, $(E^\bullet, \|\cdot\|^\bullet)$ is a Banach space, cf. [30] p. 299. We define $L^\bullet : E^\bullet \rightarrow F$ by setting $L^\bullet(x^\bullet) := L(x)$ when $x \in E$. L^\bullet is linear and continuous. L^\bullet is clearly injective. We define the following abridgement of L^\bullet : the mapping $K : E^\bullet \rightarrow \text{Im } L$ defined by $K(x^\bullet) := L^\bullet(x^\bullet)$. K remains linear, continuous, injective and moreover K is surjective. And so we can define the inverse of K , $K^{-1} : \text{Im } L \rightarrow E^\bullet$, which is linear.

i) \implies ii). Since $\text{Im } L$ is closed in the Banach space F , it is a Banach space. Since E^\bullet is a Banach space, we can use the Inverse Mapping Theorem of Banach and obtain that K^{-1} is continuous. Therefore we have

$$\exists c_0 > 0, \forall y \in \text{Im } L, \quad \|K^{-1}(y)\|^\bullet \leq c_0 \|y\|.$$

Let y an element of $\text{Im } L$. We arbitrarily fix $x_0 \in K^{-1}(y)$. When $y \neq 0$, we have $\|K^{-1}(y)\|^\bullet < 2c_0 \|y\|$ which means that $\inf\{\|u\| : u \in x_0 + \text{Ker } L\} < 2c_0 \|y\|$, so there exists $x \in x_0 + \text{Ker } L$ such that $\|x\| \leq 2c_0 \|y\|$; and moreover $L(x) = y$. When $y = 0$ we take $x = 0$ and we obtain the same inequality. Therefore ii) is proven with $c = 2c_0$.

ii) \implies i). Let $y \in \text{Im } L$. From ii) we know that there exists $x \in L^{-1}(\{y\}) = K^{-1}(y)$ such that $\|x\| \leq c \|y\|$ that implies: $\|x^\bullet\|^\bullet \leq \|x\| \leq c \|y\|$. And so we have proven

$$\exists c > 0, \forall y \in \text{Im } L, \quad \|K^{-1}(y)\|^\bullet \leq c \|y\|.$$

Since K^{-1} is linear, this last assertion means that K^{-1} is continuous. Let $(y_k)_k$ be a Cauchy sequence in $\text{Im } L$. Since K^{-1} is Lipschitzian, $(K^{-1}(y_k))_k$ is also a Cauchy sequence in E^\bullet . Since E^\bullet is complete, there exists $z^\bullet \in E^\bullet$ such $(K^{-1}(y_k))_k$ converges toward z^\bullet . Since K is continuous $(y_k)_k = (K(K^{-1}(y_k)))_k$ converges toward $K(z^\bullet)$ into $\text{Im } L$. And so $\text{Im } L$ is complete and therefore it is closed. \square

Remark 2.3. The previous proof shows that Theorem 2.1 is a consequence of Inverse Mapping Theorem of Banach. In fact, Theorem 2.1 is equivalent to Inverse Mapping Theorem of Banach. If we assume that Theorem 2.1 is true, and if we consider $L \in \mathcal{L}(E, F)$, where E and F are Banach spaces and L is bijective, since L is surjective, we have $\text{Im } L = F$ is closed in F and consequently, using Theorem 2.1, we obtain: $\exists c > 0$, $\forall y \in F$, $\|L^{-1}(y)\| \leq c \|y\|$, which means the continuity of the inverse L^{-1} .

We can extend Theorem 2.1 to unbounded closed linear operators.

Theorem 2.4. *Let E and F be Banach spaces and $T : \mathfrak{D}(T) \subset E \rightarrow F$ be a unbounded linear operator. We assume that T is closed. Then the two following assertions are equivalent.*

i) $\text{Im } T$ is closed into F .

ii) $\exists c > 0$, $\forall y \in \text{Im } T$, $\exists x \in \mathfrak{D}(T)$ such that $T(x) = y$ and $\|x\|_E \leq c \|y\|_F$.

Proof. First recall that $\|x\|_{\mathcal{G}} := \|x\|_E + \|T(x)\|_F$ defines a norm on $\mathfrak{D}(T)$ and the closeness of T implies that $(\mathfrak{D}(T), \|\cdot\|_{\mathcal{G}})$ is complete (cf. [26] p.164). We define the following abridgement of T : the mapping $L : \mathfrak{D}(T) \rightarrow F$ defined by $L(x) := T(x)$ for $x \in \mathfrak{D}(T)$. Since, $\forall x \in \mathfrak{D}(T)$, $\|L(x)\|_F = \|T(x)\|_F \leq \|x\|_{\mathcal{G}}$, we have $L \in \mathfrak{L}((\mathfrak{D}(T), \|\cdot\|_{\mathcal{G}}), (F, \|\cdot\|_F))$. We introduce the following new condition.

$$(2.1) \quad \exists d > 0, \forall y \in \text{Im } L, \exists x \in \mathfrak{D}(T) \text{ such that } L(x) = y \text{ and } \|x\|_{\mathcal{G}} \leq d \|y\|_F.$$

The equivalence i) \iff (2.1) is a consequence of Theorem 2.1 applied on the continuous linear mapping $L \in \mathfrak{L}((\mathfrak{D}(T), \|\cdot\|_{\mathcal{G}}), (F, \|\cdot\|_F))$ by remarking that $\text{Im } L = \text{Im } T$.

Let us prove the equivalence (2.1) \iff ii). For the implication (2.1) \implies ii) it suffices to take $c = d$ and to note that $\|x\|_E \leq \|x\|_{\mathcal{G}}$. Now we prove the reciprocal implication. Let $y \in \text{Im } T$. Then, from ii), $\exists x \in \mathfrak{D}(T)$ such that $T(x) = y$ and $\|x\|_E \leq c \|y\|_F$ which imply $\|x\|_{\mathcal{G}} = \|x\|_E + \|T(x)\|_F = \|x\|_E + \|y\|_F \leq (c + 1) \|y\|_F$. Setting $d := c + 1$ we obtain $\|x\|_{\mathcal{G}} \leq d \|y\|_F$. And so we have proven the following assertion ii) \implies (2.1). \square

3. GENERAL CLOSED SUMS

In this section we provide necessary and sufficient conditions to have the closeness of the sum of two closed vector subspaces in a Banach space.

Theorem 3.1. *Let E be a Banach space, and M and N be two closed vector subspaces of E . Then the three following conditions are equivalent.*

i) $M + N$ is closed into E .

ii) $\exists c > 0$, $\forall z \in M + N$, $\exists(x, y) \in M \times N$ such that $x + y = z$ and $\|x\| \leq c \|z\|$.

Proof. Let us consider the following assertion

$$(3.1) \quad \exists d > 0, \forall z \in M + N, \exists(x, y) \in M \times N \text{ s.t. } x + y = z, \|x\| \leq d \|z\|, \|y\| \leq d \|z\|.$$

The equivalence i) \iff (3.1) is a consequence of Theorem 2.1 by using the continuous linear mapping $L : M \times N \rightarrow E$ defined by $L(x, y) := x + y$ and endowing the product $M \times N$ with the norm $\|(x, y)\| = \max\{\|x\|, \|y\|\}$. The implication (3.1) \implies ii) is obvious with $c = d$ and for the reciprocal implication, note that if $\|x\| \leq c \|z\|$, we have

$$\|y\| - \|z\| \leq \|y - z\| = \|x\| \leq c \|z\| \implies \|y\| \leq (c + 1) \|z\|,$$

then the implication ii) \implies (3.1) is proved with $d = c + 1$. \square

Remark 3.2. There exists another proof of Theorem 3.1 due to Zheng and Ding in [35]. The proof of i) \implies (3.1) is given in [12] (Theorem 2.10, p. 37). A result as Theorem 3.1 is established by Kober [27] under the additional assumption $M \cap N = \{0\}$.

4. CLOSENESS OF THE SUM TWO VECTOR SUBSPACES OF A BANACH SPACE OF BOUNDED FUNCTIONS

In this section U is a set and X is a Banach space. $B(U, X)$ denotes the Banach space of all bounded functions from U into X , equipped with the norm $\|f\|_\infty = \sup_{u \in U} \|f(u)\|$ for $f \in B(U, X)$. Here we establish sufficient conditions to have the closeness of the sum of two closed subspaces of the Banach space of bounded functions using a result on the radial retraction.

Theorem 4.1. *Let \mathcal{F}_1 and \mathcal{F}_2 be two closed vector subspaces of $B(U, X)$ satisfying the following hypothesis:*

- i) *The map $\Phi : X \rightarrow X$ is Lipschitz and $f \in \mathcal{F}_1$, then $\Phi \circ f \in \mathcal{F}_1$.*
- ii) *$g \in \mathcal{F}_1$ and $h \in \mathcal{F}_2$ such that $\forall u \in U$, $\|g(u)\| \leq \|h(u)\|$, then $g \in \mathcal{F}_2$.*

Then $\mathcal{F}_1 + \mathcal{F}_2$ is a Banach space endowed with the norm $\|\cdot\|_\infty$.

For the proof of Theorem 4.1, we use the radial retraction P_R on the ball of radius R and with the center at the origin.

Lemma 4.2. *Let $R \geq 0$. We define the function $P_R : X \rightarrow X$ by setting*

$$(4.1) \quad P_R(x) = \begin{cases} x & \text{if } \|x\| \leq R \\ \frac{R}{\|x\|}x & \text{if } \|x\| > R. \end{cases}$$

Then the following assertions hold:

- i) *P_R is 2-Lipschitz: $\forall x_1, x_2 \in X$, $\|P_R(x_1) - P_R(x_2)\| \leq 2\|x_1 - x_2\|$.*
- ii) *For all $x \in X$, $\|P_R(x)\| \leq R$.*

Proof. i) If

$$(4.2) \quad \|x_1\| \leq R \leq \|x_2\|,$$

then

$$\|P_R(x_1) - P_R(x_2)\| = \left\| x_1 - \frac{R}{\|x_2\|}x_2 \right\| \leq \|x_1 - x_2\| + \left\| x_2 - \frac{R}{\|x_2\|}x_2 \right\|,$$

from $\left\| x_2 - \frac{R}{\|x_2\|}x_2 \right\| = \left| \|x_2\| - R \right|$ and (4.2) we obtain

$$\left\| x_2 - \frac{R}{\|x_2\|}x_2 \right\| \leq \|x_2\| - \|x_1\| \leq \|x_1 - x_2\|,$$

therefore

$$(4.3) \quad \|P_R(x_1) - P_R(x_2)\| \leq 2\|x_1 - x_2\|.$$

If $\|x_1\| \leq R$ and $\|x_2\| \leq R$, then we have (4.3), since $P_R(x_1) = x_1$ and $P_R(x_2) = x_2$. Dunkl and Williams have proved in [21] that for all nonzero x_1 and $x_2 \in X$

$$(4.4) \quad \left\| \frac{x_1}{\|x_1\|} - \frac{x_2}{\|x_2\|} \right\| \leq \frac{4}{\|x_1\| + \|x_2\|} \|x_1 - x_2\|.$$

From

$$\|P_R(x_1) - P_R(x_2)\| = \left\| \frac{R}{\|x_1\|}x_1 - \frac{R}{\|x_2\|}x_2 \right\|$$

and (4.4) it can be seen that (4.3) is fulfilled if $\|x_1\| \geq R$ and $\|x_2\| \geq R$. This ends the proof.

ii) From (4.1) we deduce ii). □

Proof of Theorem 4.1. We use Theorem 3.1 by setting $E = B(U, X)$, $M = \mathcal{F}_1$ and $N = \mathcal{F}_2$. Let us prove that

$$(4.5) \quad \forall f \in \mathcal{F}_1 + \mathcal{F}_2, \quad \exists g^* \in \mathcal{F}_1, \quad \exists h^* \in \mathcal{F}_2 \text{ such that } f = g^* + h^* \text{ and } \|g^*\|_\infty \leq \|f\|_\infty.$$

Let $f \in \mathcal{F}_1 + \mathcal{F}_2$. Then there exist $g \in \mathcal{F}_1$ and $h \in \mathcal{F}_2$ such that $f = g + h$. Define g^* by setting $g^* = P_R \circ g$, with $R := \|f\|_\infty$ and P_R the radial retraction of Lemma 4.2. Define h^* by setting $h^* = f - g^*$. Then we have $f = g^* + h^*$.

Since $P_R : X \rightarrow X$ is Lipschitz and $g \in \mathcal{F}_1$, from Hypothesis i), we obtain $P_R \circ g = g^* \in \mathcal{F}_1$. Inequality $\|g^*\|_\infty \leq R = \|f\|_\infty$ results of ii) of Lemma 4.2.

Since $\|f(u)\| \leq R := \|f\|_\infty$ for all $u \in U$ and from the definition of P_R , we deduce that $P_R \circ f = f$; therefore $\|h^*(u)\| = \|f(u) - g^*(u)\| = \|P_R(f(u)) - P_R(g(u))\|$. The radial retraction P_R being 2-Lipschitz, we obtain that

$$\forall u \in U, \quad \|h^*(u)\| = \|P_R(f(u)) - P_R(g(u))\| \leq 2\|f(u) - g(u)\| = 2\|h(u)\|.$$

From $h \in \mathcal{F}_2$ and Hypothesis ii), we deduce that $h^* \in \mathcal{F}_2$; so (4.5) is fulfilled, which is assertion ii) of Theorem 3.1 with $c = 1$.

By using Theorem 3.1 with $E = B(U, X)$, $M = \mathcal{F}_1$, $N = \mathcal{F}_2$ and $c = 1$, we obtain the closeness of the vector subspace $\mathcal{F}_1 + \mathcal{F}_2$ in the Banach space $(B(U, X), \|\cdot\|_\infty)$. □

Remark 4.3. Under the hypotheses of Theorem 4.1, if in addition we assume that $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$, the proof of this theorem is more simple. In this case, the decomposition (which becomes unique) $f = g + h$ with $g \in \mathcal{F}_1$ and $h \in \mathcal{F}_2$, we obtain $\|g\|_\infty \leq \|f\|_\infty$ from (4.5). This fact and the closeness of \mathcal{F}_1 and \mathcal{F}_2 imply the completeness of the sum.

5. SPACES OF μ -PSEUDO ALMOST PERIODIC AND AUTOMORPHIC FUNCTIONS

In this section X is a Banach space.

5.1. Almost periodic case. The pseudo almost periodic functions were introduced in the literature in the early nineties by Zhang [31, 32], as a natural generalization of the classical almost periodic functions in the sense of Bohr. The notion of μ -pseudo almost periodicity initiated by Blot, Cieutat and Ezzinbi [10] is a generalization of the weighted pseudo almost periodicity due to Diagana [16, 17] which generalizes the one of pseudo almost periodicity.

Definition 5.1. [15]. $f \in C(\mathbb{R}, X)$ (continuous) is said to be *almost periodic* (in the Bohr sense) if for all $\varepsilon > 0$, there exists $\ell > 0$, such that for all $\alpha \in \mathbb{R}$, there exists $\tau \in [\alpha, \alpha + \ell]$ with

$$\sup_{t \in \mathbb{R}} \|f(t + \tau) - f(t)\| < \varepsilon.$$

We denote the space of all such functions by $AP(\mathbb{R}, X)$.

We denote by \mathcal{B} the Lebesgue σ -field of \mathbb{R} and by \mathcal{M} the set of all positive measures μ on \mathcal{B} satisfying $\mu(\mathbb{R}) = +\infty$ and $\mu([a, b]) < +\infty$, for all $a, b \in \mathbb{R}$ ($a \leq b$). $BC(\mathbb{R}, X)$ denotes the Banach space of all continuous and bounded functions from \mathbb{R} to X , equipped with the norm

$$\|f\|_\infty = \sup_{t \in \mathbb{R}} \|f(t)\|.$$

Definition 5.2. [9, 10]. Let $\mu \in \mathcal{M}$. $f \in BC(\mathbb{R}, X)$ is said to be μ -ergodic if

$$\lim_{r \rightarrow +\infty} \frac{1}{\mu([-r, r])} \int_{[-r, r]} \|f(t)\| d\mu(t) = 0.$$

We denote the space of all such functions by $\mathcal{E}(\mathbb{R}, X, \mu)$.

Definition 5.3. [10]. Let $\mu \in \mathcal{M}$. $f \in BC(\mathbb{R}, X)$ is said to be μ -pseudo almost periodic if

$$f = g + \phi \quad \text{where } g \in AP(\mathbb{R}, X), \quad \phi \in \mathcal{E}(\mathbb{R}, X, \mu).$$

We denote the space of all such functions by $PAP(\mathbb{R}, X, \mu)$.

Remark 5.4. In general the sum $PAP(\mathbb{R}, X, \mu) = AP(\mathbb{R}, X) + \mathcal{E}(\mathbb{R}, X, \mu)$ is not direct ([10], Remark 2.26).

Theorem 5.5. Let $\mu \in \mathcal{M}$. Then $(PAP(\mathbb{R}, X, \mu), \|\cdot\|_\infty)$ is a Banach space.

Proof. We use Theorem 4.1 by setting $U = \mathbb{R}$, $\mathcal{F}_1 = AP(\mathbb{R}, X)$ and $\mathcal{F}_2 = \mathcal{E}(\mathbb{R}, X, \mu)$.

i) $AP(\mathbb{R}, X)$ is a closed vector subspace of $BC(\mathbb{R}, X)$ ([15], Theorem 6.1 and 6.2, pp. 138-139), then a closed vector subspace of $B(\mathbb{R}, X)$. Moreover if $\Phi \in C(X, X)$ and $f \in AP(\mathbb{R}, X)$ then $\Phi \circ f \in AP(\mathbb{R}, X)$ ([2], VII, p. 5); so Hypothesis i) is satisfied.

ii) $\mathcal{E}(\mathbb{R}, X, \mu)$ endowed with the norm $\|\cdot\|_\infty$ is a Banach space ([10], Proposition 2.11), then a closed vector subspace of $B(\mathbb{R}, X)$. By using Definition 5.2, we deduce that the pair $AP(\mathbb{R}, X)$ and $\mathcal{E}(\mathbb{R}, X, \mu)$ satisfies Hypothesis ii).

By Theorem 4.1, we obtain that $PAP(\mathbb{R}, X, \mu) = AP(\mathbb{R}, X) + \mathcal{E}(\mathbb{R}, X, \mu)$ is a Banach space endowed with the norm $\|\cdot\|_\infty$. \square

Remark 5.6. With restrictive conditions on the measure μ , this result is proved ([10], Corollary 2.31).

Remark 5.7. Theorem 5.5 generalizes Corollary 2.4 in [34] of Zheng and Ding from the space of weighted pseudo almost periodic functions to the space of μ -pseudo almost periodic functions.

5.2. Almost automorphic case. The notion of μ -pseudo almost automorphy due to Blot, Cieutat and Ezzinbi [9] is a generalization of the weighted pseudo almost automorphy initiated by Blot, Mophou, N'Guérékata and Pennequin [11] which generalizes the one of pseudo almost automorphy introduced by Liang, N'Guérékata, Xiao and Zhang [28].

Definition 5.8. [29]. $f \in C(\mathbb{R}, X)$ is said to be *almost automorphic* if for every sequence of real numbers $(s'_n)_n$ there exists a subsequence of $(s'_n)_n$ denoted by $(s_n)_n$ such that

$$g(t) = \lim_{n \rightarrow \infty} f(t + s_n) \text{ exists for all } t \text{ in } \mathbb{R}$$

and

$$\lim_{n \rightarrow \infty} g(t - s_n) = f(t) \text{ for all } t \text{ in } \mathbb{R}.$$

We denote the space of all such functions by $AA(\mathbb{R}, X)$.

Definition 5.9. [9]. Let $\mu \in \mathcal{M}$. $f \in BC(\mathbb{R}, X)$ is said to be μ -pseudo almost automorphic if

$$f = g + \phi \quad \text{where } g \in AA(\mathbb{R}, X), \quad \phi \in \mathcal{E}(\mathbb{R}, X, \mu).$$

We denote the space of all such functions by $PAA(\mathbb{R}, X, \mu)$.

Remark 5.10. In general the sum $PAA(\mathbb{R}, X, \mu) = AA(\mathbb{R}, X) + \mathcal{E}(\mathbb{R}, X, \mu)$ is not direct ([9], Remark 4.4). A sufficient condition for $PAA(\mathbb{R}, X, \mu) = AA(\mathbb{R}, X) \oplus \mathcal{E}(\mathbb{R}, X, \mu)$ is that $PAA(\mathbb{R}, X, \mu)$ be translation invariant.

Theorem 5.11. Let $\mu \in \mathcal{M}$. Then $(PAA(\mathbb{R}, X, \mu), \|\cdot\|_\infty)$ is a Banach space.

Proof. We use Theorem 4.1 by setting $U = \mathbb{R}$, $\mathcal{F}_1 = AA(\mathbb{R}, X)$ and $\mathcal{F}_2 = \mathcal{E}(\mathbb{R}, X, \mu)$.

i) $AA(\mathbb{R}, X)$ is a closed vector subspace of $BC(\mathbb{R}, X)$ ([29], Theorem 2.3, p. 11 and Theorem 2.10, p. 16), then a closed vector subspace of $B(\mathbb{R}, X)$. Moreover if $\Phi \in C(X, X)$ and $f \in AA(\mathbb{R}, X)$ then $\Phi \circ f \in AA(\mathbb{R}, X)$ ([29], Theorem 2.5, p. 13); so Hypothesis i) is satisfied.

ii) is similar to ii) in the proof of Theorem 5.5.

By Theorem 4.1, we deduce that $PAA(\mathbb{R}, X, \mu) = AA(\mathbb{R}, X) + \mathcal{E}(\mathbb{R}, X, \mu)$ is a Banach space endowed with the norm $\|\cdot\|_\infty$. \square

Remark 5.12. In the particular case where $PAA(\mathbb{R}, X, \mu)$ is translation invariant, this result is proved in ([9], Theorem 4.9).

Remark 5.13. Theorem 5.11 generalizes Theorem 2.3 in [34] of Zheng and Ding from the space of weighted pseudo almost automorphic functions to the space of μ -pseudo almost automorphic functions.

6. SPACES OF SQUARE-MEAN μ -PSEUDO ALMOST PERIODIC AND AUTOMORPHIC PROCESSES

Throughout this section, we denote by H a Banach space, (Ω, \mathcal{A}, P) a probability space and $L^2(\Omega, P, H)$ the space of all H -valued random variables x with a finite quadratic-mean:

$$E \|x\|^2 = \int_{\Omega} \|x\|^2 dP < +\infty.$$

When $x \in L^2(\Omega, P, H)$, we set $\|x\|_{L^2} = (E \|x\|^2)^{\frac{1}{2}}$. Endowed with the norm $\|\cdot\|_{L^2}$, $L^2(\Omega, P, H)$ is a Banach space.

Definition 6.1. [7]. Let $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ be a stochastic process.

i) x is said to be *stochastically continuous* if

$$\forall t \in \mathbb{R}, \quad \lim_{s \rightarrow t} E \|x(s) - x(t)\|^2 = 0.$$

ii) x is said to be *stochastically bounded* if

$$\sup_{t \in \mathbb{R}} E \|x(t)\|^2 < +\infty.$$

Remark 6.2. It is easy to verify that a stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ is stochastically continuous and bounded if and only if x is a continuous and bounded map from \mathbb{R} to the Banach space $X = L^2(\Omega, P, H)$ in the ordinary sense, i.e. $x \in B(\mathbb{R}, X)$.

The square-mean almost periodic stochastic processes were introduced in the literature by Bezandry and Diagana [7]. They established the existence and uniqueness of square-mean almost periodic mild solutions to some stochastic differential equations and some functional integro-differential stochastic evolution equations in [7, 8]. Then Fu and Liu generalize this one to the square-mean almost automorphy in [24]. The notion of square-mean μ -pseudo almost periodic and automorphic processes due to Diop, Ezzinbi and Mbaye [18, 19] is a generalization of the pseudo almost automorphic stochastic processes [14] and the weighted pseudo almost automorphy [13] due to Chen and Lin.

The definition of the square-mean μ -pseudo almost automorphy given in [19] is slightly different from the one given by Bedouhene, Challali, Mellah, Raynaud de Fitte and Smaali in [6]. A continuous and bounded stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ is square-mean μ -ergodic in [19] if

$$\lim_{r \rightarrow +\infty} \frac{1}{\mu([-r, r])} \int_{[-r, r]} E \|x(t)\|^2 d\mu(t) = 0$$

and in [6] if

$$\lim_{r \rightarrow +\infty} \frac{1}{\mu([-r, r])} \int_{[-r, r]} (E \|x(t)\|^2)^{\frac{1}{2}} d\mu(t) = 0.$$

We will see that these definitions are equivalent (cf. Proposition 6.6 and 6.12).

6.1. Almost periodic case.

Definition 6.3. [7]. A continuous stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ is said to be *square-mean almost periodic* if for all $\varepsilon > 0$, there exists $\ell > 0$, such that for all $\alpha \in \mathbb{R}$, there exists $\tau \in [\alpha, \alpha + \ell]$ with

$$\sup_{t \in \mathbb{R}} E \|x(t + \tau) - x(t)\|^2 < \varepsilon.$$

Remark 6.4. It is easy to verify that a stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ is square-mean almost periodic if and only if x is an almost periodic function from \mathbb{R} to the Banach space $X = L^2(\Omega, P, H)$ in the Bohr sense, i.e. $x \in AP(\mathbb{R}, X)$ (cf. Definition 5.1).

As in Section 5, we denote by \mathcal{B} the Lebesgue σ -field of \mathbb{R} and by \mathcal{M} the set of all positive measures μ on \mathcal{B} satisfying $\mu(\mathbb{R}) = +\infty$ and $\mu([a, b]) < +\infty$, for all $a, b \in \mathbb{R}$ ($a \leq b$).

Definition 6.5. [19] Let $\mu \in \mathcal{M}$. Let $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ be a continuous and bounded stochastic process.

i) x is said to be *square-mean μ -ergodic* if

$$\lim_{r \rightarrow +\infty} \frac{1}{\mu([-r, r])} \int_{[-r, r]} E \|x(t)\|^2 d\mu(t) = 0.$$

ii) x is said to be *square-mean μ -pseudo almost periodic* if

$$x = y + z$$

where y is square-mean almost periodic and z square-mean μ -ergodic.

Proposition 6.6. Let $\mu \in \mathcal{M}$. Let $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ be a stochastic process.

i) x is square-mean μ -ergodic if and only if x is a μ -ergodic function from \mathbb{R} to the Banach space $X = L^2(\Omega, P, H)$ in the sense of Definition 5.2, i.e. $x \in \mathcal{E}(\mathbb{R}, X, \mu)$.

ii) x is square-mean μ -pseudo almost periodic if and only if x is a μ -pseudo almost periodic function from \mathbb{R} to the Banach space $X = L^2(\Omega, P, H)$ in the sense of Definition 5.3, i.e. $x \in PAP(\mathbb{R}, X, \mu)$.

Proof. Recall that the Banach space $X = L^2(\Omega, P, H)$ is endowed with the norm $\|\cdot\|_{L^2}$ defined by $\|x\|_{L^2} = (E \|x\|^2)^{\frac{1}{2}}$ for $x \in X$.

i) Let $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ be a stochastic process. It suffices to prove that x is square-mean μ -ergodic if and only if

$$(6.1) \quad \lim_{r \rightarrow +\infty} \frac{1}{\mu([-r, r])} \int_{[-r, r]} (E \|x(t)\|^2)^{\frac{1}{2}} d\mu(t) = 0.$$

First from Cauchy-Schwartz inequality, it follows

$$\int_{[-r, r]} \|x(t)\|_{L^2} d\mu(t) \leq \left(\int_{[-r, r]} \|x(t)\|_{L^2}^2 d\mu(t) \right)^{\frac{1}{2}} \left(\int_{[-r, r]} 1^2 d\mu(t) \right)^{\frac{1}{2}},$$

then

$$\frac{1}{\mu([-r, r])} \int_{[-r, r]} (E \|x(t)\|^2)^{\frac{1}{2}} d\mu(t) \leq \left(\frac{1}{\mu([-r, r])} \int_{[-r, r]} E \|x(t)\|^2 d\mu(t) \right)^{\frac{1}{2}},$$

consequently if x is square-mean μ -ergodic, then x satisfies (6.1).

Secondly x is stochastically bounded: there exists $M > 0$ such that $E \|x(t)\|^2 \leq M$, then

$$E \|x(t)\|^2 = \|x(t)\|_{L^2}^2 \leq \|x(t)\|_{L^2} \sup_{t \in \mathbb{R}} \|x(t)\|_{L^2} \leq M^{\frac{1}{2}} (E \|x(t)\|^2)^{\frac{1}{2}},$$

therefore

$$(6.2) \quad \frac{1}{\mu([-r, r])} \int_{[-r, r]} E \|x(t)\|^2 d\mu(t) \leq \frac{M^{\frac{1}{2}}}{\mu([-r, r])} \int_{[-r, r]} (E \|x(t)\|^2)^{\frac{1}{2}} d\mu(t),$$

consequently if x satisfies (6.1), then x is square-mean μ -ergodic.

ii) results of i) and Remark 6.4. □

Remark 6.7. A consequence of Proposition 6.6 is that all results of [10] on μ -pseudo almost periodic functions are directly applicable to square-mean μ -pseudo almost periodic stochastic processes.

In view of Proposition 6.6, we denote by $PAP(\mathbb{R}, L^2(\Omega, P, H), \mu)$ the space of all square-mean μ -pseudo almost periodic stochastic processes $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$. We denote by $\|\cdot\|_\infty$ the norm of $PAP(\mathbb{R}, L^2(\Omega, P, H), \mu)$ defined by $\|x\|_\infty = \sup_{t \in \mathbb{R}} (E \|x(t)\|^2)^{\frac{1}{2}}$.

Theorem 6.8. *Let $\mu \in \mathcal{M}$. Then $(PAP(\mathbb{R}, L^2(\Omega, P, H), \mu), \|\cdot\|_\infty)$ is a Banach space.*

Proof. It is a consequence of Theorem 5.5 and Proposition 6.6. \square

6.2. Almost automorphic case.

Definition 6.9. [24]. A continuous stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ is said to be *square-mean almost automorphic* if for every sequence of real numbers $(s'_n)_n$ there exists a subsequence of the sequence $(s'_n)_n$ denoted by $(s_n)_n$ such that for some stochastic process $y : \mathbb{R} \rightarrow L^2(\Omega, P, H)$

$$\forall t \in \mathbb{R}, \quad \lim_{t \rightarrow \infty} E \|x(t + s_n) - y(t)\|^2 = 0$$

and

$$\forall t \in \mathbb{R}, \quad \lim_{t \rightarrow \infty} E \|y(t - s_n) - x(t)\|^2 = 0.$$

Remark 6.10. It is easy to verify that a stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ is square-mean almost automorphic if and only if x is an almost automorphic function from \mathbb{R} to the Banach space $X = L^2(\Omega, P, H)$ in the sense of Definition 5.8, i.e. $x \in AA(\mathbb{R}, X)$.

Definition 6.11. [19] Let $\mu \in \mathcal{M}$. A continuous and bounded stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ is said to be *square-mean μ -pseudo almost automorphic* if

$$x = y + z$$

where y is square-mean almost automorphic and z square-mean μ -ergodic.

From i) of Proposition 6.6 and Remark 6.10, we obtain the following result:

Proposition 6.12. *Let $\mu \in \mathcal{M}$. Let $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$ be a stochastic process. x is square-mean μ -pseudo almost automorphic if and only if x is a μ -pseudo almost automorphic function from \mathbb{R} to the Banach space $X = L^2(\Omega, P, H)$ in the sense of Definition 5.9, i.e. $x \in PAA(\mathbb{R}, X, \mu)$.*

Remark 6.13. A consequence of Proposition 6.12 is that all results of [9] on μ -pseudo almost automorphic functions are directly applicable to square-mean μ -pseudo almost automorphic stochastic processes.

In view of Proposition 6.12, we denote by $PAA(\mathbb{R}, L^2(\Omega, P, H), \mu)$ the space of all square-mean μ -pseudo almost automorphic stochastic processes $x : \mathbb{R} \rightarrow L^2(\Omega, P, H)$. We denote by $\|\cdot\|_\infty$ the norm of $PAA(\mathbb{R}, L^2(\Omega, P, H), \mu)$ defined by $\|x\|_\infty = \sup_{t \in \mathbb{R}} (E \|x(t)\|^2)^{\frac{1}{2}}$.

Theorem 6.14. *Let $\mu \in \mathcal{M}$. Then $(PAA(\mathbb{R}, X, \mu), \|\cdot\|_\infty)$ is a Banach space.*

Proof. It is a consequence of Theorem 5.11 and Proposition 6.12. \square

7. SPACES OF μ -PSEUDO ALMOST PERIODIC FUNCTIONS DEFINED ON THE HALF-LINE

In this section X is a Banach space. We study the case where the functions are defined only on the half-line.

Definition 7.1. $f \in C(\mathbb{R}^+, X)$ (continuous) is said to be *almost periodic* (in the Bohr sense) if for all $\varepsilon > 0$, there exists $\ell > 0$, such that for all $\alpha \geq 0$, there exists $\tau \in [\alpha, \alpha + \ell]$ with

$$\sup_{t \geq 0} \|f(t + \tau) - f(t)\| < \varepsilon.$$

We denote the space of all such functions by $AP(\mathbb{R}^+, X)$.

$BC(\mathbb{R}^+, X)$ denotes the Banach space of all continuous and bounded functions from \mathbb{R}^+ to X , equipped with the norm

$$\|f\|_\infty = \sup_{t \geq 0} \|f(t)\|.$$

One should point out that combining the extension theorem of an almost periodic function on \mathbb{R}^+ to an almost periodic function on \mathbb{R} ([4], Proposition 4.7.1, p. 305) and the recurrence property of an almost periodic function on \mathbb{R} which is a consequence of the definition of the almost periodicity: there exists a sequence of real numbers $(t_n)_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow +\infty} t_n = +\infty$ and $\lim_{n \rightarrow +\infty} f(t + t_n) = f(t)$, one can prove that the restriction operator

$$(7.1) \quad R : AP(\mathbb{R}, X) \rightarrow AP(\mathbb{R}^+, X) \quad \text{defined by } R(f) = f|_{\mathbb{R}^+}$$

is well-defined, maps $AP(\mathbb{R}, X)$ onto $AP(\mathbb{R}^+, X)$ and satisfies for $f \in AP(\mathbb{R}, X)$

$$\|f\|_{AP(\mathbb{R}, X)} = \sup_{t \in \mathbb{R}} \|f(t)\| = \sup_{t \geq 0} \|f(t)\| = \|R(f)\|_{AP(\mathbb{R}^+, X)}.$$

Similarly as for μ -pseudo almost periodic functions defined on the whole real line, we define below the μ -pseudo almost periodic functions defined on the half line.

We denote by \mathcal{B}_+ the Lebesgue σ -field of \mathbb{R}_+ and by \mathcal{M}_+ the set of all positive measures μ on \mathcal{B}_+ satisfying $\mu(\mathbb{R}^+) = +\infty$ and $\mu([a, b]) < +\infty$, for all $0 \leq a \leq b$.

Definition 7.2. Let $\mu \in \mathcal{M}_+$. $f \in BC(\mathbb{R}^+, X)$ is said to be μ -ergodic if

$$\lim_{r \rightarrow +\infty} \frac{1}{\mu([0, r])} \int_{[0, r]} \|f(t)\| d\mu(t) = 0.$$

We denote the space of all such functions by $\mathcal{E}(\mathbb{R}^+, X, \mu)$.

Theorem 7.3. Let $\mu \in \mathcal{M}_+$. Then $(\mathcal{E}(\mathbb{R}^+, X, \mu), \|\cdot\|_\infty)$ is a Banach space.

Proof. It is a slight adaptation of the proof of the completeness of the space of μ -ergodic functions defined on the whole real line: $(\mathcal{E}(\mathbb{R}, X, \mu), \|\cdot\|_\infty)$ ([10], Proposition 2.11). \square

Definition 7.4. Let $\mu \in \mathcal{M}_+$. $f \in BC(\mathbb{R}^+, X)$ is said to be μ -pseudo almost periodic if

$$f = g + \phi \quad \text{where } g \in AP(\mathbb{R}^+, X), \quad \phi \in \mathcal{E}(\mathbb{R}^+, X, \mu).$$

We denote the space of all such functions by $PAP(\mathbb{R}^+, X, \mu)$.

Theorem 7.5. Let $\mu \in \mathcal{M}_+$. Then $(PAP(\mathbb{R}^+, X, \mu), \|\cdot\|_\infty)$ is a Banach space.

Proof. We use Theorem 4.1 by setting $U = \mathbb{R}^+$, $\mathcal{F}_1 = AP(\mathbb{R}^+, X)$ and $\mathcal{F}_2 = \mathcal{E}(\mathbb{R}^+, X, \mu)$.

i) $AP(\mathbb{R}^+, X)$ is isometrically isomorphic to the Banach space $AP(\mathbb{R}, X)$, then it is a Banach space. Let $\Phi \in C(X, X)$ and $g \in AP(\mathbb{R}^+, X)$. First there exists $f \in AP(\mathbb{R}, X)$ such that $R(f) = g$ (see (7.1)). Secondly by using the definition of the restriction operator R we obtain $\Phi \circ g = \Phi \circ [R(f)] = R(\Phi \circ f)$. From the fact that $\Phi \circ f \in AP(\mathbb{R}, X)$ (see proof of Theorem 5.5) we deduce that $\Phi \circ g \in AP(\mathbb{R}^+, X)$; so Hypothesis i) is satisfied.

ii) By using Definition 7.2, we deduce that the pair $AP(\mathbb{R}^+, X)$ and $\mathcal{E}(\mathbb{R}^+, X, \mu)$ satisfies Hypothesis ii).

By Theorem 4.1, we obtain that $PAP(\mathbb{R}^+, X, \mu) = AP(\mathbb{R}^+, X) + \mathcal{E}(\mathbb{R}^+, X, \mu)$ is a Banach space endowed with the norm $\|\cdot\|_\infty$. \square

The following definition is due to Maurice Fréchet [22, 23].

Definition 7.6. $f \in BC(\mathbb{R}^+, X)$ is said to be *asymptotically almost periodic* if

$$f = g + \phi \quad \text{where } g \in AP(\mathbb{R}^+, X), \quad \phi \in C(\mathbb{R}^+, X) \text{ and } \lim_{t \rightarrow +\infty} \phi(t) = 0.$$

We denote the space of all such functions by $AAP(\mathbb{R}^+, X)$.

We will see that the following well-known result is a simple corollary of our main result.

Theorem 7.7. $(AAP(\mathbb{R}^+, X), \|\cdot\|_\infty)$ is a Banach space.

Proof. We use Theorem 4.1 by setting $U = \mathbb{R}^+$, $\mathcal{F}_1 = AP(\mathbb{R}^+, X)$ and $\mathcal{F}_2 = C_0(\mathbb{R}^+, X)$ where $C_0(\mathbb{R}^+, X) = \{\phi \in C(\mathbb{R}^+, X); \lim_{t \rightarrow +\infty} \phi(t) = 0\}$

i) is proved in i) of the proof of Theorem 7.5.

ii) It is obvious.

By Theorem 4.1, we deduce that $AAP(\mathbb{R}^+, X) = AP(\mathbb{R}^+, X) + C_0(\mathbb{R}^+, X)$ is a Banach space endowed with the norm $\|\cdot\|_\infty$. Note that this sum is direct: $AAP(\mathbb{R}^+, X) = AP(\mathbb{R}^+, X) \oplus C_0(\mathbb{R}^+, X)$. \square

8. SPACES OF WEIGHTED PSEUDO ALMOST PERIODIC AND AUTOMORPHIC SEQUENCES

In this Section we provide results of completeness on the space of weighted pseudo almost periodic sequences and of the space of weighted pseudo automorphic sequences without any assumption on the weight of the ergodic part of these sequences.

8.1. Almost periodic case. The notion of weighted pseudo almost periodicity of sequences is a generalization of pseudo almost periodicity of sequences ([32], Definition 7.11, p. 84). The notion of weighted pseudo almost periodicity of sequences is due Zhang and Li [33].

Denote by $X^{\mathbb{Z}}$ the set of all two-sided sequences $u = (u_n)_{n \in \mathbb{Z}}$ with values in the Banach space X . $\ell^\infty(\mathbb{Z}, X)$ denotes the Banach space of all bounded two-sided sequences of $X^{\mathbb{Z}}$ equipped with the norm

$$\|u\|_\infty = \sup_{n \in \mathbb{Z}} \|u_n\|.$$

Definition 8.1. ([15], p. 45). $u = (u_n)_{n \in \mathbb{Z}} \in X^{\mathbb{Z}}$ is said to be *almost periodic* if for all $\varepsilon > 0$, there exists $N \in \mathbb{N}^*$, such that for all $m \in \mathbb{Z}$, there exists $p \in \{m, m+1, \dots, m+N\}$ with

$$\sup_{n \in \mathbb{Z}} \|u_{n+p} - u_n\| < \varepsilon.$$

We denote the space of all such sequences by $AP_d(\mathbb{Z}, X)$.

The following well-known statement ([15], Theorem 1.27, p. 47) clarifies the relation between almost periodic sequences and functions: a necessary and sufficient condition for a sequence $u = (u_n)_{n \in \mathbb{Z}} \in AP_d(\mathbb{Z}, X)$ is the existence of $f \in AP(\mathbb{R}, X)$ such that $u_n = f(n)$ for all $n \in \mathbb{Z}$. This permits us to assert that the restriction operator

$$(8.1) \quad R : AP(\mathbb{R}, X) \rightarrow AP_d(\mathbb{Z}, X) \quad \text{defined by } R(f) = f|_{\mathbb{Z}}$$

is well-defined and maps $AP(\mathbb{R}, X)$ onto $AP_d(\mathbb{Z}, X)$. Let $Q = AP(\mathbb{R}, X)/\text{Ker } R$ and let $\Pi_0 : AP(\mathbb{R}, X) \rightarrow Q$ be the quotient map, so the surjective linear map R induces a unique isomorphism

$$\tilde{R} : Q \rightarrow AP_d(\mathbb{Z}, X) \quad \text{defined by } R = \tilde{R} \circ \Pi_0$$

satisfying for $f \in AP(\mathbb{R}, X)$

$$\|\Pi_0(f)\|_Q = \inf \left\{ \sup_{t \in \mathbb{R}} \|f(t) + g(t)\| ; g \in \text{Ker } R \right\} = \sup_{n \in \mathbb{Z}} \|f(n)\| = \left\| \tilde{R}(\Pi_0(f)) \right\|_{AP_d(\mathbb{Z}, X)},$$

therefore $Q = AP(\mathbb{R}, X)/\text{Ker } R$ is isometrically isomorphic to $AP_d(\mathbb{Z}, X)$.

We denote by \mathcal{M}_d the set of all two-sided sequences $p = (p_n)_{n \in \mathbb{Z}} \in \mathbb{R}^{\mathbb{Z}}$ satisfying $p_n \geq 0$, for all $n \in \mathbb{Z}$ and $\sum_{n \in \mathbb{Z}} p_n = +\infty$.

Definition 8.2. [1, 33] Let $p = (p_n)_{n \in \mathbb{Z}} \in \mathcal{M}_d$. $u = (u_n)_{n \in \mathbb{Z}} \in \ell^\infty(\mathbb{Z}, X)$ is said to be *weighted ergodic* if

$$\lim_{N \rightarrow +\infty} \frac{1}{\sum_{n=-N}^N p_n} \sum_{n=-N}^N \|u_n\| p_n = 0.$$

We denote the space of all such functions by $\mathcal{E}_d(\mathbb{Z}, X, p)$.

Proposition 8.3. Let $p \in \mathcal{M}_d$. Then $(\mathcal{E}_d(\mathbb{Z}, X, p), \|\cdot\|_\infty)$ is a Banach space.

Proof. It is enough to prove that $\mathcal{E}_d(\mathbb{Z}, X, p)$ is closed in $\ell^\infty(\mathbb{Z}, X)$. Let $(u^n)_n$ be a sequence in $\mathcal{E}_d(\mathbb{Z}, X, p)$ such that $\lim_{n \rightarrow +\infty} u^n = u$ (cv in $\ell^\infty(\mathbb{Z}, X)$). Then we have

$$\sum_{m=-N}^N \|u_m\| p_m \leq \sum_{m=-N}^N \|u_m - u_m^n\| p_m + \sum_{m=-N}^N \|u_m^n\| p_m$$

we deduce that

$$\frac{1}{\sum_{n=-N}^N p_n} \sum_{m=-N}^N \|u_m\| p_m \leq \sup_{m \in \mathbb{Z}} \|u_m - u_m^n\| + \frac{1}{\sum_{n=-N}^N p_n} \sum_{m=-N}^N \|u_m^n\| p_m,$$

it follows

$$\limsup_{N \rightarrow +\infty} \frac{1}{\sum_{n=-N}^N p_n} \sum_{m=-N}^N \|u_m\| p_m \leq \sup_{m \in \mathbb{Z}} \|u_m - u_m^n\| \quad \text{for all } n \in \mathbb{N}.$$

Since $\lim_{n \rightarrow +\infty} \|u - u^n\|_{\ell^\infty(\mathbb{Z}, X)} = \lim_{n \rightarrow +\infty} \left(\sup_{m \in \mathbb{Z}} \|u_m - u_m^n\| \right) = 0$, we deduce that

$$\lim_{N \rightarrow +\infty} \frac{1}{\sum_{n=-N}^N p_n} \sum_{m=-N}^N \|u_m\| p_m = 0.$$

□

Definition 8.4. [33] Let $p = (p_n)_{n \in \mathbb{Z}} \in \mathcal{M}_d$. $u = (u_n)_{n \in \mathbb{Z}} \in \ell^\infty(\mathbb{Z}, X)$ is said to be *weighted pseudo almost periodic* if

$$u = v + w \quad \text{where } v \in AP_d(\mathbb{Z}, X), \quad w \in \mathcal{E}_d(\mathbb{Z}, X, p).$$

We denote the space of all such functions by $PAP_d(\mathbb{Z}, X, p)$.

Remark 8.5. Contrarily to the pseudo almost periodic case: $p_n = 1$ for all $n \in \mathbb{Z}$, in general the sum is not direct for the weighted pseudo almost periodic case. For example if $p = (p_n)_{n \in \mathbb{Z}}$ is defined by $p_{2n} = 0$, $p_{2n+1} = 1$ and $u = (u_n)_{n \in \mathbb{Z}}$ by $u_{2n} = 1$, $u_{2n+1} = 0$ for all $n \in \mathbb{Z}$, then $u \in AP_d(\mathbb{Z}, \mathbb{R}) \cap \mathcal{E}_d(\mathbb{Z}, \mathbb{R}, p)$.

Theorem 8.6. Let $p \in \mathcal{M}_d$. Then $(PAP_d(\mathbb{Z}, X, p), \|\cdot\|_\infty)$ is a Banach space.

Proof. We use Theorem 4.1 by setting $U = \mathbb{Z}$, $\mathcal{F}_1 = AP_d(\mathbb{Z}, X)$ and $\mathcal{F}_2 = \mathcal{E}_d(\mathbb{Z}, X, p)$. Remark that the Banach space $B(\mathbb{Z}, X)$ used in Theorem 4.1 is $\ell^\infty(\mathbb{Z}, X)$.

i) $AP_d(\mathbb{Z}, X)$ is isometrically isomorphic to the Banach space $AP(\mathbb{R}, X)/\text{Ker } R$, then it is a Banach space. Let $\Phi \in C(X, X)$ and $u = (u_n)_{n \in \mathbb{Z}} \in AP_d(\mathbb{Z}, X)$. First there exists $f \in AP(\mathbb{R}, X)$ such that $R(f) = u$ (see (8.1)). Secondly by using the definition of the restriction operator R we obtain $\Phi \circ u = \Phi \circ [R(f)] = R(\Phi \circ f)$. From the fact that $\Phi \circ f \in AP(\mathbb{R}, X)$ (see proof of Theorem 5.5) we deduce that $\Phi \circ u \in AP_d(\mathbb{Z}, X)$; so Hypothesis i) is satisfied.

ii) By using Definition 8.2, we deduce that the pair $AP_d(\mathbb{Z}, X)$ and $\mathcal{E}_d(\mathbb{Z}, X, p)$ satisfies Hypothesis ii).

By Theorem 4.1, we obtain that $PAP_d(\mathbb{Z}, X, p) = AP_d(\mathbb{Z}, X) + \mathcal{E}_d(\mathbb{Z}, X, p)$ is a Banach space endowed with the norm $\|\cdot\|_\infty$. □

8.2. Almost automorphic case.

Definition 8.7. [3]. $u = (u_n)_{n \in \mathbb{Z}} \in X^{\mathbb{Z}}$ is said to be *almost automorphic* if for every sequence of integer numbers $(k'_n)_{n \in \mathbb{N}} \in \mathbb{Z}^{\mathbb{N}}$ there exists a subsequence of $(s'_n)_n$ denoted by $(s_n)_n$ such that

$$v_p = \lim_{n \rightarrow \infty} u_{p+k_n} \text{ exists for all } p \text{ in } \mathbb{Z}$$

and

$$\lim_{n \rightarrow \infty} v_{p-k_n} = u_p \text{ for all } p \text{ in } \mathbb{Z}.$$

We denote the space of all such functions by $AA_d(\mathbb{Z}, X)$.

The following definition is due to Abbas in [1] but it is not proved that this space is closed in $\ell^\infty(\mathbb{Z}, X)$.

Definition 8.8. Let $p = (p_n)_{n \in \mathbb{Z}} \in \mathcal{M}_d$. $u = (u_n)_{n \in \mathbb{Z}} \in \ell^\infty(\mathbb{Z}, X)$ is said to be *weighted pseudo almost automorphic* if

$$u = v + w \quad \text{where } v \in AA_d(\mathbb{Z}, X), \quad w \in \mathcal{E}_d(\mathbb{Z}, X, p).$$

We denote the space of all such functions by $PAA_d(\mathbb{Z}, X, p)$.

Theorem 8.9. Let $p \in \mathcal{M}_d$. Then $(PAA_d(\mathbb{Z}, X, p), \|\cdot\|_\infty)$ is a Banach space.

Proof. We use Theorem 4.1 by setting $U = \mathbb{Z}$, $\mathcal{F}_1 = AA_d(\mathbb{Z}, X)$ and $\mathcal{F}_2 = \mathcal{E}_d(\mathbb{Z}, X, p)$.

i) $AA_d(\mathbb{Z}, X)$ is a closed vector subspace of $\ell^\infty(\mathbb{Z}, X)$. If $\Phi \in C(X, X)$ and $v \in AA_d(\mathbb{Z}, X)$ then $\Phi \circ v \in AA(\mathbb{Z}, X)$ [3]; so Hypothesis i) is satisfied.

ii) is similar to ii) in the proof of Theorem 8.6.

By Theorem 4.1, we obtain that $PAA_d(\mathbb{Z}, X, p) = AA_d(\mathbb{Z}, X) + \mathcal{E}_d(\mathbb{Z}, X, p)$ is a Banach space endowed with the norm $\|\cdot\|_\infty$. \square

REFERENCES

- [1] S. Abbas, *Weighted pseudo almost automorphic sequences and their applications*, Electron. J. Differential Equations **121** (2010), 1-14.
- [2] L. Amerio, G. Prouse, *Almost periodic functions and functional equations*, Van Nostrand Reinhold Comp., New York, 1971.
- [3] D. Araya, R. Castro, C. Lizama, *Almost automorphic solutions of difference equations*, Adv. Difference Equ. (2009), Art. ID 591380.
- [4] W. Arendt, C. Batty, M. Hieber, F. Neubrander, *Vector-valued Laplace Transforms and Cauchy Problems*, Monographs in Mathematics, Birkhauser, 2001.
- [5] M. Bachir, J. Blot, *Infinite dimensional infinite-horizon Pontryagin principles for discrete-time problems*, Set-Valued Var. Anal. **23** (2015), 43-54.
- [6] F. Bedouhene, N. Challali, O. Mellah, P. Raynaud de Fitte, M. Smaali, *Almost automorphy and various extensions for stochastic processes*, J. Math. Anal. Appl. **429** (2015), 1113-1152.
- [7] P. Bezandry, T. Diagana, *Square-mean almost periodic solutions nonautonomous stochastic differential equations*, Electron. J. Differential Equations **117** (2007), 1-10.
- [8] P. Bezandry, T. Diagana, *Existence of almost periodic solutions to some stochastic differential equations*, Appl. Anal. **86** (2007), 819-827.
- [9] J. Blot, P. Cieutat, K. Ezzinbi, *Measure theory and pseudo almost automorphic functions, new developments and applications*, Nonlinear Anal. **75** (2012), 2426-2447.
- [10] J. Blot, P. Cieutat, K. Ezzinbi, *New approach for weighted pseudo-almost periodic functions under the light of measure theory, basic theory and applications*, Appl. Anal. **92** (2013), 493-526.
- [11] J. Blot, G.M. Mophou, G.M. N'Guérékata, D. Pennequin, *Weighted pseudo almost automorphic functions and applications to abstract differential equations*, Nonlinear Anal. **71** (2009), 903-909.
- [12] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Springer Science + Business Media LLC, New York, 2011.
- [13] Z. Chen, W. Lin, *Square-mean pseudo almost automorphic process and its application to stochastic evolution equations*, J. Funct. Anal. **261**(2011), 69-89.
- [14] Z. Chen, W. Lin, *Square-mean weighted pseudo almost automorphic solutions for non-autonomous stochastic evolution equations*, J. Math. Pures Appl. **100** (2013), 476-504.
- [15] C. Corduneanu, *Almost periodic functions*, Wiley, New York, 1968. Reprinted, Chelsea, New York, 1989.
- [16] T. Diagana, *Weighted pseudo almost periodic functions and applications*, C. R. Math. Acad. Sci. Paris, **343** (2006), 643-646.
- [17] T. Diagana, *Weighted pseudo-almost periodic solutions to some differential equations*, Nonlinear Anal. **68** (2008), 2250-2260.
- [18] M.A. Diop, K. Ezzinbi, M.M. Mbaye, *Existence and global attractiveness of a pseudo almost periodic solution in p-th mean sense for stochastic evolution equation driven by a fractional Brownian*

- motion*, Stochastics. An International Journal of Probability and Stochastic Processes (2015), DOI 10.1080/17442508.2015.1026345.
- [19] M.A. Diop, K. Ezzinbi, M.M. Mbaye, *Measure theory and S^2 -pseudo almost periodic and automorphic process: application to stochastic evolution equations*, Afr. Mat. **26** (2015), 779-812.
 - [20] N. Dunford, J.T. Schwartz, *Linear operators, Part I: general theory*, Interscience Publishers, Inc., New York, 1958.
 - [21] C. F. Dunkl and K. S. Williams, *A simple norm inequality*, Amer. Math. Monthly **71** (1964), 53-54.
 - [22] M. Fréchet, *Les fonctions asymptotiquement presque-périodiques continues*, C. R. Math. Acad. Sci. Paris, **213** (1941), 520-522 (in French).
 - [23] M. Fréchet, *Sur le théorème ergodique de Birkhoff*, C. R. Math. Acad. Sci. Paris, **213** (1941), 607-609 (in French).
 - [24] M. Fu, Z. Liu, *Square-mean almost automorphic solutions for some stochastic differential equations*, Proc. Amer. Math. Soc. **138** (2010), 3689-3701.
 - [25] C. Ganessa Moorthy, P.S. Johnson, *Composition of closed range operators*, J. Anal. **12** (2004), 165-169.
 - [26] T. Kato, *Perturbation theory for linear operators*, Second edition, Springer-Verlag, Berlin, 1980.
 - [27] H. Kober, *A theorem on Banach spaces*, Compositio Math. **7** (1940), 135-140.
 - [28] J. Liang, G.M. N'Guérékata, T-J. Xiao, J. Zhang, *Some properties of pseudo-almost automorphic functions and applications to abstract differential equations*, Nonlinear Anal. **70** (2009), 2731-2735.
 - [29] G. M. N'Guérékata, *Almost Automorphic and Almost Periodic Functions in Abstract Spaces*, Kluwer Academic/Plenum Publishers, New York, Boston, Moscow, London, 2001.
 - [30] L. Schwartz, *Analyse I; théorie des ensembles et topologie*, Hermann, Paris, 1991.
 - [31] C. Zhang, *Pseudo almost periodic solutions of differential equations*, J. Math. Anal. **181** (1994), 62-76.
 - [32] C. Zhang, *Almost periodic type functions and ergodicity*, Science Press, Beijing; Kluwer Academic Publishers, Dordrecht, 2003.
 - [33] L.-L. Zhang, H-X. Li, *Weighted pseudo almost periodic solutions of second-order neutral-delay differential equations with piecewise constant argument*, Comput. Math. Appl. **62** (2011), 4362-4376.
 - [34] Z.-M. Zheng, H.-S. Ding, *On completeness of the space of weighted pseudo almost automorphic functions*, J. Funct. Anal. **268** (2015), 3211-3218.
 - [35] Z.-M. Zheng, H.-S. Ding, *A note on closedness of the sum of two closed subspaces in a Banach space*, arXiv:1509.06445v1 [math.FA] 22 Sep 2015.