



HAL
open science

Critical Velocity and Dissipation of an ultracold Bose-Fermi Counterflow

Marion Delehaye, Sébastien Laurent, Igor Ferrier-Barbut, Shuwei Jin, Frédéric Chevy, Christophe Salomon

► **To cite this version:**

Marion Delehaye, Sébastien Laurent, Igor Ferrier-Barbut, Shuwei Jin, Frédéric Chevy, et al.. Critical Velocity and Dissipation of an ultracold Bose-Fermi Counterflow. 2015. hal-01214160v1

HAL Id: hal-01214160

<https://hal.science/hal-01214160v1>

Preprint submitted on 9 Oct 2015 (v1), last revised 20 Nov 2015 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Critical Velocity and Dissipation of an ultracold Bose-Fermi Counterflow

Marion Delehaye,^{1,*} Sébastien Laurent,^{1,*} Igor Ferrier-Barbut,^{1,†}
Shuwei Jin,¹ Frédéric Chevy,¹ and Christophe Salomon¹

¹*Laboratoire Kastler Brossel, ENS-PSL, CNRS, UPMC-Sorbonne Universités and Collège de France, Paris, France*

We study the dynamics of counterflowing bosonic and fermionic lithium atoms. First, by tuning the interaction strength we measure the critical velocity v_c of the system in the BEC-BCS crossover in the low temperature regime and we compare it to the recent prediction of Castin *et al.*, *Comptes Rendus Physique*, **16**, 241 (2015). Second, raising the temperature of the mixture slightly above the superfluid transitions reveals an unexpected phase-locking of the oscillations of the clouds. We interpret this observation as resulting from a Zeno-like effect induced by dissipation.

PACS numbers: 03.75.Kk, 03.75.Ss, 37.10.Gh

Superconductivity and superfluidity are spectacular macroscopic manifestations of quantum physics at low temperature. Besides liquid helium 4 and helium 3, dilute quantum gases have emerged over the years as a versatile tool to probe superfluid properties in diverse and controlled situations. Frictionless flows have been observed with both bosonic and fermionic atomic species, in different geometries and in a large range of interaction parameters from the weakly interacting Bose gas to strongly correlated fermionic systems [1–5]. Several other hallmarks of superfluidity such as quantized vortices or second sound were also observed in cold atoms [6–8].

A peculiar feature of superfluid flows is the existence of a critical velocity above which dissipation arises. In Landau’s original argument, this velocity is associated with the threshold for creation of elementary excitations in the superfluid: for a linear dispersion relation, it predicts that the critical velocity is simply given by the sound velocity in the quantum liquid. This critical velocity has been measured both in superfluid helium and ultracold atoms [1, 3–5, 9]. However the recent production of a Bose-Fermi double superfluid [10] raised new questions on Bose-Fermi mixtures [11–14] and interrogations on the validity of Landau’s argument in the case of superfluid counterflow [15–20].

In this letter, we study the dynamics of a Bose-Fermi superfluid counterflow in the BEC-BCS crossover and at finite temperature. We show how friction arises when the relative velocity of the Bose and Fermi clouds increases and we confirm that damping occurs only above a certain critical relative velocity v_c . We compare our measurements to Landau’s prediction and its recent generalization $v_c = c_s^F + c_s^B$ where c_s^F and c_s^B are the sound velocities of the fermionic and bosonic components respectively [16]. Finally, we study finite temperature damping

of the counterflow and we show that the system can be mapped onto a Caldeira-Leggett-like model [21] of two quantum harmonic oscillators coupled to a bath of excitations. This problem has been recently studied as a toy model for decoherence in quantum networks [22] or for heat transport in crystals [23] and we show here that the emergence of dissipation between the two clouds leads to a Zeno effect which locks their relative motions.

Our Bose and Fermi double-superfluid set-up was previously described in [10]. We prepare vapors of bosonic (B) ^7Li atoms spin-polarized in the second-to-lowest energy state and fermionic (F) ^6Li atoms prepared in a balanced mixture of the two lowest spin states noted $|\uparrow\rangle$, $|\downarrow\rangle$. The two species are kept in the same cigar-shaped hybrid magnetic-optical trap in which evaporative cooling is performed in the vicinity of the 832 G ^6Li Feshbach resonance [24]. The final number of fermions $N_F = 2.5 \times 10^5$ greatly exceeds that of the bosons $N_B \sim 2.5 \times 10^4$ and the temperature of the sample is adjusted by stopping the evaporation at different trap depths. The thermal pedestal surrounding the ^7Li Bose-Einstein Condensate (BEC) provides a convenient low temperature thermometer for both species after sufficiently long thermalization time (~ 1 second). The lowest temperature achieved in this study corresponds to almost entirely superfluid clouds with $T/T_{c,\alpha=B,F} \leq 0.5$, where $T_{c,\alpha}$ is the superfluidity transition temperature of species α .

The magnetic field values used in the experiment (780 – 880 G) enable us to scan the fermion-fermion interaction within a range $-0.5 \leq 1/k_F a_F \leq 1$. Here, a_F is the s-wave scattering length between $|\uparrow\rangle$ and $|\downarrow\rangle$ fermions and the Fermi momentum k_F is defined by $\hbar^2 k_F^2 / 2m_F = \hbar \bar{\omega} (3N_F)^{1/3}$ with $\bar{\omega}$ the geometric mean of the trap frequencies, and N_F the total number of fermions of mass m_F . In our shallowest traps, typical trap frequencies for ^6Li are $\omega_x = \omega_y = 2\pi \times 550$ Hz and $\omega_z = 2\pi \times 17$ Hz. Since the bosonic and fermionic isotopes experience the same trapping potentials, the oscillation frequencies of the two species are within a ratio $\sqrt{6/7} \simeq 0.9$.

We excite the dipole modes of the system by displac-

*These two authors contributed equally to this work

†Present Address: 5. Physikalisches Institut and Center for Integrated Quantum Science and Technology, Universität Stuttgart, Pfaffenwaldring 57, 70550 Stuttgart, Germany




FIG. 1: Center-of-mass oscillations of bosons (blue, top) and fermions (red, bottom), for different sets of parameters at unitarity. Solid lines: fits using Eq.1 for the bosons and a similar equation for the fermions. (a) $T/T_F = 0.03$, $T/T_{c,b} \leq 0.5$, $z_0 = 10 \mu\text{m}$. Superfluid regime, no damping is observed and $\omega_B = 2\pi \times 15.41(1) \text{ Hz} \approx \sqrt{6/7} \omega_F$. The observed beating at $\omega_F - \omega_B$ is due to coherent energy exchange between the clouds. (b) $T/T_F = 0.03$ and $z_0 = 150 \mu\text{m}$. For a larger initial displacement, initial damping ($\gamma_B = 2.4 \text{ s}^{-1}$) is followed by steady-state evolution. $\omega_B = 2\pi \times 14.2(1) \text{ Hz} \approx \sqrt{6/7} \omega_F$. (c) $T/T_F = 0.4$ and $z_0 = 80 \mu\text{m}$. At higher temperature, phase-locking of the two frequencies is observed with $\omega_F \approx \omega_B = 2\pi \times 17.9(3) \text{ Hz}$ and $\gamma_B = \gamma_F = 1.4(5) \text{ s}^{-1}$.

ing adiabatically the centers of mass of the clouds from their initial position by a distance z_0 along the weakly confined z direction, and abruptly releasing them in the trap. The two clouds evolve for a variable time t before *in situ* absorption images perpendicular to z direction are taken. The measurement of their doubly integrated density profiles gives access to axial positions and atom




FIG. 2: Damping rate of the center of mass oscillations versus maximal relative velocity in the BEC-BCS crossover in unit of the Fermi velocity v_F . Dark blue dots: BEC side (780 G) $1/k_F a_F = 0.68$, red squares: unitarity (832.2 G) $1/k_F a_F = 0$, light blue diamonds: BCS side (880 G) $1/k_F a_F = -0.42$. Power law fits with thresholds provide the critical velocity (Solid lines).

numbers of both species. Typical time evolutions of the centers of mass are shown in Fig.1 for different parameter values. Since the Bose and Fermi components oscillate at different frequencies, they oscillate in quadrature after a few periods. By changing z_0 , we can thus tune the maximum relative velocity between the two clouds and probe the critical superfluid counterflow.

As shown in Fig.1 a, the superfluid counterflow exhibits no visible damping on a $\simeq 5 \text{ s}$ time scale for very low temperature and small initial displacement. A striking feature is the beat note on the ${}^7\text{Li}$ oscillation amplitude due to the coherent mean-field coupling to the ${}^6\text{Li}$ cloud [10]. For larger relative velocities, ${}^7\text{Li}$ oscillations are initially damped (Fig.1 b) until a steady-state regime as in Fig.1 a is reached. We fit the time evolution of the cloud position using the phenomenological law

$$z_B(t) = d(t) [a \cos(\omega_B t) + b \cos(\omega_F t)], \quad (1)$$

$$d(t) = d_1 + d_2 \exp(-\gamma_B t).$$

We measure the damping rate γ_B as a function of relative velocity for six different values of magnetic field, exploring a large region of the crossover going from the BCS ($1/k_F a_F = -0.42$, $B = 880 \text{ G}$) to the BEC side ($1/k_F a_F = 0.68$, $B = 780 \text{ G}$), see Fig.2. For these magnetic field values, the Bose gas remains in the weakly interacting (repulsive) regime and the Bose-Fermi scattering length is $a_{BF} \simeq 41a_0$, constant in this magnetic field range, and equal for both $|\uparrow\rangle$ and $|\downarrow\rangle$ spin states.

We extract the critical velocity v_c using an *ad-hoc* power-law fitting function $\gamma_B = A\Theta(v-v_c)((v-v_c)/v_F)^\alpha$, where θ is the Heaviside function and v_F is the Fermi velocity given by $v_F = \hbar k_F/m_F$. For details see [28]. v_c in the BEC-BCS crossover is displayed in Fig.3 (red dots)




FIG. 3: Critical velocity of the Bose-Fermi superfluid counterflow in the BEC-BCS crossover normalized to the Fermi velocity v_F . Red dots: measurements. Red dot-dashed line: sound velocity c_s^F of an elongated homogeneous Fermi superfluid deduced from its equation of state [25, 26] and measured in [27]. Blue bars: calculated sound velocity c_s^B of the elongated ${}^7\text{Li}$ BEC for each magnetic field, (880 G, 860 G, 832 G, 816 G, 800 G, 780 G). Green squares indicate the prediction $v_c = c_s^F + c_s^B$. Error bars and c_s^B are discussed in [28].

and compared to Landau's and Castin *et al.* [16] predictions. In this later work, dissipation arises by creation of excitation pairs and yields a critical velocity $v_c = \text{Min}_{\substack{\mathbf{p} \\ \sigma=f,b}} \left(\frac{\epsilon^B(\mathbf{p}) + \epsilon_\sigma^F(\mathbf{p})}{p} \right)$. In this expression, $\epsilon^B(\mathbf{p})$ denotes the dispersion relation of excitations in the BEC and $\epsilon_\sigma^F(\mathbf{p})$ refers to the two possible branches of the Fermi superfluid, phonon-like ($\sigma = b$) and threshold for pair breaking excitations ($\sigma = f$) [29]. For homogeneous gases, at unitarity and on the BEC side of the crossover, this critical relative velocity turns out to be simply the sum of the respective sound velocities of the Bose and Fermi superfluids, $v_c = c_s^F + c_s^B$. We thus plot in Fig.3 the calculated sound velocities of both superfluids in an elongated geometry [25, 26, 30, 31] (red dashed line c_s^F , blue bars c_s^B). Typically c_s^B contributes $\simeq 20 - 25\%$ to the sum shown as blue dots in Fig.3. Around unitarity and on the BCS side of the resonance, our experimental data is consistent with this interpretation as well as with a critical velocity $v_c = c_s^F$ that one would expect by considering the BEC as a single impurity moving inside the fermionic superfluid. By contrast, we clearly exclude the bosonic sound velocity as a threshold for dissipation.

Our measured critical velocities are significantly higher than those previously reported in pure fermionic systems which, for all interaction strengths, were lower than Landau's criterion [3, 5]. The main difference with our study is the use of focused laser beams instead of a BEC as a moving obstacle. In [5], the laser beam is piercing the whole cloud including its non-superfluid part where the




FIG. 4: Power spectrum of the oscillations for different temperatures, obtained using the Lomb-Scargle algorithm of the center-of-mass displacement. Above $T \approx T_{c,B} \approx 0.34 T_F > T_{c,F}$, oscillations of the Bose and Fermi clouds become locked together at ω_F . A value of 10 for the power represents typically a significance of 0.002 [28].

density is low, and its potential may create a strong density modulation of the superfluid. These effects make a direct comparison to Landau criterion difficult [32]. On the contrary, in our system the size of the BEC (Thomas Fermi radii of 73, 3, 3 μm) is much smaller than the typical size of the Fermi cloud (350, 13, 13 μm around unitarity). For oscillation amplitudes up to $\pm 200 \mu\text{m}$ the BEC probes only the superfluid core of the fermionic cloud. As the mean-field interaction between the two clouds is very small [28] our BEC acts as a finite mass weakly interacting probe.

On the BEC side of the resonance (780 G) however, we observe a strong reduction of the measured critical velocity compared to the predicted values. This anomalously small value for positive scattering lengths is consistent with previous measurements [3, 5]. Its origin is still unclear but several explanations can be put forward [32]. Firstly, it is well known that vortex shedding can strongly reduce superfluid critical velocity. However, this mechanism requires a strong perturbation and the mean-field interaction between the two clouds is probably too small for vortex generation. Secondly, the critical velocity is given by the sound velocity of the cloud if the dispersion relation is *convex*. For a weakly-interacting, homogeneous Bose gas described by the Bogoliubov mean-field theory, this is indeed the case. However the convexity of the low-energy dispersion relation is inverted in a confined geometry [30] and this anomalous behaviour lowers the critical velocity [33].

We now present results of experiments performed at higher temperature ($0.03 \lesssim T/T_F \lesssim 0.5$) for $B = 835$ G. For low temperatures ($T/T_F \leq 0.2$), the two clouds re-

main weakly coupled and, as observed in the power spectrum of Fig.4, the bosonic and fermionic components oscillate at frequencies in the expected ratio $\simeq 0.9 \simeq \sqrt{6/7}$. A new feature emerges for $T \gtrsim T_{c,B} \approx 0.34 T_F > T_{c,F}$ where both gases are in the normal phase. In this “high” temperature regime, the two clouds are locked in phase: ${}^7\text{Li}$ oscillates at ${}^6\text{Li}$ frequency (Fig.4) and the two components are equally damped (Fig.1.c). This remarkable behavior can be understood as a Zeno effect arising from the increased dissipation between the two components. Indeed the system can be described as a set of two harmonic oscillators describing respectively the macroscopic motion of the global center of mass of the system (Kohn’s mode [34]) and the relative motion of the two clouds [28]. These two degrees of freedom are themselves coupled to the “bath” of the internal excitations of the two clouds (breathing mode, quadrupole modes, pair breaking excitations...).

In the spirit of the dressed-atom picture, we can represent the state of the two harmonic oscillators by the “radiative” cascade of Fig.5. Here the states $|N, n\rangle$ are labeled by the quantum numbers associated to Kohn’s mode (N) and relative motion (n) of the two clouds and we trace-out the degrees of freedom of the bath. On the one hand, Kohn’s mode is not an eigenstate of the system for fermions and bosons of different masses; center-of-mass and relative-motion modes are coupled and this coherent coupling is responsible for the dephasing of the oscillations of the two clouds in the weakly interacting regime. On the other hand interspecies interactions do not act on the center of mass of the whole system, owing to Kohn’s theorem, but on the contrary lead to an irreversible “radiative” decay of the *relative* motion at a rate γ .

In our experiments, the initial state is a pure center of mass excitation $|N, 0\rangle$. If we neglect the interspecies coupling, the system evolves in the subspace spanned by $|N - n, n\rangle_{n=0,\dots,N}$ of the two coupled oscillators and the system oscillates at a frequency $\delta\omega \simeq \omega_B - \omega_F$ as the centers of mass of the Bose and Fermi clouds dephase. If we now consider the opposite limit where the decay rate γ is larger than the dephasing frequency $\delta\omega$, the strong coupling to the bath prevents the conversion of the center of mass excitations into relative motion. As soon as the system is transferred into $|N - 1, 1\rangle$ it decays towards state $|N - 1, 0\rangle$. Similarly to optical pumping in quantum optics, we can eliminate adiabatically the excited states of the relative motion and restrict the dynamics of the system to the subspace $|N, 0\rangle_{N=0,\dots,\infty}$ of Kohn’s excitations. This situation is akin to Zeno effect where strong coupling to the environment inhibits coherent evolution. It is also reminiscent of the synchronization of two spins immersed in a thermal bath predicted in [35] and to a phenomenological classical two-coupled oscillators model.

In this letter we have investigated how a Bose-Fermi superfluid flow is destabilized by temperature or relative




FIG. 5: Radiative cascade of the center of mass motion. In $|N, n\rangle$, N (resp. n) refers to the center of mass (resp. relative) motion of the two clouds (see text). When the decay rate of the relative motion is larger than the oscillation frequency difference between the two species, the dynamics is restricted to the center of mass degree of freedom: in this Zeno-like process, dissipation prevents excitation of the relative motion and the center of mass modes of the Bose and Fermi gases do not dephase.

velocity between the two clouds. In the limit of very low temperature the measured critical velocity for superfluid counterflow slightly exceeds the speed of sound of the elongated Fermi superfluid and decreases sharply towards the BEC side of the BEC-BCS crossover. In a future study, we will investigate the role of the confining potential and of the accelerated motion of the two clouds [32] that should provide a more accurate model for the damping rate versus velocity. Identifying the quasiparticles responsible for dissipation would also be a decisive step towards a full understanding of Bose-Fermi counterflows.

The authors acknowledge support from Région Ile de France (Atomix Project), ERC (ThermoDynaMix Project) and Institut de France (Louis D. Prize). They thank I. Danaila, N. Proukakis, K.L. Lee and M. Pierce for insightful comments and discussions, and J. Dalibard, Y. Castin, S. Nascimbène and T. Yesfah for critical reading of the manuscript.

-
- [1] C. Raman, M. Köhl, R. Onofrio, D. S. Durfee, C. E. Kuklewicz, Z. Hadzibabic, and W. Ketterle. Evidence for a critical velocity in a Bose-Einstein condensed gas. *Phys. Rev. Lett.*, 83:2502–2505, Sep 1999.
 - [2] C. Ryu, M. F. Andersen, P. Cladé, Vasant Natarajan, K. Helmerson, and W. D. Phillips. Observation of persistent flow of a Bose-Einstein condensate in a toroidal trap. *Phys. Rev. Lett.*, 99:260401, Dec 2007.
 - [3] D. E. Miller, J. K. Chin, C. A. Stan, Y. Liu, W. Setiawan, C. Sanner, and W. Ketterle. Critical velocity for superfluid flow across the BEC-BCS crossover. *Phys. Rev. Lett.*, 99:070402, Aug 2007.
 - [4] Rémi Desbuquois, Lauriane Chomaz, Tarik Yefsah, Julian Léonard, Jérôme Beugnon, Christof Weitenberg, and Jean Dalibard. Superfluid behaviour of a two-

- dimensional Bose gas. *Nature Physics*, 8(9):645–648, 2012.
- [5] Wolf Weimer, Kai Morgener, Vijay Pal Singh, Jonas Siegl, Klaus Hueck, Niclas Luick, Ludwig Mathey, and Henning Moritz. Critical velocity in the BEC-BCS crossover. *Phys. Rev. Lett.*, 114:095301, Mar 2015.
- [6] KW Madison, F Chevy, W Wohlleben, and J Dalibard. Vortex formation in a stirred Bose-Einstein condensate. *Physical Review Letters*, 84(5):806, 2000.
- [7] M.W. Zwierlein, J.R. Abo-Shaer, A. Schirotzek, C.H. Schunck, and W. Ketterle. Vortices and superfluidity in a strongly interacting Fermi gas. *Nature*, 435(7045):1047–1051, 2005.
- [8] Leonid A Sidorenkov, Meng Khoon Tey, Rudolf Grimm, Yan-Hua Hou, Lev Pitaevskii, and Sandro Stringari. Second sound and the superfluid fraction in a Fermi gas with resonant interactions. *Nature*, 498(7452):78–81, 2013.
- [9] D. Dries, S. E. Pollack, J. M. Hitchcock, and R. G. Hulet. Dissipative transport of a Bose-Einstein condensate. *Phys. Rev. A*, 82:033603, Sep 2010.
- [10] Igor Ferrier-Barbut, Marion Delehaye, Sebastien Laurent, Andrew T Grier, Matthieu Pierce, Benno S Rem, Frédéric Chevy, and Christophe Salomon. A mixture of Bose and Fermi superfluids. *Science*, 345(6200):1035–1038, 2014.
- [11] Tomoki Ozawa, Alessio Recati, Marion Delehaye, Frédéric Chevy, and Sandro Stringari. Chandrasekhar-Clogston limit and critical polarization in a Fermi-Bose superfluid mixture. *Physical Review A*, 90(4):043608, 2014.
- [12] Ren Zhang, Wei Zhang, Hui Zhai, and Peng Zhang. Calibration of the interaction energy between Bose and Fermi superfluids. *Physical Review A*, 90(6):063614, 2014.
- [13] Xiaoling Cui. Atom-dimer scattering and stability of Bose and Fermi mixtures. *Physical Review A*, 90(4):041603, 2014.
- [14] J. J. Kinnunen and G. M. Bruun. Induced interactions in a superfluid Bose-Fermi mixture. *Phys. Rev. A*, 91:041605, Apr 2015.
- [15] Abad, Marta, Recati, Alessio, Stringari, Sandro, and Chevy, Frédéric. Counter-flow instability of a quantum mixture of two superfluids. *Eur. Phys. J. D*, 69(5):126, 2015.
- [16] Yvan Castin, Igor Ferrier-Barbut, and Christophe Salomon. La vitesse critique de Landau d’une particule dans un superfluide de fermions. *Comptes Rendus Physique*, 16:241, 2015.
- [17] Wei Zheng and Hui Zhai. Quasiparticle lifetime in a mixture of Bose and Fermi superfluids. *Physical Review Letters*, 113(26):265304, 2014.
- [18] Linghua Wen and Jinghong Li. Structure and dynamics of a rotating superfluid Bose-Fermi mixture. *Physical Review A*, 90(5):053621, 2014.
- [19] F. Chevy. Counterflow in a doubly superfluid mixture of bosons and fermions. *Phys. Rev. A*, 91:063606, Jun 2015.
- [20] Huitao Shen and Wei Zheng. Landau damping in a mixture of Bose and Fermi superfluids. *Phys. Rev. A*, 92:033620, Sep 2015.
- [21] Amir O Caldeira and Anthony J Leggett. Path integral approach to quantum brownian motion. *Physica A: Statistical mechanics and its Applications*, 121(3):587–616, 1983.
- [22] Chung-Hsien Chou, Ting Yu, and B. L. Hu. Exact master equation and quantum decoherence of two coupled harmonic oscillators in a general environment. *Phys. Rev. E*, 77:011112, Jan 2008.
- [23] U. Zürcher and P. Talkner. Quantum-mechanical harmonic chain attached to heat baths. I. Equilibrium properties. *Phys. Rev. A*, 42:3267–3277, Sep 1990.
- [24] G. Zürn, T. Lompe, A. N. Wenz, S. Jochim, P. S. Julienne, and J. M. Hutson. Precise characterization of ^6Li Feshbach resonances using trap-sideband-resolved RF spectroscopy of weakly bound molecules. *Phys. Rev. Lett.*, 110:135301, Mar 2013.
- [25] N. Navon, S. Nascimbène, F. Chevy, and C. Salomon. The equation of state of a low-temperature Fermi gas with tunable interactions. *Science*, 328(5979):729–732, 2010.
- [26] GE Astrakharchik. Quantum Monte Carlo study of ultracold gases (PhD thesis). *arXiv preprint arXiv:1412.4529*, 2014.
- [27] J. Joseph, B. Clancy, L. Luo, J. Kinast, A. Turlapov, and J. E. Thomas. Measurement of sound velocity in a Fermi gas near a Feshbach resonance. *Phys. Rev. Lett.*, 98:170401, Apr 2007.
- [28] See supplementary material.
- [29] R Combescot, M Yu Kagan, and S Stringari. Collective mode of homogeneous superfluid Fermi gases in the BEC-BCS crossover. *Physical Review A*, 74(4):042717, 2006.
- [30] S Stringari. Dynamics of Bose-Einstein condensed gases in highly deformed traps. *Phys. Rev. A*, 58(3):2385, 1998.
- [31] L. Luo, B. Clancy, J. Joseph, J. Kinast, and J.E. Thomas. Measurement of the entropy and critical temperature of a strongly interacting Fermi gas. *Phys. Rev. Lett.*, 98(8):80402, 2007.
- [32] Vijay Pal Singh, Wolf Weimer, Kai Morgener, Jonas Siegl, Klaus Hueck, Niclas Luick, Henning Moritz, and Ludwig Mathey. Probing superfluidity of Bose-Einstein condensates via laser stirring. *arXiv preprint arXiv:1509.02168*, 2015.
- [33] PO Fedichev and GV Shlyapnikov. Critical velocity in cylindrical Bose-Einstein condensates. *Physical Review A*, 63(4):045601, 2001.
- [34] Walter Kohn. Cyclotron resonance and de Haas-van Alphen oscillations of an interacting electron gas. *Phys. Rev.*, 123:1242–1244, Aug 1961.
- [35] Peter P. Orth, David Roosen, Walter Hofstetter, and Karyn Le Hur. Dynamics, synchronization, and quantum phase transitions of two dissipative spins. *Phys. Rev. B*, 82:144423, Oct 2010.
- [36] N.R. Lomb. Least-squares frequency analysis of unequally spaced data. *Astrophysics and Space Science*, 39(2):447–462, 1976.
- [37] J.D. Scargle. Studies in astronomical time series analysis. II - Statistical aspects of spectral analysis of unevenly spaced data. *Astrophysical Journal*, 263(835):835–853, 1982.

SUPPLEMENTAL INFORMATION

DAMPING RATES IN THE BEC-BCS CROSSOVER

All the center of mass (CoM) damping rates measured in the BEC-BCS crossover and their respective fitting functions to extract v_c are shown in Fig. 6. As indicated in the main text the fit function is $\gamma_B = A\Theta(v - v_c)((v - v_c)/v_F)^\alpha$, where θ is the Heaviside function and v_F is the Fermi velocity. The χ^2 test reveals that most of our data is consistent with $\alpha = 1$ as in [3, 5]. Nevertheless allowing α to vary between 0.5 and 2 induces a systematic correlation between the extracted α and v_c that we include in our error bars on v_c shown in Fig. 3. The fit results for $\alpha = 1$ are displayed in Tab. I along with experimental parameters to produce Fig. 3. The error bars given for v_c give the span of v_c when changing α from 0.5 to 2 in the fit function. Note the strong decrease of the damping rate towards the BCS regime.

In addition, c_s^B depends on the Bose-Bose scattering length which varies with magnetic field and in particular diverges for a magnetic field of 845.5 G (corresponding to $1/k_F a_F = -0.13$). We therefore show only the values of c_s^B at the six magnetic field values used in our experiments.

Following [15] the Bose-Fermi coupling can be characterized by the quantity $\Delta = \sqrt{\frac{\partial\mu_B}{\partial n_F} \frac{\partial\mu_F}{\partial n_B} / \frac{\partial\mu_B}{\partial n_B} \frac{\partial\mu_F}{\partial n_F}}$. As Δ increases, the interspecies interactions affect more and more the properties of the system and for $\Delta = 1$, the mixture is dynamically unstable at rest and demixes. For measurements presented here we typically have $\Delta \simeq 15\%$.

LOMB-SCARGLE ALGORITHM

We use the fit-free Lomb-Scargle periodogram - or Least Square Spectral Analysis - to extract the spectral components of the oscillations for different temperatures [36, 37]. This method is an adaptation of the Fourier transform to the case of unevenly spaced data. For N data points $\{h_i = h(t_i)\}_{i=1,\dots,N}$ taken at times $\{t_i\}$, the periodogram is defined as

$$P_N(\omega) = \frac{1}{2\sigma^2} \left\{ \frac{[\sum_j (h_j - \bar{h}) \cos \omega(t_j - \tau)]^2}{\sum_j \cos^2 \omega(t_j - \tau)} + \frac{[\sum_j (h_j - \bar{h}) \sin \omega(t_j - \tau)]^2}{\sum_j \sin^2 \omega(t_j - \tau)} \right\} \quad (2)$$

where τ is defined as $\tan(2\omega\tau) = \frac{\sum_j \sin 2\omega t_j}{\sum_j \cos 2\omega t_j}$ making the periodogram independent of the time origin. $\bar{h} = \frac{1}{N} \sum_{i=1}^N h_i$ and $\sigma = \frac{1}{N-1} \sum_{i=1}^N (h_i - \bar{h})^2$ are the mean and the variance of $\{h_i\}_i$. The periodogram, or power spectrum, gives access to the statistical significance (*ie* the probability of rejecting the null hypothesis when it is true) of each of the evaluated frequencies: noting $P_{\max} = \max_{\omega} P_N(\omega)$, the significance is proportional to $e^{-P_{\max}}$, and here a value of 10 for the power represents typically a significance of 0.002.

RADIATIVE CASCADE MODEL

Consider a mixture of two atomic species labeled by $\alpha = 1, 2$ of identical masses m and confined in identical harmonic traps. According to Kohn's theorem [34], the single-species Hamiltonian can always be written as

$$H_\alpha = \frac{P_\alpha^2}{2M_\alpha} + \frac{M_\alpha \omega^2 X_\alpha^2}{2} + H_{\text{int}}^{(\alpha)}, \quad (3)$$

where P_α is the total momentum of cloud α , X_α the position of its center of mass and $M_\alpha = N_\alpha m$ its total mass (N_α being the number of atoms). $H_{\text{int}}^{(\alpha)}$ acts only on the internal excitation modes of the cloud and commutes with the center-of-mass variables.

Neglecting interspecies interactions, the total Hamiltonian of the system can be written as $H = H_1 + H_2$. Introducing the center of mass/relative variables, we have then

$$H = H_1 + H_2 = \frac{P^2}{2M} + \frac{M\omega^2 X^2}{2} + \frac{p^2}{2\mu} + \frac{\mu\omega^2 x^2}{2} + H_{\text{int}}^{(1)} + H_{\text{int}}^{(2)}, \quad (4)$$

with the usual definitions $P = P_1 + P_2$, $X = (M_1 X_1 + M_2 X_2)/M$, $p = \mu(P_1/M_1 - P_2/M_2)$, $x = x_1 - x_2$, $M = M_1 + M_2$ and $\mu = M_1 M_2 / M$. The dynamics of the center of mass and relative variables are described by independent harmonic oscillators decoupled from the internal degrees of freedom. The decoupled base can therefore be written as $|N, n, \varphi\rangle$, where N (resp. n) is the excitation number of the center-of-mass (resp. relative) motion, and φ describes the state of the internal excitation modes.

Let's now add the interspecies interactions and the mass difference between the two species.

1. *Interspecies coupling*: interactions between the two species are described by the Hamiltonian

$$H_{1,2} = \sum_{i \leq N_1, j \leq N_2} U(x_{1,i} - x_{2,j}). \quad (5)$$




FIG. 6: Damping rates of the center of mass oscillations versus maximal relative velocity in the BEC-BCS crossover in unit of the Fermi velocity v_F . Red line: fit with $\alpha = 1$. Orange zone: region spanned by the fitting function when varying α from 0.5 to 2.

B (G)	780	800	816	832	860	880
$a_F(a_0)$	6.4×10^3	11.3×10^3	24.0×10^3	∞	-16.5×10^3	-10.3×10^3
$1/k_F a_F$	0.68 ± 0.07	0.39 ± 0.01	0.18 ± 0.02	0 ± 0.002	-0.26 ± 0.05	-0.42 ± 0.03
$a_B(a_0)$	21.3	30.8	43.3	69.5	76.0	259
$c_B(10^{-2}v_F)$	9.6 ± 1.4	9.4 ± 0.14	11.0 ± 1.6	11.1 ± 1.7	11.4 ± 1.7	15.1 ± 2.2
v_c/v_F	$0.17^{+0.06}_{-0.10}$	$0.38^{+0.02}_{-0.04}$	$0.35^{+0.04}_{-0.11}$	$0.42^{+0.08}_{-0.14}$	$0.54^{+0.02}_{-0.06}$	$0.40^{+0.10}_{-0.20}$
$A(\text{s}^{-1})$	14.8 ± 1.4	85 ± 32	24.6 ± 4.3	17.3 ± 3.6	30 ± 11	2.9 ± 0.5
v_c/c_s^F	$0.53^{+0.19}_{-0.31}$	$1.11^{+0.06}_{-0.12}$	$0.99^{+0.11}_{-0.31}$	$1.17^{+0.22}_{-0.39}$	$1.46^{+0.05}_{-0.16}$	$1.05^{+0.26}_{-0.53}$

TABLE I: Experimental parameters, sound velocity at the center of the Bose gas in an elongated geometry $c_B = \sqrt{\mu_B/2m_B}$, critical velocity v_c/v_F , damping rate $A(\text{s}^{-1})$, and v_c/c_s^F for $\alpha = 1$ in the BEC-BCS crossover. The typical number of bosons and fermions are constant in the crossover and are respectively $2.5 \pm 0.5 \times 10^4$ and $2.5 \pm 0.5 \times 10^5$.

where $x_{i,\alpha}$ is the position of the i -th particle of species α and U is the interspecies interaction potential.

Owing to Kohn's theorem, this Hamiltonian commutes with P and X and therefore couples the internal degrees of freedom only to the relative variables (x, p) .

2. *Mass difference*: assume that species α has a mass $m_\alpha = m + \epsilon_\alpha \delta m/2$, with $\epsilon_1 = 1$ and $\epsilon_2 = -1$. This mass difference adds to the kinetic energy a term

$$\delta H_K = -\frac{\delta m}{2m} \sum_{i,\alpha} \epsilon_\alpha \frac{p_{i,\alpha}^2}{2m}. \quad (6)$$

As before, we can isolate the center of mass contri-

bution and write

$$\delta H_K = -\frac{\delta m}{2m} \left[\frac{P_1^2}{2M_1} - \frac{P_2^2}{M_2} \right] + \delta H_{K,\text{int}}, \quad (7)$$

where $\delta H_{K,\text{int}}$ contributes to the internal energies of the clouds and commutes with the center-of-mass degrees of freedom.

Let's now insert these two contributions in the total Hamiltonian. We have

$$H = H_{\text{CoM}} + H_{\text{rel}} + H'_{\text{int}} + H_{1,2} + H_{\text{coh}}. \quad (8)$$

with

$$H_{\text{CoM}} = \frac{P^2}{2M} \left(1 - \rho \frac{\delta m}{2m}\right) + \frac{M\omega^2}{2} X^2 \quad (9)$$

$$H_{\text{rel}} = \frac{p^2}{2\mu} \left(1 + \rho \frac{\delta m}{2m}\right) + \frac{\mu\omega^2}{2} x^2 \quad (10)$$

$$H'_{\text{int}} = H_{\text{int}}^{(1)} + H_{\text{int}}^{(2)} + \delta H_{K,\text{int}}, \quad (11)$$

$$H_{\text{coh}} = \frac{\delta m}{m} \left(\frac{P \cdot p}{M} \right), \quad (12)$$

and $\rho = (M_1 - M_2)/M$.

This hamiltonian describes two harmonic oscillators (H_{CoM} and H_{rel}) coupled to a thermal bath (H'_{int}). The coupling is ensured by the $P \cdot p$ terms which couples only the center-of-mass and relative degrees of freedom, and $H_{1,2}$ that commutes with H_{CoM} , owing to Kohn's theorem, and couples the relative motion to the internal degrees of freedom of the two clouds.

The interaction between the relative degrees of freedom and the internal thermal bath described by $H_{\text{int}}^{1,2}$ leads to an irreversible decay of the relative motion. By contrast, H_{coh} reflects the coherent beating existing between the relative oscillations of the two species due to their oscillation-frequency difference. It can be expressed using the annihilation operators a and b for the center of mass and relative motions respectively. We have then

$$H_{\text{coh}} = -\frac{\delta m}{m} \hbar\omega \sqrt{\frac{\mu}{M}} (a - a^\dagger) (b - b^\dagger). \quad (13)$$

Using the rotating wave approximation, one can eliminate the non resonant terms and one finally gets

$$H_{\text{coh}} \simeq \frac{\delta m}{2m} \hbar\omega \sqrt{\frac{\mu}{M}} (a^\dagger b + ab^\dagger). \quad (14)$$

This Hamiltonian is similar to the generalized Caldeira-Leggett model [21] used in solid state physics to study heat transport by phonons in a crystal. The absence of coupling between the bath and Kohn's mode generalizes to the decomposition used in [22] for a bilinear coupling between the harmonic oscillators and the bath.

In the experiment, we excite the center of mass motion and the initial state is $|N, 0, \varphi\rangle$. The coherent coupling transfers the system to the state $|N - 1, 1, \varphi\rangle$. If the coupling to the bath is strong, the relative motion decays very fast and the system falls into a state $|N - 1, 0, \varphi'\rangle$ (actually, since several states of the bath are involved, it is more appropriate to describe the state of the two harmonic oscillators by a density matrix rather than a well-defined quantum state). In this case, just like for optical pumping, we can adiabatically eliminate the intermediate state $|N - 1, 1, \varphi\rangle$ and consider that the dynamics occurs only in the sub-space $|N, 0, \varphi\rangle$, where the relative motion is never excited and the centers of mass of the two clouds are locked. In some sense, this freezing of the system state in a pure motion of its center of mass can be considered as a manifestation of the quantum Zeno effect.