



HAL
open science

On Existence and Uniqueness of Solutions for Semilinear Fractional Wave Equations

Yavar Kian, Masahiro Yamamoto

► **To cite this version:**

Yavar Kian, Masahiro Yamamoto. On Existence and Uniqueness of Solutions for Semilinear Fractional Wave Equations. *Fractional Calculus and Applied Analysis*, 2017, 20 (1), pp.117-138. 10.1515/fca-2017-0006 . hal-01214747

HAL Id: hal-01214747

<https://hal.science/hal-01214747>

Submitted on 13 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON EXISTENCE AND UNIQUENESS OF SOLUTIONS FOR SEMILINEAR FRACTIONAL WAVE EQUATIONS

YAVAR KIAN AND MASAHIRO YAMAMOTO

ABSTRACT. Let Ω be a C^2 -bounded domain of \mathbb{R}^d , $d = 2, 3$, and fix $Q = (0, T) \times \Omega$ with $T \in (0, +\infty]$. In the present paper we consider a Dirichlet initial-boundary value problem associated to the semilinear fractional wave equation $\partial_t^\alpha u + \mathcal{A}u = f_b(u)$ in Q where $1 < \alpha < 2$, ∂_t^α corresponds to the Caputo fractional derivative of order α , \mathcal{A} is an elliptic operator and the nonlinearity $f_b \in C^1(\mathbb{R})$ satisfies $f_b(0) = 0$ and $|f_b'(u)| \leq C|u|^{b-1}$ for some $b > 1$. We first provide a definition of local weak solutions of this problem by applying some properties of the associated linear equation $\partial_t^\alpha u + \mathcal{A}u = f(t, x)$ in Q . Then, we prove existence of local solutions of the semilinear fractional wave equation for some suitable values of $b > 1$. Moreover, we obtain an explicit dependence of the time of existence of solutions with respect to the initial data that allows longer time of existence for small initial data.

1. INTRODUCTION

1.1. Statement of the problem. Let Ω be a C^2 -bounded domain of \mathbb{R}^d with $d = 2, 3$. In what follows, we define \mathcal{A} by the differential operator

$$\mathcal{A}u(x) = - \sum_{i,j=1}^d \partial_{x_i} (a_{ij}(x) \partial_{x_j} u) + V(x)u(x), \quad x \in \Omega,$$

where $a_{ij} = a_{ji} \in C^1(\overline{\Omega})$ and $V \in L^\kappa(\Omega)$, for some $\kappa > d$, satisfy

$$\sum_{i,j=1}^d a_{ij}(x) \xi_i \xi_j \geq c|\xi|^2, \quad x \in \overline{\Omega}, \quad \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$$

and $V \geq 0$ a.e. in Ω .

We set $T \in (0, +\infty]$, $\Sigma = (0, T) \times \partial\Omega$ and $Q = (0, T) \times \Omega$. We consider the following initial-boundary value problem (IBVP in short) for the fractional semilinear wave equation

$$\begin{cases} \partial_t^\alpha u + \mathcal{A}u = f_b(u), & (t, x) \in Q, \\ u(t, x) = 0, & (t, x) \in \Sigma, \\ u(0, x) = u_0(x), \quad \partial_t u(0, x) = u_1(x), & x \in \Omega, \end{cases} \quad (1.1)$$

where $1 < \alpha < 2$, ∂_t^α denotes the Caputo fractional derivative with respect to t ,

$$\partial_t^\alpha u(t, x) := \frac{1}{\Gamma(2-\alpha)} \int_0^t (t-s)^{1-\alpha} \partial_s^2 u(s, x) ds, \quad (t, x) \in Q,$$

$b > 1$ and $f_b \in C^1(\mathbb{R})$ satisfies $f_b(0) = 0$ and

$$|f'_b(u)| \leq C |u|^{b-1}, \quad u \in \mathbb{R}.$$

The main purpose of this paper is to give a suitable definition of solutions of (1.1) and to study the well-posedness of this problem.

1.2. Physical motivations and known results. Recall that equation (1.1) is associated to anomalous diffusion phenomenon. More precisely, for $1 < \alpha < 2$, the linear part of equation (1.1) is frequently used for super-diffusive model of anomalous diffusion such as diffusion in heterogeneous media. In particular, in the linear case (i.e., $f_b \equiv 0$), some physical background is found in Sokolov, Klafter and Blumen [24]. As for analytical results in the case of $1 < \alpha < 2$, we refer to Mainardi [16] as one early work, and also to §6.1 in Kilbas, Srivastava and Trujillo [10], §10.10 in Podlubny [20]. For $0 < \alpha < 1$, we define $\partial_t^\alpha u$ by $\partial_t^\alpha u(t, x) := \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-s)^{-\alpha} \partial_s u(s, x) ds$, and there are works in view of the theory of partial differential equations (e.g., Beckers and Yamamoto [1], Luchko [15], Sakamoto and Yamamoto [21]). Such researches are rapidly developing and here we do not intend to give any comprehensive lists of references.

In contrast to the wave equation, even linear fractional wave equations are not well studied. In fact, few authors treated the well-posedness of the linear IBVP associated to (1.1) and to our best knowledge even the definition of weak solutions does not allow source term with low regularity. For a general study of the linear fractional wave equation and the regularity of solutions we refer to [21]. When we consider e.g., reaction effects in a super-diffusive model, we have to introduce a semilinear term.

To the best knowledge of the authors, there are no publications on fractional semilinear wave equations by the Strichartz estimate which is a common technique for semilinear wave and Schrödinger equations. In fact, for the wave equation ($\alpha = 2$), the well-posedness of problem (1.1) has been studied by various authors. In the case $\Omega = \mathbb{R}^k$ with $k \geq 3$ and $\mathcal{A} = -\Delta$, the global well-posedness has been proved both in the subcritical case $1 < b < 1 + \frac{4}{k-2}$ by Ginibre and Velo [3], and in the critical case $b = 1 + \frac{4}{k-2}$ by Grillakis [5] and, Shatah and Struwe [22, 23]. For $\Omega = \mathbb{R}^2$, Nakamura and Ozawa [18, 19] proved global well-posedness with exponentially growing nonlinearity. Without being exhaustive, for other results related to regularity of solutions or existence of solutions for more general semilinear hyperbolic equations we refer to [5, 4, 8, 9, 13]. In the case of Ω a smooth bounded domain of \mathbb{R}^3 , [2] proved the global well-posedness in the critical case $b = 5$. In addition, following the strategy set by [2], [7] treated the case of Ω a smooth bounded domain of \mathbb{R}^2 with exponentially growing nonlinearity.

1.3. Main results. In order to give a suitable definition of solutions of (1.1) we first need to consider the IBVP associated to the linear fractional wave equation

$$\begin{cases} \partial_t^\alpha u + \mathcal{A}u = f(t, x), & (t, x) \in Q, \\ u(t, x) = 0, & (t, x) \in \Sigma, \\ u(0, x) = u_0(x), \partial_t u(0, x) = u_1(x), & x \in \Omega. \end{cases} \quad (1.2)$$

The present paper contains three main results. Our two first main results are related to properties of solutions of (1.2), while our last result concerns the nonlinear problem (1.1).

Let us first remark that in contrast to usual derivatives, there is no exact integration by parts formula for fractional derivatives. Therefore, it is difficult to introduce the definition of weak solutions of (1.2) in the sense of distributions. To overcome this gap we give the following definition of weak solutions of (1.2). Let $\mathbf{1}_{(0,T)}(t)$ be the characteristic function of $(0, T)$.

Definition 1.1. *Let $u_0 \in L^2(\Omega)$, $u_1 \in H^{-1}(\Omega)$ and $f \in L^1(0, T; L^2(\Omega))$. We say that problem (1.2) admits a weak solution if there exists $v \in L_{loc}^\infty(\mathbb{R}^+; L^2(\Omega))$ such that:*

- 1) $v|_Q = u$ and $\inf\{\varepsilon > 0 : e^{-\varepsilon t} v \in L^1(\mathbb{R}^+; L^2(\Omega))\} = 0$,
- 2) for all $p > 0$ the Laplace transform $V(p) = \int_0^{+\infty} e^{-pt} v(t, \cdot) dt$ with respect to t of v solves

$$\begin{cases} (\mathcal{A} + p^\alpha)V(p) = F(p) + p^{\alpha-1}u_0 + p^{\alpha-2}u_1, & \text{in } \Omega, \\ V(p) = 0, & \text{on } \partial\Omega, \end{cases}$$

where $F(p) = \mathcal{L}[f(t, \cdot)\mathbf{1}_{(0,T)}(t)](p) = \int_0^T e^{-pt} f(t, \cdot) dt$.

Remark 1. *Recall (e.g. formula (2.140) page 80 of [20]) that for $h \in \mathcal{C}^2(\mathbb{R}^+)$ satisfying $\inf\{\varepsilon > 0 : e^{-\varepsilon t} h^{(k)} \in L^1(\mathbb{R}^+), k = 0, 1, 2\} = \varepsilon_0$ we have*

$$\mathcal{L}[\partial^\alpha h](p) = p^\alpha H(p) - p^{\alpha-1}h(0) - p^{\alpha-2}h'(0), \quad p > \varepsilon_0,$$

where $H(p) = \mathcal{L}[h](p) = \int_0^{+\infty} e^{-pt} h(t) dt$. Therefore, for sufficiently smooth data u_0, u_1, f (e.g. [21]) one can check that problem (1.2) admits a unique strong solution which is also a weak solution of (1.2).

Consider the operator A acting on $L^2(\Omega)$ with domain $D(A) = \{g \in H_0^1(\Omega) : \mathcal{A}g \in L^2(\Omega)\}$ defined by $Au = \mathcal{A}u$, $u \in D(A)$. Recall that in view of the Sobolev embedding theorem (e.g. [6, Theorem 1.4.4.1]) the multiplication operator $u \mapsto Vu$ is bounded from $H^1(\Omega)$ to $L^2(\Omega)$. Thus, we have $D(A) = H^2(\Omega) \cap H_0^1(\Omega)$. Moreover, by $V \geq 0$ in Ω , the operator A is a strictly positive selfadjoint operator with a compact resolvent. Therefore, the spectrum of A consists of a non-decreasing sequence of strictly positive eigenvalues $(\lambda_n)_{n \geq 1}$. Let us also introduce an orthonormal basis in the Hilbert space $L^2(\Omega)$ of eigenfunctions $(\varphi_n)_{n \geq 1}$ of A associated to the non-decreasing sequence of eigenvalues $(\lambda_n)_{n \geq 1}$.

From now on, by $\langle \cdot, \cdot \rangle$, we denote the scalar product of $L^2(\Omega)$. For all $s \geq 0$, we denote by A^s the operator defined by

$$A^s h = \sum_{n=1}^{+\infty} \langle h, \varphi_n \rangle \lambda_n^s \varphi_n, \quad h \in D(A^s) = \left\{ h \in L^2(\Omega) : \sum_{n=1}^{+\infty} |\langle h, \varphi_n \rangle|^2 \lambda_n^{2s} < \infty \right\}$$

and consider on $D(A^s)$ the norm

$$\|h\|_{D(A^s)} = \left(\sum_{n=1}^{+\infty} |\langle h, \varphi_n \rangle|^2 \lambda_n^{2s} \right)^{\frac{1}{2}}, \quad h \in D(A^s).$$

By duality, we can also set $D(A^{-s}) = D(A^s)'$ by identifying $L^2(\Omega)' = L^2(\Omega)$ which is a Hilbert space with the norm

$$\|h\|_{D(A^{-s})} = \left(\sum_{n=1}^{\infty} \langle h, \varphi_n \rangle_{-2s} \lambda_n^{-2s} \right)^{\frac{1}{2}}.$$

Here $\langle \cdot, \cdot \rangle_{-2s}$ denotes the duality bracket between $D(A^{-s})$ and $D(A^s)$. Since $D(A^{1/2}) = H_0^1(\Omega)$, we identify $H^{-1}(\Omega)$ with $D(A^{-1/2})$.

Using eigenfunction expansions we show our first main result where we state existence and uniqueness of weak solutions of (1.2).

Theorem 1.2. *Let $u_0 \in L^2(\Omega)$, $u_1 \in H^{-1}(\Omega) = D(A^{-\frac{1}{2}})$, $f \in L^1(0, T; L^2(\Omega))$. Then, problem (1.2) admits a unique weak solution $u \in C([0, T]; L^2(\Omega))$ satisfying*

$$\|u\|_{C([0, T]; L^2(\Omega))} \leq C(\|u_0\|_{L^2(\Omega)} + \|u_1\|_{H^{-1}(\Omega)} + \|f\|_{L^1(0, T; L^2(\Omega))}). \quad (1.3)$$

Moreover, assuming that there exists $0 < r < \frac{1}{4}$ such that $u_0 \in H^{2r}(\Omega)$, we have $u \in W^{1,1}(0, T; L^2(\Omega))$ and

$$\|u\|_{W^{1,1}(0, T; L^2(\Omega))} \leq C(\|u_0\|_{H^{2r}(\Omega)} + \|u_1\|_{H^{-1}(\Omega)} + \|f\|_{L^1(0, T; L^2(\Omega))}). \quad (1.4)$$

Recall that for $\gamma, r, s \geq 0$, $1 \leq p, q, \tilde{p}, \tilde{q} \leq \infty$, Strichartz estimates for solutions u of (1.2) denotes estimates of the form

$$\|u\|_{C([0, T]; H^{2r}(\Omega))} + \|u\|_{L^p(0, T; L^q(\Omega))} \leq C(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)} + \|f\|_{L^{\tilde{p}}(0, T; L^{\tilde{q}}(\Omega))}).$$

It is well known that these estimates, introduced by [25] and extended to the endpoints by [11] for both wave and Schrödinger equations, are important tools in the study of well-posedness of nonlinear equations (e.g. [2, 5, 4, 7, 9]). In the present paper we prove these estimates for solutions of (1.2). For this purpose, we consider $1 \leq p, q \leq \infty$ and $0 < \gamma < 1$ satisfying:

- 1) $q = \infty$, for $\frac{d}{4} < \gamma < 1$,
 - 2) $2 < q < \infty$, for $\gamma = \frac{d}{4}$,
 - 3) $q = \frac{2d}{d-4\gamma}$, for $0 < \gamma < \frac{d}{4}$.
- (1.5)

$$\begin{aligned} 1) \quad & p < \frac{1}{1-\alpha(1-\gamma)}, \quad \text{for } \gamma > 1 - \frac{1}{\alpha}, \\ 2) \quad & p = \infty, \quad \text{for } \gamma \leq 1 - \frac{1}{\alpha}. \end{aligned} \tag{1.6}$$

Then, our second main result can be stated as follows.

Theorem 1.3. (*Strichartz estimates*) Assume that $1 \leq p, q \leq \infty$ and $0 < \gamma < 1$ fulfill (1.5), (1.6) and set

$$s = \max\left(0, \gamma - \frac{1}{\alpha}\right), \quad r = \min\left(1 - \frac{1}{\alpha}, \gamma\right).$$

Let $u_0 \in D(A^\gamma)$, $u_1 \in D(A^s)$, $f \in L^1(0, T; L^2(\Omega))$. Then, the unique weak solution u of problem (1.2) is lying in $L^p(0, T; L^q(\Omega)) \cap C([0, T]; H^{2r}(\Omega))$ and fulfills estimate

$$\|u\|_{C([0, T]; H^{2r}(\Omega))} + \|u\|_{L^p(0, T; L^q(\Omega))} \leq C(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)} + \|f\|_{L^1(0, T; L^2(\Omega))}). \tag{1.7}$$

Here the constant C takes the form

$$C = C_0(1 + T)^\delta, \tag{1.8}$$

where

$$\delta = \begin{cases} \max(\alpha(1-\gamma) - 1, 1 - \alpha(\gamma - s), 1 - \alpha(r - s), \alpha(1-r) - 1), & \text{for } p = \infty, \\ \max\left(\frac{1}{p}, 1 - \alpha(\gamma - s) + \frac{1}{p}, 1 - \alpha(r - s), \alpha(1-r) - 1, \alpha(1-\gamma) - 1 + \frac{1}{p}\right), & \text{for } p < \infty \end{cases} \tag{1.9}$$

and C_0 depends only on $\Omega, \gamma, d, \alpha, p$.

In the last section we apply estimates (1.7) to prove our last result which is related to the existence and uniqueness of local solutions of (1.1). For this purpose, we first need to define local solutions of (1.1). In section 2 (see also [21]), using the eigenfunction expansions we introduce the operators

$$\begin{aligned} S_1(t)h &= \sum_{k=1}^{\infty} E_{\alpha,1}(-\lambda_k t^\alpha) \langle h, \varphi_k \rangle \varphi_k, \quad h \in L^2(\Omega), \\ S_2(t)h &= \sum_{k=1}^{\infty} t E_{\alpha,2}(-\lambda_k t^\alpha) \langle h, \varphi_k \rangle \varphi_k, \quad h \in L^2(\Omega), \\ S_3(t)h &= \sum_{k=1}^{\infty} t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_k t^\alpha) \langle h, \varphi_k \rangle \varphi_k, \quad h \in L^2(\Omega), \end{aligned}$$

where for all $\alpha > 0$, $\beta \in \mathbb{R}$, $E_{\alpha,\beta}$ denotes the Mittag-Leffler function given by

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}.$$

It is well known (e.g. [1, 20, 15, 21]) that for all $t > 0$ we have $S_j(t) \in B(L^2(\Omega))$, $j = 1, 2, 3$. Moreover, in view of Theorem 1.2, for $u_0, u_1 \in L^2(\Omega)$ and $f \in L^1(0, T; L^2(\Omega))$, the unique weak solution of (1.2) is given by

$$u(t) = S_1(t)u_0 + S_2(t)u_1 + \int_0^t S_3(t-s)f(s)ds. \tag{1.10}$$

For all $T > 0$, we introduce the space

$$X_T = C([0, T]; L^2(\Omega)) \cap L^b(0, T; L^{2b}(\Omega))$$

with the norm

$$\|v\|_{X_T} = \|v\|_{\mathcal{C}([0,T];L^2(\Omega))} + \|v\|_{L^b(0,T;L^{2b}(\Omega))}.$$

Recall that, by applying the Hölder inequality, one can check that for all $u, v \in X_T$ we have $f_b(u), f_b(v) \in L^1(0, T; L^2(\Omega))$ with

$$\|f_b(u)\|_{L^1(0,T;L^2(\Omega))} \leq C_b \|u\|_{L^b(0,T;L^{2b}(\Omega))}^b \leq C_b \|u\|_{X_T}^b \quad (1.11)$$

and

$$\|f_b(u) - f_b(v)\|_{L^1(0,T;L^2(\Omega))} \leq C_b \|u - v\|_{X_T} (\|u\|_{X_T}^{b-1} + \|v\|_{X_T}^{b-1}), \quad (1.12)$$

where the constant $C_b > 0$ depends only on b, f_b . Therefore, in view of Theorem 1.2, the map \mathcal{H}_b defined by

$$\mathcal{H}_b u(t) = \int_0^t S_3(t-s) f_b(u(s)) ds, \quad u \in X_T$$

is locally Lipschitz from X_T to $\mathcal{C}([0, T]; L^2(\Omega))$.

Definition 1.4. Let $u_0, u_1 \in L^2(\Omega)$ and $T > 0$. We say that (1.1) admits a weak solution on $(0, T)$ if the map $\mathcal{G}_b : X_T \rightarrow \mathcal{C}([0, T]; L^2(\Omega))$ defined by

$$\mathcal{G}_b u(t) = S_1(t)u_0 + S_2(t)v_2 + \int_0^t S_3(t-s) f_b(u(s)) ds$$

admits a fixed point $u \in X_T$. Such a fixed point $u \in X_T$ is called a weak solution to (1.1) on $(0, T)$. We say that problem (1.1) admits a local weak solution if there exists $T > 0$, depending on u_0, u_1 , such that problem (1.1) admits a weak solution on $(0, T)$.

Now we can state our result of existence and uniqueness of local solutions for (1.1). We recall that $\delta > 0$ is given in (1.9).

Theorem 1.5. Let $b > 1$ satisfy

$$\frac{d\alpha}{d\alpha + 4(1-\alpha)} < b < \frac{d\alpha + 4}{d\alpha + 4(1-\alpha)} \quad (1.13)$$

and let

$$\gamma = \frac{d(b-1)}{4b}, \quad q = 2b, \quad s = \max(0, \gamma - \frac{1}{\alpha}), \quad r = \min(1 - \frac{1}{\alpha}, \gamma), \quad 1 \leq \ell < \frac{1}{2-\alpha}. \quad (1.14)$$

Then, we can choose $p \in \left(b, \frac{1}{1-\alpha(1-\gamma)}\right)$ such that for $u_0 \in D(A^\gamma)$, $u_1 \in D(A^s)$, $T_0 > 0$, problem (1.1) admits a local weak solution u lying in $L^p(0, T; L^q(\Omega)) \cap \mathcal{C}([0, T]; H^{2r}(\Omega)) \cap W^{1,\ell}(0, T; L^2(\Omega))$ for some $T \leq T_0$ that takes the form

$$T = \min \left[\left(\tilde{C} (\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}) \right)^{-\frac{p(b-1)}{p-b}}, T_0 \right], \quad (1.15)$$

where we set

$$\tilde{C} = \tilde{C}_0 (1 + T_0)^{\frac{\delta}{b-1}}, \quad (1.16)$$

and \tilde{C}_0 depends only on f_b , Ω , α , b , p and d . Moreover, this local weak solution u is a unique local weak solution of (1.1) lying in $L^p(0, T; L^q(\Omega))$ and satisfies

$$\|u\|_{\mathcal{C}([0, T]; H^{2r}(\Omega))} + \|u\|_{L^p(0, T; L^q(\Omega))} + \|u\|_{W^{1, \ell}(0, T; L^2(\Omega))} \leq C(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}). \quad (1.17)$$

Here the constant $C > 0$ depends on d , Ω , f_b , b , T_0 , p , α .

A direct consequence of Theorem 1.5 is the following.

Corollary 1.6. *Assume that conditions (1.13) and (1.14) are fulfilled. Let $u_0 \in D(A^\gamma)$, $u_1 \in D(A^s)$ satisfy*

$$\left[\tilde{C}_0(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}) \right]^{-\frac{p(b-1)}{p(1+\delta)-b}} > 1$$

for some $b < p < \frac{1}{1-\alpha(1-\gamma)}$, where the constant \tilde{C}_0 is introduced in (1.16). Then, for any $T > 0$ satisfying

$$T < \left[\tilde{C}_0(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}) \right]^{-\frac{p(b-1)}{p(1+\delta)-b}}, \quad (1.18)$$

problem (1.1) admits a unique weak solution u on $(0, T)$ lying in $L^p(0, T; L^q(\Omega)) \cap \mathcal{C}([0, T]; H^{2r}(\Omega)) \cap W^{1, \ell}(0, T; L^2(\Omega))$.

This last result means that for smaller initial data we obtain longer time of existence of weak solutions.

Let us remark that, this paper seems to be the first where the Definition 1.1 of weak solutions of (1.2) is considered. The main contribution of Definition 1.1 comes from the fact that it allows well-posedness of (1.2) with weak conditions. Indeed, in contrast to other definitions of weak solutions for (1.2) (e.g. [21, Definition 2.1] used by [21] to prove existence of weak solutions of (1.2) with $f \in L^2(Q)$, $u_0 \in L^2(\Omega)$, $u_1 = 0$ in [21, Corollary 2.5, 2.6]), applying Definition 1.1 we can show well-posedness of (1.2) with $f \in L^1(0, T; L^2(\Omega))$, $u_0 \in L^2(\Omega)$ and $u_1 \in H^{-1}(\Omega)$. The choice of Definition 1.1 is inspired both by the analysis of [20] and the connection between elliptic equations and fractional diffusion equations used by [12]. Note also that Definition 1.1 plays an important role in the Definition 1.4 of weak solutions of (1.1).

Let us observe that in contrast to the wave equation the solution of (1.2) are not described by a semigroup. Therefore, we can not apply many arguments that allow to improve the Strichartz estimates (1.7) such as the TT^* method of [11]. Nevertheless, we prove local existence of solution of (1.1) with estimates (1.7). Note also that estimates (1.7) are derived from suitable estimates of Mittag-Leffler functions.

To our best knowledge this paper is the first treating well-posedness for semilinear fractional wave equations. Contrary to semilinear wave equations, it seems difficult to give a suitable definition of the energy for (1.1). This is mainly due to that fact that, once again, there is no exact integration

by parts formula for fractional derivatives as well as properties of composition and conjugation of the fractional Caputo derivative ∂_t^α (e.g. [20, Section 2]). For this reason, it seems complicate to derive global well-posedness from local well-posedness. However, using the explicit dependence with respect to T of the constant in (1.7) we can establish an explicit dependence of the time of existence T of (1.1) with respect to the initial conditions u_0, u_1 . From this result, we prove long time of existence for small initial data (see Corollary 1.6).

1.4. Outline. The paper is composed of four sections. In Section 2, we treat the well-posedness of the linear problem (1.2) and we show Theorem 1.2. Then, in Section 3 we prove the Strichartz estimates associated to these solutions and given by Theorem 1.3. Finally, in Section 4 we prove the local existence of solutions stated in Theorem 1.5 and Corollary 1.6.

2. THE LINEAR EQUATION

The goal of this section is to prove Theorem 1.2. For this purpose, for $k \geq 1$ we introduce $u_k \in \mathcal{C}(\mathbb{R}^+)$ defined by

$$u_k(t) = E_{\alpha,1}(-t^\alpha \lambda_k) u_{0,k} + t E_{\alpha,2}(-t^\alpha \lambda_k) u_{1,k} + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(-(t-s)^\alpha \lambda_k) f_k(s) ds, \quad t > 0, \quad (2.1)$$

where $u_{0,k} = \langle u_0, \varphi_k \rangle$, $u_{1,k} = \langle u_1, \varphi_k \rangle_{-1}$, $f_k(s) = \langle f(s), \varphi_k \rangle \mathbf{1}_{(0,T)}(s)$. We will show that $\sum_{k \geq 1} u_k(t) \varphi_k(x)$ converge to a weak solution of (1.2) and this weak solution is unique. Let us first recall the following estimates of the behavior of the Mittag-Leffler function.

Lemma 2.1. (Theorem 1.6, [20]) *If $0 < \alpha < 2$, $\beta \in \mathbb{R}$, $\pi\alpha/2 < \mu < \min(\pi, \pi\alpha)$, then*

$$|E_{\alpha,\beta}(z)| \leq \frac{C}{1+|z|}, \quad z \in \mathbb{C}, \quad \mu \leq |\arg z| \leq \pi,$$

where the constant $C > 0$ depends only on α, β, μ .

Applying Lemma 2.1, one can check that, for all $t > 0$ and all $m, n \in \mathbb{N}^*$, we have

$$\begin{aligned} \left\| \sum_{k=m}^n u_k(t) \varphi_k \right\|_{L^2(\Omega)} &\leq C \left\| \sum_{k=m}^n u_{0,k} \varphi_k \right\|_{L^2(\Omega)} + C t^{1-\frac{\alpha}{2}} \left\| \sum_{k=m}^n \frac{(\lambda_k t^\alpha)^{\frac{1}{2}}}{1 + \lambda_k t^\alpha} \lambda_k^{-\frac{1}{2}} u_{1,k} \varphi_k \right\|_{L^2(\Omega)} \\ &\quad + C t^{\alpha-1} \int_0^T \left\| \sum_{k=m}^n f_k(s) \varphi_k \right\|_{L^2(\Omega)} ds. \end{aligned}$$

Thus, for all $T_1 > 0$ we obtain

$$\begin{aligned} \sup_{t \in (0, T_1)} \left\| \sum_{k=m}^n u_k(t) \varphi_k \right\|_{L^2(\Omega)} &\leq C \left\| \sum_{k=m}^n u_{0,k} \varphi_k \right\|_{L^2(\Omega)} + C (T_1)^{1-\frac{\alpha}{2}} \left\| \sum_{k=m}^n \lambda_k^{-\frac{1}{2}} u_{1,k} \varphi_k \right\|_{L^2(\Omega)} \\ &\quad + C (T_1)^{\alpha-1} \int_0^T \left\| \sum_{k=m}^n f_k(s) \varphi_k \right\|_{L^2(\Omega)} ds \end{aligned}$$

and it follows that

$$\lim_{m,n \rightarrow \infty} \sup_{t \in (0, T_1)} \left\| \sum_{k=m}^n u_k(t) \varphi_k \right\|_{L^2(\Omega)} = 0.$$

Therefore, for any $T_1 > 0$ the serie $\sum_{k \geq 1} u_k(t) \varphi_k$ converge uniformly in $t \in (0, T_1)$ to $v \in \mathcal{C}(\mathbb{R}^+, L^2(\Omega))$.

In addition, for all $N \in \mathbb{N}^*$ and $t > 0$, we have

$$\left\| \sum_{k=1}^N u_k(t) \varphi_k \right\|_{L^2(\Omega)} \leq C \left(\|u_0\|_{L^2(\Omega)} + t^{1-\frac{\alpha}{2}} \|u_1\|_{D(A^{-\frac{1}{2}})} + Ct^{\alpha-1} \|f\|_{L^1(0, T; L^2(\Omega))} \right). \quad (2.2)$$

Here and henceforth \mathbb{N}^* denotes the set of all the natural number > 0 . Therefore, we deduce

$$\inf\{\varepsilon > 0 : e^{-\varepsilon t} v \in L^1(\mathbb{R}^+; L^2(\Omega))\} = 0$$

and (2.2) implies that, for all $N \in \mathbb{N}^*$, $t > 0$ and $p > 0$, we obtain

$$\left\| \sum_{k=1}^N e^{-pt} u_k(t) \varphi_k \right\|_{L^2(\Omega)} \leq C \left(e^{-pt} \|u_0\|_{L^2(\Omega)} + e^{-pt} t^{1-\frac{\alpha}{2}} \|u_1\|_{D(A^{-\frac{1}{2}})} + C e^{-pt} t^{\alpha-1} \|f\|_{L^1(0, T; L^2(\Omega))} \right).$$

Then, an application of Lebesgue's dominated convergence for functions taking values in $L^2(\Omega)$ yields

$$V(p, \cdot) = \mathcal{L}[v(t, \cdot)](p) = \sum_{k=1}^{\infty} \mathcal{L}[u_k](p) \varphi_k = \sum_{k=1}^{\infty} U_k(p, \cdot)$$

with $U_k(p, \cdot) = \mathcal{L}[u_k](p) \varphi_k$. Moreover, the properties of the Lalpace transform of the Mittag-Leffler function (e.g. formula (1.80) pp 21 of [20]) imply

$$U_k(p) = \frac{p^{\alpha-1} u_{0,k} + p^{\alpha-2} u_{1,k} + F_k(p)}{p^\alpha + \lambda_k} \varphi_k = (A + p^\alpha)^{-1} \left[\langle p^{\alpha-1} u_0 + F(p), \varphi_k \rangle + \langle p^{\alpha-2} u_1, \varphi_k \rangle_{-1} \right] \varphi_k$$

with $F_k(p) = \mathcal{L}[f_k](p) = \langle F(p), \varphi_k \rangle$. Thus $U_k(p, \cdot)$ solves

$$\begin{cases} (A + p^\alpha) U_k(p) = \left(\langle p^{\alpha-1} u_0 + F(p), \varphi_k \rangle + \langle p^{\alpha-2} u_1, \varphi_k \rangle_{-1} \right) \varphi_k, & \text{in } \Omega, \\ U_k(p) = 0, & \text{on } \partial\Omega. \end{cases}$$

Combining this with the fact that $u_0, u_1, F(p, \cdot) \in H^{-1}(\Omega) = D(A^{-\frac{1}{2}})$, we deduce that $\sum_{k \geq 1} U_k(p, \cdot)$ converge in $H_0^1(\Omega)$ to $V(p, \cdot)$ and $\sum_{k \geq 1} (A + p^\alpha) U_k(p)$ converge in $H^{-1}(\Omega)$ to $F(p) + p^{\alpha-1} u_0 + p^{\alpha-2} u_1$.

Therefore, $V(p, \cdot)$ solves

$$\begin{cases} (A + p^\alpha) V(p) = F(p) + p^{\alpha-1} u_0 + p^{\alpha-2} u_1, & \text{in } \Omega, \\ V(p) = 0, & \text{on } \partial\Omega. \end{cases} \quad (2.3)$$

Thus, $u = v|_{\mathcal{Q}}$ is a weak solution of (1.2). This proves the existence of weak solutions lying in $\mathcal{C}([0, T]; L^2(\Omega))$ and by the same way we obtain estimate (1.3). It remains to show that this solution is unique and, when $u_0 \in H^{2r}(\Omega)$, that it is lying in $W^{1,1}(0, T; L^2(\Omega))$ and that it fulfills (1.4).

We first prove the uniqueness of solutions. Let v_1, v_2 be two weak solutions of (1.2). Then, for $j = 1, 2$, there exist $w_j \in L_{\text{loc}}^\infty(\mathbb{R}^+; L^2(\Omega))$ such that: $w_j|_{\mathcal{Q}} = v_j$, $\inf\{\varepsilon > 0 : e^{-\varepsilon t} w_j \in L^1(\mathbb{R}^+; L^2(\Omega))\} = 0$

and, for all $p > 0$, the Laplace transform $W_j(p)$ with respect to t of w_j solves (2.3). Let $p > 0$ and set $W(p) = W_1(p) - W_2(p) \in L^2(\Omega)$ and note that $W(p)$ solves

$$\begin{cases} (\mathcal{A} + p^\alpha)W(p) = 0, & \text{in } \Omega, \\ W(p) = 0, & \text{on } \partial\Omega. \end{cases}$$

The uniqueness of the solution of this elliptic problem implies that $W(p) = 0$. Therefore, for all $p > 0$ we have $W_1(p) = W_2(p)$ which implies that $w_1 = w_2$ and by the same way $v_1 = w_{1|Q} = w_{2|Q} = v_2$. This proves the uniqueness.

From now on we assume that $u_0 \in H^{2r}(\Omega)$, for $r \in (0, 1/4)$, and we will show that $u \in W^{1,1}(0, T; L^2(\Omega))$ and that it fulfills (1.4). For this purpose, we establish the following lemmata. Here we recall that $u_k, f_k, u_{0,k}, u_{1,k}$ appear in (2.1).

Lemma 2.2. *For $\lambda > 0$, $\alpha > 0$ and positive integer $m \in \mathbb{N}^*$, we have*

$$\frac{d^m}{dt^m} E_{\alpha,1}(-\lambda t^\alpha) = -\lambda t^{\alpha-m} E_{\alpha, \alpha-m+1}(-\lambda t^\alpha), \quad t > 0$$

and

$$\frac{d}{dt} (t E_{\alpha,2}(-\lambda t^\alpha)) = E_{\alpha,1}(-\lambda t^\alpha), \quad t > 0.$$

Proof. The power series defining $E_{\alpha,1}(-\lambda t^\alpha)$ and $t E_{\alpha,2}(-\lambda t^\alpha)$ for $t > 0$ admit the termwise differentiation any times, and the termwise differentiation yields the conclusions. \square

Lemma 2.3. *For all $k \geq 1$ and $1 \leq \ell < \frac{1}{2-\alpha}$, we have $u_k \in W^{1,\ell}(0, T)$ and*

$$\partial_t u_k(t) = -\lambda_k t^{\alpha-1} E_{\alpha,\alpha}(-t^\alpha \lambda_k) u_{0,k} + E_{\alpha,1}(-t^\alpha \lambda_k) u_{1,k} + \int_0^t (t-s)^{\alpha-2} E_{\alpha,\alpha-1}(-(t-s)^\alpha \lambda_k) f_k(s) ds, \quad (2.4)$$

for a.e. $t \in (0, T)$.

Proof. First we consider the case $f_k = 0$. Then, we have

$$u_k(t) = E_{\alpha,1}(-t^\alpha \lambda_k) u_{0,k} + t E_{\alpha,2}(-t^\alpha \lambda_k) u_{1,k}, \quad t > 0.$$

In view of Lemma 2.2, we see that $u_k \in C^1([0, T])$ and (2.4) is fulfilled.

Second we consider the the case $u_{0,k} = u_{1,k} = 0$. Introduce, for all $\varepsilon > 0$ the function

$$u_k^\varepsilon(t) = \int_0^{t-\varepsilon} (t-s)^{\alpha-1} E_{\alpha,\alpha}(-(t-s)^\alpha \lambda_k) f_k(s) ds, \quad 0 < t < T.$$

In view of Lemma 2.2, we have $u_k^\varepsilon \in W^{1,\ell}(0, T)$ and

$$\partial_t u_k^\varepsilon(t) = \varepsilon^{\alpha-1} E_{\alpha,\alpha}(-\lambda_k \varepsilon^\alpha) f_k(t-\varepsilon) + \int_0^{t-\varepsilon} (t-s)^{\alpha-2} E_{\alpha,\alpha-1}(-(t-s)^\alpha \lambda_k) f_k(s) ds, \quad a.a. t \in (0, T).$$

On the other hand, one can easily check that $(u_k^\varepsilon)_{\varepsilon>0}$ converge to u_k as $\varepsilon \rightarrow 0$ in $D'(0, T)$ and $(\partial_t u_k^\varepsilon)_{\varepsilon>0}$ converge to

$$t \mapsto \int_0^t (t-s)^{\alpha-2} E_{\alpha, \alpha-1}(-(t-s)^\alpha \lambda_k) f_k(s) ds$$

as $\varepsilon \rightarrow 0$ in $D'(0, T)$, where $D'(0, T)$ is the space of distributions in $(0, T)$. Therefore, in the sense of $D'(0, T)$ we have

$$\partial_t u_k(t) = \int_0^t (t-s)^{\alpha-2} E_{\alpha, \alpha-1}(-(t-s)^\alpha \lambda_k) f_k(s) ds, \quad 0 < t < T,$$

which implies (2.4). In addition, applying (2.1), we obtain

$$|\partial_t u_k(t)| \leq C \int_0^t (t-s)^{\alpha-2} |f_k(s)| ds.$$

Then, according to the Young inequality, we deduce that $\partial_t u_k \in L^1(0, T)$. Therefore, we have $u_k, \partial_t u_k \in L^1(0, T)$, which means that $u_k \in W^{1, \ell}(0, T)$. Combining these two cases, we complete the proof of Lemma 2.3. \square

Let us remark that, using the fact that $0 < 2r < \frac{1}{2}$ and $D(A^{\frac{1}{2}}) = H_0^1(\Omega)$, one can check by interpolation that $u_0 \in H^{2r}(\Omega) = H_0^{2r}(\Omega) = D(A^r)$ (e.g. [14, Chapter 1, Theorems 11.1 and 11.6]) and

$$\sum_{k=1}^{\infty} \lambda_k^{2r} |u_{0,k}|^2 \leq C \|u_0\|_{H^{2r}(\Omega)}^2. \quad (2.5)$$

In view of (2.4), applying our previous arguments, for all $m, n \in \mathbb{N}^*$, we obtain

$$\begin{aligned} \left\| \sum_{k=m}^n \partial_t u_k \varphi_k \right\|_{L^1(0, T; L^2(\Omega))} &\leq C \left\| \sum_{k=m}^n \frac{(\lambda_k t^\alpha)^{1-r}}{1 + (\lambda_k^{\frac{1}{\alpha}} t)^\alpha} t^{\alpha r - 1} \lambda_k^r u_{0,k} \varphi_k \right\|_{L^1(0, T; L^2(\Omega))} \\ &\quad + C \left\| \sum_{k=m}^n \frac{(\lambda_k t^\alpha)^{\frac{1}{2}}}{1 + (\lambda_k^{\frac{1}{\alpha}} t)^\alpha} t^{-\frac{\alpha}{2}} \lambda_k^{-\frac{1}{2}} u_{1,k} \varphi_k \right\|_{L^1(0, T; L^2(\Omega))} \\ &\quad + C \int_0^T \int_0^t (t-s)^{\alpha-2} \left\| \sum_{k=m}^n f_k(s) \varphi_k \right\|_{L^2(\Omega)} ds dt. \end{aligned}$$

The Young inequality implies

$$\begin{aligned} \left\| \sum_{k=m}^n \partial_t u_k \varphi_k \right\|_{L^1(0, T; L^2(\Omega))} &\leq C \frac{T^{\alpha r}}{\alpha^r} \left\| \sum_{k=m}^n \lambda_k^r u_{0,k} \varphi_k \right\|_{L^2(\Omega)} + C \frac{T^{1-\frac{\alpha}{2}}}{1-\frac{\alpha}{2}} \left\| \sum_{k=m}^n \lambda_k^{-\frac{1}{2}} u_{1,k} \varphi_k \right\|_{L^2(\Omega)} \\ &\quad + C \frac{T^{\alpha-1}}{\alpha-1} \left\| \sum_{k=m}^n f_k(s) \varphi_k \right\|_{L^1(0, T; L^2(\Omega))}. \end{aligned}$$

Thus, we have

$$\lim_{m, n \rightarrow +\infty} \left\| \sum_{k=m}^n \partial_t u_k \varphi_k \right\|_{L^1(0, T; L^2(\Omega))} = 0,$$

which means that $\sum_{k=1}^n \partial_t u_k(t) \varphi_k(x)$ is a Cauchy sequence and a convergent sequence in $L^1(0, T; L^2(\Omega))$. Since $\sum_{k=1}^n u_k(t) \varphi_k(x)$ converge to u in $\mathcal{C}([0, T]; L^2(\Omega))$, combining this with (2.2), we deduce that

$\sum_{k \geq 1} u_k(t) \varphi_k(x)$ converge to u in $W^{1,1}(0, T; L^2(\Omega))$. Finally, repeating our previous arguments and applying (2.5), for all $N \in \mathbb{N}^*$, we find

$$\left\| \sum_{k=1}^N u_k \varphi_k \right\|_{W^{1,1}(0, T; L^2(\Omega))} \leq C(\|u_0\|_{H^{2r}(\Omega)} + \|u_1\|_{H^{-1}(\Omega)} + \|f\|_{L^1(0, T; L^2(\Omega))}).$$

Then, combining this estimate with (2.2) and taking the limit $N \rightarrow \infty$, we deduce (1.3). Thus, the proof of Theorem 1.2 is completed.

3. STRICHARTZ ESTIMATES

The goal of this section is to show Theorem 1.3. We divide the proof of Theorem 1.3 into two steps. First we prove estimates (1.7) for the weak solution u of (1.2) with $f = 0$ and then for $u_0 = u_1 = 0$. Henceforth $C > 0$ denotes generic constants which are dependent only on $\Omega, d, \alpha, \gamma$.

First step: Let $f = 0$ and let $1 \leq p, q \leq \infty, 0 < \gamma < 1$ fulfill (1.5) and (1.6). Then, (1.10) implies that

$$u(t) = S_1(t)u_0 + S_2(t)u_1.$$

Applying estimate Lemma 2.1, we deduce that for $t \mapsto S_1(t)u_0 \in \mathcal{C}([0, T]; D(A^\gamma)) \subset \mathcal{C}([0, T]; H^{2\gamma}(\Omega))$ with

$$\|S_1(t)u_0\|_{H^{2\gamma}(\Omega)} \leq C \|S_1(t)u_0\|_{D(A^\gamma)} \leq C \|u_0\|_{D(A^\gamma)} \leq C \|u_0\|_{H^{2\gamma}(\Omega)}, \quad 0 < t < T. \quad (3.1)$$

We have $0 \leq \gamma - s < 1$ by the definition of γ, s . Therefore, in the same way, Lemma 2.1 yields that, for all $0 < t < T$, we have

$$\lambda_k^{2\gamma} |tE_{\alpha, 2}(-\lambda_k t^\alpha) \langle u_1, \varphi_k \rangle|^2 \leq C t^{2(1-(\gamma-s)\alpha)} \lambda_k^{2s} |\langle u_1, \varphi_k \rangle|^2 \left(\frac{(\lambda_k t^\alpha)^{\gamma-s}}{1 + \lambda_k t^\alpha} \right)^2.$$

Thus, for all $0 < t < T$, we deduce that $S_2(t)u_1 \in D(A^\gamma) \subset H^{2\gamma}(\Omega)$ with

$$\|S_2(t)u_1\|_{H^{2\gamma}(\Omega)} \leq C t^{1-(\gamma-s)\alpha} \|u_1\|_{H^{2s}(\Omega)}, \quad 0 < t < T. \quad (3.2)$$

By the Sobolev embedding theorem, for all $0 < t < T$, we have $u(t, \cdot) \in H^{2\gamma}(\Omega) \subset L^q(\Omega)$ and

$$\|u(t, \cdot)\|_{L^q(\Omega)} \leq C \|u(t, \cdot)\|_{H^{2\gamma}(\Omega)} \leq C \max\left(t^{1-(\gamma-s)\alpha}, 1\right) (\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}).$$

On the other hand, we have $1 - (\gamma - s)\alpha \geq 0$ and so $u \in L^\infty(0, T; L^q(\Omega))$ and

$$\|u\|_{L^p(0, T; L^q(\Omega))} \leq C(1 + T)^{1-(\gamma-s)\alpha + \frac{1}{p}} (\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}). \quad (3.3)$$

In the same way, we have

$$\begin{aligned} \|S_1(t)u_0\|_{H^{2r}(\Omega)} &\leq C \|S_1(t)u_0\|_{H^{2\gamma}(\Omega)} \leq C \|u_0\|_{H^{2\gamma}(\Omega)}, \quad 0 < t < T, \\ \|S_2(t)u_1\|_{H^{2r}(\Omega)} &\leq C(1 + T)^{1-(r-s)\alpha} \|u_1\|_{H^{2s}(\Omega)}, \quad 0 < t < T. \end{aligned} \quad (3.4)$$

Combining these two estimates in (3.4) with (3.3), we deduce (1.7) for $f = 0$.

Second step: Let $u_0 = u_1 = 0$. In view of Lemma 2.1, for all $0 < t < T$, we have

$$\lambda_k^{2\gamma} |t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_k t^\alpha) \langle f, \varphi_k \rangle|^2 \leq t^{2(\alpha(1-\gamma)-1)} |\langle f, \varphi_k \rangle|^2 \left(\frac{(\lambda_k t^\alpha)^\gamma}{1 + \lambda_k t^\alpha} \right)^2.$$

Thus, for all $0 < t < T$ and $h \in L^2(\Omega)$, we deduce that $S_3(t)h \in D(A^\gamma) \subset H^{2\gamma}(\Omega)$ with

$$\|S_3(t)h\|_{H^{2\gamma}(\Omega)} \leq C t^{\alpha(1-\gamma)-1} \|h\|_{L^2(\Omega)}, \quad 0 < t < T.$$

By the Sobolev embedding theorem, for all $0 < t < T$, we have $S_3(t)h \in H^{2\gamma}(\Omega) \subset L^q(\Omega)$ with

$$\|S_3(t)h\|_{L^q(\Omega)} \leq C \|S_3(t)h\|_{H^{2\gamma}(\Omega)} \leq C t^{\alpha(1-\gamma)-1} \|h\|_{L^2(\Omega)}.$$

Applying this estimate, we obtain

$$\|u(t, \cdot)\|_{L^q(\Omega)} \leq \int_0^t \|S_3(t-s)f(s)\|_{L^q(\Omega)} ds \leq C \int_0^t (t-s)^{\alpha(1-\gamma)-1} \|f(s)\|_{L^2(\Omega)} ds.$$

By $t \mapsto t^{\alpha(1-\gamma)-1} \in L^p(0, T)$, the Young inequality yields

$$\|u\|_{L^p(0, T; L^q(\Omega))} \leq C \frac{T^{\alpha(1-\gamma)-1 + \frac{1}{p}}}{(p(\alpha(1-\gamma) - 1) + 1)^{1/p}} \|f\|_{L^1(0, T; L^2(\Omega))}. \quad (3.5)$$

Repeating the above arguments, we deduce that

$$\|u(t, \cdot)\|_{H^{2r}(\Omega)} \leq C \int_0^t (t-s)^{\alpha(1-r)-1} \|f(s)\|_{L^2(\Omega)} ds.$$

Then, since $\alpha(1-r) - 1 \geq \alpha(1 - (1 - \alpha^{-1})) - 1 = 0$, we deduce from the Young inequality that

$$\|u(t, \cdot)\|_{H^{2r}(\Omega)} \leq C T^{\alpha(1-r)-1} \|f\|_{L^1(0, T; L^2(\Omega))}.$$

Combining this estimate with (3.3) - (3.5), we deduce (1.7) for $u_0 = u_1 = 0$. This completes the proof of Theorem 1.3.

4. LOCAL SOLUTIONS OF (1.1)

In this section we will apply the results of the previous section to prove Theorem 1.5 and Corollary 1.6.

Proof of Theorem 1.5 Note first that for γ and b given by (1.13) and (1.14), we have $\gamma < \frac{d}{4}$ and

$$\frac{d}{d-4\gamma} = b > \frac{d\alpha}{d\alpha + 4(1-\alpha)},$$

which implies by $1 < \alpha < 2$ and $d = 2, 3$ that

$$\gamma > 1 - \frac{1}{\alpha}. \quad (4.1)$$

On the other hand, for $1 - \frac{1}{\alpha} < \gamma < \frac{d}{4}$, one can check that

$$\gamma < \frac{d\alpha}{4+d\alpha} \iff \frac{d(b-1)}{4b} < \frac{d\alpha}{4+d\alpha} \iff b < \frac{d\alpha+4}{d\alpha+4(1-\alpha)}. \quad (4.2)$$

Therefore, γ given by (1.14) fulfills

$$1 - \frac{1}{\alpha} < \gamma < \frac{d\alpha}{4+d\alpha},$$

which yields

$$\frac{1}{1 - \alpha \left(1 - \frac{d\alpha}{4+d\alpha}\right)} < \frac{1}{1 - \alpha(1 - \gamma)}.$$

Therefore, we can choose p satisfying

$$b < \frac{d\alpha + 4}{d\alpha + 4(1 - \alpha)} = \frac{1}{1 - \alpha \left(1 - \frac{d\alpha}{4+d\alpha}\right)} < p < \frac{1}{1 - \alpha(1 - \gamma)}.$$

Moreover, for q given by (1.14) we have $q = \frac{2d}{d-4\gamma}$. Thus, for q, γ given by (1.14) and $b < p < \frac{1}{1 - \alpha(1 - \gamma)}$, p, q, γ fulfill conditions (1.5) and (1.6) with $p > b$. Provided that $0 < T \leq T_0$ and $M > 0$ will be chosen suitably later, we set $Y_T = L^p(0, T; L^q(\Omega)) \cap \mathcal{C}([0, T]; H^{2r}(\Omega))$ and $B_M = \{u \in Y_T : \|u\|_{Y_T} \leq M\}$. Moreover, we set

$$\|u\|_{Y_T} = \|u\|_{L^p(0, T; L^q(\Omega))} + \|u\|_{\mathcal{C}([0, T]; H^{2r}(\Omega))}.$$

We fix the constant $C'_b > 0$ which appears in estimates (1.11) and (1.12). We note that C'_b is independent of T . We put $C' = C_0(1 + T_0)^\delta$, where the constants C_0, δ are introduced in (1.8), (1.9) and are independent of T . Finally we fix $C = C'(1 + C'_b) + 1$. Since $p > b$, for all $u \in Y_T$ we have $u \in L^b(0, T; L^{2b}(\Omega))$. Therefore, in view of Theorem 1.3 and estimates (1.7), (1.8) and (1.11), we have $\mathcal{G}_b(u) \in Y_T$ and

$$\begin{aligned} \|\mathcal{G}_b(u)\|_{Y_T} &\leq C'(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)} + \|f_b(u)\|_{L^1(0, T; L^2(\Omega))}) \\ &\leq C'(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)} + C_b \|u\|_{L^b(0, T; L^{2b}(\Omega))}^b) \\ &\leq C(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)} + \|u\|_{L^b(0, T; L^{2b}(\Omega))}^b). \end{aligned} \quad (4.3)$$

On the other hand, by the Hölder inequality one can check that

$$\int_0^T \|u(t, \cdot)\|_{L^q(\Omega)}^b dt \leq \left(\int_0^T \|u(t, \cdot)\|_{L^q(\Omega)}^p dt \right)^{\frac{b}{p}} T^{1 - \frac{b}{p}},$$

which implies

$$\|u\|_{L^b(0, T; L^{2b}(\Omega))} \leq T^{\frac{p-b}{bp}} \|u\|_{L^p(0, T; L^q(\Omega))}. \quad (4.4)$$

Applying this estimate to (4.3), we obtain

$$\|\mathcal{G}_b(u)\|_{Y_T} \leq C(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)} + T^{\frac{p-b}{p}} \|u\|_{Y_T}^b). \quad (4.5)$$

We set $M = 2C(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)})$ and $T = \min\left((3CM^{b-1})^{-\frac{p}{p-b}}, T_0\right)$. With these values of M and T , one can easily verify that (4.5) implies

$$\|\mathcal{G}_b u\|_{Y_T} \leq M, \quad u \in B_M.$$

In the same way, applying estimates (1.12) and (1.7) in

$$\mathcal{G}_b u - \mathcal{G}_b v = \int_0^t S_3(t-s)[f_b(u(s)) - f_b(v(s))] ds,$$

we obtain

$$\|\mathcal{G}_b u - \mathcal{G}_b v\|_{Y_T} \leq C \|u - v\|_{L^b(0,T;L^{2b}(\Omega))} (\|u\|_{L^b(0,T;L^{2b}(\Omega))}^{b-1} + \|v\|_{L^b(0,T;L^{2b}(\Omega))}^{b-1}).$$

Then, (4.4) and the choice of T imply that for every $u, v \in B_M$, we have

$$\begin{aligned} \|\mathcal{G}_b u - \mathcal{G}_b v\|_{Y_T} &\leq CT^{\frac{p-b}{p}} \|u - v\|_{L^p(0,T;L^q(\Omega))} (\|u\|_{L^p(0,T;L^q(\Omega))}^{b-1} + \|v\|_{L^p(0,T;L^q(\Omega))}^{b-1}) \\ &\leq 2CM^{b-1} T^{\frac{p-b}{p}} \|u - v\|_{Y_T} \\ &\leq \frac{2}{3} \|u - v\|_{Y_T}. \end{aligned}$$

Therefore, \mathcal{G}_b is a contraction from B_M to B_M . Consequently \mathcal{G}_b admits a unique fixed point $u \in B_M$ which is a local weak solution of (1.1). Moreover, from our choice of M and T we deduce (1.15) and (1.16).

Now we show that this solution is unique in $L^p(0, T; L^q(\Omega))$. For this purpose, consider the space $Z_T = \mathcal{C}([0, T]; L^2(\Omega)) \cap L^p(0, T; L^q(\Omega))$ with the norm

$$\|v\|_{Z_T} = \|v\|_{\mathcal{C}([0,T];L^2(\Omega))} + \|v\|_{L^p(0,T;L^q(\Omega))}, \quad v \in Z_T.$$

Repeating our previous arguments we can show that \mathcal{G}_b is a contraction from B'_M to B'_M with $B'_M = \{u \in Z_T : \|u\|_{Z_T} \leq M\}$. Therefore, the fixed point $u \in B_M$ of \mathcal{G}_b is a unique local weak solution of (1.1) lying in $L^p(0, T; L^q(\Omega))$. Now let us show that the unique weak solution of (1.1) lying in $L^p(0, T; L^q(\Omega))$ is also lying in $W^{1,\ell}(0, T; L^2(\Omega))$ and it fulfills (1.17). Since $\|u\|_{Z_T} \leq M$ and $T = \min\left((3CM^{b-1})^{-\frac{p}{p-b}}, T_0\right)$, by (1.11), we obtain that $f_b(u) \in L^1(0, T; L^2(\Omega))$ satisfies

$$\|f_b(u)\|_{L^1(0,T;L^2(\Omega))} \leq C(\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}). \quad (4.6)$$

Now let us set

$$f_k(t) = \langle f_b(u(t)), \varphi_k \rangle, \quad u_{0,k} = \langle u_0, \varphi_k \rangle, \quad u_{1,k} = \langle u_1, \varphi_k \rangle.$$

Then, in view of Lemma 2.3, $u_k(t) = \langle u(t), \varphi_k \rangle \in W^{1,\ell}(0, T)$ fulfills (2.4). Repeating the arguments used in the last part of the proof of Theorem 1.2, we obtain

$$\begin{aligned} \left\| \sum_{k=m}^n (\partial_t u_k) \varphi_k \right\|_{L^\ell(0,T;L^2(\Omega))} &\leq C \left\| \sum_{k=m}^n \frac{(\lambda_k t^\alpha)^{1-\gamma}}{1 + (\lambda_k^\alpha t)^\alpha} t^{\alpha\gamma-1} \lambda_k^\gamma u_{0,k} \varphi_k \right\|_{L^\ell(0,T;L^2(\Omega))} + CT^{\frac{1}{2}} \left\| \sum_{k=m}^n u_{1,k} \varphi_k \right\|_{L^2(\Omega)} \\ &\quad + C \left\| \int_0^t (t-s)^{\alpha-2} \left\| \sum_{k=m}^n f_k(s) \varphi_k \right\|_{L^2(\Omega)} ds \right\|_{L^\ell(0,T)} \end{aligned}$$

for all $m, n \in \mathbb{N}^*$. In view of (4.1), we have

$$\ell(\alpha\gamma - 1) > \frac{\alpha - 2}{2 - \alpha} > -1.$$

Therefore, the Young inequality yields

$$\begin{aligned} \left\| \sum_{k=m}^n (\partial_t u_k) \varphi_k \right\|_{L^\ell(0,T;L^2(\Omega))} &\leq C \left(\frac{T^{\ell(\alpha\gamma-1)+1}}{\ell(\alpha\gamma-1)+1} \right)^{\frac{1}{\ell}} \left\| \sum_{k=m}^n u_{0,k} \varphi_k \right\|_{L^2(\Omega)} + CT^{\frac{1}{\ell}} \left\| \sum_{k=m}^n u_{1,k} \varphi_k \right\|_{L^2(\Omega)} \\ &+ C \left(\frac{T^{\ell(\alpha-2)+1}}{\ell(\alpha-2)+1} \right)^{\frac{1}{\ell}} \left\| \sum_{k=m}^n f_k(s) \varphi_k \right\|_{L^1(0,T;L^2(\Omega))}. \end{aligned}$$

Thus, we have

$$\lim_{m,n \rightarrow +\infty} \left\| \sum_{k=m}^n (\partial_t u_k) \varphi_k \right\|_{L^\ell(0,T;L^2(\Omega))} = 0,$$

which means that $\sum_{k \geq 1} (\partial_t u_k)(t) \varphi_k(x)$ is a Cauchy sequence and is a convergent sequence in $L^\ell(0, T; L^2(\Omega))$.

Combining this with the fact that $\sum_{k \geq 1} u_k(t) \varphi_k(x)$ converge to u in $\mathcal{C}([0, T]; L^2(\Omega))$, we deduce that

$\sum_{k \geq 1} u_k(t) \varphi_k(x)$ converge to u in $W^{1,\ell}(0, T; L^2(\Omega))$. Finally, for all $N \in \mathbb{N}^*$, we find

$$\left\| \sum_{k=1}^N u_k \varphi_k \right\|_{W^{1,\ell}(0,T;L^2(\Omega))} \leq C (\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)} + \|f_b(u)\|_{L^1(0,T;L^2(\Omega))}).$$

Combining this estimate with (4.6) and letting $N \rightarrow \infty$, we deduce (1.17). Thus, the proof of Theorem 1.5 is completed. \square

Proof of Corollary 1.6. Let $T > 0$ fulfill (1.18) and set $T_0 = T$. Without lost of generality we can assume that $T \geq 1$. Then, we have

$$\left(\tilde{C}_0 T_0^{\frac{\delta}{b-1}} (\|u_0\|_{H^{2\gamma}(\Omega)} + \|u_1\|_{H^{2s}(\Omega)}) \right)^{-\frac{p(b-1)}{p-b}} > T_0.$$

Since $T_0 \geq 1$ we can replace T_0 by $T_0 + 1$ in condition (1.15). Therefore, with this value of T_0 , condition (1.15) holds. Thus, according to Theorem 1.5, problem (1.1) admits a unique weak solution u on $(0, T)$ lying in $L^p(0, T; L^q(\Omega)) \cap \mathcal{C}([0, T]; H^{2r}(\Omega)) \cap W^{1,\ell}(0, T; L^2(\Omega))$. \square

Acknowledgements. The second author is partially supported by Grant-in-Aid for Scientific Research (S) 15H05740 of Japan Society for the Promotion of Science.

REFERENCES

- [1] S. BECKERS AND M. YAMAMOTO, *Regularity and uniqueness of solution to linear diffusion equation with multiple time-fractional derivatives*, International Series of Numerical Mathematics, **164** (2013), 45-55.
- [2] N. BURQ, G. LEBEAU AND F. PLANCHON, *Global existence for energy critical waves in 3-D domains*, J. Amer. Math. Soc, **21** (3) (2008), 831-845.
- [3] J. GINIBRE AND G. VELO, *The global Cauchy problem for nonlinear Klein-Gordon equation*, Math. Z, **189**, (1985) 487-505.
- [4] J. GINIBRE AND G. VELO, *Regularity of solutions of critical and subcritical nonlinear wave equations*, Nonlinear Anal., **22** (1994), 1-19.
- [5] M. G. GRILLAKIS, *Regularity and asymptotic behavior of the wave equation with a critical non-linearity*, Ann. of Math., **132** (1990), 485-509.

- [6] P. GRISVARD, *Elliptic problems in nonsmooth domains*, Pitman, London, 1985.
- [7] S. IBRAHIM AND R. JRAD, *Strichartz type estimates and the well-posedness of an energy critical 2D wave equation in a bounded domain*, J. Differ. Equ., **250** (9) (2011), 3740-3771.
- [8] L. KAPITANSKI, *Weak and yet weaker solutions of semi-linear wave equations*, Comm. Partial Differential Equation, **19** (1994), 1629-1676.
- [9] Y. KIAN, *Cauchy problem for semilinear wave equation with time-dependent metrics*, Nonlinear Anal., **73** (2010), 2204-2212.
- [10] A.A. KILBAS, H.M. SRIVASTAVA AND J.J. TRUJILLO, *Theory and applications of fractional differential equations*, Elsevier, Amsterdam, 2006.
- [11] M. KEEL AND T.TAO, *Endpoint Strichartz estimates*, Amer. J. Math., **120** (1998), 955-980.
- [12] Z. LI, O. Y. IMANUVILOV AND M. YAMAMOTO, *Uniqueness in inverse boundary value problems for fractional diffusion equations*, preprint, arXiv:1404.7024.
- [13] H. LINDELBAG AND C. D. SOGGE, *On existence and scattering with minimal regularity for the semilinear wave equation*, J. Funct. Anal., **130** (1995), 357-426.
- [14] J.-L. LIONS AND E. MAGENES, *Non-homogeneous Boundary Value Problems and Applications*, Vol. I, Springer-Verlag, Berlin, 1972.
- [15] Y. LUCHKO, *Some uniqueness and existence results for the initial-boundary-value problems for the generalized time-fractional diffusion equation*, Computers and Mathematics with Applications, **59** (2010), 1766-1772.
- [16] F. MAINARDI, *On the initial value problem for the fractional diffusion-wave equation*, in: S. Rionero, T. Ruggeri (Eds.), *Waves and Stability in Continuous Media*, World Scientific, Singapore, 1994, pp. 246-251.
- [17] C. MIAO AND B. ZHANG, *H^s -global well-posedness for semilinear wave equations*, J. Math. Anal. Appl, **283** (2003), 645-666.
- [18] M. NAKAMURA AND T. OZAWA, *Global solutions in the critical Sobolev space for the wave equations with nonlinearity of exponential growth*, Math. Z, **231** (1999), 479-487.
- [19] M. NAKAMURA AND T. OZAWA, *The Cauchy problem for nonlinear wave equations in the Sobolev space of critical order*, Discrete and Continuous Dynamical Systems, **5** N. 1 (1999), 215-231.
- [20] I. PODLUBNY, *Fractional differential equations*, Academic Press, San Diego, 1999.
- [21] K. SAKAMOTO AND M. YAMAMOTO, *Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems*, J. Math. Anal. Appl., **382** (2011), 426-447.
- [22] J. SHATAH AND M. STRUWE, *Regularity results for nonlinear wave equations*, Ann. of Math, **2** (138) (1993), 503-518.
- [23] J. SHATAH AND M. STRUWE, *Well-Posedness in the energy space for semilinear wave equation with critical growth*, IMRN, **7** (1994), 303-309.
- [24] I.M. SOKOLOV, J. KLAFTER AND A. BLUMEN, *Fractional kinetics*, Physics Today, **55** (2002), 48-54.
- [25] R. STRICHARTZ, *A priori estimates for the wave equation and some applications*, J. Funct. Anal., **5** (1970), 218-235.

AIX-MARSEILLE UNIVERSITÉ, CNRS, CPT UMR 7332, 13288 MARSEILLE, FRANCE & UNIVERSITÉ DE TOULON,
CNRS, CPT UMR 7332, 83957 LA GARDE, FRANCE

E-mail address: yavar.kian@univ-amu.fr

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-9 KOMABA MEGURO, TOKYO
153-8914, JAPAN

E-mail address: myama@ms.u-tokyo.ac.jp