

Probability that the maximum of the reflected Brownian motion over a finite interval [0; t] is achieved by its last zero before t

Agnès Lagnoux, Sabine Mercier, Pierre Vallois

To cite this version:

Agnès Lagnoux, Sabine Mercier, Pierre Vallois. Probability that the maximum of the reflected Brownian motion over a finite interval [0; t] is achieved by its last zero before t. Electronic Communications in Probability, 2015, $10.1214/ECP.vVOL-PID$. hal-01214773

HAL Id: hal-01214773 <https://hal.science/hal-01214773v1>

Submitted on 13 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Electron. Commun. Probab. **0** (2012), no. 0, 1–9. DOI: [10.1214/ECP.vVOL-PID](http://dx.doi.org/10.1214/ECP.vVOL-PID) ISSN: [1083-589X](http://ecp.ejpecp.org/)

ELECTRONIC COMMUNICATIONS in PROBABILITY

Probability that the maximum of the reflected Brownian motion over a finite interval $[0, t]$ is achieved by its last zero before t

Agnès Lagnoux * Sabine Mercier† Pierre Vallois‡

Abstract

We calculate the probability p_c that the maximum of a reflected Brownian motion U is achieved on a complete excursion, i.e. $p_c := P(\overline{U}(t) = U^*(t))$ where $\overline{U}(t)$ (respectively $U^*(t)$) is the maximum of the process U over the time interval [0, t] (resp. $[0, g(t)]$ where $g(t)$ is the last zero of U before t).

Keywords: Reflected Brownian motion ; Bessel process ; Brownian meander and Brownian bridge ; Brownian excursions ; Gamma function ; local score.. **AMS MSC 2010:** 60 G 17 ; 60 J 25 ; 60 J 65.. Submitted to ECP on May 2015, final version accepted on .

1 Introduction

1.1 Motivation. The local score of a biological sequence is its "best" segment with respect to some scoring scheme (see e.g. [12] for more details) and the knowledge of its distribution is important (see e.g. [8], [9]). Let us briefly recall the mathematical setting while biological interpretations can be found in [5]. Let $S_n := \epsilon_1 + \cdots + \epsilon_n$ be the random walk generated by the sequence of the independent and identically distributed random variables $(\epsilon_i, \, i \in \mathbb{N})$ that are centered with unit variance. The local score is the process: $U_n := S_n - \min_{0 \leq i \leq n} S_i$, where $n \geq 0$. The path of $(U_n, n \in \mathbb{N})$ is a succession of 0 and excursions above 0. In [5], the authors only took into consideration complete excursions up to a fixed time n and so considered the maximum U_n^\ast of the heights of all the complete excursions up to time n instead of the maximum \overline{U}_n of the path until time n. They also introduced the random time θ^* of the length of the segment that realizes $U_n^*.$ Since it is easy to simulate $(S_k, 0 \leq k \leq n)$, for any n not too large, we get an approximation of the law of (U_n^*,θ_n^*) for a given $n.$ Simulations have shown that for an important proportion of sequences, \overline{U}_n is realized during the last incomplete excursion. As expected, the number of excursions naturally increases when the length of the sequence growths, however the proportion of sequences that achieve their maximum on a complete excursion remains strikingly constant. The main goal of this study is to explain these observations and to calculate this probability when n is large, see Proposition 1.2 below.

^{*}Institut de Mathématiques de Toulouse, UMR 5219, Université Toulouse 2, France.

E-mail: lagnoux@univ-tlse2.fr

[†] Institut de Mathématiques de Toulouse, UMR 5219, Université Toulouse 2, France.

E-mail: mercier@univ-tlse2.fr

[‡] Institut de Mathématiques Elie Cartan, INRIA-BIGS, CNRS UMR 7502, Université de Lorraine, France. E-mail: pierre.vallois@univ-lorraine.fr

Maximum of the reflected Brownian motion before its last zero

1.2 Link with the Brownian motion. According to the functional convergence theorem of Donsker, the random walk $(S_k, 0 \le k \le n)$ (resp. $(U_k, 0 \le k \le n)$) normalized by the factor $1/\sqrt{n}$ converges in distribution, as $n\to\infty$, to the Brownian motion (resp. the reflected Brownian motion). We prove (see Theorem 1.1 for a precise formulation) that the probability that the maximum of a reflected Brownian motion over a finite interval $[0, t]$ is achieved on a complete excursion is around 30% and is thus independent of t. This result permits to answer to the two questions asked in the discrete setting, when n is large.

Let U be the reflected Brownian motion started at 0, i.e. $U(t) = |B(t)|$ where $(B(t), t > 0)$ is the standard one-dimensional Brownian motion started at 0. In Chabriac et al. [5], the authors have considered two maxima: $\overline{U}(t)$ and $U^*(t)$, the first (resp. second) one being the maximum of U up to time t (resp. the last zero before t), namely $U(t):=\max\limits_{0\leqslant s\leqslant t}U(s)$

and $U^*(t) = \overline{U}(g(t))$, where $g(t) := \sup\{s \leqslant t, U(s) = 0\}$. In [5], the density function of the pair $(U^*(t),\theta^*(t))$ has been calculated where $\theta^*(t)$ is the length of time segment that realizes $\dot{U}^*(t)$, i.e. the first hitting time of level $U^*(t)$ by the process $(U(s), 0 \leqslant s \leqslant g(t)).$ Here we only deal with $U^*(t)$ and $\overline{U}(t)$.

It is clear that $\overline{U}(t)=U^*(t)$ if and only if $U^{**}(t)\leqslant \overline{U}(t)$, where

$$
U^{**}(t) := \max_{g(t) \leq s \leq t} U(s).
$$
 (1.1)

In that case, the maximum of U over $[0, t]$ is the maximum of all the complete excursions of U which hold before t. We introduce the probability p_c that the maximum of U over $[0, t]$ is achieved on a complete excursion:

$$
p_c = \mathbb{P}\left(\overline{U}(t) = U^*(t)\right) = \mathbb{P}\left(U^*(t) > U^{**}(t)\right). \tag{1.2}
$$

Let ψ be the logarithmic derivative of the Gamma function:

$$
\psi(x) := \Gamma'(x) / \Gamma(x). \tag{1.3}
$$

The main result of our study is

Theorem 1.1. The probability p_c equals $\psi(1/4) - \psi(1/2) + 1 + \pi/2 \approx 0.3069$.

1.3 Back to discrete sequences. We now go back to the setting of random walks introduced in paragraph 1.1. Let $p_c^{(n)}$ be the probability that the maximum of $(U_k, 0\leqslant$ $k \leqslant n)$ is achieved on a complete excursion, namely

$$
p_c^{(n)} := P\Big(\max_{0 \le k \le n} U_k = \max_{0 \le k \le g_n} U_k\Big) \quad \text{where } g_n := \max\{k \le n, U_k = 0\}.
$$
 (1.4)

Proposition 1.2. $p_c^{(n)}$ converges to p_c as $n \to \infty.$

The convergence of $p^{(n)}_{c}$ can be obtained from Theorem 3.3 in [5] and the fact that the event N defined by (4.23) in [5] is actually included in $\Big\{\max_{0\leqslant k\leqslant n}U_k=\max_{0\leqslant k\leqslant g_n}U_k\Big\}.$ 1.4 Main steps of the proof. We now consider the Brownian motion setting. The density function of $\overline{U}(t)$ is known (see either Subsection 2.11 in [3] or Lemma 3.2 in [11]) and the one of $U^*(t)$ has been calculated in [5]. Obviously, the knowledge of the distributions of $\overline{U}(t)$ and $U^*(t)$ is not sufficient to determine $p_c.$ The trajectory of $(U(s), 0 \leq s \leq t)$ naturally splits in two parts before and after the random time $q(t)$ which is not a stopping time. Although $(U(s), 0 \leq s \leq g(t))$ and $(U(s), g(t) \leq s \leq t)$ are not independent, the scaling with $q(t)$ leads to independence. Indeed,

$$
\left(\frac{1}{\sqrt{g(t)}}U(sg(t)),\ 0\leqslant s\leqslant 1\right),\ \left(\frac{1}{\sqrt{t-g(t)}}\left|B(g(t)+s(t-g(t))\right|,\ 0\leqslant s\leqslant 1\right),\ g(t)\tag{1.5}
$$

ECP **0** [\(2012\), paper 0.](http://dx.doi.org/10.1214/ECP.vVOL-PID)

Maximum of the reflected Brownian motion before its last zero

are independent. Moreover each part of the above triplet has a known distribution. The process $(g(t)^{-1/2}B(g(t)s),\ 0\leqslant s\leqslant 1)$ is distributed as the Brownian bridge $(b(s),\ 0\leqslant s\leqslant 1)$ $s \leq 1$), (see e.g. [1]) and the second component in (1.5) is the Brownian meander denoted m. The scaling property of the Brownian motion implies that $g(t)$ is distributed as $tg(1)$ while the distribution of $g(1)$ is the arcsine one (see again [1]):

$$
\mathbb{P}(g(1) \in dx) = \frac{1}{\pi \sqrt{x(1-x)}} \mathbb{1}_{[0,1]}(x) dx.
$$
 (1.6)

Consequently,

$$
(U^*(t), U^{**}(t)) \stackrel{(d)}{=} \left(\sqrt{tg(1)}b^*, \sqrt{t(1-g(1))} \max_{0 \le u \le 1} m(u)\right) \tag{1.7}
$$

where $b^* := \sup_{0 \le s \le 1} |b(s)|$. Its distribution function is given by the Kolmogorov-Smirnov formula (see e.g. [10]):

$$
\mathbb{P}(b^* > x) = 2\sum_{k \ge 1} (-1)^{k-1} e^{-2k^2 x^2}, \quad x > 0.
$$
 (1.8)

Formula (1.7) permits to determine the law of $(U^*(t),U^{**}(t))$, once we know the distribution of $\max\limits_{0\leqslant u\leqslant 1}m(u).$ But by [2], for any bounded Borel function f ,

$$
\mathbb{E}[f(m(u), 0 \le u \le 1)] = \sqrt{\frac{\pi}{2}} \mathbb{E}\left[\frac{1}{R(1)} f(R(u), 0 \le u \le 1)\right],
$$
\n(1.9)

where $(R(u), 0 \leq u \leq 1)$ stands for a 3-dimensional Bessel started at 0. Due to the scaling property (1.7), we deduce that p_c does not depend on t and

$$
p_c = \sqrt{\frac{\pi}{2}} \mathbb{E}\left[F\left(b^* \sqrt{\frac{g(1)}{1 - g(1)}}\right)\right]
$$
\n(1.10)

where

$$
F(x) := \mathbb{E}\left[\frac{1}{R(1)}1\!\!1_{\{\max_{0\leq u\leq 1} R(u)< x\}}\right].\tag{1.11}
$$

According to (1.10) we have first to determine the function F (see Lemma 2.1 below), second the distribution function of $b^* \sqrt{\dfrac{g(1)}{1-g(1)}}$ (see Proposition 2.2) and third to calculate the expectation. The details are given in Section 2.

2 Proof of Theorem 1.1

Lemma 2.1. For any $x > 0$,

$$
F(x) = \sqrt{\frac{2}{\pi}} \sum_{k \in \mathbb{Z}} \left\{ e^{-2k^2 x^2} - e^{-(2k+1)^2 x^2/2} \right\} = \frac{4}{x} \sum_{k \ge 0} \exp \left\{ -\frac{(2k+1)^2 \pi^2}{2x^2} \right\}.
$$
 (2.1)

Proof. First, by [4, formula 1.1.8, p317],

$$
\mathbb{P}\left(\max_{0 \le u \le 1} R(u) < y, \ R(1) \in dz \, \middle| \, R(0) = x\right) = \frac{z}{x\sqrt{2\pi}} \times S \times 1_{\{y > x, z < y\}} dz\tag{2.2}
$$

ECP **0** [\(2012\), paper 0.](http://dx.doi.org/10.1214/ECP.vVOL-PID)

where

$$
S = \sum_{k \in \mathbb{Z}} \left[\exp \left\{ -\frac{(z - x + 2ky)^2}{2} \right\} - \exp \left\{ -\frac{(z + x + 2ky)^2}{2} \right\} \right].
$$

A Taylor expansion of $x \mapsto \exp \left\{-\frac{(z \pm x + 2ky)^2}{2}\right\}$ $\left\{\frac{+2ky)^2}{2}\right\}$ at $x=0$ leads to

$$
\mathbb{P}\left(\max_{0\leq u\leq 1} R(u) < y, \ R(1) \in dz \, \bigg| R(0) = 0\right) = \frac{2z}{\sqrt{2\pi}} \left[\sum_{k\in\mathbb{Z}} (z + 2ky) \exp\left\{-\frac{(z + 2ky)^2}{2}\right\} \right] \, \mathbb{1}_{\{z < y\}} dz.
$$

As a consequence,

$$
F(y) = \frac{2}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \int_0^y (z + 2ky) \exp\left\{-\frac{(z + 2ky)^2}{2}\right\} dz
$$

= $\frac{2}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\exp\left\{-2k^2y^2\right\} - \exp\left\{-\frac{(2k+1)^2y^2}{2}\right\} \right].$

The second equality in (2.1) is a direct consequence of the Poisson summation formula (see, e.g. [7, Chap. XIX p.630]:

$$
\frac{1}{\sqrt{2\pi t}}\sum_{k\in\mathbb{Z}}\left[\exp\left\{-\frac{1}{2t}\left(x_0+2k(\beta_0-\alpha_0)\right)^2\right\}-\exp\left\{-\frac{1}{2t}\left(2\beta_0-x_0+2k(\beta_0-\alpha_0)\right)^2\right\}\right]
$$
\n
$$
=\frac{1}{\beta_0-\alpha_0}\sum_{k\geqslant 1}\left[\left(\cos\left(\frac{k\pi x_0}{\beta_0-\alpha_0}\right)-\cos\left(\frac{k\pi(2\beta_0-x_0)}{\beta_0-\alpha_0}\right)\right)\exp\left\{-\frac{k^2\pi^2t}{2(\beta_0-\alpha_0)^2}\right\}\right]
$$
\napplied with $t=1$, $x_0=0$, $\beta_0=x/2$ and $\alpha_0=-x/2$.

applied with $t = 1$, $x_0 = 0$, $\beta_0 = x/2$ and $\alpha_0 = -x/2$.

Proposition 2.2. For any $x > 0$,

$$
\mathbb{P}\left(b^*\sqrt{\frac{g(1)}{1-g(1)}} > x\right) = \frac{2}{\pi} \int_0^\infty A(u)e^{-2x^2u} \frac{du}{\sqrt{u}},\tag{2.3}
$$

where

$$
A(u) := \sum_{k \geqslant 1} (-1)^{k-1} \frac{k}{k^2 + u}.
$$
\n(2.4)

Proof. We introduce a cut-off $0 < \varepsilon < 1$ and we define:

$$
\varphi_{\varepsilon}(x) := \mathbb{P}\left(g(1) < 1 - \varepsilon, \ b^* \sqrt{\frac{g(1)}{1 - g(1)}} > x\right). \tag{2.5}
$$

Using the independence between $g(1)$ and b^* , (1.6) and (1.8), we deduce:

$$
\varphi_{\varepsilon}(x) = \frac{1}{\pi} \int_0^{1-\varepsilon} \frac{1}{\sqrt{y(1-y)}} \mathbb{P}\left(b^* > x\sqrt{\frac{1-y}{y}}\right) dy = \frac{2}{\pi} \sum_{k \geqslant 1} I_k(\varepsilon)
$$

where $I_k(\varepsilon) := (-1)^{k-1} \int^{1-\varepsilon}$ 0 1 $\sqrt{y(1-y)}$ $\exp \left\{-\frac{2k^2x^2(1-y)}{x^2}\right\}$ \hat{y} $\partial_t dy$. The inversion of the sum and the integral is available since $(1-y)/y \geqslant \varepsilon' > 0$ where $\varepsilon' := \varepsilon/(1-\varepsilon)$. Making the change of variables $(1 - y)/y = u/k^2$ leads to:

$$
I_k(\varepsilon) = \frac{(-1)^{k-1}}{k} \int_{\varepsilon' k^2}^{\infty} \frac{1}{\sqrt{u}(1+u/k^2)} \exp\left\{-2x^2u\right\} du.
$$

ECP **0** [\(2012\), paper 0.](http://dx.doi.org/10.1214/ECP.vVOL-PID)

The identity $\frac{1}{1+u/k^2} = 1 - \frac{u}{k^2(1+u)}$ $\frac{1}{k^2(1+u/k^2)}$ allows to invert the sum and the integral. Finally we get:

$$
\varphi_{\varepsilon}(x) = \frac{2}{\pi} \int_{\varepsilon'}^{\infty} \left(\sum_{\varepsilon' k^2 \leq u} \frac{(-1)^{k-1}}{k} \frac{1}{1+u/k^2} \right) \exp\left\{-2x^2 u\right\} \frac{du}{\sqrt{u}} = \frac{2}{\pi} \int_{\varepsilon'}^{\infty} S_{n(\varepsilon', u)}(u) \frac{\exp\left\{-2x^2 u\right\}}{\sqrt{u}} du
$$

with $n(\varepsilon', u) = \lfloor \sqrt{u/\varepsilon'} \rfloor$, $S_n(u) := \sum^{n}$ $k=1$ $(-1)^{k-1}\phi(1/k, u)$ and $\phi(y, u) := y/(1 + uy^2)$.

Note that $\Big|$ $\partial \phi$ ∂y $\begin{array}{c} \hline \end{array}$ $\leqslant 1$, then, considering $n = 2m$ and $n = 2m + 1$ and using the mean value inequality we obtain:

$$
|S_{2m}(u)| = \left|\sum_{k'=1}^{m} \phi\left(\frac{1}{2k'-1}, u\right) - \phi\left(\frac{1}{2k'}, u\right)\right| \leqslant \sum_{k'=1}^{m} \left|\frac{1}{2k'-1} - \frac{1}{2k'}\right| \leqslant \sum_{k'=1}^{\infty} \frac{1}{2k'(2k'-1)} < \infty. \tag{2.6}
$$

Similarly, $|S_{2m+1}(u)| < \infty$. Since $\varepsilon' \to 0$ and $n(\varepsilon', u) \to 0$ as ε goes to 0, then, identity (2.3) is a direct consequence of the Lebesgue dominated convergence theorem. \square

Lemma 2.1 and Proposition 2.2 allow to obtain a new integral form for p_c . **Lemma 2.3.** One has

$$
p_c = 8 \int_0^\infty \frac{uA(u^2)}{\sinh(2\pi u)} du.
$$

Proof. We deduce easily from (2.4) that

$$
A(u) = \sum_{k' \geq 1} \phi\left(\frac{1}{2k'-1}, u\right) - \phi\left(\frac{1}{2k'}, u\right),
$$

where $\phi(y, u) := y/(1 + uy^2)$. Then inequality (2.6) implies that $\sup_{u \geq 0} A(u) < \infty$. By Lemma 2.1, Proposition 2.2, the definition (2.4) of A and the Fubini theorem, we get

$$
p_c = \sqrt{\frac{\pi}{2}} \frac{32}{\pi} \sum_{k \ge 0} \int_0^\infty \sqrt{u} A(u) \left(\int_0^\infty \exp \left\{ -\frac{(2k+1)^2 \pi^2}{2x^2} - 2x^2 u \right\} dx \right) du.
$$

But making $s = 2x^2u$ and letting $z = 2(2k+1)\pi\sqrt{u}$, we get:

$$
\int_0^\infty \exp\left\{-\frac{(2k+1)^2\pi^2}{2x^2} - 2x^2u\right\}dx = \frac{1}{\sqrt{2u}}\left(\frac{z}{2}\right)^{1/2}K_{1/2}(z)
$$

where (cf [13, Formula (15) p183])

$$
K_{\nu}(x) = \frac{1}{2} \left(\frac{x}{2}\right)^{\nu} \int_0^{\infty} \exp\left\{-s - \frac{x^2}{4s}\right\} ds.
$$

Recall that $K_{\nu}=K_{-\nu}$ [13, Formula (8) p79] and $K_{-1/2}(x)=\sqrt{\frac{\pi}{2}}$ $\frac{\pi}{2x}e^{-x}$ [13, Formula (13) p80] . It follows that

$$
p_c = 8 \sum_{k \ge 0} \int_0^\infty A(u) e^{-2(2k+1)\pi\sqrt{u}} du = 4 \int_0^\infty \frac{A(u)}{\sinh\left(2\pi\sqrt{u}\right)} du = 8 \int_0^\infty \frac{v A(v^2)}{\sinh\left(2\pi v\right)} dv.
$$

ECP **0** [\(2012\), paper 0.](http://dx.doi.org/10.1214/ECP.vVOL-PID)

Maximum of the reflected Brownian motion before its last zero

We now focus on the function A . Our method is based on the crucial fact that A can be expressed with the function ψ defined by (1.3).

Lemma 2.4. 1. We have:

$$
A(u) = \frac{1}{4} \left[\psi\left(\frac{i\sqrt{u}}{2}\right) + \psi\left(-\frac{i\sqrt{u}}{2}\right) - \psi\left(\frac{1+i\sqrt{u}}{2}\right) - \psi\left(\frac{1-i\sqrt{u}}{2}\right) \right], \quad u \ge 0.
$$
\n(2.7)

2. There exists $a, b > 0$ such that

$$
|\psi(z)| \leqslant a + b |z|^2, \quad \forall \ z \in \mathbb{C}, \ |Im \ z| \geqslant 1. \tag{2.8}
$$

Consequently, $p_c = I_2 - I_1$, where

$$
I_k := 2 \int_0^\infty \frac{v F_k(v)}{\sinh(2\pi v)} dv, \ k = 1, 2 \tag{2.9}
$$

and $F_1(v) := \psi\left(\frac{1+iv}{2}\right) + \psi\left(\frac{1-iv}{2}\right)$, $F_2(v) := \psi\left(\frac{iv}{2}\right) + \psi\left(-\frac{iv}{2}\right)$.

Proof. Formula (2.7) and inequality (2.8) are a direct consequence of identity (3), p 15 in [6], i.e.

$$
\psi(z) = -\gamma - \frac{1}{z} + \sum_{n \ge 1} \frac{z}{n(z+n)} = -\gamma - \frac{1}{z} + \left(\sum_{n \ge 1} \frac{1}{n^2}\right)z - \sum_{n \ge 1} \frac{z^2}{n^2(z+n)}\tag{2.10}
$$

with
$$
\gamma := \lim_{m \to \infty} \left(\sum_{k=1}^{m} \frac{1}{k} - \ln m \right)
$$
.

Due to the form of the functions F_1 and F_2 , the integrals I_1 and I_2 can be viewed as integrals over a straight line in the plane. More precisely, we have:

Lemma 2.5. I_1 and I_2 can be written as:

$$
I_1 = -8i \int_{\Delta_{1/2}} \frac{z - 1/2}{\sin(4\pi z)} \psi(z) dz, \quad I_2 = -8i \int_{\Delta_0} \frac{z\psi(z+1)}{\sin(4\pi z)} dz
$$
 (2.11)

where, for any $a \in \mathbb{R}$, Δ_a is the line with parametrization by $z = a + it$, $t \in \mathbb{R}$.

Proof. 1) We note that $sin(2\pi iv) = i sinh(2\pi v)$. Thus

$$
2\int_0^\infty \frac{v}{\sinh(2\pi v)} \psi\left(\frac{1+iv}{2}\right) dv = \frac{8}{i} \int_0^\infty \frac{\frac{1+iv}{2} - \frac{1}{2}}{\sin\left(4\pi i \left(\frac{1+iv}{2}\right)\right)} \psi\left(\frac{1+iv}{2}\right) d\left(\frac{1+iv}{2}\right)
$$

$$
= -8i \int_{\Delta'_{1/2}} \frac{z - 1/2}{\sin(4\pi z)} \psi(z) dz
$$

where Δ'_a is the half-line: $z = a + it$, $t \ge 0$. Similarly,

$$
2\int_0^\infty \frac{v}{\sinh(2\pi v)}\psi\left(\frac{1-iv}{2}\right)dv = -8i\int_{\Delta_{1/2}''} \frac{z-1/2}{\sin(4\pi z)}\psi(z) dz
$$

where $\Delta''_a := \{z = a + it, t \leq 0\}$ and $a \in \mathbb{R}$. This implies the value of I_1 given by (2.11).

ECP **0** [\(2012\), paper 0.](http://dx.doi.org/10.1214/ECP.vVOL-PID)

Page 6/9

2) Formula (2.10) tells us that $\hat{\psi}(z) := \psi(z) + \frac{1}{z}$ has no singularity at $z = 0$. Thus we study

$$
2\int_0^\infty \frac{v}{\sinh\left(2\pi v\right)}\hat{\psi}\left(\frac{iv}{2}\right)dv = \frac{8}{i}\int_0^\infty \frac{iv/2}{\sin\left(4\pi \frac{iv}{2}\right)}\hat{\psi}\left(\frac{iv}{2}\right)d\left(\frac{iv}{2}\right) = -8i\int_{\Delta_0'}\frac{z}{\sin\left(4\pi z\right)}\hat{\psi}\left(z\right)dz
$$

and similarly

$$
2\int_0^\infty \frac{v}{\sinh\left(2\pi v\right)}\hat{\psi}\left(-\frac{iv}{2}\right)dv = 2\int_0^\infty \frac{iv}{\sin\left(2\pi iv\right)}\hat{\psi}\left(\frac{iv}{2}\right)dv = -8i\int_{\Delta_0^{\prime\prime}}\frac{z}{\sin\left(4\pi z\right)}\hat{\psi}\left(z\right)dz.
$$

The identity (formula (8) p 16 in [6]) :

$$
\hat{\psi}(z) = \psi(z) + \frac{1}{z} = \psi(z+1)
$$
\n(2.12)

implies $\psi\left(\frac{iv}{\delta}\right)$ $+\psi\left(-\frac{iv}{\delta}\right)$ $\bigg\} = \hat{\psi}\left(\frac{iv}{\delta}\right)$ $+\hat{\psi}\left(-\frac{iv}{\delta}\right)$ $\big)$ and finally (2.11). \Box 2 2 2 2

We show in the following lemma that I_1 and I_2 are integrals over the vertical line. **Lemma 2.6.** Let $0 < \varepsilon < 1/4$, then

$$
I_1 = -8i \int_{\Delta_{1/2+\varepsilon}} \frac{z - 1/2}{\sin(4\pi z)} \psi(z) dz, \quad I_2 = -8i \int_{\Delta_{1/2+\varepsilon}} \frac{z\psi(z+1)}{\sin(4\pi z)} dz + \psi(1/4) - 2\psi(1/2).
$$
\n(2.13)

Proof. We only deal with I_2 , the proof related to I_1 is similar and easier. The quantity $\sin(4\pi z)$ cancels at $z = k/4$ for every $k \in \mathbb{Z}$, the zeros are simple. From (2.10), we deduce that $h(z) := z\psi(z+1)/\sin(4\pi z)$ is meromorphic in $\{z \in \mathbb{C}; -1/4 < Re z < 3/4\}$ with poles at $1/4$ and $1/2$. We introduce the contour defined in Figure 2.

Then the residue theorem gives

$$
\int_{C_nB_n} h(z)dz - \int_{D_nA_n} h(z)dz + \int_{D_nC_n} h(z)dz - \int_{B_nA_n} h(z)dz = 2i\pi \left\{ Res\left(h, 1/4\right) + Res\left(h, 1/2\right) \right\}.
$$
\n(2.14)

The residual at $1/4$ is given by

$$
Res(h, 1/4) = \lim_{z \to 1/4} \left\{ h(z)(z - 1/4) \right\} = \frac{1}{4} \psi (5/4) \left(\lim_{z \to 1/4} \frac{z - 1/4}{\sin(4\pi z) - \sin(4\pi 1/4)} \right) = -\frac{1}{16\pi} \psi (5/4)
$$

Using (2.12) with $z = 1/4$, we get $Res(h, 1/4) = -\frac{1}{4\pi} \left(1 + \frac{1}{4} \psi (1/4) \right)$. Similarly, $Res(h, 1/2) = \frac{1}{4\pi} (1 + \psi (1/2) / 2)$.

ECP **0** [\(2012\), paper 0.](http://dx.doi.org/10.1214/ECP.vVOL-PID)

Now, we let n goes to infinity. Inequality (2.8) implies

$$
\lim_{n \to \infty} \int_{A_n B_n} h(z) dz = \lim_{n \to \infty} \int_{C_n D_n} h(z) dz = 0.
$$

Indeed, it follows from the parametrization of A_nB_n of the type $z = in + t$, inequality $|in + t| \le n + 1$ is valid for $0 \le t \le \varepsilon + 1/2$, and

$$
|\sin(4\pi(in+t))|^2 = \sinh(4\pi n)^2 \cos(4\pi t)^2 + \cosh(4\pi n)^2 \sin(4\pi t)^2
$$

\n
$$
\ge \min \left\{ \sinh(4\pi n)^2, \cosh(4\pi n)^2 \right\}.
$$

We proceed analogously on C_nD_n . As a consequence, letting $n \to \infty$ in (2.14), we get

$$
I_2 = -8i \left\{ \int_{\Delta_{1/2+\varepsilon}} h(z)dz - 2i\pi \left[-\frac{1}{4\pi} \left(1 + \psi \left(1/4 \right) / 4 \right) + \frac{1}{4\pi} \left(1 + \psi \left(1/2 \right) / 2 \right) \right] \right\}
$$

= $-8i \int_{\Delta_{1/2+\varepsilon}} h(z)dz + \psi \left(1/4 \right) - 2\psi \left(1/2 \right).$

 \Box

Bringing together Lemmas 2.4, 2.6 leads to

$$
p_c = -8i \int_{\Delta_{1/2+\varepsilon}} \frac{1}{\sin(4\pi z)} \left(z\psi(z+1) - (z-\frac{1}{2})\psi(z) \right) dz + \psi(1/4) - 2\psi(1/2).
$$

Setting $z = 1/2 + u$ and using identity (2.12) with $u + 1/2$ instead of z gives:

$$
p_c = -8i \int_{\Delta_{\varepsilon}} h^+(u) du + \psi(1/4) - 2\psi(1/2), \qquad (2.15)
$$

where $h^{\pm}(z) := \frac{1}{\sin{(4\pi z)}}$ $\left(1 + \frac{1}{2}\right)$ $\frac{1}{2}\psi(\frac{1}{2})$ $\frac{1}{2} \pm z)$.

We are not able to calculate directly $\overline{}$ $Δ_ε$ $h^+(u)du$, however it is possible for $Δ_ε$ $h^+(u)du \pm$ Z $h^-(u)du$.

Lemma 2.7. For any $\varepsilon \in]0, 1/4[$,

∆^ε

$$
\int_{\Delta_{\varepsilon}} h^{+}(u) du + \int_{\Delta_{\varepsilon}} h^{-}(u) du = \frac{i}{2} (1 + \psi(1/2)/2)
$$
\n(2.16)

$$
\int_{\Delta_{\varepsilon}} h^{+}(u) du - \int_{\Delta_{\varepsilon}} h^{-}(u) du = \frac{\pi}{2} \int_{\Delta_{\varepsilon}} \frac{\tan(\pi z)}{\sin(4\pi z)} dz.
$$
\n(2.17)

Proof. 1) We begin proving (2.16). The function h^+ is meromorphic in $\{z \in \mathbb{C}; -1/4 < Re\ z < 1/4\}$ with unique pole $z = 0$. Then the residue theorem yields

$$
\int_{\Delta_{\varepsilon}} h^+(z)dz = \int_{\Delta_{-\varepsilon}} h^+(z)dz + 2i\pi Res(h^+,0) = -\int_{\Delta_{\varepsilon}} h^-(z)dz + \frac{i}{2}(1+\psi(1/2)/2).
$$

2) Formula (2.17) is a direct consequence of formula 11 p16 in [6]: $\psi\left(\frac{1}{2}+z\right)-\psi\left(\frac{1}{2}-z\right)=$ $\pi \tan(\pi z)$.

It is easy to deduce from (2.16) and (2.17) that:

$$
2\int_{\Delta_{\varepsilon}} h^+(z)dz = \frac{\pi}{2} \int_{\Delta_{\varepsilon}} \frac{\tan(\pi z)}{\sin(4\pi z)} dz + \frac{i}{2} (1 + \psi(1/2)/2).
$$

ECP **0** [\(2012\), paper 0.](http://dx.doi.org/10.1214/ECP.vVOL-PID)

Relation (2.15) implies directly that p_c equals $\psi(1/4) - \psi(1/2) + 2 - 2\pi i\alpha(\epsilon)$, where $\alpha(\varepsilon) :=$ ∆^ε $\tan(\pi z)$ $\frac{\sinh(n\pi)}{\sin(4\pi z)}$ dz. Since the real number p_c does not depend on ε , letting $\varepsilon \to 0$ leads to:

$$
p_c = \psi(1/4) - \psi(1/2) + 2 - 2\pi i\alpha(0+),
$$
\n(2.18)

.

where $\alpha(0+) = \alpha$ Δ_0 $\tan(\pi z)$ $\frac{\sin(kz)}{\sin(4\pi z)}$ dz. This integral is easy to calculate.

$$
\alpha(0+) = i \int_{\mathbb{R}} \frac{\tanh(\pi x)}{\sinh(4\pi x)} dx = \frac{i}{4} \int_{\mathbb{R}} \frac{dx}{\cosh^2(\pi x) \cosh(2\pi x)}
$$

We make the change of variable $u = \tanh(\pi x)$:

$$
\alpha(0+) = \frac{i}{2\pi} \int_0^1 \frac{1-u^2}{1+u^2} du = \frac{i}{2\pi} \left(-1 + 2 \int_0^1 \frac{du}{1+u^2} \right) = \frac{i}{2\pi} \left(-1 + \frac{\pi}{2} \right).
$$

Theorem 1.1 follows from (2.18) and the above result.

References

- [1] J. Bertoin. *Lévy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- [2] Ph. Biane, J.-F. Le Gall, and M. Yor. Un processus qui ressemble au pont brownien. In Séminaire de Probabilités, XXI, volume 1247 of Lecture Notes in Math., pages 270–275. Springer, Berlin, 1987.
- [3] P. Billingsley. Convergence of probability measures. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons Inc., New York, second edition, 1999. A Wiley-Interscience Publication.
- [4] A.N. Borodin and P. Salminen. Handbook of Brownian motion—facts and formulae. Probability and its Applications. Birkhäuser Verlag, Basel, second edition, 2002.
- [5] C. Chabriac, A. Lagnoux, S. Mercier, and P. Vallois. Elements related to the largest complete excursion of a reflected Brownian motion stopped at a fixed time. Application to local score. Stochastic Processes and their Applications, 124(12), 2014.
- [6] A. Erdélyi, W. Magnus, F. Oberhettinger, and F.G. Tricomi. Higher transcendental functions. Vol. I. Robert E. Krieger Publishing Co., Inc., Melbourne, Fla., 1981. Based on notes left by Harry Bateman, With a preface by Mina Rees, With a foreword by E. C. Watson, Reprint of the 1953 original.
- [7] W. Feller. An introduction to probability theory and its applications. Vol. II. Second edition. John Wiley & Sons, Inc., New York-London-Sydney, 1971.
- [8] S. Karlin and S.-F. Altschul. Methods for assessing the statistical significance of molecular sequence features by using general scoring schemes. PNAS, 87:2264–2268, 1990.
- [9] S. Mercier and J.J. Daudin. Exact distribution for the local score of one i.i.d. random sequence. Jour. Comp. Biol, 8(4):373–380, 2001.
- [10] J. Pitman and M. Yor. On the distribution of ranked heights of excursions of a Brownian bridge. Ann. Probab., 29(1):361–384, 2001.
- [11] B. Roynette, P. Vallois, and M. Yor. Penalisations of Brownian motion with its maximum and minimum processes as weak forms of Skorokhod embedding. Theory Stoch. Process., 14(2):116–138, 2008.
- [12] M.S. Waterman. Introduction to Computational Biology: Maps, Sequences and Genomes. Chapman & Hall, 1995.
- [13] G.N. Watson. A treatise on the theory of Bessel functions. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1995. Reprint of the second (1944) edition.