



HAL
open science

The 1-loop self-energy of an electron in a strong external magnetic field revisited

Bruno Machet

► **To cite this version:**

Bruno Machet. The 1-loop self-energy of an electron in a strong external magnetic field revisited. 2015. hal-01215168v1

HAL Id: hal-01215168

<https://hal.science/hal-01215168v1>

Preprint submitted on 13 Oct 2015 (v1), last revised 13 Nov 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**THE 1-LOOP SELF-ENERGY OF AN ELECTRON
IN A STRONG EXTERNAL MAGNETIC FIELD REVISITED**

B. Machet ^{1 2 3 4}

Abstract: I revisit the 1-loop self-energy of an electron in a strong, constant and uniform external magnetic field B . First, I show, after Tsai [1], how, for an electron in the lowest Landau level, Schwinger's techniques [2] explained by Dittrich and Reuter [3] lead to the same integral deduced by Demeur [4] and used later by Jancovici [5]. Then, I calculate the Demeur-Jancovici integral in the range $75 \leq L \equiv \frac{|e|B}{m^2} \leq 10\,000$, which yields $\delta m \simeq \frac{\alpha m}{4\pi} \left[\left(\ln L - \gamma_E - \frac{3}{2} \right)^2 - \frac{9}{4} + \frac{\pi}{\beta-1} + \frac{\pi^2}{6} + \frac{\pi \Gamma[1-\beta]}{L^{\beta-1}} + \frac{1}{L} \left(\frac{\pi}{2-\beta} - 5 \right) + \mathcal{O}\left(\frac{1}{L^{\geq 2}}\right) \right]$, $\beta \simeq 1.175$, close to Jancovici's last estimate $\delta m \simeq \frac{\alpha m}{4\pi} \left[\left(\ln 2L - \gamma_E - \frac{3}{2} \right)^2 + A + \dots \right]$ with $A \simeq 3.5$ (previously undetermined). The term proportional to $(\ln \frac{|e|B}{m^2})^2$ can never be considered to be leading and gets largely damped, in particular by the large $\ln \frac{|e|B}{m^2}$ which arises from the counterterm implementing suitable renormalization conditions. The addition with respect to existing literature is small but some consequences may be worth deeper investigations.

PACS: 12.15.Lk 12.20.Ds

¹Sorbonne Université, UPMC Univ Paris 06, UMR 7589, LPTHE, F-75005, Paris, France

²CNRS, UMR 7589, LPTHE, F-75005, Paris, France.

³Postal address: LPTHE tour 13-14, 4^{ème} étage, UPMC Univ Paris 06, BP 126, 4 place Jussieu, F-75252 Paris Cedex 05 (France)

⁴machet@lpthe.jussieu.fr

1 Generalities

I shall be concerned in this short ⁵ note, with the self-energy of an electron at 1-loop in the presence of a strong external, constant and uniform, magnetic field B . The electron propagator is described by the sum of the 2 diagrams




Fig. 1: 1-loop radiative correction to the mass of an electron.

in which the double horizontal lines, external as well as internal, stand for an electron of mass m in an external B . The self-energy that we shall calculate is the second diagram. For the sake of simplicity, we shall restrict external electrons to lie in the lowest Landau level. This does not apply to the internal electron propagator, which includes a summation over all Landau levels.

1.1 Historical remarks

To my knowledge, the uses ⁶ of the self-energy of an electron in a strong external B rely on the $\left(\ln \frac{|e|B}{m^2}\right)^2$ that has been extracted in 1969 by Jancovici [5] from a general formula deduced by Demeur in 1953 [4] and, as far as I could see, Demeur's calculations, performed with techniques which are unfamiliar today, have not been reproduced ⁷. Despite the presence of potentially large corrections was mentioned at the end of Jancovici's paper [5] (in there, one constant could not be determined), all terms but the $(\ln)^2$ have been dropped, later, with the argument that they are "non-leading". I shall show that this is untenable.

An alternate way is the one pioneered by Schwinger in the late 1940's [2]. Calculations have been explained in details in the book by Dittrich and Reuter [3] in 1985, which includes a long list of references. One finds there, in particular, the expression for the renormalized 1-loop mass operator $\Sigma(\pi)$ for an electron in an external B , as deduced in 1974 by Tsai [1], which will be our starting point. At the end of his paper, Tsai mentions that his calculation, which uses the techniques and results of Schwinger, yields, when projected on the ground state of the electron, "...the known result of Demeur" (this correspondence is the subject of section 2).

1.2 The procedure

I go along Schwinger's path and make use of Demeur's technique [4] to sandwich the mass operator $\Sigma(\pi)$ between 2 "privileged" electron states $|\psi\rangle$ (to reproduce the terminology of Demeur and previous authors, in particular Luttinger [15]), on mass-shell. This restricts, but greatly simplifies the calculations. This matrix element corresponds to δm of the electron at 1-loop in the presence of B , the electron mass being defined as the pole of its propagator (subsection 2.2). The privileged state, that always exists in the presence of B , is the one with energy m . In our present terminology, it corresponds to the Lowest Landau Level (LLL) and, on mass shell, it satisfies the Dirac equation $(\gamma\pi + m)|\psi\rangle = 0$, $\pi = p - eA$ ^{8 9}.

Then, I show how changes of variables cast δm in the form deduced by Demeur [4] and used by Jancovici [5]. It is a convergent double integral that only depends on $\frac{|e|B}{m^2}$. A fully analytical evaluation lies beyond my ability. However,

⁵This is why I do not pay a fair enough tribute to the many authors that contributed to this subject, and I apologize for this. I will instead insist on very small details, generally not mentioned, that can help the reader.

⁶Some important steps can be found in [6] [7] [8], [9], [10], in the book [11] and in the review [12] in which one can find a large amount of other important references.

⁷They have been critically examined and completed by Newton [14] at small values of $\frac{|e|B}{m^2}$, but this path seems to have then been abandoned.

⁸ $\gamma\pi$ stands for $\gamma_\mu\pi^\mu$.

⁹We use Schwinger's metric $(-, +, +, +)$.

a trick due to M.I. Vysotsky in his study the screening of the Coulomb potential in an external magnetic field [16] comes to the rescue: the part of the integrand that resists analytical integration can be nearly perfectly fitted inside the range of integration by a simpler function that can be analytically integrated.

2 The self-energy Σ in external B of an electron in the lowest Landau level; equivalence between the calculations of Schwinger and Demeur

2.1 The general formula for the electron self-energy operator at 1-loop

From now onwards we rely on the operatorial expression of the self-energy of an electron in an external B deduced by Tsai [1]¹⁰ in the formalism of Schwinger (in the whole paper “c-term” stands for “counterterm”)

$$\begin{aligned} \Sigma(\pi) = \frac{\alpha m}{2\pi} \int_0^\infty \frac{ds}{s} \int_0^1 du e^{-isu^2 m^2} \left\{ \frac{1}{\sqrt{\Delta}} e^{-is\Phi} \left[1 + e^{-2i\sigma^3 Y} + (1-u)e^{-2i\sigma^3 Y} \frac{\not{Y}}{m} \right. \right. \\ \left. \left. + (1-u) \left(\frac{1-u}{\Delta} + \frac{u}{\Delta} \frac{\sin Y}{Y} e^{-i\sigma^3 Y} - e^{-2i\sigma^3 Y} \right) \frac{\not{Y}_\perp}{m} \right] \right. \\ \left. \underbrace{-(1+u) - (m + \not{Y}) \left[\frac{1-u}{m} - 2imu(1-u^2)s \right]}_{c\text{-term}} \right\} \end{aligned} \quad (1)$$

in which the notations are the following^{11 12}

$$\begin{aligned} Y &= eB s u, \\ \Delta &= (1-u)^2 + 2u(1-u) \frac{\sin Y \cos Y}{Y} + u^2 \left(\frac{\sin Y}{Y} \right)^2, \\ \Phi &= u(1-u)[m^2 - (\gamma\pi)^2] + \frac{u}{Y}[\beta - (1-u)Y]\pi_\perp^2 - u^2 \frac{e}{2} \sigma_{\mu\nu} F^{\mu\nu}, \\ \tan \beta &= \frac{(1-u) \sin Y}{(1-u) \cos Y + u \frac{\sin Y}{Y}}. \end{aligned} \quad (2)$$

This formula has been obtained with an internal photon in the Feynman gauge (like Demeur [4]) and internal electron propagator inside an external B as determined by Schwinger¹³ (it includes all Landau levels).

The variables of integration s and u are deduced from the 2 Schwinger’s parameters s_1 and s_2 , respectively for the electron and for the photon¹⁴ and both integrated from 0 to ∞ , by the change $s_1 = su$, $s_2 = s(1-u)$; this manifestly requires cutting the extremities 0 and 1 off the u -interval of integration, values at which respectively s_1 and s_2 stay vanishing (or, at best, are undefined).

The counterterm is determined by the 2 equations (3.39) and (3.40) of [3]

$$\lim_{\gamma\pi \rightarrow -m} \lim_{B \rightarrow 0} \Sigma(\pi) = 0, \quad \lim_{\gamma\pi \rightarrow -m} \lim_{B \rightarrow 0} \frac{\partial \Sigma(\pi)}{\partial \not{Y}} = 0. \quad (3)$$

They ensure that, after turning off B , ($\gamma\pi \rightarrow \gamma p$), the renormalization conditions $\Sigma(\not{p} = -m) = 0$ and $\frac{\partial \Sigma}{\partial \not{p}}(\not{p} = -m) = 0$ are fulfilled¹⁵. The renormalized electron mass (pole of its propagator), is then (i.e. at $B = 0$) defined by

$$m = m_0 + \delta m, \quad \delta m = \Sigma(\gamma p + m = 0). \quad (4)$$

¹⁰This is eq. (3.44) p.52 of Dittrich-Reuter [3]. The expressions for Φ and Δ are given in their equations (3.38b) and (3.38c) (see also footnote 12).

¹¹ e stands here for the charge of the electron $e < 0$.

¹²There is a sign misprint in the definition (3.38b) of Φ in [3], which has been corrected here. The correct sign is the one in eq. (3.35) of [3].

¹³See for example [1].

¹⁴For example $\frac{1}{k^2 - i\epsilon} = i \int_0^\infty ds_2 e^{-is_2(k^2 - i\epsilon)}$.

¹⁵These renormalization conditions are carefully explained in p. 38-41 of [3].

2.2 Defining the electron mass in the presence of B

The propagator of a Dirac electron in an external field A^μ is

$$G = \frac{1}{\gamma\pi + m_0 + \Sigma(\pi)}, \quad \pi = p - eA. \quad (5)$$

We define, in analogy with eq. (4), the mass of the electron as the pole of its propagator by

$$m = m_0 + \Sigma(\gamma\pi + m = 0) \Leftrightarrow \delta m = \Sigma(\gamma\pi + m = 0). \quad (6)$$

δm is expected to depend on the external field. Note that the mass-shell is defined by $(\gamma\pi)^2 \equiv -\pi^2 + \frac{e}{2} \sigma_{\mu\nu} F^{\mu\nu} = m^2 \neq p^2$.

2.3 Projecting $\Sigma(\pi)$ on the ‘‘privileged state’’: δm for the lowest Landau level

The spectrum of a Dirac electron in a pure magnetic field directed along z is [17]

$$\epsilon_n^2 = m^2 + p_z^2 + (2n + 1 + \sigma_z) |e|B \quad (7)$$

in which $\sigma_z = \pm 1$ is $2 \times$ the spin projection of the electron on the z axis. So at $n = 0, \sigma_z = -1, p_z = 0, \epsilon_n = m$: this is the privileged state, which is nothing more than the lowest Landau level.

We can consider $A_\mu = \begin{pmatrix} A_0 = 0 \\ A_x = 0 \\ A_y = xB \\ A_z = 0 \end{pmatrix}$ such that $F_{12} = B$ is the only non-vanishing component of the classical external $F_{\mu\nu}$. Then, the wave function of the privileged state of energy m writes [15] [11]

$$\psi_{n=0, s=-1, p_y=p_z=0} = \frac{1}{\sqrt{N}} \left(\frac{|e|B}{\pi} \right)^{1/4} e^{-\frac{|e|B}{2} x^2} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad N \stackrel{[11]}{=} L_y L_z. \quad (8)$$

According to (6), in order to determine δm for the (on mass-shell) LLL, we shall sandwich the general self-energy operator (1) between two states $|\psi\rangle$ defined in (8) and satisfying $(\gamma\pi + m)|\psi\rangle = 0$.

The expression (1) involves \not{A} that we shall replace by $-m$, Δ that needs not be transformed, and Φ which involves $m^2 - (\gamma\pi)^2, \pi_\perp^2$ and $\sigma_{\mu\nu} F^{\mu\nu}$. The only non-vanishing component of $F^{\mu\nu}$ being $F^{12} = B, \sigma_{\mu\nu} F^{\mu\nu} = \sigma_{12} F^{12} + \sigma_{21} F^{21} = 2\sigma_{12} F^{12} \equiv 2\sigma_3 B$. Since the electron is an eigenstate of the Dirac equation in the presence of $B, m^2 - (\gamma\pi)^2$ can be taken to vanish. $\pi_\perp^2 \equiv \pi_1^2 + \pi_2^2$ is also identical, since the privileged state has $p_z = 0$ and we work in a gauge with $A_z = 0$, to $\bar{\pi}^2 \equiv \pi^2 + \pi_0^2$. One has $(\gamma\pi)^2 = -\pi^2 + \frac{e}{2} \sigma_{\mu\nu} F^{\mu\nu}$ such that $\pi_\perp^2 = -(\gamma\pi)^2 + \pi_0^2 + \sigma_3 eB$. Since our gauge for the external B has $A_0 = 0, \pi_0^2 = p_0^2$, which is the energy squared of the electron, identical to m^2 for the privileged state. Therefore, on mass-shell, $\pi_\perp^2 = \sigma_3 eB$, with [3] $\sigma^{12} = \sigma^3 = \frac{i}{2} [\gamma^1, \gamma^2] = \text{diag}(1, -1, 1, -1)$. When sandwiched between privileged states,

$$\langle \psi | \sigma^3 | \psi \rangle = \begin{pmatrix} 0 & 1 & 0 & 0 \end{pmatrix} \text{diag}(1, -1, 1, -1) \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} = -1 \text{ such that } \sigma^3 \text{ can be replaced by } (-1) \text{ and } \Phi(u, Y)$$

shrinks to $u eB \left(1 - \frac{\beta}{Y}\right)$. σ^3 can also be replaced by (-1) in the exponentials of (1).

$\Sigma(\pi)$ in (1) also involves a term proportional to \not{k}_\perp . Since the privileged state has $p_z = 0$ and we work at $A_z = 0$, this is also equal to $\vec{\gamma} \cdot \vec{\pi} = \gamma^\mu \pi_\mu - \gamma^0 \pi_0 = \gamma\pi + \gamma^0 p^0$. $\langle \psi | \gamma\pi | \psi \rangle = -m$ such that $\langle \psi | \not{k}_\perp | \psi \rangle = \langle \psi | -m + \gamma^0 p^0 | \psi \rangle$. $\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and using (8) yields therefore $\langle \psi | \not{k}_\perp | \psi \rangle = -m + p^0$. The energy p^0 of the privileged state $|\psi\rangle$ being equal to m , this term vanishes.

Gathering all information and simplifications leads finally to

$$\delta m_{LLL} \equiv \Sigma(\not{k} + m = 0) \stackrel{LLL}{=} \frac{\alpha m}{2\pi} \int_0^\infty \frac{ds}{s} \int_0^1 du e^{-isu^2 m^2} \left[\frac{e^{-is\Phi(u,Y)}}{\sqrt{\Delta(u,Y)}} (1 + ue^{2iY}) - \underbrace{(1+u)}_{c\text{-term}} \right],$$

$$Y = eBsu,$$

$$\Phi(u, Y) = ueB \left(1 - \frac{\beta(u, Y)}{Y} \right) = ueB - \frac{\beta(u, Y)}{s}, \quad (9)$$

$$\Delta(u, Y) = (1-u)^2 + 2u(1-u) \frac{\sin Y \cos Y}{Y} + u^2 \left(\frac{\sin Y}{Y} \right)^2,$$

$$\sin \beta(u, Y) \stackrel{p.49 \text{ of [3]}}{=} \frac{(1-u) \sin Y}{\sqrt{\Delta(u, Y)}}, \quad \cos \beta(u, Y) \stackrel{(3.31) \text{ of [3]}}{=} \frac{(1-u) \cos Y + u \frac{\sin Y}{Y}}{\sqrt{\Delta(u, Y)}}.$$

or, equivalently

$$\delta m_{LLL} \equiv \Sigma(\not{k} + m = 0) \stackrel{LLL}{=} \frac{\alpha m}{2\pi} \int_0^\infty \frac{ds}{s} \int_0^1 du e^{-isu^2 m^2} \left[\frac{e^{i[-sueB + \beta(u, Y)]} + ue^{i[sueB + \beta(u, Y)]}}{\sqrt{\Delta(u, Y)}} - \underbrace{(1+u)}_{c\text{-term}} \right] \quad (10)$$

which is the expression that we have to evaluate.

2.4 A few remarks

* At $B \rightarrow 0, Y \rightarrow 0, \beta \sim (1-u)Y + \mathcal{O}(Y^2)$ yields $\Phi(B=0) = 0$. One also has $\Delta(B=0) = 1$ such that $\Sigma(\pi)_{B=0} = \frac{\alpha m}{2\pi} \int_0^\infty \frac{ds}{s} \int_0^1 du e^{-isu^2 m^2} [(1+u) - (1+u)] = 0$. This agrees with the renormalization condition (3).

* Δ , which occurs by its square root, is a seemingly naughty denominator. Its zeroes u_\pm can be written $u_+ = u_-^* = \frac{1 - \frac{\sin Y}{Y} e^{iY}}{\xi(Y)}$, with $\xi(Y) = 1 - 2 \frac{\sin Y \cos Y}{Y} + \left(\frac{\sin Y}{Y} \right)^2$. The real zeroes $u_+ = 1 = u_-$ are degenerate and are located at $Y = n\pi, n \neq 0$, values at which $\beta = 0$. Fortunately, as mentioned in subsection 2.1, $u = 0$ and $u = 1$ should be cut off the u -interval of integration.

* The renormalized δm given by (9) is finite. The contribution $\propto (1+u)$ from the counterterm is tailored for this.

* The (infinite) counterterm does not *seem* to depend on B ¹⁶.

2.5 Changing variables; the Demeur-Jancovici integral [4] [5]

We first perform the change of variables¹⁷

$$(u, s) \rightarrow (u, Y \equiv eBsu) \Rightarrow \frac{du ds}{s} = \frac{du dY}{Y}. \quad (11)$$

In Dittrich-Reuter [3], e stands for the (negative) charge of the electron¹⁸. Therefore, $Y < 0$, too, and $\int_0^\infty \frac{ds}{s} = \int_0^{-\infty} \frac{dY}{Y}$. Then, δm in (10) becomes

$$\delta m_{LLL} = \frac{\alpha m}{2\pi} \int_0^{-\infty} \frac{dY}{Y} \int_0^1 du e^{-iuY \frac{m^2}{eB}} \left[\frac{e^{i[\beta(u, Y) - Y]} + ue^{i[\beta(u, Y) + Y]}}{\sqrt{\Delta(u, Y)}} - \underbrace{(1+u)}_{\text{from } c\text{-term}} \right], \quad (12)$$

¹⁶It is evaluated in pp. 53-56 of [3]: $\delta m_{B=0} = \lim_{s_0 \rightarrow 0} \frac{3\alpha m}{4\pi} \left(-\gamma_E + \ln \frac{1}{im^2 s_0} + \frac{5}{6} \right)$, where s_0 is the lower limit of integration for the Schwinger parameter s_1 attached to the electron propagator. It coincides with the result given by Ritus in [13].

¹⁷It is legitimate to integrate Y up to ∞ because $u = 0$ is cut off the u -interval of integration (see subsection 2.1).

¹⁸unlike in [1] in which, like in Schwinger, both q and e are introduced. In there, e has the meaning of the elementary charge $e > 0$.

which is seen to only depend on $\frac{eB}{m^2}$. The divergence of δm occurs now at $Y \rightarrow 0$. The change (11) introduces a dependence of the (infinite, therefore undefined¹⁹) counterterm on $\frac{|e|B}{m^2}$.

To summarize in a symbolic way, this change of variables amounts to rewriting $\delta m_{LLL} \equiv \left(\infty + \eta\left(\frac{|e|B}{m^2}\right)\right) - \infty = \left(\infty + \eta\left(\frac{|e|B}{m^2}\right) + \zeta\left(\frac{|e|B}{m^2}\right)\right) - \left(\infty + \zeta\left(\frac{|e|B}{m^2}\right)\right)$. Then, we shall get rid of infinities via a regularization and calculate $\eta + \zeta$ and $-\zeta$ which give respectively the $\left(\ln \frac{|e|B}{m^2}\right)^2$ and $\ln \frac{|e|B}{m^2}$ terms.

It is interesting to expand the sole $e^{i\beta}$ into $\cos \beta + i \sin \beta$, to use the expressions (9) of $\cos \beta$ and $\sin \beta$, to cast δm in the form

$$\delta m_{LLL} = \frac{\alpha m}{2\pi} \int_0^{-\infty} \frac{dY}{Y} \int_0^1 du e^{-iuY \frac{m^2}{eB}} \left[(1 + u e^{2iY}) \frac{1 - u + u \frac{\sin Y}{Y} e^{-iY}}{\Delta(u, Y)} - (1 + u) \right] \quad (13)$$

and to notice that $\Delta(u, Y) = (1 - u + u \frac{\sin Y}{Y} e^{+iY})(1 - u + u \frac{\sin Y}{Y} e^{-iY})$ to simplify the previous expression into

$$\delta m_{LLL} = \frac{\alpha m}{2\pi} \int_0^{-\infty} \frac{dY}{Y} \int_0^1 du e^{-iuY \frac{m^2}{eB}} \left[\frac{1 + u e^{2iY}}{1 - u + u \frac{\sin Y}{Y} e^{+iY}} - (1 + u) \right]. \quad (14)$$

Expressing $\sin Y$ in the denominator in terms of complex exponentials gives

$$\delta m_{LLL} = \frac{\alpha m}{2\pi} \int_0^{-\infty} dY \int_0^1 du e^{-iuY \frac{m^2}{eB}} \left[\frac{2i(1 + u e^{2iY})}{2iY(1 - u) + u(e^{2iY} - 1)} - \frac{1 + u}{Y} \right]. \quad (15)$$

Going to $t = -iY$ yields

$$\delta m_{LLL} = \frac{\alpha m}{2\pi} \int_0^{+i\infty} dt \int_0^1 du e^{ut \frac{m^2}{eB}} \left[\frac{2(1 + u e^{-2t})}{2t(1 - u) + u(1 - e^{-2t})} - \frac{1 + u}{t} \right]. \quad (16)$$

Last, we change to $z = ut \Rightarrow du dt = \frac{du dz}{u}$ and get

$$\begin{aligned} \delta m_{LLL} &= \frac{\alpha m}{2\pi} \int_0^{+i\infty} dz \int_0^1 du e^{z \frac{m^2}{eB}} \left[\frac{2(1 + u e^{-2z/u})}{2z(1 - u) + u^2(1 - e^{-2z/u})} - \frac{1 + u}{z} \right] \\ &= \frac{\alpha m}{2\pi} \int_0^{+i\infty} dz \int_0^1 du e^{-z \frac{m^2}{|e|B}} \left[\frac{2(1 + u e^{-2z/u})}{2z(1 - u) + u^2(1 - e^{-2z/u})} - \frac{1 + u}{z} \right] \end{aligned} \quad (17)$$

which still differs from eq. 3 of Jancovici [5] by the 2 following points:

* that we have $e^{+z \frac{m^2}{eB}}$ instead of $e^{-z \frac{m^2}{eB}}$ is due to $e > 0$ in [5], while, here, $e < 0$;

* we have $\int_0^{i\infty} dt$ instead of $\int_0^{\infty} dt$. $\int_0^{+i\infty} + \int_{1/4 \text{ infinite circle}} + \int_{\infty}^0 = 2i\pi \sum \text{residues}$. Because of $e^{-z \frac{m^2}{|e|B}}$ the contribution on the infinite 1/4 circle is vanishing. That the residue at $z = 0$ vanishes is easily seen because, as already mentioned (subsection 2.1), u must never come down strictly to 0. Other poles (now considering (16)) can only occur at $2t(1 - u) + u(1 - e^{-2t}) = 0 \Leftrightarrow u = \frac{2t}{2t + e^{-2t} - 1}$. That u must be real requires $t \in \mathbb{R}$ or $t = n i\pi$. For $t \in \mathbb{R}$, $u = \frac{2t}{2t + e^{-2t} - 1} > 1$ (except at $t = 0$ but we have seen that there is no problem at $t = 0$), thus outside the domain of integration for u . The most delicate poles occur at $t = n i\pi$, in which case $u_{pole} = 1$ is indeed real but at the border of the u -domain of integration. Fortunately we do not have to take them into account since $u = 1$ should be excluded, too (subsection 2.1). We can therefore safely make the Wick rotation, which yields

$$\delta m_{LLL} = \frac{\alpha m}{4\pi} 2 \int_0^{\infty} dz \int_0^1 du e^{-z \frac{m^2}{|e|B}} \left[\frac{2(1 + u e^{-2z/u})}{2z(1 - u) + u^2(1 - e^{-2z/u})} - \underbrace{\frac{1 + u}{z}}_{\text{from } c\text{-term}} \right]. \quad (18)$$

It is now the same as Jancovici's eq. 3 [5] (see eqs. (20,21) below). This proves in particular that the latter (and therefore Demeur's calculation [4]) satisfy the same renormalization conditions (3), which was not evident in [4].

¹⁹See footnote 16.

3 Calculating Jancovici's integral [5]

3.1 Generalities and definition

Along with Jancovici [5], let us write the rest energy of the electron

$$E_0 = m(c^2) \left(1 + \frac{\alpha}{4\pi} I(L) \right), \quad L = \frac{(\hbar)|e|B}{(c^3)m^2} \quad (19)$$

in which, at all orders in B

$$I(L) = 2 \int_0^\infty dz e^{-z/L} \int_0^1 dv \left(\frac{2(1+v)e^{-2z/v}}{2z(1-v) + v^2(1-e^{-2z/v})} - \frac{1+v}{z} \right) = 2 \int_0^\infty dz e^{-z/L} \int_0^1 dv f(v, z), \quad (20)$$

$$f(v, z) = \frac{2(1+v)e^{-2z/v}}{2z(1-v) + v^2(1-e^{-2z/v})} - \frac{1+v}{z}.$$

Jancovici [5] defines accordingly (we set hereafter $\hbar = 1 = c$)

$$\delta m = \frac{\alpha m}{4\pi} I(L). \quad (21)$$

$I(L)$ has been obtained from Demeur's original integral [4]^{20 21}

$$D(L) = \int_0^1 dv (1+v) \int \frac{dw}{w} \frac{w}{|w|} e^{i v w} \frac{2iLw(v e^{2iLw} + 1)}{(1+v)[v e^{2iLw} + 2iLw(1-v) - v]} \quad (22)$$

after subtracting its value at $B = 0 \Leftrightarrow L = 0$ and after the change of variables $z = -iLv w$. Therefore, (21) corresponds to the magnetic radiative corrections to the electron mass, after subtracting the self-energy of the “free” (i.e. at $B = 0$) electron²². The latter corresponds to the term $\propto \frac{1+v}{z}$ in the integrand of (20). Accordingly, (21) satisfies $\delta m \xrightarrow{B \rightarrow 0} 0$. Demeur's calculation concerns what he calls, after Luttinger [15], the “privileged state” of the electron which has energy m .

We want an analytical expression for $I(L)$ valid for large values of the magnetic field, say $\frac{|e|B}{m^2} > 75$. That $I(L)$ can easily be integrated numerically makes checks easy.

3.2 First steps: a simple convergent approximation for $L \equiv eB/m^2 > 75$

The 2 integrals in (20) both diverge at $z \rightarrow 0$. The cancellation of the divergences is ensured by the first renormalization condition (3), but its practical implementation needs a kind of regularization. Following Jancovici [5], one splits $I(L)$ into $\int_0^\infty dz = \int_0^a dz + \int_a^\infty dz$, with a large enough such that $e^{-2z/v} \ll 1$ can be neglected inside $f(v, z)$. Since $v \in [0, 1]$, this requires at least $a \geq 1$, that we check numerically. $I(L)$ can then be approximated by

$$I(L) \approx 2 \int_0^a dz e^{-z/L} \int_0^1 dv f(v, z) + 2 \int_a^\infty dz e^{-z/L} \int_0^1 dv \left(\frac{2}{v^2 + 2z(1-v)} - \frac{1+v}{z} \right), \quad (23)$$

in which the second integral is manifestly convergent. We focus on the first one, which includes the two canceling divergences. It turns out, as in [5], that, for L large enough, for example $L > 75$, its numerical value decreases with a and that one can go very safely down to $a = 1$ at which it is totally negligible with respect to the value of the full I ²³.

²⁰It is eq. (21) of § 8: “La self-énergie de l'électron”, p. 78 of [4].

²¹It has been manifestly obtained with an internal photon in the Feynman gauge (see eq. (1) p. 56 of [4]).

²²See Demeur [4] chapitre III “Les corrections radiatives magnétiques”, § 1 “La self-énergie”, p.55

²³We proceed as follows. Though $f(0, z) = \frac{1}{z} - \frac{1}{z} = 0$, $f(v, z)$ cannot be integrated $\int_0^1 dv$ at small z because its expansion $f(v, z) \xrightarrow{z \rightarrow 0} -(1-v) + z(-\frac{5}{3} + \frac{4}{3v} + v) + z^2(-\frac{7}{3} - \frac{1}{v^2} + \frac{7}{3v} + v) + \dots$ has (fake) poles at $v = 0$. Even numerical integration becomes hazardous. To achieve it safely, we regularize the first integral in (23) by introducing a small parameter ϵ , replace $\int_0^1 dv f(v, z)$ with $\int_\epsilon^1 dv f(v, z)$, then decrease $\epsilon = 10^{-3}, 10^{-6}, 10^{-9} \dots$ while checking stability.

We thus approximate, for $L \geq 75$ ²⁴

$$I(L) \stackrel{L \geq 75}{\approx} 2 \int_{a=1}^{\infty} dz e^{-z/L} \int_0^1 dv \left(\frac{2}{v^2 + 2z(1-v)} - \frac{1+v}{z} \right). \quad (24)$$

The second contribution to (24), which comes from the counterterm, is easily integrated, and one gets

$$I(L) \stackrel{L \geq 75}{\approx} 2 \underbrace{\int_1^{\infty} dz e^{-z/L} \int_0^1 dv \frac{2}{v^2 + 2z(1-v)}}_{J(L)} - 3 \Gamma(0, 1/L) = 2 J(L) - 3 \Gamma(0, 1/L) \quad (25)$$

in which $\Gamma(0, z)$ is the incomplete Gamma function $\Gamma(0, z) = \int_z^{\infty} \frac{e^{-t}}{t} dt$. $\int_0^1 dv \frac{2}{v^2 + 2z(1-v)}$ can be easily integrated, too, leading to

$$I(L) \stackrel{L \geq 75}{\approx} 2 \underbrace{\int_1^{\infty} dz e^{-z/L} \frac{\ln \left(z - 1 + \sqrt{z(z-2)} \right)}{\sqrt{z(z-2)}}}_{J(L)} - \underbrace{3 \Gamma(0, 1/L)}_{\text{from } c\text{-term}}. \quad (26)$$

The result of the change of variables done in subsection 2.5 associated with the regularization-approximation just performed is a sum of 2 finite integrals. The most peculiar and also the most important for our purposes is the second one which originates from the counter-term and includes the large $\ln \frac{|e|B}{m^2}$ generally ignored. The occurrence of a dependence on $\frac{|e|B}{m^2}$ in the counterterm is non-trivial since, naively, before regularization, it looks (infinite and) independent of it (see footnote 16).

3.3 Further evaluation

$J(L) \equiv \int_1^{\infty} dz g(z)$, $g(z) = \frac{\ln(z-1+\sqrt{z(z-2)})}{\sqrt{z(z-2)}}$ cannot be integrated exactly but, following [16], one can find an accurate approximation for the integrand

$$g_{app}(z) \approx \frac{\ln z}{z} + \frac{\pi}{2} \frac{1}{z^\beta}, \quad \beta = 1.175 \quad (27)$$

as shown on Fig. 2 below where the 2 curves for the exact g (blue) and the approximate g_{app} (yellow) are practically indistinguishable.




Fig. 2: exact (blue) and approximate (yellow) values for the integrand $g(z)$ of $J(L)$.

Without the $\frac{\pi}{2} \frac{1}{z^\beta}$, g would go to 0 instead of $\frac{\pi}{2}$ at $z = 1$. This term yields in particular the term $\propto \frac{1}{L^{\beta-1}}$ in the expansion of J_{app} at $L \rightarrow \infty$. The integral can now be done analytically, leading to

$$J_{app}(L) = \int_1^{\infty} dz e^{-z/L} \left(\frac{\ln z}{z} + \frac{\pi}{2} \frac{1}{z^\beta} \right) = \underbrace{\frac{\pi}{2} \text{ExpIntegralE}[\beta, \frac{1}{L}]}_{\text{from } \frac{\pi}{2} \frac{1}{z^\beta}} + \underbrace{\text{MeijerG}[\{((), (1, 1)\}, \{(0, 0, 0), (,)\}, \frac{1}{L}]}_{\text{from } \frac{\ln z}{z}}. \quad (28)$$

We compare in Fig. 3 the integrals $J(L)$ (blue) and $J_{app}(L)$ (yellow), which prove extremely close.

²⁴The term $\frac{1+v}{z}$ was neglected in eq. (4) of [5], where only \ln^2 are focused on.




Fig. 3: exact (blue) and approximate (yellow) values for $J(L)$.

3.4 Final result

The final result is obtained by expanding $J_{app}(L)$ and $\Gamma(0, 1/L)$ at large L

$$\begin{aligned}
J_{app} \stackrel{L \rightarrow \infty}{\simeq} \frac{1}{L^\beta} \left(\frac{\pi}{2} L \Gamma[1 - \beta] + \mathcal{O}\left(\frac{1}{L^2}\right) \right) + \frac{\gamma_E^2}{2} + \frac{\pi}{12} \left(\frac{6}{\beta - 1} + \pi \right) - \frac{1}{2} \ln L (2\gamma_E - \ln L) + \frac{-1 + \frac{\pi}{4-2\beta}}{L} + \mathcal{O}\left(\frac{1}{L^2}\right), \\
\Gamma(0, 1/L) \stackrel{L \rightarrow \infty}{\simeq} -\gamma_E + \ln L + \frac{1}{L} + \mathcal{O}\left(\frac{1}{L^2}\right) \quad (\text{comes from the counterterm})
\end{aligned} \tag{29}$$

which yields for $I(L)$

$$\begin{aligned}
I_{app}(L, \beta) \stackrel{L \geq 75}{\approx} \gamma_E^2 \underbrace{+3\gamma_E}_{\text{from c-term}} + \frac{\pi}{\beta - 1} + \frac{\pi^2}{6} + \frac{\pi \Gamma[1 - \beta]}{L^{\beta-1}} - \ln L \left(2\gamma_E + \underbrace{3}_{\text{from c-term}} \right) + (\ln L)^2 \\
+ \frac{1}{L} \left(\frac{\pi}{2 - \beta} - 2 \underbrace{-3}_{\text{from c-term}} \right) + \mathcal{O}\left(\frac{1}{L^2}\right) \\
= \left(\ln L - \gamma_E - \frac{3}{2} \right)^2 - \frac{9}{4} + \frac{\pi}{\beta - 1} + \frac{\pi^2}{6} + \frac{\pi \Gamma[1 - \beta]}{L^{\beta-1}} + \frac{1}{L} \left(\frac{\pi}{2 - \beta} - 5 \right) + \mathcal{O}\left(\frac{1}{L^2}\right).
\end{aligned} \tag{30}$$

The terms under-braced “from c-term” result from the subtraction of the electron self-energy at $B = 0$; they include a large $-3(\ln L - \gamma_E)$, which therefore originates from the counterterm (together with part of the constant term in δm).

At $L \geq 75$ the term $\propto 1/L$ can be very safely neglected and one can approximate

$$I_{app}(L, \beta) \stackrel{L \geq 75}{\approx} \left(\ln L - \gamma_E - \frac{3}{2} \right)^2 - \frac{9}{4} + \frac{\pi}{\beta - 1} + \frac{\pi^2}{6} + \frac{\pi \Gamma[1 - \beta]}{L^{\beta-1}} + \mathcal{O}\left(\frac{1}{L}\right), \quad \beta \approx 1.175 \tag{31}$$

which is very different, as we shall see, from the brutal approximation $I_{app} \approx (\ln L)^2$ that has been systematically used in the following years. At $\beta = 1.175$, one gets explicitly

$$I_{app}(L, \beta = 1.175) \stackrel{L \geq 75}{\approx} 21.6617 - \frac{20.4164}{L^{0.175}} + \ln L (\ln L - 4.15443) + \mathcal{O}\left(\frac{1}{L^{\geq 1}}\right). \tag{32}$$

We plot in Fig. 4 the different contributions to the Demeur-Jancovici integral: the yellow curve is the constant term, the green one is the inverse power, the red one the \ln contribution, the violet one the $(\ln)^2$, and the blue curve is the global result. The comparison between the violet and blue curve is that between the systematically used $(\ln)^2$ approximation and our accurate evaluation (31). A large cancellation between $(\ln)^2$ and \ln terms²⁵ makes in particular the role of the large constant important.

²⁵They exactly cancel at $\ln L \approx 4.15443 \Leftrightarrow B \approx 63 B_0$, where $B_0 \equiv \frac{m^2}{|e|}$ is the “Schwinger critical field”.




Fig. 4: contributions to the Demeur-Jancovici integral; constant term (yellow), inverse power (green), \ln (red), \ln^2 (violet), sum of all (blue).

One needs $L > 2 \cdot 10^{17}$ for the relative error $\left| \frac{I_{app}(L) - (\ln L)^2}{I_{app}(L)} \right| < \frac{1}{10}$, which is a totally unrealistic value of the magnetic field.

Jancovici mentioned at the end of his work [5] a refined estimate $I(L) \simeq (\ln 2L - \gamma_E - \frac{3}{2})^2 + A$ with $-6 \leq A \leq +7$. Actually, the value $A = 3.5$ yields a good agreement with our calculation in the range $75 \leq L \leq 100\,000$, as shown in Figs. 5.




Fig. 5: comparison between the present calculation of $I(L)$ (blue) and Jancovici's final refined estimate with $A = 3.5$, $I(L) \simeq (\ln 2L - \gamma_E - \frac{3}{2})^2 + 3.5$ (yellow).

The inverse power term, only present in our calculation, only plays a substantial role at the lowest values of L (see the green curve on Fig. 4); at very large L , a fit “à la Jancovici” with only $(\ln)^2$, \ln and cst terms is in practice enough. This explains the slight disagreement between the two curves of Fig. 5, which cannot perfectly match in the whole range of L that we are considering.

3.5 Concluding remarks and two challenges

In view of these results, we conclude that one cannot reasonably approximate the integral of Demeur-Jancovici (nor the corresponding δm of the electron) by the sole term proportional to $(\ln \frac{|e|B}{m^2})^2$; at least, the large single $\ln \frac{|e|B}{m^2}$ (with opposite sign) and the large constant should be included in all estimates.

Renormalization conditions are of the uttermost importance. We have seen that, at order α , if one forgets about the counterterm, one skips, among others, the large $\ln \frac{|e|B}{m^2}$, which leads to erroneous results. Counterterms have to be determined order by order in the perturbative series. This constitutes the first challenge when operating a resummation, which becomes necessary at very large values of $\ln \frac{|e|B}{m^2}$ or for theories more strongly coupled than standard QED. We have no proof that this has been achieved yet. The second challenge concerns extending the present calculation to electrons lying in higher Landau levels. Both tasks look very non-trivial.

Though it is premature to make any prospect, the sharp damping of δm that we have found with respect to previous approximations nevertheless suggests that physical consequences should also be substantially weakened. This is left for later investigations.

References

- [1] W.-y. TSAI: “Modified electron propagation function in strong magnetic fields”, *Phys. Rev. D* 10 (1974) 1342.
- [2] J. SCHWINGER: “Quantum Electrodynamics. II. Vacuum Polarization and Self-Energy”, *Phys. Rev.* 75 (1949) 651.
- [3] W. DITTRICH, M. REUTER: “Effective Lagrangians in Quantum Electrodynamics”, Springer-Verlag, *Lecture Notes in Physics* 220 (1985).
- [4] M. DEMEUR: “Étude de l’interaction entre le champ propre d’une particule et un champ electro-magnétique homogène et constant”, *Acad. Roy. Belg., Classe Sci., Mem.* 28, No 1643 (1953-55) 651 C.
- [5] B. JANCOVICI: “Radiative Correction to the Ground-State Energy of an Electron in an Intense Magnetic Field”, *Phys. Rev.* 187 (1969) 2275.
- [6] Yu.M. LOSKUTOV, B.B. SKOBELEV: “Radiative corrections to the electron mass operator in the two-dimensional approximation of quantum electrodynamics”, *Teor. Mat. Fiz.* 38 (1979) 195-200.
- [7] Yu.M. LOSKUTOV, B.B. SKOBELEV: “Behavior of the mass operator in a superstrong magnetic field: summation of the perturbation theory diagrams”, *Teor. Mat. Fiz.* 48 (1981) 44-48.
- [8] Yu.M. LOSKUTOV, B.B. SKOBELEV: “Massovii operator : adnolarifmitcheskaja polevaja asimptotika”, *Vestn. Mosk. Un-Ta. Ser. 3 Fisica Atronomia*, 1983, T.24 No 6, p.95.
- [9] V.P. GUSYNIN, A.V. SMILGA: “Electron self-energy in strong magnetic field: summation of double logarithmic terms”, *Phys. Lett. B* 450 (1999) 267-274.
- [10] A.V. KUZNETSOV, N.V. MIKHEEV & M.V. OSIPOV: “Electron mass operator in a strong magnetic field”, *Mod. Phys. Lett. A* 17 (2002) 231-235.
- [11] A. KUZNETSOV, N. MIKHEEV: “Electroweak Processes in External Electromagnetic Fields”, *Springer Tracts in Modern Physics* 197 (2004).
- [12] V.A. MIRANSKY, I.A. SHOVKOVY: “Quantum field theory in a magnetic field; from quantum chromodynamics to graphene and Dirac semimetals”, *arXiv:1503.00732*, *Phys. Rept.* 576 (2015) 1-209.
- [13] V.I. RITUS: “Lagrangian of an intense electromagnetic field and quantum electrodynamics at short distances”, *Zh. Ekps. Teor. Fiz.* 69, 1517-1535 (1975), *Sov. Phys. JETP* 42, 774 (1976)
- [14] R.G. NEWTON: “Radiative Effects in a Constant Field”, *Phys. Rev.* 96 (1954) 523.
- [15] J.M. LUTTINGER: “A Note on the Magnetic Moment of the Electron”, *Phys. Rev.* 74 (1948) 893.
- [16] M.I. VYSOTSKY: “Atomic levels in superstrong magnetic fields and $D = 2$ QED of massive electrons: screening”, *Pis'ma v ZhETF* 92 (2010) 22-26.
- [17] V.B. BERETSETSKII, E.M. LIFSHITZ, & L.P. PITAYEVSKY, “Quantum Electrodynamics”, *Theoretical Physics Vol. IV (Fizmatlit, Moscow, 2001)* [“Quantum Electrodynamics”, *Course of Theoretical Physics Vol. 4*, translated from russian by J.B. Sykes and J.S. Bell (Pergamon, New York, 1982)].