



HAL
open science

Frasnian (Late Devonian) conodonts and environment at the northern margin of the Algerian Sahara platform: the Ben Zireg section

Abdessamed Mahboubi, Raimund Feist, Jean-Jacques Cornee, Abdelkader Ouali Mehadji, Catherine Girard

► To cite this version:

Abdessamed Mahboubi, Raimund Feist, Jean-Jacques Cornee, Abdelkader Ouali Mehadji, Catherine Girard. Frasnian (Late Devonian) conodonts and environment at the northern margin of the Algerian Sahara platform: the Ben Zireg section. *Geological Magazine*, 2015, 152 (5), pp.844-857. 10.1017/S0016756814000715 . hal-01217151

HAL Id: hal-01217151

<https://hal.science/hal-01217151>

Submitted on 30 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3 **Frasnian (Late Devonian) conodonts and environment at the northern margin**
4
5
6 **of the Algerian Sahara platform: the Ben Zireg section**
7
8
9

10
11 **Category: Stratigraphy**
12
13

14
15 Abdessamed Mahboubi¹, Raimund Feist¹, Jean-Jacques Cornée², Abdelkader Ouali Mehadji³,
16
17
18 Catherine Girard¹
19
20

21
22 ¹ UMR 5554 CNRS, Université Montpellier, Place Eugène Bataillon, Montpellier (France)
23
24

25 ² UMR 5243 CNRS, Université Montpellier 2, Geosciences Montpellier, Place Eugène
26
27 Bataillon, 34095 Montpellier Cedex 05 (France)
28
29

30 ³ Département des Sciences de la Terre, Université d'Oran, Es Sénia, Oran (Algérie)
31
32
33

34
35 Corresponding author: Catherine Girard (catherine.girard@univ-montp2.fr)
36
37
38

39
40 **Short title: Frasnian biostratigraphy at Ben Zireg (Algeria)**
41
42
43

44 **Abstract**
45

46 The Ben Zireg anticline NW of Bechar yields the so far most important Frasnian succession
47
48 on the northern margin of the Algerian Sahara. It represents 26.5 m of calcilutites which are
49
50 attributed to the middle through late Frasnian. As the early Frasnian is not represented, the
51
52 succession rests conformably on an indifferentiated, probably late Givetian substrate. Fine-
53
54 scaled conodont biostratigraphy reveals a continuous sequence of Montagne Noire Zones 5
55
56
57
58
59
60

1
2
3 to 13, superseded by the earliest Famennian *triangularis* Zone. The deposits are organized
4
5 into 15 sedimentary cycles that are initially condensed and become dilated upwards.
6
7 Conodont biofacies indicate an overall deepening towards the top of the section with an
8
9 episode of slight shallowing at the MN11/12 transition marked by an increase of
10
11 Ancyrodellids which is also seen in the Marhouma section of the Ougarta region (SW
12
13 Algeria). Homogeneous calcareous microfacies with only a few shaly intercalations
14
15 determine a predominantly oxygenated depositional environment on an outer platform,
16
17 submarine rise or ramp setting. This is also indicated by relatively low sedimentation rates
18
19 which are similar in selected sections from the Tafilalet platform, and contrast with those
20
21 from the Marhouma trough and the Maider basin. Unlike other Frasnian successions S of the
22
23 Atlas fault, but similar to the Moroccan Meseta, the Upper Kellwasser horizon is clearly
24
25 discriminated at Ben Zireg by an outstanding occurrence of black shales on top of the
26
27 oxygenated Latest Frasnian deposits.
28
29
30
31
32
33
34
35

36 Key-words: Frasnian, conodont biostratigraphy, biofacies, depositional environment,
37
38 northwestern Algerian Sahara
39
40
41
42

43 **1. Introduction**

44
45
46
47

48 The Ben Zireg area at 50 km NE of Bechar township constitutes the north-eastern most
49
50 exposure of Saharan Palaeozoic deposits adjacent to the Main Atlas fault. They differ from
51
52 all neighbouring outcrops of the Bechar basin such as Soltane el Betoum and Maider el
53
54 Mhadjib to the N and W of Djebel Antar (Weyant, 1988) (Fig. 1a) in comprising a well
55
56 exposed, almost complete succession of marine sediments from Cambrian-Ordovician
57
58
59
60

1
2
3 through Early Carboniferous. As such the Ben Zireg section is a reference for the northern
4
5 margin of epicontinental Sahara in SW Algeria, at 200 km in distance from the Palaeozoic of
6
7 the Ougarta region further to the South (Fig. 1a). A major contribution to its structure and
8
9 stratigraphy by Pareyn (1961) was mainly devoted to the Carboniferous, whereas apart from
10
11 a few scattered data, the Devonian, and Frasnian in particular, remained largely unstudied
12
13 regards biostratigraphical subdivision and sedimentary development.
14
15

16
17 After the first recognition of Devonian fossils at Ben Zireg by Flamand (1911), Menchikoff
18
19 (1936) established the existence of Late Devonian series based on cephalopods. Among
20
21 them *Pharciceras* was considered at that time representative of the early Frasnian. This
22
23 taxon, however, is now restricted to the Latest Givetian since the definition of Frasnian stage
24
25 boundaries by SDS/IUGS (Klapper, Feist & House, 1987, 1993). The presence of Frasnian
26
27 strata was definitely established when Massa et al. (1965) recovered the first conodonts,
28
29 notably *Ancyrognathus asymmetricus* and *Ancyrodella curvata*.
30
31
32

33
34 Since then, and in contrast to numerous conodont-based biostratigraphical investigations in
35
36 the neighbouring Tafilalet area (e.g. Belka et al., 1999), no more work was carried out to
37
38 evaluate the extent and completeness of the early Upper Devonian in the Algerian part of
39
40 northern Sahara.
41
42

43
44 This study aims establishing fine-scaled conodont biostratigraphy and biofacies as well as
45
46 sedimentary dynamics of the Ben Zireg section to emphasise the importance of its pivotal
47
48 position between Frasnian sections on both sides of the Main Atlas fault, i.e. the
49
50 allochthonous Hercynian Meseta of Central Morocco, and their eastern prolongation into
51
52 the Algerian Hercynian Tifrit Horst, and cratonic Gondwana sections both in SW Algeria
53
54 (Marhouma) and Tafilalet / Maider in SE Morocco (Fig. 1a).
55
56
57
58
59
60

2. Geological Setting

In contrast to Palaeozoic rocks north of the alpine Main Atlas fault which were affected by complex tangential tectonics and deformation during the Variscan orogeny (e.g. Allary et al., 1976), the Palaeozoic deposits of the western Sahara platform north of the Precambrian Reguibat shield (Fig. 1a) remained generally only weakly deformed to wide synclinal and narrower anticlinal structures, where Devonian sedimentation rates reflect basin and ridge depending depositional environments (Elmi & Ameur, 1984; Conrad, Massa & Weyant, 1986; Wendt et al., 2006). Exposures of the north-eastern Bechar basin close to the Atlas fault are more intensely folded to form narrow W-E trending anticlinal structures, often over-thrusted to the N, and these features along with the occurrence of synorogenic wildflysch sediments with olistolithes of various ages characterising the Dinantian of Ben Zireg (Pareyn, 1961) might indicate dynamic relations to the Maghrebien variscan belt in the north.

In this context the Ben Zireg Palaeozoic forms an acute, narrow anticlinal structure with subvertical axial plane, opening to the ENE under 50°, before being unconformably covered by continental Permo-Triassic sediments and marine Liassic and Cenomanian-Turonian deposits (Fig. 1b). The structure is crossed by numerous vertical faults that run sub-parallel to the Main Atlas fault and, affecting also the Cretaceous cover, may belong to the alpine transform system. The eastern end of the structure is framed by a prominent ridge of some 200m of Middle through Late Devonian carbonates easily accessible from the S where the road-truck from Ben Zireg station to Taghit runs along the Oued Ben Zireg shortly before it bends to the S (Fig.1b). The investigated section in the Frasnian is situated on the steep southern flank of the Devonian ridge along a southerly directed, deeply incised gorge (coordinates: N 31° 54' 39.4" W 001° 47' 58.8", Fig. 1b).

3. Material and Methods

One single section was logged in detail and thirty-seven oriented samples were taken both for sedimentological investigations and conodont research. Thin slices from all samples were cut to investigate the corresponding microfacies. Microfacies were determined by using optical observations under photonic microscope. Depositional settings result in zonations proposed by Wright & Burchette (1996) and Flügel (2004). Noticeable is a general recrystallization of micrite into microsparite in all samples, and the presence of pressure-dissolution surfaces indicative of a discrete cleavage. These feature are related to the Variscan deformation and do not allow observations of the primary matrix of the rocks.

For conodont extraction, the calcareous samples were dissolved in 10% acetic acid. The sulphic acid (10%) was used to dissolve iron-and pyrite-rich samples (beds BZ1A and 1C and BZ2). The insoluble residue was then rinsed through two sieves (1mm and 100 μ m). Residue smaller than 1 mm were dried and handpicked under a stereomicroscope. Specimens were photographed with SEM (JEOL 5600). All platform (P1) elements were picked (2660 elements) (Table 1). The limestone samples yielded relatively abundant conodont elements (mean of 115/kg). Some of these elements were identified at the specific level for biostratigraphy. In this paper the biostratigraphic zonations are those of Klapper (1989, 2007a) for the Frasnian and Klapper (2007b) and Schülke (1999) for the lowermost Famennian.

Among the platform P1 elements, different genera can be recognised that are supposed to have different ecological preferences (e.g. Seddon & Sweet, 1971; Sandberg, 1976). The changing distribution patterns of conodont abundance (called biofacies) in space and time can therefore provide some information about the depositional environment. The biofacies is

1
2
3 named after the one or the two most abundant genera, and trends in the variations of the
4
5 relative proportion of the different genera are assumed to provide indirect water depth proxy
6
7 through time.
8

9
10 During the Frasnian period, five conodont genera dominated the assemblages. *Icriodus* is
11
12 considered as characteristic of shallow environments, whereas *Palmatolepis* should prefer
13
14 deeper conditions. *Ancyrodella* and *Ancyrognathus* are grouped under the term
15
16 ancyrodellids. Together with *Polygnathus* they are considered as ubiquitous, with a slight
17
18 preference for deep environments for *Polygnathus* whereas Ancyrodellids would prefer
19
20 shallow ones (Sandberg et al., 1992; Sobstel et al., 2006). The variations of relative
21
22 abundances of these 5 different genera were investigated through the Ben Zireg section. The
23
24 relative abundances of the four groups (*Palmatolepis*, *Icriodus*, *Polygnathus* and
25
26 ancyrodellids) are represented as percentage of the total assemblage.
27
28
29
30
31
32
33

34 **4. Results**

35 *4.a. Lithostratigraphy*

36
37 The Frasnian deposits at Ben Zireg section are 26.5 metres thick, between sample BZ1
38
39 (Givetian) and sample BZ16 (early Famennian) (Fig. 2). Sedimentary rocks display a rhythmic
40
41 pattern characterized by cm to dm-thick alternations (couplets) of hard, fine-grained
42
43 limestones and softer, argillaceous fine-grained limestones or claystones. In the field, these
44
45 alternations clearly display a rather constant organization which indicates sedimentary
46
47 cycles and subcycles (Fig. 3). Sedimentary cycles are limited at top and bottom by thick,
48
49 argillaceous carbonates or dark claystones. Between, twenty couplets are generally found,
50
51 organized into four lithological subcycles, each composed of five couplets. Sedimentary
52
53
54
55
56
57
58
59
60

1
2
3 subcycles are defined by rhythmic facies and bed-thickness changes. In the present-day of
4
5 our knowledge, it is not possible to assign a clear significance to these sedimentary cycles.
6

7
8 The Frasnian deposits comprise 15 sedimentary cycles. From bottom to top, a general
9
10 increase in thickness of sedimentary cycles is observed through the Frasnian deposits: cycles
11
12 1 to 9 and 11 to 12 are 1 to 2m thick, whilst the cycles 10 and 13 to 15 are 2.5 to 3.5 m thick.
13

14
15 The facies are rather homogenous with rare macrofossil remains, but slight differences can
16
17 be depicted, from bottom to top:
18

19
20 1) Above sample BZ1 is a cm-thick iron crust. Between 0.6 and 4m above the base of the
21
22 section, there are interbedded cm to dm thick massive, greyish to brownish micritic
23
24 limestones and cm to dm thick argillaceous limestones. The first beds above the hard-ground
25
26 are cherty and ferruginous (BZ 1b). The massive limestones are tentaculid-rich wackestones.
27
28 The top of some beds can be coated by iron hydroxydes. Limestones display a discrete
29
30 nodular structure and some pyrite crystals.
31
32

33
34 2) Between 4 and 10.7 m above the base of the section are interbedded ferruginous
35
36 limestones and argillaceous limestones (Fig. 4a). Ferruginous limestones display a
37
38 pronounced pseudo-nodular structure underlined by iron films around micritic nodules
39
40 because of pressure-dissolution process of tectonic origin or/and bioturbation (=bioerosion;
41
42 Elmi & Ameur, 1984). Limestones are mudstones to wackestones with tentaculids (Fig. 4d),
43
44 some entomozoan crustaceans and pelagic molluscs ("filaments"). Pyrite crystals are present
45
46 and sparse phosphate grains occur. Most of the beds are coated by thin iron coatings.
47
48

49
50 Argillaceous limestones also display nodular structure and iron films.
51

52
53 3) Between 10.7 and 15.5 m above the base of the section, massive and argillaceous
54
55 limestones are found. These are bioturbated mudstones with rare pelagic organisms
56
57
58
59
60

1
2
3 (tentaculids, entomozoan crustaceans, pelagic molluscs). They also yielded foraminifers in
4
5 residues obtained for conodont extraction. Some cm-thick blackish shales are also
6
7 interbedded.
8
9

10 4) Between 15.5 and 19.5 m above the base of the section are massive, greyish pseudo-
11
12 nodular limestones. These are mudstones with some pelagic organisms (tentaculids,
13
14 entomozoan crustaceans, pelagic molluscs) and some benthic ostracods (Fig. 4e).
15
16

17 5) Between 19.5 and 26.5m above the base of the section deposits are dominant pinkish to
18
19 greyish pseudo-nodular limestones and interbedded argillaceous micrites (Fig. 4b).
20
21

22 Limestones are tentaculid mudstones, with some entomozoan crustaceans, pelagic molluscs
23
24 and goniatites. They also yielded rare benthic ostracods, foraminifers, and rare fragments of
25
26 brachiopods and trilobites; the latter exclusively occur in dish-like flattened mud pebbles
27
28 occurring on the surface of a single bed (Bed 14 d with *Acuticryphops acuticeps*; Fig. 4f). The
29
30 uppermost part of the Frasnian deposits consists of a 30 cm thick black shale level overlain
31
32 by 35 cm bed of plane-laminated, grain-sorted, pinky calcisiltite between samples BZ15c and
33
34 BZ16 (Fig. 4c). The significance of this reddish calcisiltite remains enigmatic. It could have
35
36 resulted in turbidite current activity carrying detrital carbonates and iron oxides of possibly
37
38 of continental origin into the offshore domain.
39
40
41
42
43
44
45

46 *4.b. Biostratigraphy*

47

48
49 Distribution of conodont platform elements is given on Fig. 2, and some biostratigraphic
50
51 species are presented on the Fig. 5. Genera abundance per level is listed on Table 1.
52
53
54
55
56
57
58
59
60

1
2
3 The first level (BZ1) delivered *Polygnathus linguiformis linguiformis* (gamma morphotype), *Po.*
4 *linguiformis mucronatus*, *Po. linguiformis weddigei*, *Po. pseudofoliatus* and *Po. latifossatus*
5
6
7 indicating the Givetian (Aboussalam, 2003).
8
9

10 The following level BZ1A indicates MN Zone 5 due to the occurrence of *Palmatolepis*
11 *punctata*. This level coincides with the base of the Middle Frasnian (Becker & House, 2000a).
12
13 *Polygnathus dubius* and *Po. decorosus* also occur within this zone.
14
15
16

17 The MN Zone 6 is identified with the entry of *Ancyrodella gigas* form 2 in bed BZ2. *Icriodus*
18 *symmetricus*, *Po. decorosus*, *Po. dubius* are also present in this zone.
19
20
21

22 The occurrences of both *An. lobata* and *An. curvata* (early form) in bed BZ4B.1 mark the base
23
24 of the MN Zone 7.
25
26

27 The MN Zone 8 is identified by the co-occurrence of *Pa. hassi* and *Ag. coeni*.
28
29

30 An undifferentiated MN Zone 9-10 is discriminated from bed BZ7 to bed BZ7A by the presence
31
32 of *Pa. ljaschenkoae* together with *Pa. kireevae*. The conodont association is composed by *Pa.*
33 *amplificata* in bed BZ7, *Po. zinaidae* and *Po. aequalis* in bed BZ 7A which have their first
34
35 occurrences during this interval.
36
37
38

39 The base of the MN Zone 11 is defined by the first occurrence of the marker *Pa. feisti* in bed
40
41 BZ8 and corresponds to the base of the Upper Frasnian substage (Becker & House 1998).
42
43
44

45 The presence of the MN Zone 12 is attested in bed BZ12A with the first appearance of *Pa.*
46
47 *winchelli*, index-species of the zone.
48
49

50 The presence of the marker *Pa. bogartensis* in bed BZ13 allows discriminating the basis of MN
51
52 Zone 13. Thereafter, the level BZ13A.1 yields *Pa. ederi* and *Ag. asymmetricus*, and *Pa. brevis*
53
54 and *Pa. boogaardi* occur in bed BZ14. The co-occurrence of *Pa. beckeri* and *Pa. juntianensis* is
55
56 recorded in bed BZ15 at MN Zone 13.
57
58
59
60

1
2
3 The occurrences of *Pa. ultima* and *Pa. triangularis* in Bed BZ16 indicate the Famennian stage
4
5 (early *triangularis* Zone) (Klapper, 2007b). Other species present are *Pa. subperlobata*, *Pa. sp.*
6
7 B, *Pa. quadrantinodosalobata praeterita*, *Po. angustidiscus*, *Po. cf. politus* and *lc. alternatus*. In
8
9 our record, *Pa. cf. spathula* occurs slightly before the first known appearance of *Pa. spathula*
10
11 (late *triangularis* zone, Schülke 1999).
12
13
14
15
16
17

18 *4.c. Temporal calibration of terminal Frasnian black shales*

19
20 Temporary dysoxic events are recorded by several thin intervals of black shale deposits that
21
22 occur throughout the section, especially at the top of the succession where a 30 cm thick
23
24 discrete black shale level (between BZ 15 C and 15D) is developed. The Lower Kellwasser is
25
26 known to occur at the boundary between the MN zone 12 and MN zone 13 (Girard &
27
28 Renaud 2007). At Ben Zireg the Lower Kellwasser event is located at the end of sedimentary
29
30 cycle 12 and beginning of cycle 13, coinciding with the entry of the marker *Palmatolepis*
31
32 *bogartensis*. It is not defined by a distinct horizon but rather constitutes a two metres-thick
33
34 succession of dark micritic limestones and few black shales. In this interval the percentage of
35
36 *Icriodus* elements increases (Fig. 6). The Upper Kellwasser horizon that comprises at Ben
37
38 Zireg black shale deposits in its lower part and reddish calcisiltites in its upper part (BZ 15C
39
40 and D), represents the uppermost part of MN Zone 13 before being superseded by basal
41
42 Famennian strata (BZ 16). As the most basal part of the lower *triangularis* Zone has not been
43
44 established, a small depositional gap might exist. This gap might be related to a sea-level
45
46 instability, materialized by the pinkish detrital level. Such instability is often recorded at the
47
48 top of the Upper Kellwasser horizon event (i.e. Girard & Renaud, 2007; Schindler, 1990).
49
50
51
52
53
54
55
56
57
58
59
60

4.d. Biofacies

Numerous studies using the biofacies approach were performed through the Frasnian, from very different parts of the world, showing that the assemblages were dominated by *Polygnathus* during the early and middle Frasnian and then by *Palmatolepis* at the end of the Frasnian (e.g. Morocco: Lazreq, 1999; China: Ji & Ziegler 1993; USA: Sandberg et al., 1989; Russia: Ovnatanova et al., 1999). At Ben Zireg, three biofacies have been distinguished from bottom to top (Fig. 6): polygnathid-icriodid, polygnathid-ancyrodellid and then palmatolepid-polygnathid. Several peaks of *Icriodus* occur successively in Zone MN5 and through MN8 to MN10 zones. Ancyrodellids are never dominant but their percentage can reach more than 20% of the assemblage at the end of Zone MN11 and during Zone MN12.

The change from relatively shallow-water polygnathid-icriodid and polygnathid-ancyrodellid to deep-water palmatolepid-polygnathid biofacies through the Frasnian correlates with conclusions of Johnson, Klapper & Sandberg (1985) and trends in conodont biofacies that have been observed globally.

5. Interpretations and Discussion

5.a. Biostratigraphy and depositional environment

The fine-scaled conodont stratigraphy in the Ben Zireg section bears witness to a considerable stratigraphic hiatus at the base of the section. The underlying bed (BZ1) yields a Givetian, probably late Givetian conodont association where no typical Frasnian elements are found. This age attribution is emphasised by the occurrence of *Pharciceras* at Ben Zireg mentioned by Menchikoff (1936) but not recovered since (Massa et al., 1965). As the superseding bed (BZ1B) already yields *Pa. punctata*, the entire early Frasnian is missing (i.e. MN Zone 1 to MN Zone 4).

1
2
3 The interval of Zones MN5 to MN10 is only 9 m thick, attesting to severe condensation of
4
5 deposits under slow sedimentation rates. Conversely, sedimentation rates increased
6
7 considerably thereafter. Indeed, sedimentary cycles 10 to 15 comprise an interval of 17 m in
8
9 thickness that represents two-thirds of the entire Frasnian deposits. The sedimentary cycles
10
11 increase in importance from bottom to top of the section. Deposits of Zone MN13 reach a
12
13 thickness of 9 metres that correspond to one third of the total thickness of the section.
14
15

16
17 The resumption of the Middle Frasnian sedimentation is neither marked by erosional feature
18
19 nor transgressive deposits. Noticeable is only the presence of ferruginous cherty beds just
20
21 above the iron crust (bed BZ 1b). Sedimentary rocks are condensed in the lower part of the
22
23 section (cycles 1 to 6 with frequent iron coatings as micro-hard-grounds, phosphatic grains).
24
25 They display a rhythmic pattern dominated by micritic deposits. Detrital deposits are
26
27 restricted to a few shaly interbeds. Sediment texture is widely dominated by mudstones and
28
29 fewer wackestones. The fauna is poorly diversified, mostly represented by pelagic organisms
30
31 (tentaculites, pelagic bivalves, entomozoan crustaceans, goniatites and conodonts)
32
33 associated with some autochthonous benthic foraminifers and benthic ostracods, and rare
34
35 fragments of trilobites and brachiopods in flat mudstone pebbles. These criteria indicate that
36
37 the Frasnian deposits were emplaced on a distal ramp or distal platform setting (*e.g.* Franke
38
39 and Walliser, 1983), below wave action base and very far from emerged areas. Condensation
40
41 in intervals 1) and lower 2) could indicate a distal submarine rise. The interval 3) between
42
43 10.7 and 15.5m yield bioturbation features, rare pelagic organisms, abundant foraminifers,
44
45 benthic ostracods and numerous *Icriodus*. These indicate that the corresponding sediments
46
47 were probably deposited at shallower depth than both the overlying and underlying ones.
48
49 The interval 4) during the MN 13 Zone interval indicates a return to deeper outer ramp
50
51
52
53
54
55
56
57
58
59
60

1
2
3 setting. Noticeable is the presence of some soft pebbles levels composed of mudstones with
4
5 trilobites, indicative of downslope transport.
6
7
8
9

10 *5.b. Conodont distribution patterns*

11
12 In order to elucidate the significance of the conodont biofacies, assemblages from Ben Zireg
13
14 were compared with those established from the Mahrouma section (Ougarta area) by
15
16 Mahboubi *et al.* (2013a and b; Fig. 6).
17
18
19

20 The Marhouma section is distinct from Ben Zireg in the lower part of the Middle Frasnian
21
22 where a pure *Polygnathus* biofacies prevails. Thereafter, the Marhouma pattern is consistent
23
24 with the trends of the Ben Zireg section: first by a clear increase of the percentage of
25
26 ancyrodellids during the MN11 Zone (base of the Upper Frasnian), second by an obvious
27
28 increase of *Palmatolepis* till the Frasnian/Famennian boundary.
29
30
31

32 In detail, a slight shallowing is indicated during early late Frasnian (MN Zones 11-12) in both
33
34 sections due to the obvious increase of Ancyrodellids. As depositional environments are
35
36 different, *i.e.* submarine rise vs subsiding basin, this trend seems independent from local
37
38 setting dynamics.
39
40
41
42
43

44 *5.c. Gaps at the Givetian/Frasnian transition*

45
46 Occurrences of gaps in the later Devonian, and at the Givetian-Frasnian transition in
47
48 particular, are quite common in sections South of the Atlas Fault. This is most obvious in
49
50 Devonian outcrops of the Bechar Basin N and W of Ben Zireg where Lower Visean
51
52 conformably supersedes various levels from Tournaisian to Late Emsian (Weyant, 1988),
53
54 bearing witness of the marked diachronism of the omission. Whether the latter is due to
55
56
57
58
59
60

1
2
3 syntectonic movements as Weyant (1988) claims or results from current induced erosion or
4
5 influence of differentiated submarine scenery into sea-mounts and depressions is up to
6
7 debate.
8
9

10 The Ben Zireg section can be compared with other conodont-dated sections in Algeria and
11
12 Morocco (Fig. 6). Whereas in the basinal Marhouma section the Givetian-Frasnian transition
13
14 is presumably complete (Göddertz, 1987; Mahboubi et al., 2003 a), various sections on the
15
16 Tafilalet platform exhibit hiatuses (Becker & House, 2000b): Jebel Mech Irdane (*latifossatus-*
17
18 *semialternans* – MN 4 zones, Ebert, 1993; Walliser, 2000; Aboussalam, 2003), Jebel Amelane
19
20 (MN 1-6 zones, Becker & House 2000a), Bou Tchrafine (lower part of the MN zone 5, Becker
21
22 & House, 2000b; Aboussalam, 2003) and Hamer El Khdad section (*disparilis*-MN zone 4,
23
24 Gouwy, Haydukiewicz & Bultynck, 2007). In these areas, the extent of the gap is variable and
25
26 may range between the Givetian *disparilis* zone and the Frasnian MN zone 6. Hiatuses in the
27
28 same time interval were also noted in northern Morocco, France, Germany and Austria,
29
30 implying a supra-regional distribution (summarized in Table 1 of Hüneke, 2006). These
31
32 hiatuses have been interpreted as the result of a strong current-induced erosion, often
33
34 occurring within lower parts of fossil contourite units (Hüneke, 1995; 2006).
35
36
37 However, possible contourite deposits were not yet identified at Ben Zireg, and current
38
39 activity is not obvious as no sorting of conodont assemblages were observed, and
40
41 accumulations of benthic biodetritus do not occur. Instead, we favour the interpretation that
42
43 these hiatuses and condensation features resulted from depositional conditions on off-shore
44
45 submarine rises under slow sedimentation rates with numerous episodes of non-
46
47 sedimentation, sub-marine dissolution and abrasion of bedding surfaces marked by residual
48
49 iron-rich incrustations and hard-grounds.
50
51
52
53
54
55
56
57
58
59
60

5.d. Accumulation rates

Interbasinal correlation with the South Marhouma section reveals that the Middle Frasnian interval is at least four times thicker than the studied section (Fig. 6). This suggests an important accumulation rate that reflects high subsidence in the Ougarta trough, while the Bechar basin suffered condensation (Fig. 7). This is presumably due to voluminous fine-grained detrital influx from the far highlands (*i.e.* Reguibat shield) into the Ougarta sill.

At Ben Zireg condensation progressively reversed during Zones 11-13. This was also the case in Marhouma and in the Anti-Atlas and Meseta domains (Wendt and Belka, 1991, Gouwy, Haydukiewicz & Bultynck, 2007; Lazreq, 1999) (Fig. 7). At the end of the Upper Frasnian, deposits tend to homogenize in thickness over wide areas, which may reflect more uniform depositional conditions and less differentiated subsidence rates, both in platform and basin domains.

5.e. Late Frasnian Kellwasser facies

The terminal Frasnian Upper Kellwasser horizon is defined by an outstanding level of oxygen depleted carbon-rich black shales that occur intercalated in a succession of normally oxygenated sediments. This is generally the case in European sections of offshore cephalopod limestones (Schindler, 1990) but also in sections characterising the Central Moroccan Hercynian Meseta near Mrirt (Lazreq, 1999).

In contrast, South of the Main atlas Fault, the upper Frasnian both in platform and basin settings of the Tafilalet and Maider regions as well as in the South Marhouma area are exclusively composed of black cephalopod-rich limestones and shales. In these, the two Kellwasser horizons are lithologically difficult to discriminate but are discernable by

1
2
3 conodont and goniatite biostratigraphy. This lithological unit of Kellwasser-like facies that
4
5 starts diachronously at the base or in the lower part of the late Frasnian and extends into the
6
7 lower Famennian is termed by Wendt and Belka (1991) "Upper Kellwasser member".
8
9

10 The Ben Zireg section differs from all other known sections of SE Morocco and adjacent north-
11
12 western Algerian Sahara in the persistence of well-oxygenated carbonate sedimentation
13
14 throughout the Late Frasnian and Early Famennian in which, similarly to the situation in the
15
16 Moroccan Meseta, the intercalation of a clearly defined Upper Kellwasser horizon is obvious.
17
18 Additionally, the presence of the trilobite *Acuticryphops acuticeps*, index of pre-Upper
19
20 Kellwasser MN-Zone 13 in European and Central Moroccan sections, is exceptional as it is
21
22 unknown from contemporaneous dysoxic deposits of Kellwasser facies in Southern Morocco
23
24 and Algeria. Its occurrence shortly before the Upper Kellwasser horizon witnesses for normal
25
26 oxygenated level bottom conditions at Ben Zireg.
27
28
29

30
31 These "Hercynian" aspects of late Frasnian lithology and fauna bear evidence of closer
32
33 palaeogeographic relations to the Frasnian of the Maghrebid hercynides than any other area
34
35 South of the Atlas Fault.
36
37
38
39

40 **Conclusions**

41
42 The Ben Zireg section is representative for Frasnian deposits on the northern edge of the
43
44 Algerian Sahara platform where the early Upper Devonian is generally largely incomplete or
45
46 even absent in the Bechar Basin. Though the entire early Frasnian is missing at Ben Zireg, this
47
48 section is the most complete as, conformably superseding early mid-Devonian, all conodont
49
50 zones of the middle and late Frasnian are represented below the earliest Famennian. Such a
51
52 fine-scaled biozonation representing MN zones 5-13 is obtained for the first time in Frasnian
53
54 sequences of the Algerian Sahara.
55
56
57
58
59
60

1
2
3 The lithological succession is one of largely oxygenated, fine-grained carbonates of the off-
4
5 shore realm that are condensed in its lower part to continuously gain in importance when
6
7 sedimentation rates increase. The latest Frasnian MN Zone 13 is the thickest one. Initial
8
9 depositional environment was that of an offshore submarine rise subsequently smoothed to
10
11 bottom level at the top of the section.
12
13

14 Both the Lower and Upper Kellwasser events are recorded, the latter being represented by an
15
16 outstanding horizon of anoxic black-shale deposits intercalated in the normally oxygenated
17
18 carbonate sequence. This disposition is unique among all other known terminal Frasnian
19
20 sections S of the Atlas Fault where the Kellwasser horizons cannot be distinguished
21
22 lithologically within a uniform shale/limestone succession of Kellwasser-like kind. Conversely,
23
24 it is identical with equivalent sections characterizing terrains of the European and Maghrebian
25
26 Variscids. In particular, evidence of instability in the topmost part of the Upper Kellwasser
27
28 horizon has also been observed in many European sections.
29
30
31

32
33 Frasnian conodont biofacies patterns at Ben Zireg, *i.e.* the notable increase of Ancyrodellids at
34
35 the beginning of the late Frasnian, follows the global trend independently from local
36
37 conditions linked to subsidence or paleogeographic constraints.
38
39
40
41
42
43
44
45
46
47

48 Acknowledgements

49
50 The authors are very grateful to Abess Kada, M'hamed Mahboubi, Tadjedinne Hassen, Brigitte
51
52 Meyer Berthaud for their help in the field. Gilbert Klapper is deeply acknowledged for
53
54 checking some Frasnian conodonts. The first fieldwork was supported by the ANR Palasiafrica
55
56
57
58
59
60

1
2
3 (ANR- 08-JCJC-0017). Doriane Delmas and Christophe Nevado are acknowledged for the thin
4
5 sections. ISEM Contribution 2014-XXX.
6
7
8
9

10 References

- 11
12 ABOUSSALAM, Z. S. 2003. *Das "Taghanic-Event" im höheren Mittel-Devon von West-Europa und*
13 *Marokko*. Thesis, Geologisch-Paläontologisches Institut der Universität Münster 1–332.
14
15 ALLARY, A., LAVENU, A. & RIBEYROLLES, M., 1976. Etude tectonique et microtectonique d'un
16 segment de chaîne hercynienne dans la partie sud-orientale du Maroc central. *Notes et Mémoires*
17 *Service géologique Maroc* **261**, 113–66.
18
19
20 BECKER, R. T. & HOUSE, M. R. 2000a. Devonian ammonoid succession at Jebel Amelane (western
21 Tafilalt, southern Morocco). *Notes et Mémoires du Service Géologique du Maroc* **399**, 49– 56.
22
23 BECKER, R. T. & HOUSE, M. R. 2000b. Late Givetian and Frasnian ammonoid succession at Bou
24 Tchrafine (Anti Atlas, southern Morocco). *Notes et Mémoires du Service Géologique du Maroc*
25 **399**, 27–36.
26
27
28 BEŁKA, Z., KLUG, C., KAUFMANN, B., KORN, D., DÖRING, S., FEIST, R. & WENDT, J. 1999. Devonian
29 conodont and ammonoid succession of the eastern Tafilalt (Ouidane Chebbi section), Anti-Atlas,
30 Morocco. *Acta Geologica Polonica* **49**, 1–23.
31
32
33 BRANSON, E. B. & MEHL, M. G. 1934. Conodonts from the Grassy Creek Shale of Missouri. The
34 University of Missouri Studies **8**, 171–59.
35
36
37 BULTYNCK, P. & JACOBS, L. 1981. Conodontes et sédimentologie des couches de passage du Givétien
38 au Frasnian dans le Nord du Tafilalt et dans le Ma' der (Maroc présaharien). *Bulletin de l'Institut*
39 *royal des Sciences naturelles de Belgique, Sciences de la Terre* **53**, 1–24.
40
41
42 BULTYNCK, P. 1986. Accuracy and reliability of conodont zones: the *Polygnathus asymmetricus*
43 "zone" and the Givetian–Frasnian boundary. *Bulletin de l'Institut royal des Sciences naturelles de*
44 *Belgique, Sciences de la Terre* **56**, 269–80.
45
46
47 BULTYNCK, P. 1987. Pelagic and neritic conodont successions from the Givetian of the pre-Sahara
48 Morocco and the Ardennes). *Bulletin de l'Institut royal des Sciences naturelles de Belgique,*
49 *Sciences de la Terre* **57**, 149–81.
50
51
52 CONRAD, J., MASSA, D. & WEYANT, M. 1986. Late Devonian regression and Early Carboniferous
53 transgression on the Northern African platform. *Annales de la Société Géologique de Belgique*
54 **109**, 113–22.
55
56
57
58
59
60

- 1
2
3 EBERT, J. 1993. Globale events im Grenz-Bereich Mittel-/Oberdevon. *Göttinger Arbeiten Zur Geologie*
4 *und Paläontologie* **59**, 1–106.
5
6 ELMI, S. & AMEUR, M. 1984. Quelques environnements des faciès noduleux mésogéens. *Geologica*
7 *Romana* **23**, 13-22.
8
9
10 FLAMAND, G. B. M. 1911. Recherches géologiques et géographiques sur le Haut-Pays de l'Oranie et
11 sur le Sahara (Algérie et Territoires du Sud). Thèse Sciences, Lyon, n° 47, Rey édition. , pp.1–1001.
12
13 FLÜGEL, E. 2004. *Microfacies of carbonate rocks. Analysis, Interpretation and Application*. Berlin,
14 Heidelberg, NewYork: Springer Verlag.
15
16
17 FRANKE, W & WALLISER, O. H. 1983. Pelagic carbonates in the Variscan belt – their sedimentary and
18 tectonic environments. In *Intracontinental fold belts* (eds H. Martin & F. W, Eder), pp. 77–92.
19 Springer, Berlin & Heidelberg.
20
21
22 GIRARD, C. & RENAUD, S. 2007. Quantitative conodont-based approaches for correlation of the Late
23 Devonian Kellwasser anoxic events. *Palaeogeography, Palaeoclimatology, Palaeoecology* **250**,
24 114–25.
25
26
27 GÖDDERTZ, B. 1987. Devonische Goniatiten aus SW-Algerien und ihre stratigraphische Einordnung in
28 die Conodonten-Abfolge. *Palaeontographica Abteilung A* **A197**, 127–220.
29
30
31 GOUWY, S., HAYDUKIEWICZ, J. & BULTYNCK, P. 2007. Conodont-based graphic correlation of upper
32 Givetian-Frasnian sections of the Eastern Anti-Atlas (Morocco). *Geological Quarterly* **51**, 375–92.
33
34
35 HINDE, G. J. 1879. On Conodonts from the Chazy and Cincinnati Group of the Cambro-Silurian, and
36 from the Hamilton and Genesee-Shale Divisions of the Devonian, in Canada and the United States.
37 *Quarterly Journal of the Geological Society* **35**, 351–69.
38
39
40 HÜNEKE, H. 1995. Early Devonian (Emsian) to Late Devonian (Famennian) stratigraphy and conodonts
41 of the Antoinettenweg section in the Lower Harz Mountains (Germany). *Courier*
42 *Forschungsinstitut Senckenberg* **188**, 99–131.
43
44
45 HÜNEKE, H. 2006. Erosion and deposition from bottom currents during the Givetian and Frasnian:
46 Response to intensified oceanic circulation between Gondwana and Laurussia. *Palaeogeography,*
47 *Palaeoclimatology, Palaeoecology* **234**, 146–67.
48
49
50 JI, Q., ZIEGLER, W. 1993. Lali section: an excellent reference section for Upper Devonian in south
51 China. *Courier Forschungsinstitut Senckenberg* **157**, 1–183.
52
53
54 JOHNSON, J. G., KLAPPER, G. & SANDBERG, C. A. 1985. Devonian eustatic fluctuations in Euramerica.
55 *Geological Society of America Bulletin* **96**, 567-87.
56
57
58 KLAPPER, G., FEIST, R. & HOUSE, M.R. 1987. Decision on the boundary stratotype for the
59 Middle/Upper Devonian Series boundary. *Episodes* **16**, 433–41.
60

- 1
2
3 KLAPPER, G. 1989. The Montagne Noire Frasnian (Upper Devonian) conodont succession, In *Devonian*
4 *of the World*. (eds N. J. McMillan, A. F. Embry, A. F. Glass), pp. 449–68. Calgary. Canadian Society
5 of Petroleum Geologists, 14 (3).
6
7
8 KLAPPER, G. 1990. Frasnian species of the Late Devonian conodont genus *Ancyrognathus*. *Journal of*
9 *Paleontology* **64**, 998–1025.
10
11 KLAPPER, G., FEIST, R., BECKER, R. & HOUSE, M.R. 1993. Definition of the Frasnian / Famennian stage
12 boundary. *Episodes* **16**, 433–41.
13
14 KLAPPER, G. & FOSTER, C. T. 1993. Shape analysis of Frasnian species of the Late Devonian conodont
15 genus *Palmatolepis*. *Journal of Paleontology* **67**, 1–33.
16
17
18 KLAPPER, G. 2007a. Frasnian (Upper Devonian) conodont succession at Horse Spring and correlative
19 sections, Canning Basin, Western Australia. *Journal of Paleontology* **81**, 513–37.
20
21
22 KLAPPER, G. 2007b. Conodont taxonomy and the recognition of the Frasnian/Famennian (Upper
23 Devonian) Stage Boundary. *Stratigraphy* **4**, 67–76.
24
25
26 LAZREQ, N. 1999. Biostratigraphie des conodontes du Givétien au Famennien du Maroc central -
27 Biofacies et événement Kellwasser. *Courier Forschungsinstitut Senckenberg* **214**, pp. 1–111.
28
29 MAHBOUBI, A., GIRARD, C., CORNÉE, J. J & FEIST, R. 2013a. Late Devonian sequences and
30 biostratigraphy of the South Marhouma region, Beni Abbès, SW Algeria - Preliminary results. In
31 *International field symposium « The Devonian and Carboniferous of Northern Gondwana »*, SDS
32 joint field meeting, (eds El Hassini, A., Becker, R. T, Tahiri, A.), pp. 80. Document de l'institut
33 Scientifique, Rabat, Mémoire 27.
34
35
36 MAHBOUBI, A., CORNÉE, J. J & GIRARD, C. 2013b. Biostratigraphie et évolution sédimentaire du
37 dévonien supérieur de la plate-forme occidentale saharienne la coupe de Marhouma sud, SO-
38 algerien. *14ème congrès français de sédimentologie*, pp. 257, Paris.
39
40
41
42 MASSA, D., COMBAZ, A. & MANDERSIED, G. 1965. Observations sur les séries Siluro – Dévoniennes
43 des confins algéro-marocains du Sud. *Notes et mémoire de la Compagnie Française du Pétrole*,
44 Paris, pp.1–188.
45
46
47 MAMET, B. & PRÉAT, A. 2006. Iron-bacterial mediation in Phanerozoic red limestones: State of the
48 art. *Sedimentary Geology* **185**, 147–157.
49
50
51 MENCHIKOFF, N. 1936. Etude géologique sur les confins algéro-marocains du Sud. *Bulletin de la*
52 *Société Géologique de France* **6**, 131–49.
53
54
55 MÜLLER, K. J & MÜLLER, E. M. 1957. Early Upper Devonian (Independence) conodonts from Iowa,
56 part 1. *Journal of Paleontology* **31**, 1069–1108.
57
58
59
60

- 1
2
3 OVNATANOVA, N. S. 1976. Novye pozdnedevonskie konodonty Russkoy Platformy.
4 *Paleontologicheskii Zhurnal* 2, 106-15. [In Russia].
5
6 PAREYN, C. 1961. Les Massifs Carbonifères du Sahara Sud-Oranais. *Publications du Centre de*
7 *Recherches Sahariennes, Série Géologie, CNRS., (ed), Paris, pp. 1–324.*
8
9
10 SANDBERG, C. A. 1976. Conodont biofacies of late Devonian *Polygnathus styriacus* Zone in western
11 United States. In *Conodont Paleoecology* (ed. C. R. Barnes), pp. 171–86. Montreal: *Geological*
12 *Association of Canada, Special Paper No. 15.*
13
14 SANDBERG, C. A., POOLE, F. G. & JOHNSON, J. G. 1989. Upper Devonian of Western United States. In
15 *Devonian of the World*, (eds N. J. McMillan, A. F. Embry & D. J. Glass). Calgary: Canadian Society of
16 Petroleum Geologists, Memoir 14 (1) 183–220.
17
18
19
20 SANDBERG, C. A., ZIEGLER, W., DREESEN, R & BUTLER, J. L. 1992. Conodont biochronology, biofacies,
21 taxonomy and event stratigraphy around Middle Frasnian Lion Mudmound (F2h), Belgium. *Courier*
22 *Forschungsinstitut Senckenberg* 150, 1–87.
23
24
25 SANNEMANN, D. 1955. Beitrag zur Untergliederung des Oberdevons nach Conodonten. *Neues*
26 *Jahrbuch für Geologie und Paläontologie, Abhandlungen* 100, 324–31.
27
28
29 SCHINDLER, E. 1990. *Die Kellwasser-Krise (höhe Frasn-Stufe, Oberdevon)*. Göttingen, Göttinger
30 Arbeiten zur Geologie und Päläontologie; 115 p.
31
32 SCHÜLKE, I. 1995. Evolutive Prozesse bei *Palmatolepis* in der frühen Famenne-Stufe (Conodonta,
33 Ober-Devon). *Göttinger Arbeiten zur Geologie und Paläontologie* 67, pp.1–108.
34
35
36 SCHÜLKE, I. 1999. Early Famennian conodont biostratigraphy of the stratotype area (Montagne
37 Noire, Southern France). *Bolletino della Società Paleontologica Italiana* 37, 375–391.
38
39 SEDDON, G. & SWEET, W. C. 1971. An ecologic model for conodonts. *Journal of Paleontology* 45(5),
40 869– 80.
41
42
43 STAUFFER, C. R. 1938. Conodonts of the Olentangy Shale. *Journal of Paleontology* 12, 411–43.
44
45 WALLISER, O. H. 2000. The Jebel Mech Irdane section. *Notes et Mémoires du Service Géologique du*
46 *Maroc* 399, 57– 62.
47
48 WENDT, J. & BELKA, Z. 1991. Age and depositional environment of upper devonian (early Frasnian to
49 early famennian) black shales and limestones (Kellwasser facies) in the eastern Anti-Atlas,
50 Morocco. *Facies* 25, 51–89.
51
52
53 WENDT, J., KAUFMANN, B., BELKA, Z., KLUG, C & LUBESSEDER, S. 2006. Sedimentary evolution of a
54 Palaeozoic basin and ridge system: the Middle and Upper Devonian of the Ahnet and Mouydir
55 (Algerian Sahara). *Geological Magazine* 143, 269–99.
56
57
58
59
60

- 1
2
3 WEYANT, M. 1988. Relationship between Devonian and Carboniferous strata near the northern
4 confines of the Bechar basin, Algeria. *Courier Forschungsinstitut Senckenberg* **100**, 235–41.
5
6 WRIGHT, V.P. & BURCHETTE, T.P. 1996. Shallow-water carbonate environments. In *Sedimentary*
7 *Environments: processes, facies, and stratigraphy* (ed H.G. Reading). Blackwell Sc., Oxford, pp.
8 325-394.
9
10 YOUNGQUIST, W. 1947. A New Upper Devonian Conodont Fauna from Iowa. *Journal of Paleontology*
11 **21**, 95–12.
12
13 ZIEGLER, W., 1975. *Polygnathus linguiformis linguiformis* (gamma morphotype). In *Catalogue of*
14 *Conodonts* (ed W. Ziegler). E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, II, pp. 463.
15
16
17
18
19
20
21
22

23 **Figure and table captions**

24
25 Figure 1. Main Palaeozoic outcrops in north-west Africa. (a) location of selected Frasnian
26 sites: 1 - Bou Terga; 2 - Jebel Atrous II; 3 - Bou Thrafine; 4 - Ouidane Chebbi; 5 – Oued Rheris;
27
28 6 – Anajdam; 7 – Bou Alzaz nord; 8 – El Mader el Madjib; 9 – Soltane el Boutoum; 10 – South
29
30 Mahrouma; (b) investigated section in the Ben Zireg anticline (after Pareyn, 1961).
31
32
33
34

35
36
37 Figure 2. Stratigraphic log with sedimentary cycles, Kellwasser events and distribution of
38 conodont species in the Givetian, Frasnian and lowermost Famennian at Ben Zireg section,
39 Bechar basin. *An-Ancyrodella*; *Ag-Ancyrognathus*; *Pa-Palmatolepis*; *Po- Polygnathus*; *Ic-*
40 *Icriodus*; *ling - linguiformis linguiformis* (gamma morphotype); *q-quadrantinodosalobata*;
41
42 LKW- Lower Kellwasser ; UKW- Upper Kellwasser; MN Montagne Noire Zone.
43
44
45
46
47
48
49
50

51 Figure 3. Exposure of the Ben Zireg succession from the Givetian to the Famennian showing
52 15 sedimentary cycles. Numbers correspond to sedimentary cycles. The Frasnian/Famennian
53
54
55
56
57
58
59
60

1
2
3 boundary is located on top of the Upper Kellwasser horizon, and coincides with the top of
4
5 cycle 15. The scale is given by B. Meyer-Berthaud who is 1.55 m-high on the up-right.
6
7
8
9

10 Figure 4. Upper Devonian facies types of Ben Zireg section (Algeria). Os – benthic ostracods;

11
12 Tr – trilobite; Ra – Radiolarian; Te – Tentaculite (*Homoctenus*).
13

14
15 (a) View of the ferruginous limestone unit (7th cycle)
16

17 (b) View of the thin to thick-bedded grey nodular limestone (13th cycle)
18

19 (c) Upper Kellwasser horizon (UKW) composed of a lower part with black shales (to the left)
20

21 and a upper part of pinkish calcisiltites (BZ15D). BZ16 correspond to the lowermost
22

23 Famennian. The UKW is 0.65 m-thick.
24

25
26 (d) Tentaculid wackestone (Bed BZ6C, MN Zone 8).
27

28 (e) Ostracods mudstone with unsculptured benthic specimens (Bed BZ13A, MN Zone 13).
29

30 (f) Surface of mudstone slab with trilobite sclerites of *Acuticryphops acuticeps* (Kayser,
31
32 1889). Pointed glabella and reduced size of eye-lobe with few prominent lenses (arrow) are
33
34 characteristic (Bed BZ14D, MN Zone 13).
35
36
37
38
39

40
41 Figure 5. Significant conodonts from the Ben Zireg section:
42

43 (a) *Ancyrodella gigas* Youngquist, 1947 form 2 (Bed BZ2); (b) *Polygnathus pseudofoliatius*
44

45 Wittekindt, 1866 (Bed BZ1); (c) *Polygnathus linguiformis linguiformis* Hinde, 1879 gamma
46

47 morphotype, Ziegler et al. 1976 (Bed BZ1); (d) *Ancyrodella curvata* (Branson & Mehl, 1934)
48

49 early form (Bed BZ5B); (e) *Ancyrodella lobata* Branson & Mehl, 1934 (Bed BZ6); (f)
50

51 *Ancyrognathus triangularis* Youngquist, 1945 (Bed BZ11); (g) *Palmatolepis kireevae*
52

53 Ovnatanova, 1976 (Bed BZ8); (h) *Palmatolepis ljaschenkoae* Ovnatanova, 1976 (Bed BZ6); (i)
54

55 *Ancyrognathus coeni* Klapper, 1990 (Bed BZ11); (j) *Palmatolepis feisti* Klapper, 2007a (Bed
56
57
58
59
60

1
2
3 BZ8); (k) *Palmatolepis punctata* (Hinde, 1879), (Bed BZ2); (l) *Palmatolepis jamieae* Ziegler &
4 Sandberg, 1990 (Bed BZ10B); (m) *Palmatolepis bogartensis* (Stauffer, 1938), (Bed BZ13); (n)
5 *Palmatolepis boogaardi* Klapper & Foster, 1993 (Bed BZ14D.1); (o) *Palmatolepis hassi* Müller
6 & Müller, 1957 (Bed BZ6B); (p) *Palmatolepis* cf. *spathula* Schülke, 1995 (Bed BZ16); (q)
7 *Palmatolepis ultima* Sannemann, 1955 (Bed BZ16); (r) *Palmatolepis subperlobata* Branson &
8 Mehl, 1934 (Bed BZ16); (s) *Palmatolepis winchelli* Stauffer, 1938 (Bed BZ10A); (t)
9 *Palmatolepis triangularis* Sannemann, 1955.
10
11
12
13
14
15
16
17
18
19
20
21

22 Figure 6. Percentage of conodont genera at Ben Zireg (left) and at the South Marhouma
23 section (right) and fine-scaled correlation between these sections based on the detailed
24 biostratigraphical frame. Po = Polygnathid biofacies; Po-Ic = Polygnathid-Icriodid biofacies;
25 Po-An = Polygnathid-Ancyrodellid biofacies; Pa-Po = Palmatolepid-Polygnathid biofacies. G =
26 Givetian; t. = *triangularis*.
27
28
29
30
31
32
33
34
35

36 Figure 7. Correlation of selected Frasnian sections from both sides of the Main Atlas Fault
37 (location see Fig. 1b): Bou Terga, Jebel Atrous II (Gouwy, Haydukiewicz & Bultynck, 2007);
38 Bou Tchrafine (Bultynck & Jacob, 1981, Bultynck, 1986, 1987; Becker & House, 2000b);
39 Ouidane Chebbi (Wendt & Belka, 1991; Belka et al. 1999), Oued Rheris (Wendt & Belka,
40 1991), Anajdam, Bou Alzaz Nord (Lazreq, 1999), South Marhouma (Mahboubi et al. 2013 a,
41 b; Mahboubi & Gatovsky, submitted).
42
43
44
45
46
47
48
49
50
51




52 Table 1. Numbers of platform conodont genera at the Ben Zireg section. The low conodont
53 abundances (<20P1) in italic are not included in the biofacies analyses. Pa-*Palmatolepis*; Po-
54 *Polygnathus*; An- *Ancyrodella* and *Ancyrognathus*; Ic-*Icriodus*.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Sample	Weight (g)	Depth (m)	Pa	Po	An	Ic	Total
BZ16B	200	27.6	52	2		2	56
BZ16A	1138	27.25	193	25		1	219
BZ16	990	26.7	178	31			209
BZ15D	200	26.5	26	5	8		39
BZ15C	827	26	70	18	9	1	98
BZ15A	100	25	21		2	1	24
BZ15	873	24.1	52	5	2	9	68
BZ14D.1	1000	23.5	296	73	9	16	394
<i>BZ14C</i>	90	22.5	4	8	1		13
BZ14	1205	22	80	22			102
BZ13B	210	21	35	13	1	4	53
BZ13A.1	1260	20.1	160	24	19	42	245
BZ13	879	18.5	62	22	1	23	108
BZ12A	751	17.6	5	37	8	4	54
BZ11	230	16	1	14	13	2	30
BZ10B	230	15.5	3	13	4	3	23
BZ10A	200	13.65	7	9	3	3	22
BZ10	671	12.82	15	32	7	9	63
BZ9	456	11.85		44	4	21	69
BZ8A.1	747	11.5	13	54	5	43	115
<i>BZ8A</i>	140	11.25	8	4		5	17
BZ8	875	8.75	12	20	7	43	82
<i>BZ7A</i>	170	9	4	7	1	3	15
BZ7	834	8.1	12	41	4	28	85
BZ6B	400	7.25	6	35	1	5	47
BZ6	620	7	4	33		8	45
<i>BZ5B</i>	150	5.25		4	4		8
BZ5	730	5.75		66		4	70
BZ4B.1	1120	5.25		79	6	40	125
<i>BZ4B</i>	267	5		14		1	15
<i>BZ3C</i>	580	3.25		6		1	7
BZ3	610	3	2	12		12	26
<i>BZ2C</i>	347	2.25		2		3	5
BZ2	1083	1.75	2	13	2	17	34
<i>BZ1C</i>	1037	1		3			3
BZ1A	715	0.5	1	48	1	4	54
<i>BZ1</i>	800	0		27		1	28


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



63x87mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




76x112mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




289x80mm (300 x 300 DPI)




194x212mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




140x180mm (170 x 170 DPI)



42x37mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



42x42mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60