



HAL
open science

On the Galois groups of generalized Laguerre Polynomials

Shanta Laishram

► **To cite this version:**

Shanta Laishram. On the Galois groups of generalized Laguerre Polynomials. Hardy-Ramanujan Journal, 2014, Volume 37 - 2014, pp.8-12. 10.46298/hrj.2014.1317 . hal-01220303

HAL Id: hal-01220303

<https://hal.science/hal-01220303>

Submitted on 27 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Galois groups of generalized Laguerre Polynomials

Shanta Laishram

Abstract. For a positive integer n and a real number α , the generalized Laguerre polynomials are defined by

$$L_n^{(\alpha)}(x) = \sum_{j=0}^n \frac{(n+\alpha)(n-1+\alpha)\cdots(j+1+\alpha)(-x)^j}{j!(n-j)!}.$$

These orthogonal polynomials are solutions to *Laguerre's Differential Equation* which arises in the treatment of the harmonic oscillator in quantum mechanics. Schur studied these Laguerre polynomials for their interesting algebraic properties. In this short article, it is shown that the Galois groups of Laguerre polynomials $L_n^{(\alpha)}(x)$ is S_n with $\alpha \in \{\pm\frac{1}{2}, \pm\frac{1}{3}, \pm\frac{2}{3}, \pm\frac{1}{4}, \pm\frac{3}{4}\}$ except when $(\alpha, n) \in \{(\frac{1}{4}, 2), (-\frac{2}{3}, 11), (\frac{2}{3}, 7)\}$. The proof is based on ideas of p -adic Newton polygons.

Keywords. Irreducibility, Laguerre Polynomials, Newton Polygons, Arithmetic Progressions, Primes.

2010 Mathematics Subject Classification. 2000 Mathematics Subject Classification: Primary 11A41, 11B25, 11N05, 11N13, 11C08, 11Z05

1. Introduction

For a positive integer n and a real number α , the generalized Laguerre polynomials are defined by

$$L_n^{(\alpha)}(x) = \sum_{j=0}^n \frac{(n+\alpha)(n-1+\alpha)\cdots(j+1+\alpha)(-x)^j}{j!(n-j)!}.$$

These orthogonal polynomials have a wide range of applications in several areas of mathematics. Not long after its appearance in the literature early in the twentieth century, it became evident, in the hands of Schur, that the generalized Laguerre polynomials also enjoy algebraic properties of great interest. In fact the irreducibility of these polynomials is connected to finding explicit examples of solutions to Hilbert's *Inverse Galois Problem*. We refer to [FKT12] for more details.

It was shown that $L^{(\alpha)}(x)$ is irreducible for $\alpha \in \{\pm\frac{1}{2}\}$ in [Sch29] and [Sch31] and for $\alpha \in \{\pm\frac{1}{3}, \pm\frac{2}{3}, \pm\frac{1}{4}, \pm\frac{3}{4}\}$ in [LaSh, Theorem 1] except when $\alpha = \frac{1}{4}, n = 2$. By using these results of irreducibility, it was shown in [SaSh15, Theorem 1.4] that the Galois group of $L^{(\alpha)}(x)$ is S_n for $n \geq n_0$ where $n_0 = 182, 876, 1325$ if $q \in \{\pm\frac{1}{2}\}$, $q \in \{\pm\frac{1}{3}, \pm\frac{2}{3}\}$ and $q \in \{\pm\frac{1}{4}, \pm\frac{3}{4}\}$, respectively. In this short note, we give a complete result for all n . Here S_n is the Symmetric Group on n symbols and A_n is the Alternating Group on n symbols. We prove

Theorem 1.1. *Let $\alpha \in \{\pm\frac{1}{2}, \pm\frac{1}{3}, \pm\frac{2}{3}, \pm\frac{1}{4}, \pm\frac{3}{4}\}$. The Galois group of Laguerre polynomials $L_n^{(\alpha)}(x)$ is S_n for every $n \geq 1$ except when $(\alpha, n) \in \{(\frac{1}{4}, 2), (-\frac{2}{3}, 11), (\frac{2}{3}, 7)\}$ where it is A_n for $(\alpha, n) \in \{(-\frac{2}{3}, 11), (\frac{2}{3}, 7)\}$ and S_1 for $(\alpha, n) = (\frac{1}{4}, 2)$.*

We give a proof of Theorem 1.1 in Section 3. The proof of Theorem 1.1 is an application of a result of Hajir [Haj05] based on p -adic Newton polygons, see Lemmas 2.1 and 2.2. The new ingredient in this paper is the clever use of Lemma 2.1 as Lemma 2.2 instead of [SaSh15, Lemma 3.3]. In fact the proof of [SaSh15, Theorem 1.4] can be much shortened by using Lemma 2.2.

2. Preliminaries

Hajir [Haj05] gave a criterion for an irreducible polynomial to have large Galois group using Newton polygons. We restate the result which is [Haj05, Lemma 3.1].

Lemma 2.1. *Let $f(x) = \sum_{j=0}^m \binom{m}{j} c_j x^j \in \mathbb{Q}[X]$ be an irreducible polynomial of degree m . Let p be a prime with $\frac{m}{2} < p < m - 2$ such that*

- (i) $\text{ord}_p(c_0) = 1$,
- (ii) $\text{ord}_p(c_j) \geq 1$ for $0 \leq j \leq m - p$,
- (iii) $\text{ord}_p(c_p) = 0$.

*Then p divides the order of Galois group of f over \mathbb{Q} . In fact, this Galois group is A_m if $\text{disc}(f) \in \mathbb{Q}^{*2}$ and S_m otherwise.*

We will be applying the above lemma to following polynomial. Let $\alpha = \frac{u}{v}$ with $u, v \in \mathbb{Z}$, $\text{gcd}(u, v) = 1$ and $v > 0$. Let

$$\begin{aligned} G(x, u, v) &:= v^n n! L\left(\frac{u}{v}\right)\left(\frac{-x}{v}\right) \\ &= \sum_{j=0}^n \binom{n}{j} (u + vn)(u + v(n-1)) \cdots (u + v(j+1)) x^j. \end{aligned} \quad (2.1)$$

In [Sch31], Schur showed that its discriminant is given by

$$D_n^{(u,v)} := \text{Disc}(G(x, u, v)) = \prod_{j=2}^n j^j \left(\frac{u}{v} + j\right)^{j-1}.$$

We write $D_n^{(u,v)} = bY^2$, $Y \in \mathbb{Q}$ with

$$b = \begin{cases} \frac{3 \cdot 5 \cdots n \cdot (u+2v)(u+4v) \cdots (u+(n-1)v)}{v^\delta} & \text{if } n \equiv 1, 3 \pmod{4} \\ \frac{3 \cdot 5 \cdots (n-1) \cdot (u+2v)(u+4v) \cdots (u+nv)}{v^\delta} & \text{if } n \equiv 0, 2 \pmod{4} \end{cases} \quad (2.2)$$

where $\delta = 0$ if $n \equiv 0, 1 \pmod{4}$ and 1 if $n \equiv 2, 3 \pmod{4}$.

We now apply Lemma 2.1 to $G(x, u, v)$. We prove

Lemma 2.2. *Let $1 \leq r < v$, $\text{gcd}(r, v) = 1$ and p be a prime with*

$$p > v, p \equiv r^{-1}u \pmod{v} \text{ and } \frac{u + v + nv}{r + v} \leq p \leq n - 3. \quad (2.3)$$

Let $G(x, u, v)$ be given by (2.1) be an irreducible polynomial of degree n . Assume that $|u| < v$. Then the Galois group of $G(x, u, v)$ is A_n or S_n according as b (given by (2.2)) is a square or not an square of an integer.

Proof. We apply Lemma 2.1 with

$$c_j = (u + vn)(u + v(n-1)) \cdots (u + v(j+1)).$$

Since $1 \leq r < v$, we have $\frac{u+v+nv}{r+v} > \frac{n}{2}$ and hence $\frac{n}{2} < p < n - 2$ is valid. It suffices to check conditions (i) – (iii) of Lemma 2.1.

Since $p \equiv r^{-1}u \pmod{v}$, we get $p|(u+iv)$ for some i . Let i_0 be the least positive integer i with this property. Then $1 \leq i_0 < p$. Further let $u + i_0v = pr_0$. Then $r_0 \equiv r \pmod{v}$. We claim that $r_0 < v$. Suppose not. Then $u + i_0v = pr_0 \geq pv \geq (i_0 + 1)v$ since $i_0 < p$ contradicting $|u| < v$. Thus $r_0 < v$. This with $r_0 \equiv r \pmod{v}$ and $1 \leq r < v$ implies $r = r_0$. Since $\frac{u+v+nv}{r+v} \leq p$, we have

$$u + v + (n - p)v \leq rp = r_0p = u + i_0v$$

giving $i_0 > n - p$. Thus $n - p < i_0 < p$. This gives $i_0 - p < 0$ and $i_0 + p > n$ and hence $u + i_0v$ is the only multiple of p in $\{u, u + v, \dots, u + nv\}$. Further $u + i_0v = pr < pv < p^2$ implying $p|(u + i_0v)$. Hence conditions (i) – (iii) of Lemma 2.1 are valid and the assertion follows.

The above Lemma contains [SaSh15, Lemma 3.3]. We also need the following result on b being a square or not.

Lemma 2.3. *Let $2 \leq n \leq 1325$,*

$$\alpha = \frac{u}{v} \in \left\{ \pm \frac{1}{2}, \pm \frac{1}{3}, \pm \frac{2}{3}, \pm \frac{1}{4}, \pm \frac{3}{4} \right\}$$

and b be given by (2.2). Then b is square only when $(u, v, n) \in \{(-2, 3, 3), (-2, 3, 11), (2, 3, 7)\}$.

Proof. First we check that for $2 \leq n \leq u + 2v$, the assertion is valid. Hence we now take $n > u + 2v$. Let

$$n_1 = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ \frac{n-1}{2} & \text{if } n \text{ is odd.} \end{cases}$$

Assume $u \neq \pm 2$ if $v = 3$. Then we see that b is divisible by every prime $p \equiv u \pmod{2v}$ with $n < p \leq u + 2vn_1$ to the first power. Hence if there is such a prime, b cannot be a square. For $u + 2v < n \leq 1325$, we check that this is true. Thus we now take $v = 3, u = \pm 2$. Let $u_1 = \frac{u}{2}$. Then

$$(u + 2v) \cdots (u + 2n_1v) = 2^{n_1}(u_1 + v)(u_1 + 2v) \cdots (u_1 + n_1v)$$

and hence b is not a square if there is a prime p with $n < p \leq u_1 + n_1v$ and $p \equiv u_1 \pmod{v}$ where $v = 3$. We check that this is the case for $u + 2v < m \leq 1325$ except when $u = -2, m \in \{5, 6, 7, 11\}$ and $u = 2, m = 19$. For $u = -2, m \in \{5, 6, 7, 11\}$ and $u = 2, m = 19$, we check that b is not a square except when $u = -2, m = 11$. Hence the assertion follows.

3. Proof of Theorem 1.1

Let

$$\alpha = \frac{u}{v} \in \left\{ \pm \frac{1}{2}, \pm \frac{1}{3}, \pm \frac{2}{3}, \pm \frac{1}{4}, \pm \frac{3}{4} \right\}.$$

As mentioned before, it was shown that $L^{(\alpha)}(x)$ is irreducible for $\alpha \in \{\pm \frac{1}{2}\}$ in [Sch29] and [Sch31] and in [LaSh] for $\alpha \in \{\pm \frac{1}{3}, \pm \frac{2}{3}, \pm \frac{1}{4}, \pm \frac{3}{4}\}$ except when $\alpha = \frac{1}{4}, n = 2$ and hence same is true for $G(x, u, v)$. For $n \leq 13$, we check in *SAGE* for $n \leq 11$ and *MAGMA* for $n = 12, 13$ that the assertion of Theorem 1.1 is valid.

Hence we may suppose that $n > 13$. Further we can take $n \leq 1325$ by [SaSh15, Theorem 1.4]. It suffices to prove that $G(x, u, v)$ has Galois group S_n . We use Lemmas 2.2 and 2.2. It suffices to find a prime p with

$$p > v, p \equiv r^{-1}u \pmod{v} \text{ and } \frac{u + v + nv}{r + v} \leq p \leq n - 3.$$

for some $r, 1 \leq r < v$, $\gcd(r, v) = 1$. Let $\alpha = \frac{u}{v} \in \{\pm\frac{1}{2}\}$. We check that there is a prime p with $\frac{2n+2+u}{3} \leq p \leq n-3$ except when $u = 1, n = 19$. We check that for $n = 19$, the Galois group of $L^{(\frac{1}{2})}(x)$ is S_n .

Let $\alpha = \frac{u}{v} \in \{\pm\frac{1}{3}, \pm\frac{2}{3}\}$. Since $1 \leq r < 3$, we need to find a prime p with

$$\frac{3n}{4} + \frac{3+u}{4} \leq p \leq n-3, p \equiv u \pmod{3}$$

or

$$\frac{3n}{5} + \frac{3+u}{5} \leq p \leq n-3, p \equiv 2u \pmod{3}.$$

Hence it suffices to find a prime p with $\frac{3n}{4} + \frac{3+u}{4} \leq p \leq n-3$ or

$$\frac{3n}{5} + \frac{3+u}{5} \leq p < \frac{3n}{4} + \frac{3+u}{4}, p \equiv 2u \pmod{4}.$$

We check that this is the case except when

$$\begin{aligned} u = -1 : n = 15 \\ u = -2 : n \in \{19\} \\ u = 1 : n \in \{18, 19\} \\ u = 2 : n \in \{14, 15, 31\}. \end{aligned}$$

For these values of $\frac{u}{v}$ and n , we check in *MAGMA* that Galois group of $L^{(\frac{u}{v})}(x)$ is S_n .

Let $\alpha = \frac{u}{v} \in \{\pm\frac{1}{4}, \pm\frac{3}{4}\}$. Since $r \in \{1, 3\}$, we need to find a prime p with

$$\frac{4n}{5} + \frac{4+u}{5} \leq p \leq n-3, p \equiv u \pmod{4}$$

or

$$\frac{4n}{7} + \frac{4+u}{7} \leq p \leq n-3, p \equiv 3u \pmod{4}.$$

Hence it suffices to find a prime p with $\frac{4n}{5} + \frac{4+u}{5} \leq p \leq n-3$ or

$$\frac{4n}{7} + \frac{4+u}{7} \leq p < \frac{4n}{5} + \frac{4+u}{5}, p \equiv 3u \pmod{4}.$$

We check that this is the case except when

$$\begin{aligned} u = -1 : n \in \{14, 15, 30, 31\} \\ u = -3 : n \in \{20, 21, 23\} \\ u = 1 : n = 19 \\ u = 3 : n \in \{14, 29, 30, 31\}. \end{aligned}$$

For these values of $\frac{u}{v}$ and n , we check in *MAGMA* that Galois group of $L^{(\frac{u}{v})}(x)$ is S_n . This completes the proof of Theorem 1.1. \square

Acknowledgements

We thank the anonymous referee for the remarks in an earlier version of this paper. We also thank DST for supporting this work under the Fast Track Project.

References

- [FKT12] M. Filaseta, T. Kidd and O. Trifonov, *Laguerre polynomials with Galois group A_m for each m* , J. Number Theory, **132** (2012), no. 4, 776–805.
- [Haj05] F. Hajir, *On the Galois group of generalized Laguerre polynomials*, J. Théor. Nombres Bordeaux, **17**(2) (2005), 517–525.
- [LaSh] S. Laishram and T. N. Shorey, *Irreducibility of generalized Hermite–Laguerre polynomials III*, Submitted.
- [SaSh15] N. Saradha and T. N. Shorey, *Squares in blocks from an arithmetic progression and Galois group of Laguerre polynomials*, Int. Jour. of Number Theory, **11**(1) (2015), 233–250.
- [Sch29] I. Schur, *Einige Sätze über Primzahlen mit Anwendungen auf Irreduzibilitätsfragen, II*, Sitzungsber. Preuss. Akad. Wiss. Berlin Phys.-Math. Kl., **14** (1929), 370–391.
- [Sch31] I. Schur, *Affektlose Gleichungen in der Theorie der Laguerreschen und Hermitschen Polynome*, J. Reine Angew. Math., **165** (1931), 52–58.

Shanta Laishram
Stat-Math Unit, Indian Statistical Institute
7 S. J. S. Sansanwal Marg
New Delhi, 110016, India.
e-mail: shanta@isid.ac.in