



Comparing bottom-up and top-down approaches at the landscape scale, including agricultural activities and water systems, at the Roskilde Fjord, Denmark

Emeline Lequy (1), Andreas Ibrom (2), Per Ambus (2), Raia-Silvia Massad (1), Stiig Markager (3), Eero Asmala (3), Josette Garnier (4), Benoit Gabrielle (1), and Benjamin Loubet (1)

(1) INRA, UMR1091 INRA-AgroParisTech Ecosys, F-78850 Thiverval-Grignon, France (emeline.lequy@grignon.inra.fr), (2) Tech Univ Denmark, Dept Chem & Biochem Engn, Ctr Ecosyst & Environm Sustainabil, DK-4000 Roskilde, Denmark, (3) Aarhus Univ, Dept Biosci, DK-4000 Roskilde, Denmark, (4) Univ Paris 06, CNRS, UMR Sisyphe 7619, F-75005 Paris, France

The greenhouse gas nitrous oxide (N₂O) mainly originates in direct emissions from agricultural soils due to microbial reactions stimulated by the use of nitrogen fertilisers. Indirect N₂O emissions from water systems due to nitrogen leaching and deposition from crop fields range between 26 and 37% of direct agricultural emissions, indicating their potential importance and uncertainty (Reay et al. 2012). The study presented here couples a top-down approach with eddy covariance (EC) and a bottom-up approach using different models and measurements. A QCL sensor at 96-m height on a tall tower measures the emissions of N₂O from 1100 ha of crop fields and from the south part of the Roskilde fjord, in a 5-km radius area around the tall tower at Roskilde, Denmark. The bottom-up approach includes ecosystem modelling with CERES-EGC for the crops and PaSIM for the grasslands, and the N₂O fluxes from the Roskilde fjord are derived from N₂O sea water concentration measurements.

EC measurements are now available from July to December 2014, and indicate a magnitude of the emissions from the crop fields around 0.2 mg N₂O-N m⁻² day⁻¹ (range -9 to 5) which is consistent with the CERES-EGC simulations and calculations using IPCC emission factors. N₂O fluxes from the Roskilde fjord in May and July indicated quite constant N₂O concentrations around 0.1 μg N L⁻¹ despite variations of nitrate and ammonium in the fjord. The calculated fluxes from these concentrations and the tall tower measurements consistently ranged between -7 and 6 mg N₂O-N m⁻² day⁻¹. The study site also contains a waste water treatment plant, whose direct emissions will be measured in early 2015 using a dynamic plume tracer dispersion method (Mønster et al. 2014). A refined source attribution methodology together with more measurements and simulations of the N₂O fluxes from the different land uses in this study site will provide a clearer view of the dynamics and budgets of N₂O at the regional scale. The complementarity between these bottom-up and top-down approaches and their usefulness to disentangle direct and indirect N₂O fluxes will also be discussed.

Acknowledgements:

This work was funded by the EU-FP7 InGOS project.

References:

- Mønster JG, Samuelsson J, Kjeldsen P, Rella CW, Scheutz C. Quantifying methane emission from fugitive sources by combining tracer release and downwind measurements – A sensitivity analysis based on multiple field surveys. *Waste Management*. 2014 Aug;34(8):1416–28.
- Reay DS, Davidson EA, Smith KA, Smith P, Melillo JM, Dentener F, et al. Global agriculture and nitrous oxide emissions. *Nature Clim Change*. 2012 Jun;2(6):410–6.