



HAL
open science

Introduction of a carboxymethylamino (or oxy or thio) group in the 3 position of 2-aryl (or arylmethyl) isoindol-1-ones from alpha hydroxylactams

Pascal Pigeon, Bernard Decroix

► To cite this version:

Pascal Pigeon, Bernard Decroix. Introduction of a carboxymethylamino (or oxy or thio) group in the 3 position of 2-aryl (or arylmethyl) isoindol-1-ones from alpha hydroxylactams. *Synthetic Communications*, 1997, 27 (8), pp.1423-1431. 10.1080/00397919708006073 . hal-01230420

HAL Id: hal-01230420

<https://hal.science/hal-01230420>

Submitted on 20 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INTRODUCTION OF A CARBOXYMETHYLAMINO(or OXY or THIO)
GROUP IN THE 3 POSITION OF 2-ARYL(or ARYLMETHYL)
ISOINDOL-1-ONES FROM α HYDROXYLACTAMS


Pascal Pigeon and Bernard Decroix*

Laboratoire de Chimie, Université du Havre, 30, rue Gabriel Péri, 76600 Le Havre, France

*To whom correspondence should be addressed

Abstract: The preparation of 3-carboxymethylamino (or oxy or thio)-2-aryl (or arylmethyl)-isindol-1-ones are described from hydroxylactams according to three different pathways.

Tetracyclic systems as **1** incorporating an isoindole moiety with its nitrogen atom as one of the two junction atoms are widely expanded. The isoindolo[1,2-b][3]benzazepine **1** ($n=2$, $m=0$, $X=CH_2$, ring D=benzene) belonging to the aporphoadane alkaloid serie^{1,2} is one example. Compounds with a second heteroatom ($X=N$, O, S) in the C ring are few reported. Structures with a second nitrogen atom ([1,3] or [2,4]benzodiazepines, $X=N$, $n=0, 1$, $m=1, 2$) are described³⁻⁸ but we have found one report⁹ concerning compounds with an oxygen atom ([3,1]benzoxazepine, $X=O$, $n=0$, $m=2$) and no report with a sulfur atom. Similarly, isoindolobenzodiazocine ($X=NH$, $n+m=3$), oxazocine ($X=O$, $n+m=3$) or thiazocine ($X=S$, $n+m=3$) are just now unknown.




$n = 0, 1, 2$ $m = 0, 1, 2$
 $X = O, S, NH, CH_2$
ring D = benzene, thiophene

As a further development of our search on tetracyclic systems we hope to synthesize polycyclic systems as **1** (X=O, S, NH, n=1, m=2). For this purpose, substituted isoindolones **6**, **7** or **8** could be the precursors and we wish to describe herein their synthesis.


Since it is well known that hydroxylactam can give a substitution product¹⁰ under an acidic catalyze in the presence of a nucleophile via a *N*-acyliminium ion, we tried three nucleophiles: glycine, glycolic acid and thioglycolic acid and their ethyl esters. Thus (Scheme 1), only the thioglycolic acid (or its ethyl ester) have reacted with the previously reported hydroxylactams **2a-d**^{11,12} (n=0, 1, Ar=Ph, thien-2(3)-yl) under these conditions of reaction. The substitution products **3b** or **6a-d** were obtained in excellent yields (81 to 94%). Glycolic acid, glycine and their corresponding esters did not react or degradation products were observed. A related compound substituted with a furfuryl group had been prepared from methyl 2-formylbenzoate, furfurylamine and methyl thioglycolate¹³. Nevertheless other aminolactams or thioalkyl(aryl)lactams could be obtained using aniline, *p*-methylaniline or isopropanethiol, thiophenol as nucleophiles¹⁴.

The requisite carboxymethoxy esters derivatives **4a-d** (X=O) were prepared by an alkylation of the hydroxylactam. Actually, compound **2a-d** treated with butyllithium in tetrahydrofuran followed by the action of ethyl bromoacetate dissolved in hexamethylphosphotriamide (without HMPA, the reaction did not occur) afforded the expected ether derivatives **4a-d**.

The amino derivatives **5a-d** (X=NH) were synthesized according to Scheme 2. First we prepared 3-amino-2-alkyl(aryl)isoindol-1-ones **10a-d** via the chlorolactams **9a-d**.



Scheme 1



Scheme 2

The hydroxylactams **2a-d** were treated with thionyl chloride in dichloromethane and the resulting chlorolactams **9a-d** submitted to the action of ammonia in dichloromethane led to the 3-amino derivatives **10a-d** in good yield (72% to 81%). Alkylation of the latter using ethyl bromoacetate gave the expected alkylated products **5a-d**. It is noteworthy that the direct amination using ethyl glycinate upon the chlorolactams **9a-d** did not occur. On the other hand amine **5d** could be prepared by the reaction of 2-cyanobenzaldehyde with ammonia followed by a selective *N*-alkylation of the resulting amide in the presence of a phase transfer catalyst¹⁵. All the esters **3, 4, 5a-d** are viscous oils and their saponification was accomplished using potassium carbonate in aqueous methanol giving the corresponding carboxylic acids **6, 7, 8a-d** as crystalline products.

The microanalysis and spectroscopic data (IR, ¹H and ¹³C NMR) confirmed the assigned structures for the various compounds obtained (details are reported in the experimental part). These results indicate that direct conversion of hydroxylactams **2a-d** to thio derivatives **3a-d** was very efficient via a *N*-acyliminium ion, while an alkylation was necessary to obtain the oxy derivatives **4a-d**, and an alkylation of the amino compounds **10a-d**, via a chlorolactam intermediate, was essential to get the alkyamino derivatives **5a-d**. All of these compounds are possible precursors of polycyclic structures and cyclization reactions are under investigation.

EXPERIMENTAL

Melting points are uncorrected. The infrared spectra of solids (potassium bromide) were recorded on a Perkin Elmer FTIR paragon 1000 spectrometer. The ¹H and ¹³C NMR spectra were recorded on a Bruker AC-200 (200 MHz) instrument in deuteriochloroform

solution and chemical shift (δ) are expressed in ppm relative to internal TMS. Ascending thin layer chromatography was performed on precoated plates of silica gel 60 F 254 (Merck) and the spots visualized using an ultraviolet lamp or iodine vapor. E. Merck silica gel 60 F (70-300 mesh) was used for column chromatography. The elemental analyses were carried out by the microanalysis laboratory of INSA at Rouen, F 76130 M^t. S^t. Aignan, France.

Preparation of acids 6a-d : general procedure.

Hydroxylactams **2a-d** (10 mmol) and thioglycolic acid (0.83 ml, 1.1 g, 12 mmol) were refluxed overnight in acetic acid. The solvent was evaporated and the solid was triturated with ether, filtered, then recrystallized from acetone.

3-(Carboxymethylthio)-2,3-dihydro-2-phenyl-1H-isoindol-1-one (6a). Yield 81%; mp 188°C; IR: 3064 (OH), 1732 (C=O), 1668 (C=O) cm⁻¹; ¹H NMR: δ 2.75 (s, 2H, SCH₂), 4.80 (s, 1H, OH), 6.25 (s, 1H, CH), 7.24-7.34 (m, 1H, H_{arom}), 7.39-7.60 (m, 5H, H_{Ph}), 7.63-7.73 (m, 2H, H_{arom}), 7.90 (d, 1H, H_{arom}, J = 7.5 Hz). Anal. Calcd. for C₁₆H₁₃NO₃S: C, 64.20; H, 4.38; N, 4.68. Found: C, 63.97; H, 4.26; N, 4.55.

3-(Carboxymethylthio)-2,3-dihydro-2-(thien-2'-ylmethyl)-1H-isoindol-1-one (6b). Yield 94%; mp 201°C; IR: 3071 (OH), 1732 (C=O), 1670 (C=O) cm⁻¹; ¹H NMR (DMSO-d₆) : δ 2.61 (d, 1H, SCH₂, J = 15.3 Hz), 2.84 (d, 1H, SCH₂, J = 15.3 Hz), 3.65 (s, 1H, OH), 4.64 (d, 1H, NCH₂, J = 15.6 Hz), 5.12 (d, 1H, NCH₂, J = 15.6 Hz), 5.74 (s, 1H, CH), 6.97 (dd, 1H, H_{4'}, J = 4.4, 3.1 Hz), 7.08 (d, 1H, H_{3'}, J = 3.1 Hz), 7.43 (d, 1H, H_{5'}, J = 4.4 Hz), 7.45-7.80 (m, 4H, H_{arom}). Anal. Calcd. for C₁₅H₁₃NO₃S₂: C, 56.41; H, 4.10; N, 4.39. Found: C, 56.27; H, 4.06; N, 4.31.

3-(Carboxymethylthio)-2,3-dihydro-2-(thien-3'-ylmethyl)-1H-isoindol-1-one (6c). Yield 87%; mp 159°C; IR: 3100 (OH), 1730 (C=O), 1669 (C=O) cm⁻¹; ¹H NMR (DMSO-d₆) : δ 2.59 (d, 1H, SCH₂, J = 15.5 Hz), 2.82 (d, 1H, SCH₂, J = 15.5 Hz), 3.67 (s, 1H, OH), 4.48 (d, 1H, NCH₂, J = 15.3 Hz), 4.99 (d, 1H, NCH₂, J = 15.3 Hz), 5.66 (s, 1H, CH), 7.01 (d, 1H, H_{4'}, J = 4.6 Hz), 7.30-7.80 (m, 6H, 4H_{arom}+H_{2',5'}). Anal. Calcd. for C₁₅H₁₃NO₃S₂: C, 56.41; H, 4.10; N, 4.39. Found: C, 56.22; H, 4.04; N, 4.35.

3-(Carboxymethylthio)-2,3-dihydro-2-(phenylmethyl)-1H-isoindol-1-one (6d). Yield 93%; mp 202°C; IR: 3063 (OH), 1732 (C=O), 1674 (C=O) cm⁻¹; ¹H NMR: δ 2.60 (d, 1H, SCH₂, J = 15.6 Hz), 2.75 (d, 1H, SCH₂, J = 15.6 Hz), 3.46 (s, 1H, OH), 4.39 (d, 1H, NCH₂, J = 15.1 Hz), 5.32 (d, 1H, NCH₂, J = 15.1 Hz), 5.34 (s, 1H, CH), 7.29 (s, 5H, H_{Ph}), 7.40-7.64 (m, 3H, H_{arom}), 7.87 (d, 1H, H_{arom}, J = 7.0 Hz). Anal. Calcd. for C₁₇H₁₅NO₃S: C, 65.16; H, 4.82; N, 4.47. Found: C, 64.88; H, 4.64; N, 4.35.

3-(Ethoxycarbonylmethylthio)-2,3-dihydro-2-(thien-2'-ylmethyl)-1H-isoindol-1-one (3b).

Hydroxylactam **2b** (2.45 g, 10 mmol), ethyl thioglycolate (1.44 g, 12 mmol) and a catalytic amount of paratoluenesulfonic acid were stirred overnight in dry dichloromethane. The solvent was evaporated to dryness and the residue was dissolved in dichloromethane. The organic layer was washed with a saturated solution of sodium hydrogenocarbonate, then dried and evaporated. The ester was recrystallized from ethanol. Yield 93%; mp 77°C; IR: 1746 (C=O), 1703 (C=O) cm^{-1} ; ^1H NMR: δ 1.13 (t, 3H, CH_3 , $J = 7.1$ Hz), 2.59 (d, 1H, SCH_2 , $J = 15.3$ Hz), 2.71 (d, 1H, SCH_2 , $J = 15.3$ Hz), 4.04 (q, 2H, OCH_2 , $J = 7.1$ Hz), 4.60 (d, 1H, NCH_2 , $J = 15.3$ Hz), 5.38 (d, 1H, NCH_2 , $J = 15.3$ Hz), 5.45 (s, 1H, CH), 6.92 (dd, 1H, H_4 , $J = 5.1, 3.5$ Hz), 7.05 (d, 1H, H_3 , $J = 3.5$ Hz), 7.19 (d, 1H, H_5 , $J = 5.1$ Hz), 7.40-7.63 (m, 3H, H_{arom}), 7.84 (d, 1H, H_{arom} , $J = 7.0$ Hz).

Preparation of aminolactams 10a-d : general procedure.

A solution of ammonia in dichloromethane was prepared by extraction of 400 ml of concentrated aqueous ammonia solution with 400 ml of dichloromethane. The aqueous layer was kept in the separatory funnel for later use and the solution of ammonia in dichloromethane was dried over magnesium sulfate then filtered.

Hydroxylactams **2a-d** (10 mmol) and thionyl chloride (1.5 ml, 21 mmol) were stirred in dry dichloromethane until all solid had disappeared, then reaction was continued for 30 minutes. This solution was poured into the previous solution of ammonia in dichloromethane and the mixture was stirred for 10 min. The solution was transferred into the separatory funnel containing the previous aqueous ammonia, then the mixture was made more basic with addition of sodium hydroxide solution. The organic layer was decanted then extracted twice with 5% HCl solution. The aqueous solutions were combined, washed with dichloromethane, made basic with sodium hydroxide solution then extracted twice with dichloromethane. The combination of organic layers was dried and evaporated. The solid was recrystallized from ethanol.

3-Amino-2,3-dihydro-2-phenyl-1H-isoindol-1-one (10a). Yield 78%; mp 112°C; IR: 3396 (NH_2), 1686 (C=O) cm^{-1} ; ^1H NMR: δ 1.90 (s, 2H, NH_2), 5.94 (s, 1H, CH), 7.17-7.30 (m, 1H, H_{arom}), 7.38-7.73 (m, 7H, $5\text{H}_{\text{Ph}}+2\text{H}_{\text{arom}}$), 7.88 (d, 1H, H_{arom} , $J = 7.0$ Hz). Anal. Calcd. for $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}$: C, 74.98; H, 5.39; N, 12.49. Found: C, 74.57; H, 5.31; N, 12.25.

3-Amino-2,3-dihydro-2-(thien-2'-ylmethyl)-1H-isoindol-1-one (10b). Yield 76%; mp 113°C; IR: 3373 (NH_2), 1667 (C=O) cm^{-1} ; ^1H NMR: δ 2.54 (s, 2H, NH_2), 4.63 (d, 1H, NCH_2 , $J = 15.5$ Hz), 5.21 (s, 1H, CH), 5.26 (d, 1H, NCH_2 , $J = 15.5$ Hz), 6.92 (dd, 1H, H_4 , $J = 5.1, 3.4$ Hz), 7.02 (d, 1H, H_3 , $J = 3.4$ Hz), 7.18 (dd, 1H, H_5 , $J = 5.1, 1.3$ Hz), 7.40-7.60 (m, 3H,

H_{arom}), 7.80 (d, 1H, H_{arom}, J = 7.0 Hz). Anal. Calcd. for C₁₃H₁₂N₂OS: C, 63.91; H, 4.95; N, 11.47. Found: C, 63.67; H, 4.86; N, 11.31.

3-Amino-2,3-dihydro-2-(thien-3'-ylmethyl)-1H-isoindol-1-one (10c). Yield 81%; mp 118°C; IR: 3372 (NH₂), 1665 (C=O) cm⁻¹; ¹H NMR: δ 1.74 (s, 2H, NH₂), 4.48 (d, 1H, NCH₂, J = 15.2 Hz), 5.09 (d, 1H, NCH₂, J = 15.2 Hz), 5.15 (s, 1H, CH), 7.03 (d, 1H, H_{4'}, J = 4.9 Hz), 7.13-7.33 (m, 2H, H_{2',5'}), 7.41-7.62 (m, 3H, H_{arom}), 7.82 (d, 1H, H_{arom}, J = 6.7 Hz). Anal. Calcd. for C₁₃H₁₂N₂OS: C, 63.91; H, 4.95; N, 11.47. Found: C, 63.61; H, 4.89; N, 11.52.

3-Amino-2,3-dihydro-2-(phenylmethyl)-1H-isoindol-1-one (10d). Yield 72%; mp 124°C; IR: 3378 (NH₂), 1676 (C=O) cm⁻¹; ¹H NMR: δ 1.70 (s, 2H, NH₂), 4.36 (d, 1H, NCH₂, J = 15.3 Hz), 5.07 (s, 1H, CH), 5.10 (d, 1H, NCH₂, J = 15.3 Hz), 7.22 (s, 5H, H_{Ph}), 7.36-7.52 (m, 3H, H_{arom}), 7.78 (d, 1H, H_{arom}, J = 6.7 Hz). Anal. Calcd. for C₁₅H₁₄N₂O: C, 75.61; H, 5.92; N, 11.76. Found: C, 75.65; H, 5.90; N, 11.68.

Preparation of glycine derivatives 8a-d

Aminolactams **10a-d** (10 mmol), potassium carbonate (6.4 g, 20 mmol), ethyl bromoacetate (2.2 ml, 20 mmol) and 20 ml of dry 1,4 dioxane were refluxed for 5 days. After removal of the solvent, water and dichloromethane were added to the residue. The organic layer was washed with 5% aqueous hydrochloric solution (no extraction of desired product), washed with water, dried, then evaporated. The residue was chromatographed with dichloromethane to afford esters **5a-d** as an oil. These crude esters, potassium carbonate (2.4 g), water (6 ml) and methanol (24 ml) were stirred under reflux for 2 hours, then the solution was concentrated under reduced pressure. Water and dichloromethane were added and the organic layer was discarded. The aqueous layer was washed with dichloromethane and acidified with hydrochloric acid (10%) to pH=2. Compounds **8a-d** were rapidly extracted with dichloromethane before crystallization occurs. After removal of the solvent, the residue was recrystallized from acetone to give pure **8a-d**.

3-(Carboxymethylamino)-2,3-dihydro-2-phenyl-1H-isoindol-1-one (8a).

Yield: 55%; mp 162°C; IR: 3365 (NH), 2914 (OH), 1727 (C=O), 1651 (C=O) cm⁻¹; ¹H NMR: δ 1.83 (m, 2H, OH+NH), 2.86 (d, 1H, CH₂COO, J = 17.7 Hz), 3.04 (d, 1H, CH₂COO, J = 17.7 Hz), 6.14 (s, 1H, CH), 7.22-7.69 (m, 8H, 3H_{arom}+5H_{Ph}), 7.92 (d, 1H, H_{arom}, J = 7.0 Hz). Anal. Calcd. for C₁₆H₁₄N₂O₃: C, 68.08; H, 5.00; N, 9.92. Found: C, 67.75; H, 5.12; N, 10.07.

3-(Carboxymethylamino)-2,3-dihydro-2-(thien-2'-ylmethyl)-1H-isoindol-1-one (8b).

Yield: 54%; mp 125°C; IR: 3422 (NH), 2943 (OH), 1718 (C=O, C=O) cm⁻¹; ¹H NMR: δ 2.93 (m, 2H, OH+NH), 2.97 (s, 2H, CH₂COO), 4.56 (d, 1H, NCH₂, J = 15.3 Hz), 5.20 (d, 1H, NCH₂, J = 15.3 Hz), 5.42 (s, 1H, CH), 6.92 (dd, 1H, H_{4'}, J = 3.8, 5.0 Hz), 7.04 (d, 1H, H_{3'}, J =

3.0 Hz), 7.21 (d, 1H, H_{5'}, J = 5.0 Hz), 7.43-7.62 (m, 3H, 3H_{arom}), 7.84 (d, 1H, H_{arom}, J = 7.0 Hz). Anal. Calcd. For C₁₅H₁₄N₂O₃S: C, 59.59; H, 4.67; N, 9.27. Found: C, 59.34; H, 4.32; N, 9.12.

3-(Carboxymethylamino)-2,3-dihydro-2-(thien-3'-ylmethyl)-1H-isoindol-1-one (8c).

Yield: 73%; mp 112°C; IR: 3386 (OH+NH), 1686 (C=O, C=O) cm⁻¹; ¹H NMR: δ 2.90 (s, 2H, CH₂COO), 4.38 (d, 1H, NCH₂, J = 15.1 Hz), 5.01 (d, 1H, NCH₂, J = 15.1 Hz), 5.35 (s, 1H, CH), 7.07 (dd, 1H, H_{4'}, J = 1.6, 4.5 Hz), 7.15-7.31 (m, 2H, H_{2',5'}), 7.39-7.59 (m, 3H, H_{arom}), 7.69 (d, 1H, H_{arom}, J = 7.0 Hz). Anal. Calcd. for C₁₅H₁₄N₂O₃S: C, 59.59; H, 4.67; N, 9.27. Found: C, 59.41; H, 4.55; N, 9.13.

3-(Carboxymethylamino)-2,3-dihydro-2-(phenylmethyl)-1H-isoindol-1-one (8d).

Yield: 54%; mp 170°C; IR: 2903 (OH+NH), 1614 (C=O, C=O) cm⁻¹; ¹H NMR: δ 2.92 (s, 2H, CH₂COO), 4.27 (d, 1H, NCH₂, J = 15.0 Hz), 5.10 (d, 1H, NCH₂, J = 15.0 Hz), 5.32 (s, 1H, CH), 6.02 (s, 2H, OH+NH), 6.89-7.68 (m, 8H, 3H_{arom}+5H_{Ph}), 7.74-7.88 (m, 1H, H_{arom}). Anal. Calcd. for C₁₇H₁₆N₂O₃: C, 68.91; H, 5.44; N, 9.45. Found: C, 68.69; H, 5.64; N, 9.77.

Preparation of glycolic derivatives 7a-d

Hydroxylactams **2a-d** (10 mmol) were dissolved in 50 ml of dry tetrahydrofuran and BuLi (7 ml, 1.6 M, 1.1 eq) was added with stirring. Then a solution of 1.7 ml (15 mmol) of ethyl bromoacetate in 10 ml of dry hexamethylphosphoramide was added. Stirring was continued overnight, then the solution was poured into water then extracted with dichloromethane. The organic layer was washed with water then evaporated. The oil was triturated several times with water and decanted. The oil was dried under reduced pressure then chromatographed with dichloromethane. The ester **4a-d** was obtained as an oil. Saponification of esters **4a-d** was performed in a similar manner as above and led to acids **7a-d** which were recrystallized from acetone.

3-(Carboxymethyloxy)-2,3-dihydro-2-phenyl-1H-isoindol-1-one (7a).

Yield 56%; mp 163°C; IR: 3069 (OH), 1758 (C=O), 1708 (C=O) cm⁻¹; ¹H NMR: δ 3.60 (d, 1H, OCH₂, J = 16.4 Hz), 3.82 (d, 1H, OCH₂, J = 16.4 Hz), 4.21 (s, 1H, OH), 6.60 (s, 1H, CH), 7.19-7.27 (m, 1H, H_{arom}), 7.38-7.47 (m, 2H, H_{arom}), 7.54-7.78 (m, 5H, H_{Ph}), 7.92 (d, 1H, H_{arom}, J = 7.0 Hz). Anal. Calcd. for C₁₆H₁₃NO₄: C, 67.84; H, 4.63; N, 4.85. Found: C, 67.42; H, 4.34; N, 4.85.

2,3-Dihydro-3-(carboxymethyloxy)-2-(thien-2'-ylmethyl)-1H-isoindol-1-one (7b).

Yield 60%; mp 136°C; IR: 3113 (OH), 1737 (C=O), 1682 (C=O) cm⁻¹; ¹H NMR: δ 3.60 (d, 1H, OCH₂, J = 16.4 Hz), 3.71 (d, 1H, OCH₂, J = 16.4 Hz), 4.60 (d, 1H, NCH₂, J = 15.3 Hz), 5.15 (d, 1H, NCH₂, J = 15.3 Hz), 5.96 (s, 1H, CH), 6.88 (dd, 1H, H_{4'}, J = 5.1, 3.3 Hz), 7.06 (d, 1H, H_{5'}, J = 3.3 Hz), 7.16 (d, 1H, H_{5'}, J = 5.1 Hz), 7.40-7.65 (m, 3H, H_{arom}), 7.84 (d, 1H, H_{arom}, J =

5.4 Hz), 8.69 (s, 1H, OH). Anal. Calcd. for C₁₅H₁₃NO₄S: C, 59.40; H, 4.32; N, 4.62. Found: C, 59.25; H, 4.28; N, 4.60.

3-(Carboxymethoxy)-2,3-dihydro-2-(thien-3'-ylmethyl)-1H-isoindol-1-one (7c). Yield 49%; mp 149°C; ¹H NMR: δ 3.53 (d, 1H, OCH₂, J = 16.2 Hz), 3.63 (d, 1H, OCH₂, J = 16.2 Hz), 4.42 (d, 1H, NCH₂, J = 14.9 Hz), 4.95 (d, 1H, NCH₂, J = 14.9 Hz), 5.86 (s, 1H, CH), 7.04 (d, 1H, H_{4'}, J = 3.8 Hz), 7.10-7.25 (m, 2H, H_{2',5'}), 7.40-7.60 (m, 3H, H_{arom}), 7.81 (d, 1H, H_{arom}, J = 5.4 Hz), 7.93 (s, 1H, OH). Anal. Calcd. for C₁₅H₁₃NO₄S: C, 59.40; H, 4.32; N, 4.62. Found: C, 59.34; H, 4.30; N, 4.65.

3-(Carboxymethoxy)-2,3-dihydro-2-(phenylmethyl)-1H-isoindol-1-one (7d). Yield 66%; mp 115°C; IR: 3447 (OH), 1734 (C=O), 1683 (C=O) cm⁻¹; ¹H NMR: δ 3.54 (d, 1H, OCH₂, J = 16.5 Hz), 3.66 (d, 1H, OCH₂, J = 16.5 Hz), 4.37 (d, 1H, NCH₂, J = 14.9 Hz), 5.05 (d, 1H, NCH₂, J = 14.9 Hz), 5.25 (s, 1H, OH), 5.87 (s, 1H, CH), 7.17-7.42 (m, 5H, H_{Ph}), 7.48-7.62 (m, 3H, H_{arom}), 7.82-7.92 (m, 1H, H_{arom}). Anal. Calcd. for C₁₇H₁₅NO₄: C, 68.68; H, 5.09; N, 4.71. Found: C, 68.47; H, 5.13; N, 4.77.

REFERENCES AND NOTES

- 1 Napolitano E., Spinelli G., Fiaschi R., Marsili A., *J. Chem. Soc., Perkin Trans. I*, **1986**, 785.
- 2 Lamas C., Saa C., Castedo L., Dominguez D., *Tetrahedron Lett.*, **1992**, 33, 5653.
- 3 Hatt H.H., Stephenson E.F.M., *J. Chem. Soc., Chem. Commun.*, **1952**, 199.
- 4 Stephenson E.F.M., *J. Chem. Soc., Chem. Commun.*, **1952**, 5024.
- 5 Eguchi S., Takeuchi H., *J. Chem. Soc., Chem. Commun.*, **1989**, 602.
- 6 Kurita J., Iwata T., Yasuike S., Tsuchiya T., *J. Chem. Soc., Chem. Commun.*, **1992**, 81.
- 7 Kurita J., Iwata T., Yasuike S., Tsuchiya T., *J. Chem. Soc., Chem. Commun.*, **1992**, 970.
- 8 Lessel J., *Pharmazie*, **1993**, 48, 812.
- 9 Baumann H., Fletcher I.J., *Eur. Pat. Appl. EP 475, 908 (18 march 1992)*, *Chem. Abstr.* **1992**, 117, 7947.
- 10 Speckamp W.N., *Recl. Trav. Chim., Pays-Bas*, **1981**, 100, 345.
- 11 Ishihara Y., Kiyota Y., Goto G., *Chem. Pharm. Bull.*, **1990**, 38, 3024.
- 12 Pigeon P., Decroix B., *J. Heterocyclic Chem.*, **1996**, 33, 129.
- 13 Liebl R., Handte R., *Liebigs Ann. Chem.*, **1985**, 1093.

- 14 Pigeon P., *Thèse-Université Le Havre, France, 1996.*
- 15 Sato R., Senzaki T., Goto T., Saito M., *Bull. Chem. Soc. Jpn.*, **1986**, 59, 2950.