



HAL
open science

Synthesis of dibenz[c,e]azepine and benzo[e]thieno[c]azepine via N-acyliminium ion cyclization

Pascal Pigeon, Bernard Decroix

► **To cite this version:**

Pascal Pigeon, Bernard Decroix. Synthesis of dibenz[c,e]azepine and benzo[e]thieno[c]azepine via N-acyliminium ion cyclization. *Tetrahedron Letters*, 1997, 38 (6), pp.1041-1042. 10.1016/S0040-4039(96)02497-5. hal-01230421

HAL Id: hal-01230421

<https://hal.science/hal-01230421>

Submitted on 4 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Tetrahedron Letters (1997), 38(6), 1041-1042.

DOI : [10.1016/S0040-4039\(96\)02497-5](https://doi.org/10.1016/S0040-4039(96)02497-5)

Synthesis of Dibenz[c,e]azepine and Benzo[e]thieno[c]azepine via *N*-Acyliminium ion Cyclization

Pascal Pigeon and Bernard Decroix*

~~Laboratoire de Chimie, Faculté des Sciences et des Techniques de l'Université du Havre,
30, rue Gabriel Péri, 76600 Le Havre, France~~

New address: Université du Havre, URCOM, 25 rue Philippe Lebon, BP 1123, 76063 Le Havre cedex, France

Abstract: Dibenz[c,e]azepine **4a** and benzo[e]thieno[c]azepine **4b** have been synthesized utilizing the acid-catalyzed cyclization of hydroxylactams **3a,b** as key step.


Over the last ten years many synthetic efforts have been directed toward synthesis of [2] or [3] benzazepines annelated to various rings (isoindole, pyrrolidine, benzene...) because a number of natural products as Chilenine ¹, Lennoxamine ², Cephalotaxine ³ contains this moiety and present potential biological activities. Any number derivatives of dibenz[c,e ; b,e ; b,f]azepines are known and present hypolipidemic activity ⁴, central nervous system activity ⁵, antiarrhythmic activity ⁶ respectively. With our interest in the synthesis of diversely substituted polycyclic systems ⁷⁻¹⁰ we wish to report herein an interesting approach to dibenzazepine or thienobenzazepine annelated to an isoindole moiety as **4a** or **4b**.

We have shown ¹¹ that 3-phenyl-2-(2'-bromothien-3'-ylmethyl)phthalimide subjected to the usual radical cyclization conditions (AIBN, Bu₃SnH, toluene) or intramolecular Heck reaction did not give the expected [2]benzazepine **4b**. On the other hand it was reported ¹² that cyclodehydration of *N*-(*o*-biphenyl)hydroxyphthalimidine under refluxing of trifluoroacetic acid occurred leading to a new isoindolophenanthridine.

From these results we suggested to use the hydroxylactam **3a** for the synthesis of the dibenzazepine derivative **4a**. *N*-acyliminium ion precursor **3a** was prepared from phthalimide derivative **1a**. A palladium (0) catalyzed cross coupling of bromo derivative **1a** with phenylboronic acid provided the biphenyl compound **2a** in good yield (76%). Reduction of

imide **2a** by sodium borohydride in the presence of acid ¹³, afforded the hydroxylactam **3a** (82%). This latter was then subjected to trifluoroacetic acid at room temperature and according to the Baldwin's rules ¹⁴ led to the expected formation of a 7-membered ring, the new dibenz[*c,e*]azepine **4a** annelated to an isoindole moiety in a yield of 95%. The structure of **4a** was supported by NMR (¹H, ¹³C) spectroscopic analyses ¹⁵. A study of the long range heteronuclear shift correlation (HETCOR) spectrum of **4a** further substantiated our conclusions. The spectrum showed seven different signals for the benzenic correlation. Such correlations would not be expected from the isomeric structure **5a**.

This promoted result allied to the fact that a *N*-acyliminium ion cyclization gave a 5 or 6-membered ring from the α position of thiophene and only a 6-membered ring from the β position of the thiophene ¹⁶ we tested the hydroxylactam **3b**. In similar conditions (CF₃COOH, room temperature), cyclization of **3b** gave a 7-membered ring (**4b**) as in the above benzene series rather than a 5-membered one (**5b**).



The structure of the thienobenzazepine **4b** was supported by NMR (¹H, ¹³C) spectroscopic analyses ¹⁷. The two protons of the thiophene ring appear as doublets (AB system) with chemical shifts of 7.13 ppm and 7.38 ppm and a coupling constant of 5 Hz characteristic of a α,β substituted thiophene.

The present investigation has thus led to a new method of synthesis of dibenzo[*c,e*]azepine or benzo[*e*]thieno[*c*]azepine annelated to an isoindole moiety starting from hydroxylactams.

REFERENCES AND NOTES

1. Mazzocchi, P.H.; King, C.R.; Ammon, H.L. *Tetrahedron Lett.*, **1987**, 28, 2473-2476.
2. Ishibashi, H.; Kawanami, H.; Iriyama, H.; Ikeda, M. *Tetrahedron Lett.*, **1995**, 36, 6733-6734.
3. Weinreb, S.M.; Auerbach, J. *J. Am. Chem. Soc.*, **1975**, 97, 2503-2506.
4. Hall, I.; Murthy, A.R.K.; Wyrick, S.D. *J. Pharm. Sci.*, **1986**, 75, 622-626.
5. Steiner, G.; Franke, A.; Haedicke, E.; Lenke, D.; Teschendorf, H.; Hofmann, H.; Kreiskott, H.; Werstmann, W. *J. Med. Chem.*, **1986**, 29, 1877-1888.
6. Wunderlich, H.; Strak, A.; Carstens, E.; Lohmann, D.; Gritsenko, A.N.; Skoldinov, A.P. *Pharmazie*, **1985**, 40, 827-830.
7. Marchalin, S.; Decroix, B. *Heterocycles*, **1995**, 41, 689-696.
8. Daich, A.; Ohier, P.; Decroix, B. *Tetrahedron Lett.*, **1995**, 36, 83-86.
9. Pigeon, P.; Decroix, B. *J. Heterocyclic Chem.*, **1996**, 33, 129-135.
10. Othman, M.; Decroix, B. *Synth. Commun.*, **1996**, 26, 2803-2809.
11. Pigeon, P., PhD University Le Havre, 1996.
12. Hitchings, G.J.; Heliwell, M.; Vernon, J.M. *J. Chem. Soc. Perkin Trans. I*, **1990**, 83-87.
13. Wijnberg, J.B.P.A.; Schoemaker, H.J.; Speckamp, W.N. *Tetrahedron*, **1978**, 34, 179-187.
14. Baldwin, J.E. *J. Chem. Soc. Chem. Comm.*, **1976**, 734-736.
15. The HETCOR spectrum of **4a** was acquired using the standard Bruker microprogram XHCORR.AU (Bax, A.; Morris, G. *J. Mag. Res.*, **1981**, 42, 501). Yield: 95%; mp: 196°C; IR: 1686 cm⁻¹ (C=O); ¹H NMR (CDCl₃): δ 3.89 (d, 1H, NCH₂, J = 14 Hz), 5.05 (d, 1H, NCH₂, J = 14 Hz), 5.31 (s, 1H, CH), 7.05 (d, 1H, H_{arom}, J = 7 Hz), 7.28 (t, 1H, H_{arom}, J = 7 Hz), 7.31-7.72 (m, 9H, H_{arom}), 7.92 (d, 1H, H_{arom}, J = 7 Hz). ¹³C NMR: δ 44.2 (CH₂), 60.2 (CH), 123.5 (CH), 123.8 (CH), 124.8 (CH), 128.0 (CH), 128.1 (2CH), 128.4 (CH), 128.5 (CH), 128.8 (2CH), 128.9 (CH), 130.6 (CH), 133.2 (C), 133.4 (C), 133.7 (C), 140.0 (C), 140.0 (C), 141.3 (C), 165.1 (CO).
16. Othman, M.; Pigeon, P.; Decroix, B. *Tetrahedron*, **1997**, 53(7), 2495-2504.
17. Yield: 88%; mp: 241°C; IR: 1684 cm⁻¹ (C=O); ¹H NMR (CDCl₃): δ 4.07 (d, 1H, NCH₂, J = 15 Hz), 5.20 (d, 1H, NCH₂, J = 15 Hz), 5.60 (s, 1H, CH), 6.99-7.77 (m, 7H, H_{arom}), 7.13 (d, 1H, H_βthiophene, J = 5 Hz), 7.38 (d, 1H, H_αthiophene, J = 5 Hz), 7.95 (d, 1H, H_{arom}, J = 6 Hz).

