



## Thin linearly piezoelectric junctions

Christian Licht, Somsak Orankitjaroen, Patcharakorn Viriyasrisuwattana,  
Thibaut Weller

### ► To cite this version:

Christian Licht, Somsak Orankitjaroen, Patcharakorn Viriyasrisuwattana, Thibaut Weller. Thin linearly piezoelectric junctions. Comptes Rendus Mécanique, 2015, 343 (4), pp.282-288. 10.1016/j.crme.2015.01.001 . hal-01232788

HAL Id: hal-01232788

<https://hal.science/hal-01232788>

Submitted on 24 Nov 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Thin linearly piezoelectric junctions

## *Jonctions minces linéairement piézoélectriques*

Christian Licht <sup>a,b</sup>, Somsak Orankitjaroen <sup>b</sup>, Patcharakorn Viriyasrisuwattana <sup>b,c</sup>,  
Thibaut Weller <sup>a</sup>

<sup>a</sup> LMGC, UMR-CNRS 5508, Université Montpellier-2, case courier 048, place Eugène-Bataillon, 34095 Montpellier cedex 5, France

<sup>b</sup> Department of Mathematics, Faculty of Science, Mahidol University, Bangkok 10400, Thailand

<sup>c</sup> Department of Mathematics, Faculty of Science and Technology, Suan Sunandha Rajabhat University, Bangkok 10300, Thailand

### A B S T R A C T

Through a rigorous mathematical analysis, we present various asymptotic models for a thin piezoelectric junction between two linearly piezoelectric or elastic bodies. Depending on the relative behavior of a stiffness parameter with respect to its thickness, the joint is replaced by either a(n) (electro)mechanical constraint or a piezoelectric material surface.

### R É S U M É

Par une analyse mathématique rigoureuse, nous présentons divers modèles asymptotiques pour une jonction mince piézoélectrique entre deux corps linéairement élastiques ou piézoélectriques. Selon l'ordre de grandeur relatif entre un paramètre de rigidité et l'épaisseur, le joint est remplacé par une liaison (électro)mécanique ou par une surface matérielle piézoélectrique.

---

### 1. Introduction

We present various asymptotic models, indexed by  $p = (p_1, p_2) \in \{1, 2, 3, 4\}^2$ , for a thin piezoelectric junction between two linearly piezoelectric ( $p_2 = 1$ ) or elastic ( $p_2 > 1$ ) bodies. Index  $p_1$  is relative to the magnitude of the piezoelectric coefficients of the adhesive, characterized by a single parameter  $\mu$ , with respect to that of the constant thickness  $2\varepsilon$  of a layer containing the adhesive. More precisely, we assume that  $h := (\varepsilon, \mu)$  takes values in a countable set with a sole cluster point  $\bar{h} \in \{0\} \times [0, +\infty)$  so that:

$$\begin{cases} p_1 = 1 : \bar{\mu}_1 := \lim_{h \rightarrow \bar{h}} (\varepsilon \mu) \in (0, +\infty) \\ p_1 = 2 : \bar{\mu}_1 := \lim_{h \rightarrow \bar{h}} (\varepsilon \mu) = 0, \quad \bar{\mu}_2 := \lim_{h \rightarrow \bar{h}} (\mu / 2\varepsilon) = +\infty \\ p_1 = 3 : \bar{\mu}_2 := \lim_{h \rightarrow \bar{h}} (\mu / 2\varepsilon) \in (0, +\infty) \\ p_1 = 4 : \bar{\mu}_2 := \lim_{h \rightarrow \bar{h}} (\mu / 2\varepsilon) = 0. \end{cases} \quad (1)$$

---

E-mail addresses: [clicht@univ-montp2.fr](mailto:clicht@univ-montp2.fr) (C. Licht), [somsak.ora@mahidol.ac.th](mailto:somsak.ora@mahidol.ac.th) (S. Orankitjaroen), [vdplek@hotmail.com](mailto:vdplek@hotmail.com) (P. Viriyasrisuwattana), [Patcharakorn.vi@ssru.ac.th](mailto:Patcharakorn.vi@ssru.ac.th) (P. Viriyasrisuwattana), [thibaut.weller@univ-montp2.fr](mailto:thibaut.weller@univ-montp2.fr) (T. Weller).

As previously said, index  $p_2$  characterizes the status of the adherents but also that of the interfaces between adherents and adhesive:

$$\begin{cases} p_2 = 1 : \text{the two interfaces are electromechanically perfectly permeable,} \\ p_2 = 2 : \text{the two interfaces are electrically impermeable,} \\ p_2 = 3 : \text{one interface is electrically impermeable while the other is electroded,} \\ p_2 = 4 : \text{the two interfaces are electroded.} \end{cases} \quad (2)$$

The space  $\mathbb{R}^3$  is assimilated with the physical Euclidean space with basis  $\{e_1, e_2, e_3\}$ . Let  $\Omega$  be a domain, with Lipschitz-continuous boundary, whose intersection  $S$  with  $\{x_3 = 0\}$  is a domain of  $\mathbb{R}^2$  of positive two-dimensional Hausdorff measure  $\mathcal{H}_2(S)$ . Let  $\Omega_{\pm} := \Omega \cap \{\pm x_3 > 0\}$  and  $\varepsilon$  be a small positive number, then adhesive and adherents occupy  $B^{\varepsilon} := S \times (-\varepsilon, \varepsilon)$ ,  $\Omega_{\pm}^{\varepsilon} := \Omega_{\pm} \pm \varepsilon e_3$ , respectively; let  $\Omega^{\varepsilon} = \Omega_{+}^{\varepsilon} \cup \Omega_{-}^{\varepsilon}$ ,  $S_{\pm}^{\varepsilon} := S \pm \varepsilon e_3$ ,  $\mathcal{O}^{\varepsilon} := \Omega^{\varepsilon} \cup B^{\varepsilon} \cup_{\pm} S_{\pm}^{\varepsilon}$ . Let  $(\Gamma_{mD}, \Gamma_{mN})$ ,  $(\Gamma_{eD}, \Gamma_{eN})$  be two partitions of  $\partial\Omega$  with  $\mathcal{H}_2(\Gamma_{mD}), \mathcal{H}_2(\Gamma_{eD}) > 0$  and  $0 < \delta := \text{dist}(\Gamma_{eD}, S)$ . For all  $\Gamma$  in  $\{\Gamma_{mD}, \Gamma_{mN}, \Gamma_{eD}, \Gamma_{eN}\}$ ,  $\Gamma_{\pm}, \Gamma_{\pm}^{\varepsilon}, \Gamma^{\varepsilon}$  denotes  $\Gamma \cap \{\pm x_3 > 0\}$ ,  $\Gamma_{\pm} \pm \varepsilon e_3$ ,  $\cup_{\pm} \Gamma_{\pm}^{\varepsilon}$ , respectively; if  $(\gamma_D, \gamma_N)$  is a partition of  $\gamma := \partial S$ , we denote  $\{\gamma_D, \gamma_N, \gamma\} \times (-\varepsilon, \varepsilon)$  by  $\{\Gamma_{DI}^{\varepsilon}, \Gamma_{NI}^{\varepsilon}, \Gamma_{lat}^{\varepsilon}\}$ . The structure made of the adhesive and the two adherents, perfectly stuck together along  $S_{\pm}^{\varepsilon}$ , is clamped on  $\Gamma_{mD}^{\varepsilon}$ , subjected to body forces of density  $f^{\varepsilon}$  and to surface forces of density  $F^{\varepsilon}$  on  $\Gamma_{mN}^{\varepsilon}$  and vanishing on  $\Gamma_{lat}^{\varepsilon}$ . Moreover, a given electric potential  $\varphi_{p_0}^h$  is applied on  $\Gamma_{DI}^{\varepsilon}$  and, when  $p_2 = 1$ , on  $\Gamma_{eD}^{\varepsilon}$ , while electric charges of density  $a^{\varepsilon}$  appear on  $\Gamma_{NI}^{\varepsilon}$  and, when  $p_2 = 1$ , on  $\Gamma_{eN}^{\varepsilon}$ .

If  $\sigma_p^h, u_p^h, e(u_p^h), D_p^h, \varphi_p^h$  stand for the fields of stress, displacement, strain, electric displacement and electric potential, respectively, the constitutive equations of the structure, for all  $p_1$  in  $\{1, 2, 3, 4\}$ , read as:

$$\begin{cases} (\sigma_p^h, D_p^h) = \mu M_I(e(u_p^h), \nabla \varphi_p^h) & \text{in } B^{\varepsilon} \quad \forall p_2 \in \{1, 2, 3, 4\}, \\ \left\{ \begin{array}{ll} (\sigma_p^h, D_p^h) = M_E^{\varepsilon}(e(u_p^h), \nabla \varphi_p^h) & \text{in } \Omega^{\varepsilon} \text{ if } p_2 = 1, \\ \sigma_p^h = a_E^{\varepsilon} e(u_p^h) & \text{in } \Omega^{\varepsilon} \text{ if } p_2 > 1 \end{array} \right. \end{cases} \quad (3)$$

where

$$(M_E^{\varepsilon}, a_E^{\varepsilon})(x) = (M_E, a_E)(x \mp \varepsilon e_3) \quad \forall x \in \Omega_{\pm}^{\varepsilon} \quad (4)$$

$$\begin{cases} (M_I, M_E) \in L^{\infty}(S \times \Omega; \text{Lin}(\mathbb{K})) \text{ such that} \\ M_P = \begin{bmatrix} a_P & -b_P \\ b_P^T & c_P \end{bmatrix}; \exists \kappa > 0, \kappa |k|^2 \leq M_P(x) k \cdot k, \quad \forall k \in \mathbb{K} := \mathbb{S} \times \mathbb{R}^3, \text{ a.e. } x \in \Omega, \forall P \in \{I, E\} \end{cases} \quad (5)$$

and  $\text{Lin}(\mathbb{S}^3)$  is the space of linear operators on the space  $\mathbb{S}^N$  of  $N \times N$  symmetric matrices whose inner product and norm are noted  $\cdot$  and  $|\cdot|$  as in  $\mathbb{R}^3$  (the same notations for the norm and inner product stand also for  $\mathbb{K}$ ).

Lastly we have to add the following conditions on  $S_{\pm}^{\varepsilon}$ :

$$\begin{cases} p_2 = 2 \quad D_p^h \cdot e_3 = 0 \quad \text{on } S_{\pm}^{\varepsilon}, \\ p_2 = 3 \quad D_p^h \cdot e_3 = 0 \quad \text{on } S_{+}^{\varepsilon}, \quad \varphi_p^h = \varphi_{p_0}^h \text{ on } S_{-}^{\varepsilon}, \\ p_2 = 4 \quad \varphi_p^h = \varphi_{p_0}^h \quad \text{on } S_{\pm}^{\varepsilon}, \end{cases} \quad (6)$$

the electric potential  $\varphi_{p_0}^h$  being given on  $S_{+}^{\varepsilon}$  or  $S_{\pm}^{\varepsilon}$ .

It will be convenient to use the following notations:

$$\begin{cases} \hat{k} := (\hat{e}, \hat{g}) \quad \hat{e} := e_{\alpha\beta}, 1 \leq \alpha, \beta \leq 2, \quad \hat{g} := (g_1, g_2), \quad \forall k = (e, g) \in \mathbb{K} \\ \tilde{e}_{\alpha\beta} = e_{\alpha\beta}, 1 \leq \alpha, \beta \leq 2, \quad \tilde{e}_{i3} = 0, 1 \leq i \leq 3, \quad \forall e \in \mathbb{S}^3 \\ k(r) = k(v, \psi) := (e(v), \nabla \psi) \quad \forall r \in H^1(\mathcal{O}; \mathbb{R}^3 \times \mathbb{R}) \\ e(v) \in \mathcal{D}'(S; \mathbb{S}^2); \quad (e(v))_{\alpha\beta} = \frac{1}{2}(\partial_{\alpha} v_{\beta} + \partial_{\beta} v_{\alpha}), 1 \leq \alpha, \beta \leq 2, \quad \forall v \in \mathcal{D}'(S; \mathbb{R}^3) \end{cases} \quad (7)$$

and the same symbol  $e(\cdot)$  shall also stand for the symmetrized gradient in the sense of distributions of  $\mathcal{D}'(\mathcal{O}; \mathbb{R}^3)$ ,  $\mathcal{O} \in \{\mathcal{O}^{\varepsilon}, \Omega, \Omega \setminus S, B^{\varepsilon}, \Omega^{\varepsilon}\}$  or  $\mathcal{D}'(S; \mathbb{R}^2)$ . An electromechanical state with vanishing electric potential on  $\Gamma_{DI}^{\varepsilon}$  and on  $\Gamma_{eD}^{\varepsilon}$  when  $p_2 = 1$  will belong to  $V_p^{\varepsilon} := H_{\Gamma_{mD}^{\varepsilon}}^1(\mathcal{O}^{\varepsilon}; \mathbb{R}^3) \times \Phi_{p_2}^{\varepsilon}$ , with

$$\begin{cases} \Phi_1^{\varepsilon} = H_{\Gamma_{DI}^{\varepsilon} \cup \Gamma_{eD}^{\varepsilon}}^1(\mathcal{O}^{\varepsilon}) \\ \Phi_2^{\varepsilon} = H_{\Gamma_{DI}^{\varepsilon}}^1(B^{\varepsilon}) \text{ if } \mathcal{H}_2(\Gamma_{DI}^{\varepsilon}) > 0, H_m^1(B^{\varepsilon}) \text{ if } \mathcal{H}_2(\Gamma_{DI}^{\varepsilon}) = 0 \\ \Phi_3^{\varepsilon} = H_{\Gamma_{DI}^{\varepsilon} \cup S_{-}^{\varepsilon}}^1(B^{\varepsilon}) \\ \Phi_4^{\varepsilon} = H_{\Gamma_{DI}^{\varepsilon} \cup \pm S_{\pm}^{\varepsilon}}^1(B^{\varepsilon}) \end{cases} \quad (8)$$

where, for any domain  $\mathcal{O}$  of  $\mathbb{R}^N$ ,  $N = 2, 3$ ,  $H_\Sigma^1(\mathcal{O}; \mathbb{R}^M)$  denotes the subspace of  $H^1(\mathcal{O}; \mathbb{R}^M)$ ,  $M = 1$  or  $3$ , of all elements with vanishing traces on a part  $\Sigma$  of the boundary of  $\mathcal{O}$ , while  $H_m^1(\mathcal{O}; \mathbb{R}^M)$  denotes the subspace of all elements with vanishing average.

We make the following assumptions on the data:

$$\left\{ \begin{array}{l} \text{Given } (f, F, d_E, d_I) \text{ in } L^2(\Omega; \mathbb{R}^3) \times L^2(\Gamma_{mN}; \mathbb{R}^3) \times L^2(\Gamma_{eN}) \times L^2(\gamma_N \times (-1, 1)) \\ \text{with } \int_{\Gamma_{\text{lat}}} d_I = 0 \text{ when } p_2 = 2 \text{ and } \mathcal{H}_2(\gamma_D \times (-1, 1)) = 0, \\ \varphi_{oI} \text{ in } H^{3/2}(\mathbb{R}) \text{ vanishing in } \{|x_3| > 1 + \delta/2\}, \text{ and } \varphi_{oE} \text{ in } H^1(\Omega) \text{ vanishing on } S, \text{ then:} \\ f^\varepsilon(x) = f(x \mp \varepsilon e_3) \text{ a.e. } x \in \Omega_{\pm}^\varepsilon, \quad f^\varepsilon(x) = 0 \text{ a.e. } x \in B^\varepsilon, \\ F^\varepsilon(x) = F(x \mp \varepsilon e_3) \text{ a.e. } x \in \Gamma_{mN\pm}^\varepsilon, \\ d^\varepsilon(x) = (\mu/\varepsilon)^{1/2} d_I(\hat{x}, x_3/\varepsilon) \text{ a.e. } x \in B^\varepsilon, \\ d^\varepsilon(x) = d_E(x \mp \varepsilon e_3) \text{ a.e. } x \in \Gamma_{eN\pm}^\varepsilon \text{ if } p_2 = 1, \\ \varphi_{p_0}^h(x) = \begin{cases} \varphi_{oE}(x \mp \varepsilon e_3) + \varepsilon^{p_{DI}} \varphi_{oI}(x \pm (1 - \varepsilon)e_3) \text{ a.e. } x \in \Omega_{\pm}^\varepsilon \\ \varepsilon^{p_{DI}} \varphi_{oI}(\hat{x}, x_3/\varepsilon) \text{ a.e. } x \in B^\varepsilon \end{cases} \end{array} \right. \quad (9)$$

where  $p_{DI}$  is such that  $p_{DI} = 0$  if  $\partial_3 \varphi_{oI} = 0$  in  $S \times (-1, 1)$ ,  $p_{DI} = 1$  if  $\partial_3 \varphi_{oI} \neq 0$  in  $S \times (-1, 1)$ . We also introduce the element  $\varphi_o$  of  $H^{1,1}(\Omega, S) := \{\psi \in H^1(\Omega) \text{ whose trace } \gamma_o(\psi) \text{ on } S \text{ belongs to } H^1(S)\}$  defined by  $\varphi_o(x) = \varphi_{oE}(x) + (1 - p_{DI})\varphi_{oI}(x \pm e_3)$  a.e.  $x \in \Omega_{\pm}$ . We note  $\bar{\varphi}_o$  the trace on  $\gamma_D$  of  $\varphi_o$  and set  $\Delta \varphi_{oI} = \frac{1}{2}(\varphi_{oI}(\cdot, 1) - \varphi_{oI}(\cdot, -1))$ .

Then, if  $\mathcal{M}_p$  and  $\mathcal{L}_p$  are defined by:

$$\left\{ \begin{array}{l} \mathcal{M}_p(s, r) := \begin{cases} \int_{\Omega^\varepsilon} M_E^\varepsilon k(s) \cdot k(r) dx + \mu \int_{B^\varepsilon} M_I k(s) \cdot k(r) dx, & \text{if } p_2 = 1 \\ \int_{\Omega^\varepsilon} a_E^\varepsilon e(u) \cdot e(v) dx + \mu \int_{B^\varepsilon} M_I k(s) \cdot k(r) dx, & \text{if } p_2 > 1 \end{cases} \\ \mathcal{L}_p(r) := \int_{\Omega} f^\varepsilon \cdot v dx + \int_{\Gamma_{mN}^\varepsilon} F^\varepsilon \cdot v d\mathcal{H}_2 + \int_{\Gamma_{DI}^\varepsilon \cup \Gamma^\varepsilon} d^\varepsilon \psi d\mathcal{H}_2 \quad \Gamma^\varepsilon = \Gamma_{eN}^\varepsilon \text{ if } p_2 = 1, \Gamma^\varepsilon = \emptyset \text{ if } p_2 > 1. \end{array} \right. \quad (10)$$

Seeking an equilibrium state leads to the problem

$$(\mathcal{P}_p^h): \quad \text{Find } s_p^h \text{ in } (0, \varphi_{p_0}^h) + V_p^\varepsilon \text{ such that } \mathcal{M}_p(s_p^h, r) = \mathcal{L}_p(r), \quad \forall r \in V_p^\varepsilon$$

which, by Stampacchia theorem, has a unique solution.

## 2. The asymptotic models

In the following  $C, C'$  will denote various constants independent of  $h$  that may vary from line to line. It will be convenient in the cases  $p_2 > 1$  to use the same symbol  $s_p^h$  for  $(u_p^h, \tilde{\varphi}_p^h)$  where  $\tilde{\varphi}_p^h$  denotes the extension into  $\Omega^\varepsilon$  of  $\varphi_p^h$  by 0. Without loss of generality, we suppose  $\mathcal{H}_2(\Gamma_{mD+}) > 0$ ; moreover, when  $p_1 = 4$ , we assume  $\mathcal{H}_2(\Gamma_{mD\pm}) > 0, \mathcal{H}_2(\Gamma_{eD\pm}) > 0$ . We recall that if the same symbol  $u_p^h$  denotes a continuous extension from  $H^1(\Omega_+^\varepsilon; \mathbb{R}^3)$  into  $H^1(\{x_3 > \varepsilon\}; \mathbb{R}^3)$  and  $\eta$  is a  $C_0^\infty(\mathbb{R})$  cut-off function such that

$$\eta = 1 \text{ on } [-\frac{\delta}{3}, \frac{\delta}{3}], \quad 0 \leq \eta \leq 1 \text{ and } 0 < \left| \frac{d\eta}{dx_3} \right| \leq \frac{4}{\delta} \text{ on } \frac{\delta}{3} \leq |x_3| \leq \frac{2\delta}{3}, \quad \eta = 0 \text{ on } \left\{ |x_3| \geq \frac{2\delta}{3} \right\} \quad (11)$$

then Korn inequality imply

$$\begin{aligned} \int_{B^\varepsilon} |\nabla u_p^h|^2 dx &\leq \int_{S \times (-\varepsilon, \delta - \varepsilon)} |\nabla \eta u_p^h|^2 dx \leq C \int_{S \times (-\varepsilon, \delta - \varepsilon)} |e(\eta u_p^h)|^2 dx \\ &\leq C' \left( |e(u_p^h)|_{L^2(B^\varepsilon; \mathbb{S}^3)}^2 + |e(u_p^h)|_{L^2(\Omega_+^\varepsilon; \mathbb{S}^3)}^2 \right) \end{aligned} \quad (12)$$

We will propose our models by studying the asymptotic behavior of  $s_p^h$  when  $h$  goes to  $\bar{h}$  in three steps.

As announced in [1], we may proceed, when  $p_1 = 1$ , similarly but by taking due account of the realistic nonvanishing electric loading on  $\Gamma_{\text{lat}}^\varepsilon$  easily handled though the standard inequalities:

$$\left\{ \begin{array}{l} \int_{\Gamma_{NI}^\varepsilon} \psi^2 d\mathcal{H}_2 \leq C \int_{B^\varepsilon} |\nabla \psi|^2 dx \quad \forall \psi \in H_{\Gamma_{NI}^\varepsilon}^1(B^\varepsilon) \text{ if } \mathcal{H}_2(\Gamma_{DI}^\varepsilon) > 0 \\ \int_{\Gamma_{\text{lat}}^\varepsilon} \psi^2 d\mathcal{H}_2 \leq C \int_{B^\varepsilon} |\nabla \psi|^2 dx \quad \forall \psi \in H_m^1(B^\varepsilon) \text{ if } \mathcal{H}_2(\Gamma_{DI}^\varepsilon) > 0 \end{array} \right. \quad (13)$$

When  $p_1 > 1$ , we proceed in the spirit of [2–4].

**Step 1 (a priori estimates):** By taking  $r = s_p^h - (0, \varphi_{p_0}^h)$  in the variational formulation of  $(\mathcal{P}_p^\varepsilon)$ , (12), (13) and

$$\int_S |s_p^h(\cdot, \varepsilon) - s_p^h(\cdot, -\varepsilon)|^2 d\hat{x} \leq 2\varepsilon \int_{B^\varepsilon} |\nabla s_p^h|^2 dx \quad (14)$$

imply:

$$\mu |k(s_p^h)|_{L^2(B^\varepsilon; \mathbb{K})}^2 + |k(s_p^h)|_{L^2(\Omega^\varepsilon; \mathbb{K})}^2 \leq C. \quad (15)$$

**Step 2 (convergence of  $(s_p^h)$ ):** The following two tools are well suitable to describe the asymptotic behavior of the electromechanical state in the adherents and adhesive, respectively. First, let  $T^\varepsilon$  be the mapping from  $H^1(\Omega^\varepsilon; \mathbb{R}^3 \times \mathbb{R})$  into  $H^1(\Omega \setminus S; \mathbb{R}^3 \times \mathbb{R})$  defined by:

$$(T^\varepsilon r)(x) = (T^\varepsilon(v, \psi))(x) = (T_1^\varepsilon v, T_2^\varepsilon \psi)(x) := (v, \psi)(x \pm \varepsilon e_3) \quad \forall x \in \Omega_\pm. \quad (16)$$

Note that  $T^\varepsilon s_p^h = (T_1 u_p^h, 0)$  if  $p_2 > 1$ . For any  $w$  in  $H^1(\Omega \setminus S; \mathbb{R}^N)$ ,  $N \in \{1, 3\}$ , if  $\gamma_o^\pm(w^\pm)$  denotes the trace on  $S$  of its restriction  $w^\pm$  to  $\Omega_\pm$ ,  $\llbracket w \rrbracket$  stands for  $\gamma_o^+(w^+) - \gamma_o^-(w^-)$ .

Next, as for all  $r$  of  $H^1(B^\varepsilon; \mathbb{R}^3 \times \mathbb{R})$ , one has

$$\int_{B^\varepsilon} \left| r(x) - \frac{1}{2\varepsilon} \int_{-\varepsilon}^{\varepsilon} r(\hat{x}, x_3) dx_3 \right|^2 dx \leq C\varepsilon^2 \int_{B^\varepsilon} |\nabla r|^2 dx \quad (17)$$

we introduce the following element of  $L^2(S; \mathbb{K})$ :

$$k_p(\varepsilon, r) := \frac{1}{(2\varepsilon)^q} \int_{-\varepsilon}^{\varepsilon} k(r)(\cdot, x_3) dx_3, \quad q = \max(2 - p_1, 0) \quad (18)$$

and, obviously, there holds

$$\widehat{k_p(\varepsilon, r)} = (e(\widehat{U_p}), \nabla \Phi_p), \quad S_p^h := (U_p^h, \Phi_p^h) := \frac{1}{(2\varepsilon)^q} \int_{-\varepsilon}^{\varepsilon} s_p^h(\cdot, x_3) dx_3. \quad (19)$$

So (12), (14) and (15) imply:

$$\begin{cases} |k(T^\varepsilon s_p^h)|_{L^2(\Omega \setminus S; \mathbb{K})} \leq C, \quad \llbracket T^\varepsilon s_p^h \rrbracket_{L^2(S; \mathbb{R}^3 \times \mathbb{R})}^2 \leq C\left(\varepsilon + \frac{\varepsilon}{\mu}\right) \\ |k_p(\varepsilon, s_p^h)|_{L^2(S; \mathbb{K})}^2 \leq C\varepsilon^{-2q} \cdot \frac{\varepsilon}{\mu} \\ |U_p^h|_{L^2(S; \mathbb{R}^3)}^2 \leq C\varepsilon^{2(1-q)} \left(1 + \frac{\varepsilon}{\mu}\right) \\ \varepsilon^{2(q-1)} |\Phi_p^h|_{L^2(S)}^2 \leq Cc^*(h) \quad c^*(h) = \left(1 + \frac{\varepsilon}{\mu}\right) \text{ if } p_2 = 1, \frac{1}{\varepsilon\mu} \text{ if } p_2 = 2, \varepsilon^2 + \frac{\varepsilon}{\mu} \text{ if } p_2 > 2 \\ |e(\widehat{U_p^h}), \nabla \Phi_p^h|_{L^2(S; \mathbb{R}^3 \times \mathbb{R}^2)}^2 \leq C \frac{1}{\varepsilon^{2q}} \cdot \frac{\varepsilon}{\mu} \\ |S_p^h - \gamma_o^\pm((T^\varepsilon s_p^h)^\pm)|_{L^2(S; \mathbb{R}^3 \times \mathbb{R})}^2 \leq C\left(\varepsilon + \frac{\varepsilon}{\mu}\right) \text{ if } p_1 = p_2 = 1 \\ |U_p^h - \gamma_o^\pm((T_1^\varepsilon u_p^h)^\pm)|_{L^2(S; \mathbb{R}^3)}^2 \leq C\left(\varepsilon + \frac{\varepsilon}{\mu}\right) \text{ if } p_1 = 1, p_2 > 1. \end{cases} \quad (20)$$

Thus, if  $a \otimes_S b$  denotes the symmetrized tensor product of  $a$  and  $b$  in  $\mathbb{R}^3$ , we deduce:

**Proposition 2.1.** There exists  $\bar{s}_p = (\bar{u}_p, \bar{\varphi}_p)$  in  $H^1_{\Gamma_{\text{mD}}}(\Omega \setminus S; \mathbb{R}^3) \times H^1_{\Gamma_{\text{eD}}}(\Omega \setminus S)$  such that  $T^\varepsilon s_p^h$  weakly converges in  $H^1(\Omega \setminus S; \mathbb{R}^3 \times \mathbb{R})$  toward some  $\bar{s}_p$ ;  $\bar{\varphi}_p = 0$  when  $p_2 > 1$  and  $\bar{s}_p$  belongs to  $H^1(\Omega; \mathbb{R}^3 \times \mathbb{R})$  when  $p_1 \leq 2$ .

When  $p_1 \neq 4$ ,  $k_p(\varepsilon, s_p^h)$  weakly converges in  $L^2(S; \mathbb{K})$  toward some  $\bar{k}_p = (\bar{e}_p, \bar{g}_p)$ , and there exists  $(\bar{U}_p, \bar{\Phi}_p)$  in  $H^1(S; \mathbb{R}^3 \times \mathbb{R})$  such that  $(\widehat{U_p^h}, \Phi_p^h)$  converges weakly in  $H^1(S; \mathbb{R}^3 \times \mathbb{R})$  toward  $(\widehat{\bar{U}_p}, \bar{\Phi}_p)$ ,  $(U_p^h)_3$  converges strongly in  $L^2(S)$  toward  $(\bar{U}_p)_3$ . Moreover

- i) when  $p_1 = 1$ ,  $\bar{U}_p = \gamma_0(\bar{u}_p)$  for all  $p_2$ , while  $\bar{\Phi}_p$  is equal to  $\gamma_0(\bar{\varphi}_p)$  when  $p_2 = 1$  or to  $\gamma_0(\varphi_0)$  when  $p_2 \geq 3$ . Furthermore, the trace on  $\gamma_D$  of  $\bar{\Phi}_p$  is equal to  $\bar{\varphi}_o$  while  $\widehat{\bar{k}_p} = (e(\widehat{\bar{U}_p}), \nabla \bar{\Phi}_p)$ ,  $(\bar{g}_p)_3 = \Delta \varphi_{0l}$  when  $p_2 = 4$ ;
- ii) when  $p_1 = 2$ ,  $\bar{U}_p = 0$ ,  $\bar{\Phi}_p = 0$  and  $\bar{k}_p = 0$ ;
- iii) when  $p_1 = 3$ ,  $\bar{U}_p = 0$  and  $\bar{e}_p = \llbracket \bar{u}_p \rrbracket \otimes_S e_3$  for all  $p_2$ , while  $\bar{\Phi}_p$  and  $\widehat{\bar{g}_p}$  vanish only when  $p_2 \neq 2$ ,  $(\bar{g}_{(3,1)})_3 = \llbracket \bar{\varphi}_p \rrbracket$ ,  $(\bar{g}_{(3,4)})_3 = 0$ .

Actually, in the next step, we will show that  $(\bar{u}_p, \bar{\varphi}_p)$  is necessarily the unique solution to a variational problem so that the whole sequences converge. To identify  $\bar{k}_p$ , when  $p_1 > 1$ , it suffices to go to the limit in the identity

$$\int_{B^\varepsilon} \left( k(s_p^h)(x) \cdot \tau^i(\hat{x}) + s_p^h(x) \cdot \operatorname{div} \tau^i(\hat{x}) \right) dx = \int_S \left( \llbracket T_1^\varepsilon u_p^h \rrbracket_i \theta_1 \right) + \left( \llbracket T_2^\varepsilon \varphi_p^h \rrbracket \theta_2 \right) d\hat{x} \quad (21)$$

with  $\tau^i = (\theta_1 e_3 \otimes_S e_i, \theta_2 e_3)$ ,  $\theta_j \in C_0^\infty(S)$ ,  $j = 1, 2$ ,  $i = 1, 2, 3$  and to use the convergence to 0 in the sense of distributions of  $\widehat{k_p(\varepsilon, s_p^h)}$  by due account of (20)!

### Step 3 (identification of $(\bar{s}_p, \bar{k}_p)$ ):

When  $p_1 > 1$ , we simply go to the limit in the variational formulation of  $(\mathcal{P}_p^h)$  by using suitable test-functions  $r_p^\varepsilon$ . For all  $w^1$  in  $H_{\Gamma_{\text{mD}}}^1(\Omega; \mathbb{R}^3)$  and all  $\zeta^1$  in  $H_{\Gamma_{\text{eD}}}^1(\Omega)$  vanishing in a neighborhood of  $\gamma_D$ , let  $(w^{1,\varepsilon}, \zeta^{1,\varepsilon})$  be defined by

$$(w^{1,\varepsilon}, \zeta^{1,\varepsilon})(x) = \begin{cases} (w^1, \zeta^1)(x \mp \varepsilon e_3) & \text{a.e. } x \in \Omega_\pm^\varepsilon \\ (w^1, \zeta^1)(\hat{x}, 0) & \text{a.e. } x \in B^\varepsilon \end{cases}$$

For all  $w^2$  in  $H_{\Gamma_{\text{mD}}}^1(\Omega \setminus S; \mathbb{R}^3)$  and all  $\zeta^2$  in  $H_{\Gamma_{\text{eD}}}^1(\Omega \setminus S)$  vanishing in a neighborhood of  $\gamma_D$ , let  $(w^{2,\varepsilon}, \zeta^{2,\varepsilon})$  be defined by

$$(w^{2,\varepsilon}, \zeta^{2,\varepsilon})(x) = \begin{cases} (w^2, \zeta^2)(x \mp \varepsilon e_3) & \text{a.e. } x \in \Omega_\pm^\varepsilon \\ (w^a, \zeta^a)(\hat{x}, x_3/\varepsilon) + \frac{|x_3|}{\varepsilon} (w^s, \zeta^s)(\hat{x}, x_3/\varepsilon) & \text{a.e. } x \in B^\varepsilon \end{cases}$$

with

$$\begin{aligned} (w^a, \zeta^a)(x) &= \frac{1}{2} [(w^2, \zeta^2)(\hat{x}, x_3) - (w^2, \zeta^2)(\hat{x}, -x_3)] \\ (w^s, \zeta^s)(x) &= \frac{1}{2} [(w^2, \zeta^2)(\hat{x}, x_3) + (w^2, \zeta^2)(\hat{x}, -x_3)] \end{aligned}$$

Note (see the proof of Lemma 4.1 of [3]) that

$$\left| \left( e(w^{2,\varepsilon}) - \frac{\llbracket w^2 \rrbracket \otimes_S e_3}{2\varepsilon}, \nabla \zeta^{2,\varepsilon} - \frac{\llbracket \zeta^2 \rrbracket e_3}{2\varepsilon} \right) \right|_{L^2(B^\varepsilon; \mathbb{K})} \leq C |(w^2, \zeta^2)|_{H^1(B^\varepsilon \setminus S; \mathbb{R}^3 \times \mathbb{R})}.$$

So  $r_p^\varepsilon = (v_p^\varepsilon, \psi_p^\varepsilon)$  reads as:

$$\begin{cases} v_p^\varepsilon = w^{\min(p_1-1, 2), \varepsilon}, & 1 \leq p_2 \leq 4 \\ \psi_p^\varepsilon = \begin{cases} \zeta^{\min(p_1-1, 2), \varepsilon} & p_2 = 1 \\ (\theta_1 + x_3 \theta_2)/\varepsilon, \quad \theta_1, \theta_2 \in C_0^\infty(S) & p_1 = 3, 4, p_2 = 2 \\ (1 + x_3/\varepsilon)\theta, \quad \theta \in C_0^\infty(S) & p_1 = 3, 4, p_2 = 3 \\ 0 & \text{if } (2 \leq p_1 \leq 4, p_2 = 4) \text{ or } (p_1 = 2, p_2 = 2, 3) \end{cases} \end{cases} \quad (22)$$

When  $p_1 = 1$ , one proceeds in two steps. First we prove

$$(M\bar{k}_p)_p^2 = 0 \quad (23)$$

where  $k_p^i$  denotes the projection on  $\mathbb{K}_p^i$  of any element  $k$  of  $\mathbb{K}$  with:

$$\begin{cases} \mathbb{K} = \mathbb{K}_p^1 \oplus \mathbb{K}_p^2 \oplus \mathbb{K}_p^3, \\ p_2 \leq 2 : \mathbb{K}_p^1 := \{(e, g) \in \mathbb{K}; e_{13} = 0, g_3 = 0\}, \mathbb{K}_p^2 := \{(e, g) \in \mathbb{K}; \hat{e} = 0, \hat{g} = 0\}, \mathbb{K}_p^3 := \{0\}, \\ p_2 > 2 : \mathbb{K}_p^1 = \{(e, g) \in \mathbb{K}; e_{13} = 0, \hat{g} = 0\}, \mathbb{K}_p^2 = \{(e, g) \in \mathbb{K}; \hat{e} = 0, g = 0\}, \\ \mathbb{K}_p^3 = \{(e, g) \in \mathbb{K}; e = 0, g_3 = 0\} \end{cases} \quad (24)$$

For that, we simply use test functions  $\rho_p^\varepsilon$  built as follows: given  $(w, \psi)$  in  $C_0^\infty(S; \mathbb{R}^3 \times \mathbb{R})$ ,

$$\rho_p^\varepsilon(x) = \begin{cases} (x_3 + \varepsilon)(w, I_{p_2}\psi)(\hat{x}) & \text{a.e. } x \in B^\varepsilon \\ 2\varepsilon(w^+, I_{p_2}\psi^+)(x - \varepsilon e_3) & \text{in } \Omega_+^\varepsilon, 0 \text{ in } \Omega_-^\varepsilon \end{cases} \quad (25)$$

where  $(w^+, \psi^+)$  is an extension into  $H_{\Gamma_{\text{mD+}}}^1(\Omega_+; \mathbb{R}^3) \times H_{\Gamma_{\text{eD+}}}^1(\Omega_+)$  and  $I_{p_2} = 1$  if  $p_2 \leq 2$ ,  $I_{p_2} = 0$  if  $p_2 > 2$ . Hence, as Proposition 2.1 yields  $(\bar{k}_p)_p^3 = 0$ , we deduce

$$(M_I \bar{k}_p)^1 = \widetilde{M}_{I,p}(\bar{k}_p)^1; \quad \widetilde{M}_{I,p} := M_{I,p}^{11} - M_{I,p}^{12}(M_{I,p}^{22})^{-1}M_{I,p}^{21} \quad (26)$$

with  $M_{I,p}^{ij}$ ,  $1 \leq i, j \leq 3$  being the decomposition of  $M_I$  in linear operators mapping  $\mathbb{K}_p^i$  into  $\mathbb{K}_p^j$ .

Next, given  $(v, \psi)$  in  $(H_{\Gamma_{\text{mD}}}^1(\Omega; \mathbb{R}^3) \times H_{\Gamma_{\text{eD}}}^1(\Omega)) \cap H^2(\Omega; \mathbb{R}^3 \times \mathbb{R})$   $\psi$  vanishing in a neighborhood of  $\gamma_D$ , we define  $r_p^\varepsilon = (v_p^\varepsilon, \psi_p^\varepsilon)$  by:

$$\begin{cases} \widehat{v_p^\varepsilon}(x) = \hat{v}(\hat{x}, 0) - x_3 \nabla v_3(\hat{x}, 0), & (v_3^\varepsilon(x) = v_3(\hat{x}, 0)) \text{ a.e. } x \in B^\varepsilon \\ v_p^\varepsilon(x) = v(x \mp \varepsilon e_3) \mp \varepsilon R^\pm(\nabla v_3(\cdot, 0), 0)(x \mp \varepsilon e_3) & \text{a.e. } x \in \Omega_\pm^\varepsilon \\ \psi_p^\varepsilon(x) = \psi(x \mp \varepsilon e_3) \text{ in } \Omega_\pm^\varepsilon, & \psi(\hat{x}, 0) \text{ in } B^\varepsilon \\ \psi_p^\varepsilon(x) = \psi(\hat{x}, 0) & \text{if } p_2 = 1 \\ \psi_p^\varepsilon(x) = 0 & \text{if } p_2 = 2 \\ \psi_p^\varepsilon(x) = 0 & \text{if } p_2 \geq 3 \end{cases} \quad (27)$$

where  $R^\pm$  is a continuous lifting operator from  $H^{1/2}(S; \mathbb{R}^3)$  into  $H_{\Gamma_{mD\pm}}^1(\Omega_\pm; \mathbb{R}^3)$ . As  $(\widetilde{e}(\hat{v}), \nabla \psi)$  belongs to  $\mathbb{K}_p^1$  almost everywhere in  $S$ , (26) yields

$$\lim_{h \rightarrow \bar{h}} \int_{B^\varepsilon} \mu M_l k(s_p^h) k(r_p^\varepsilon) dx = \int_S M_l \bar{k}_p \cdot (\widetilde{e}(\hat{v}), \nabla \psi) d\hat{x} = \int_S \widetilde{M}_{lp}(\bar{k}_p^1) \cdot (\widetilde{e}(\hat{v}), \nabla \psi) d\hat{x} \quad (28)$$

while, obviously, we have:

$$\begin{cases} \lim_{h \rightarrow \bar{h}} \int_{\Omega^\varepsilon} M_E^\varepsilon k(s_p^h) \cdot k(r_p^\varepsilon) dx = \int_\Omega M_E k(\bar{s}_p) \cdot k(v, \psi) dx & \text{if } p_2 = 1 \\ \lim_{h \rightarrow \bar{h}} \int_{\Omega^\varepsilon} a_E^\varepsilon e(u_p^h) \cdot e(v_p^\varepsilon) dx = \int_\Omega a_E e(\bar{u}_p) \cdot e(v) dx & \text{if } p_2 \geq 2 \\ \lim_{h \rightarrow \bar{h}} \mathcal{L}_{(1, p_2)}(r_{(1, p_2)}^\varepsilon) \\ := \begin{cases} \int_\Omega f \cdot v dx + \int_{\Gamma_{mN}} F \cdot v d\mathcal{H}_2 + \int_{\Gamma_{eN}} d_E \psi d\mathcal{H}_2 + (\bar{\mu}_1)^{1/2} \int_{\gamma_N} \left( \int_{-1}^1 d_l(\cdot, x_3) dx_3 \right) \psi d\mathcal{H}_2 & p_2 = 1 \\ \int_\Omega f \cdot v dx + \int_{\Gamma_{mN}} F \cdot v d\mathcal{H}_2 + (\bar{\mu}_1)^{1/2} \int_{\gamma_N} \left( \int_{-1}^1 d_l(\cdot, x_3) dx_3 \right) \psi d\mathcal{H}_2 & p_2 \geq 2 \end{cases} \end{cases} \quad (29)$$

Lastly, Jensen inequality and the previously established weak convergences achieve the proof of the following convergence result which supports our asymptotic models in the form of variational problems  $(\overline{\mathcal{P}}_p)$  where the convention  $\infty \times 0 = 0$  is understood.

**Theorem 2.1.** If  $p_2 = 1$ , when  $h$  goes to  $\bar{h}$ ,  $T^\varepsilon s_p^h$  converges strongly in  $H^1(\Omega \setminus S; \mathbb{R}^3 \times \mathbb{R})$  toward  $\bar{s}_p$  the unique solution to

$$(\overline{\mathcal{P}}_{(p_1, 1)}) : \begin{cases} \text{Find } (u, \varphi) \text{ in } (0, \varphi_0) + V_{p_1} \times \Psi_{p_1} \text{ such that} \\ \overline{\mathcal{M}}_{(p_1, 1)}((u, \varphi), (v, \psi)) = \overline{\mathcal{L}}_{(p_1, 1)}(v, \psi) \quad \forall (v, \psi) \in V_{p_1} \times \Psi_{p_1} \end{cases}$$

where

$$\begin{aligned} \overline{\mathcal{M}}_{(p_1, 1)}((u, \varphi), (v, \psi)) &:= \begin{cases} \int_\Omega M_E k(u, \varphi) \cdot k(v, \psi) dx + \bar{\mu}_1 \int_S \widetilde{M}_{lp}(e(\hat{u}), \nabla \varphi) \cdot (e(\hat{u}), \nabla \psi) d\hat{x} & p_1 = 1 \\ \int_\Omega M_E k(u, \varphi) \cdot k(v, \psi) dx + \bar{\mu}_2 \int_S M_l([u] \otimes_S e_3, [\varphi] e_3) \cdot ([v] \otimes_S e_3, [\psi] e_3) d\hat{x} & p_1 \geq 2 \end{cases} \\ \overline{\mathcal{L}}_{(p_1, 1)}(v, \psi) &:= \begin{cases} \int_\Omega f \cdot v dx + \int_{\Gamma_{mN}} F \cdot v d\mathcal{H}_2 + \int_{\Gamma_{eN}} d_E \psi d\mathcal{H}_2 + (\bar{\mu}_1)^{1/2} \int_{\gamma_N} \left( \int_{-1}^1 d_l(\cdot, x_3) dx_3 \right) \psi d\mathcal{H}_2 & p_1 = 1 \\ \int_\Omega f \cdot v dx + \int_{\Gamma_{mN}} F \cdot v d\mathcal{H}_2 + \int_{\Gamma_{eN}} d_E \psi d\mathcal{H}_2 & p_1 \geq 2 \end{cases} \end{aligned}$$

$$V_1 := \left\{ v \in H_{\Gamma_{mD}}^1(\Omega; \mathbb{R}^3); \hat{v} \in H^1(S; \mathbb{R}^2) \right\}, \quad V_2 := H_{\Gamma_{mD}}^1(\Omega; \mathbb{R}^3), \quad V_3 = V_4 := H_{\Gamma_{mD}}^1(\Omega \setminus S; \mathbb{R}^3)$$

$$\Psi_1 := \left\{ \psi \in H_{\Gamma_{eD}}^1(\Omega); \psi \in H_{\gamma_D}^1(S) \right\}, \quad \Psi_2 := H_{\Gamma_{eD}}^1(\Omega), \quad \Psi_3 = \Psi_4 := H_{\Gamma_{eD}}^1(\Omega \setminus S)$$

If  $p_2 > 1$ , when  $h$  goes to  $\bar{h}$ ,  $(T_1^\varepsilon u_p^h, \Phi_p^h)$  converges strongly in  $H^1(\Omega \setminus S; \mathbb{R}^3) \times H^1(S)$  toward  $(\bar{u}_p, \overline{\Phi}_p)$  the unique solution to

$$(\overline{\mathcal{P}}_p) : \begin{cases} \text{Find } (u, \phi) \text{ in } (0, q \gamma_0(\varphi_0)) + V_{p_1} \times \Psi_p \text{ such that} \\ \overline{\mathcal{M}}_p((u, \phi), (v, \psi)) = \overline{\mathcal{L}}_p(v, \psi) \quad \forall (v, \psi) \in V_{p_1} \times \Psi_p \end{cases}$$

where

$$\overline{\mathcal{M}}_{(1, p_2)}((u, \phi), (v, \psi)) := \begin{cases} \int_\Omega a_E e(u) \cdot e(v) dx + \bar{\mu}_1 \int_S \widetilde{M}_{lp}(e(\hat{u}), \nabla \phi) \cdot (e(\hat{u}), \nabla \psi) d\hat{x} & p_2 = 2 \\ \int_\Omega a_E e(u) \cdot e(v) dx + \bar{\mu}_1 \int_S \widetilde{M}_{lp}(e(\hat{u}), (\bar{g}_p)_3) \cdot (e(\hat{u}), 0) d\hat{x} & p_2 \geq 3 \\ \text{with } \widetilde{M}_{lp}(e(\hat{u}), (\bar{g}_p)_3) \cdot (0, e_3) = 0 \text{ if } p_2 = 3 \\ (\bar{g}_p)_3 = \Delta \varphi_{0l} \text{ if } p_2 = 4 \\ \int_\Omega a_E e(u) \cdot e(v) dx + \bar{\mu}_1 \int_S \widetilde{M}_{lp}(e(\hat{u}), \Delta \varphi_{0l}) \cdot (e(\hat{u}), 0) d\hat{x} & p_2 = 4 \end{cases}$$

$$\overline{\mathcal{L}}_{(1, p_2)}(v, \psi) := \int_\Omega f \cdot v dx + \int_{\Gamma_{mN}} F \cdot v d\mathcal{H}_2 + (\bar{\mu}_1)^{1/2} \int_{\gamma_N} \left( \int_{-1}^1 d_l(\cdot, x_3) dx_3 \right) \psi d\hat{x} \quad \forall p_2 \geq 2$$

and for  $2 \leq p_1, p_2 \leq 4$

$$\bar{\mathcal{M}}_{(p_1, p_2)}((u, \phi), (v, \psi)) := \begin{cases} \int_{\Omega} a_E e(u) \cdot e(v) dx + \bar{\mu}_2 \int_S M_I([\![u]\!] \otimes_S e_3, (\nabla \phi, (\bar{g}_p)_3)) \cdot ([\![v]\!] \otimes_S e_3, (\nabla \psi, 0)) d\hat{x} \\ \text{with } M_I([\![u]\!] \otimes_S e_3, (\nabla \phi, (\bar{g}_p)_3)) \cdot (0, e_3) = 0 \text{ if } p_2 = 2, 3 \\ (\bar{g}_p)_3 = 0 \text{ if } p_2 = 4 \text{ or } p_1 = 2 \end{cases}$$

$$\bar{\mathcal{L}}_{(p_1, p_2)}(v, \psi) = \int_{\Omega} f \cdot v dx + \int_{\Gamma_{mN}} F \cdot v d\mathcal{H}_2$$

$$\Psi_{(p_1, 2)} := H_{\gamma_D}^1(S) \text{ or } H_m^1(S) \text{ according to the positivity of the length of } \gamma_D, \quad \Psi_{(p_1, 3)} = \Psi_{(p_1, 4)} := \{0\}, \quad p_1 \neq 2$$

$$\Psi_{(2, p_2)} := \{0\}, \quad 2 \leq p_2 \leq 4$$

### 3. Concluding remarks

In the case of piezoelectric adhesive and adherents ( $p_2 = 1$ ), our results extend those obtained in elasticity (see [4–7]). The asymptotic behavior of the adhesive strongly depends on the magnitude of the stiffness compared to that of the thickness. When the magnitude of the stiffness is of the order of the inverse of the thickness, the adhesive is replaced by a *material piezoelectric surface* perfectly bonded to the adherents. When it is lesser, the adhesive is replaced by an *electromechanical constraint* between the two adherents which can be perfect adhesion, electromechanical pull-back or free separation, according to the order of magnitude of the stiffness which is, respectively, larger, equal or lower than that of the thickness. There is a large discrepancy between our results and that of [8,9] obtained by formal or questionable arguments.

Similarly, in the case of a thin piezoelectric layer embedded between two elastic adherents, depending on the magnitude of the stiffness, the adhesive is replaced by a material elastic surface perfectly bonded to the adherents or by a mechanical constraint between the adherents. In the case of electrically impermeable interfaces, the material surface has a *non-local* elastic behavior (since the additional state variable of electric nature  $\phi$  can be eliminated!), the constitutive equations being derived from the asymptotic behavior of a thin piezoelectric plate acting as a sensor (case  $p = 1$  in [10]). When one interface is electrically impermeable while the other is electroded, the material surface is an elastic membrane. When the two interfaces are electroded, the material surface is an elastic membrane with residual stress. In these last two cases, the constitutive equations are derived from the asymptotic behavior of a thin piezoelectric plate acting as an actuator (case  $p = 2$  in [10]). The mechanical constraint is perfect adhesion, elastic pull-back or free separation according to the order of magnitude of the stiffness. In the case of electrically impermeable interfaces, the elastic pull-back is of non-local nature (since the state variable of electric nature  $\phi$ , additional to the relative displacement, can be eliminated). In the two other cases, the elastic pull-back is local. When the two interfaces are electroded, it is similar to the purely elastic case, while, if only one interface is electroded, piezoelectric and dielectric coefficients enter the constitutive equations.

The realistic dual situation [11] in which an elastic layer is embedded between two piezoelectric bodies can be treated within the same framework.

### References

- [1] C. Licht, S. Orankitjaroen, P. Viriyasrisuwattana, T. Weller, Bonding a linearly piezoelectric patch on a linearly elastic body, *C. R. Mecanique* 342 (4)(2014) 234–239.
- [2] C. Licht, S. Orankitjaroen, Dynamics of elastic bodies connected by a thin soft inelastic layer, *C. R. Mecanique* 341 (3) (2013) 323–332.
- [3] C. Licht, A. Léger, S. Orankitjaroen, A. Ould Khaoua, Dynamics of elastic bodies connected by a thin soft viscoelastic layer, *J. Math. Pures Appl.* (2013) 685–703.
- [4] A. Aitmoussa, Modélisation et études des singularités de contraintes d'un joint collé très mince, Ph.D. thesis, Université Montpellier-2, France, 1989.
- [5] E. Acerbi, G. Buttazzo, D. Percivale, Thin inclusions in linear elasticity: a variational approach, *J. Reine Angew. Math.* 386 (1988) 99–115.
- [6] C. Licht, G. Michaille, A modelling of elastic adhesive bonded joints, *Adv. Math. Sci. Appl.* 7 (2) (1997) 711–740.
- [7] C. Licht, Some new mathematical modelings of junctions, *East-West J. Math.* 13 (1) (2011) 23–33.
- [8] M. Serpilli, An asymptotic model of a multimaterial with a thin piezoelectric interphase, *C. R. Mecanique* 342 (4) (2014) 258–262.
- [9] M. Serpilli, Asymptotic analysis of a multimaterial with a thin piezoelectric interphase, *Mecanica* 49 (7) (2014) 1641–1652.
- [10] T. Weller, C. Licht, Asymptotic modeling of thin piezoelectric plates, *Ann. Solid Struct. Mech.* 1 (2010) 173–188.
- [11] W. Geis, G. Mishuris, A.-M. Sändig, Asymptotic models for piezoelectric stack actuators with thin metal inclusions, preprint 2004/001, Universität Stuttgart, Germany.