



**HAL**  
open science

## **$n$ automorphism groups of affine surfaces**

Sergei Kovalenko, Alexander Perepechko, Mikhail Zaidenberg

► **To cite this version:**

Sergei Kovalenko, Alexander Perepechko, Mikhail Zaidenberg.  $n$  automorphism groups of affine surfaces. 2015. hal-01235237v1

**HAL Id: hal-01235237**

**<https://hal.science/hal-01235237v1>**

Preprint submitted on 29 Nov 2015 (v1), last revised 6 Aug 2016 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON AUTOMORPHISM GROUPS OF AFFINE SURFACES

S. KOVALENKO, A. PEREPECHKO, M. ZAIDENBERG

ABSTRACT. This is a survey on the automorphism groups in various classes of affine algebraic surfaces and the algebraic group actions on such surfaces. Being infinite-dimensional, these automorphism groups share some important features of algebraic groups. At the same time, they can be studied from the viewpoint of the combinatorial group theory, so we put a special accent on group-theoretical aspects (ind-groups, amalgams, etc.). We provide different approaches to classification, prove certain new results, and attract attention to several open problems.

## CONTENTS

1. Introduction	2
1.1. Classification according to the Makar-Limanov invariant	2
1.2. Classification according to rank	3
1.3. A general classification scheme	3
2. Ind-groups, amalgams, and all this	4
2.1. Ind-groups	4
2.2. Nested ind-groups	6
2.3. Amalgams	8
2.4. Bearable automorphism groups	8
3. Automorphism groups and amalgams: the first examples	10
4. Algebraic group actions on affine surfaces	13
4.1. Generalities	13
4.2. Quasihomogeneous affine surfaces	14
4.3. Actions with an open orbit	15
4.4. $\mathbb{G}_m$ -surfaces: Dolgachev-Pinkham-Demazure presentation	16
5. Automorphism groups of Gizatullin surfaces	18
5.1. Definition, characterizations, examples	18
5.2. One-parameter subgroups and bearability on Gizatullin surfaces	18
5.3. Standard completions and extended divisors of Gizatullin surfaces	19
5.4. Associated graph of groups	23
5.5. Amalgam structures for Gizatullin surfaces	24
6. Automorphism groups of $\mathbb{A}^1$ -fibrations	26
6.1. Generalized de Jonquières groups	26
6.2. The surface case	29
7. Formal neighborhood of a fiber in an $\mathbb{A}^1$ -fibration	29
7.1. Chain of contractions	30
7.2. Formal neighborhoods and coordinate charts	31
7.3. Arcs and multiplicities	32
7.4. Puiseux arc spaces	34
7.5. Stabilizer of a special fiber	36

---

**Acknowledgements:** The second author is supported by the SPbGU grant for postdocs no. 6.50.22.2014 and partially granted by the “Dynasty” foundation. He also expresses his gratitude to the Institut Fourier, Grenoble. The third author is grateful to the INdAM (Istituto Nazionale di Alta Matematica “F. Severi”) in Rome, where a part of the work was done. The second and the third authors are grateful to the Max-Planck Institute for Mathematics, Bonn, for hospitality and generous support.

8. Automorphism groups of $\mathbb{A}^1$ -fibrations on surfaces	39
8.1. Notation	39
8.2. Stabilizers of arc spaces	39
8.3. Automorphism groups of $\mathbb{A}^1$ -fibrations	40
References	45

## 1. INTRODUCTION

Our aim is to give a tedious survey on the automorphism groups of affine algebraic surfaces and algebraic group actions on such surfaces. We are using several different classifications of surfaces, according to the Makar-Limanov invariant, to the rank of the automorphism group, etc. However, our ultimate goal is to approach a sensible classification of the automorphism groups themselves. These groups are often infinite dimensional, and even ‘wild’ in a sense, so, we do not reach the final goal. Nevertheless, we are trying to put some order in our present knowledge on the subject, and to indicate difficult open problems.

We apologize for incompleteness of the reference list and of the overview of the cited literature. Instead, we prove several new results, especially concerning the automorphism groups of the surfaces of classes  $(ML_1)$  and  $(ML_2)$ , where one possesses by now a rather complete knowledge. In contrast, we are far from a good understanding of the automorphism groups of surfaces of class  $(ML_0)$ , that is, the Gizatullin surfaces.

Before passing to the content of the paper, we recall some general notions and facts.

### 1.1. Classification according to the Makar-Limanov invariant.

**1.1.** Let  $X$  be a normal affine variety over an algebraically closed field  $\mathbb{K}$  of characteristic zero. Sometimes further restrictions on the field are imposed; however, all results are valid over the complex number field  $\mathbb{C}$ . The *special automorphism group*  $\text{SAut } X$  is the subgroup of  $\text{Aut } X$  generated by all its one-parameter unipotent subgroups ([1]). This group is trivial if and only if  $X$  does not admit any nontrivial  $\mathbb{G}_a$ -action, and in case  $\dim X = 2$  if and only if  $X$  does not admit any  $\mathbb{A}^1$ -fibration over an affine curve, see [42, Rem. 1.7] or [33, Lem. 1.6]. The *Makar-Limanov invariant*  $ML(X) = \mathcal{O}(X)^{\text{SAut } X}$  is the subalgebra of invariants of  $\text{SAut } X$ . The normal affine varieties can be classified according to the transcendence degree of the Makar-Limanov invariant or, which is the same, according to the *Makar-Limanov complexity* of  $X$ , that is, the codimension of a general  $\text{SAut } X$ -orbit. The latter has sense since these orbits are locally closed subvarieties in  $X$  ([1]). One says that a variety  $X$  is of class  $(ML_i)$ ,  $i \in \{0, 1, \dots, \dim X\}$ , if its Makar-Limanov complexity equals  $i$ . A normal affine surface  $X$  is of class

- $(ML_2)$  if  $X$  does not admit any  $\mathbb{A}^1$ -fibration over an affine curve,
- $(ML_1)$  if  $X$  admits a unique such fibration, and
- $(ML_0)$  if  $X$  admits at least two distinct such fibrations.

The surfaces of class  $(ML_0)$  are also called *Gizatullin surfaces*, cf. Definition 5.1. The surfaces of classes  $(ML_0)$  and  $(ML_1)$  are  $\mathbb{G}_a$ -surfaces, which means that they admit an effective action of the additive group  $\mathbb{G}_a = \mathbb{G}_a(\mathbb{K})$  of the field  $\mathbb{K}$ , while the  $ML_2$ -surfaces do not admit such an action.

**1.2.** A derivation  $\partial$  of a ring  $A$  is called *locally nilpotent* if for any  $a \in A$ ,  $\partial^n a = 0$  for some  $n = n(a) \in \mathbb{N}$ . For a normal affine surface  $X$ , the coordinate ring  $\mathcal{O}_X(X)$

- does not admit any nonzero locally nilpotent derivation if  $X \in (ML_2)$ ;

- admits a unique such derivation up to a factor, which is a rational function on  $X$ , if  $X \in (\text{ML}_1)$ ;
- admits two non-proportional such derivations if  $X \in (\text{ML}_0)$ .

**1.3.** There is a combinatorial counterpart of the ML-classification, see Lemma 1.4. Let  $X$  be a normal affine surface, and let  $V$  be a completion of  $X$  with boundary divisor  $D = V \setminus X$ . Assume that  $(V, D)$  is a minimal NC-completion, that is,  $V$  is smooth near  $D$  and  $D$  is a normal crossing divisor such that no (smooth, rational)  $(-1)$ -component of  $D$  can be contracted without losing the NC-property. Let  $\Gamma = \Gamma(D)$  be the weighted dual graph of  $D$ . It is easily seen that if  $X$  admits an  $\mathbb{A}^1$ -fibration then  $(V, D)$  is in fact an SNC-completion, and  $\Gamma$  is a tree.

A linear weighted graph is called *admissible* if all its weights are  $\leq -2$ . A non-admissible linear chain can be transformed birationally into a chain with an end vertex of weight 0.

**Lemma 1.4** ([42, Rem. 1.7]). *The graph  $\Gamma$  as in 1.3*

- *has only admissible extremal linear branches for  $X \in (\text{ML}_2)$ ;*
- *is non-linear and has a non-admissible extremal linear branch for  $X \in (\text{ML}_1)$  different from  $\mathbb{A}^1 \times \mathbb{A}_*^1$ ;*
- *is a linear chain (a zigzag) non-transformable into the linear chain with weights  $[[0, 0, 0]]$  for  $X \in (\text{ML}_0)$ .*

**1.2. Classification according to rank.** The *rank* of an ind-group  $G$  acting morphically and effectively on a variety  $X$  is the maximal dimension of an algebraic torus contained in  $G$ . This rank does not exceed the dimension of  $X$ , and  $X$  is toric in the case of equality. A surface  $X$  with  $\text{rk Aut } X \geq 1$  is called a  $\mathbb{G}_m$ -surface. The rank distinguishes toric surfaces, non-toric  $\mathbb{G}_m$ -surfaces, and surfaces without  $\mathbb{G}_m$ -actions; indeed, their ranks are 2, 1, and 0, respectively.

**1.3. A general classification scheme.** The two independent classifications of normal affine surfaces, according to the rank of the automorphism group and according to the Makar–Limanov class as defined before, give altogether 9 classes of affine surfaces denoted  $(\text{ML}_i, r)$ ,  $(i, r) \in \{0, 1, 2\}^2$ , where  $r$  is the rank of  $\text{Aut } X$  and  $i$  the Makar–Limanov complexity of  $X$ .

To describe the automorphism groups of affine surfaces and the algebraic groups actions on them, one applies various means. Some of these found their place in our survey. The content of the present notes is as follows.

- In Section 2 we introduce different classes of groups: ind-groups, nested ind-groups, amalgams, bearable groups.
- In Section 3 we study classical examples of affine surfaces, including the toric surfaces, along with a presentation of their automorphism groups as amalgams.
- Section 4 contains generalities on algebraic group actions on affine surfaces;
- in Subsection 4.4 we classify affine surfaces of rank  $\geq 1$ , along with one-parameter groups acting on such surfaces, in terms of the DPD presentation.
- In Section 5 we consider the automorphism groups of the  $\text{ML}_0$ -surfaces, also called Gizatullin surfaces. We provide a classification of their one-parameter subgroups. For some particular classes of Gizatullin surfaces, we describe their automorphism groups as amalgams. The structure of the automorphism groups of general Gizatullin surfaces remains mysterious.
- In Sections 6 we study the automorphism groups of  $\mathbb{A}^1$ -fibrations as nested ind-groups.

---

<sup>1</sup>The surface  $X = \mathbb{A}^1 \times \mathbb{A}_*^1$  of class  $(\text{ML}_1)$  admits an SNC-completion  $(X, D)$  with  $\Gamma(D)$  being the linear chain with weights  $[[0, 0, 0]]$ .

- Sections 7 – 8 deal with  $\mathbb{A}^1$ -fibrations on affine surfaces. In Section 7 we introduce Puiseux arc spaces and study the actions of automorphisms on these spaces.
- In Section 8 this techniques is applied in order to describe the automorphism groups of  $\mathbb{A}^1$ -fibrations on affine surfaces in more detail. The neutral components of these automorphism groups occur to be metabelian nested ind-groups of rank at most two, while the discrete part is at most countable, see Theorems 8.12, 8.24, and Corollary 8.25.

## 2. IND-GROUPS, AMALGAMS, AND ALL THIS

**2.1. Ind-groups.** Recall (see [65, 67, 96, 97]) that an *ind-group*  $G$  is an inductive limit  $G = \varinjlim \Sigma_i$  of an increasing sequence of algebraic varieties

$$\Sigma_1 \subset \Sigma_2 \subset \dots \subset \Sigma_n \subset \dots$$

with closed embeddings  $\Sigma_i \subset \Sigma_{i+1}$ , where  $G$  is endowed with a group structure such that for each pair  $(i, j) \in \mathbb{N}^2$  the multiplication  $(f, g) \mapsto f \cdot g^{-1}$  yields a morphism  $\Sigma_i \times \Sigma_j \rightarrow \Sigma_{n(i,j)}$  with a suitable  $n(i, j) \in \mathbb{N}$ .

The *neutral component*  $G^\circ$  of an ind-group  $G = \varinjlim \Sigma_i$  is defined as the inductive limit of the connected components of  $\Sigma_i$  passing through the neutral element  $e \in G$ .

A *morphism of ind-groups*  $G = \varinjlim \Sigma_i$  and  $G' = \varinjlim \Sigma'_i$  is a group homomorphism  $\phi: G \rightarrow G'$  such that for any  $i \geq 1$ ,  $\phi(\Sigma_i) \subset \Sigma'_j$  for some  $j = j(i) \geq 1$ , and  $\phi|_{\Sigma_i}: \Sigma_i \rightarrow \Sigma'_j$  is a morphism of varieties. Clearly,  $\phi(G^\circ) \subset G'^\circ$ .

A subgroup  $H \subset G$  of an ind-group  $G = \varinjlim \Sigma_i$  is *closed* if for any  $i \geq 1$ , the intersection  $H \cap \Sigma_i$  is closed in  $\Sigma_i$ . In the latter case  $H = \varinjlim (H \cap \Sigma_i)$  is an ind-group.

One says that an ind-group  $G = \varinjlim \Sigma_i$  *acts morphically* on a variety  $X$  if there is an action  $G \times X \rightarrow X$  of  $G$  on  $X$  such that for each  $i \in \mathbb{N}$  the restriction  $\Sigma_i \times X \rightarrow X$  is a morphism of algebraic varieties. The following lemma is well known in the affine case, see, e.g., [65, Prop. 2.5]; the first (unpublished) proof is due to Bialynicki-Birula, see [58, Rem. after Cor. 1.2].

**Lemma 2.1.** *For a quasi-affine algebraic variety  $X$ , the automorphism group  $\text{Aut } X$  has a natural structure of an ind-group acting morphically on  $X$ .*

*Proof.* Choose a finitely generated subalgebra  $A = \mathbb{K}[a_1, \dots, a_n]$  of  $\mathcal{O}_X(X)$  such that  $X$  embeds as a dense, open subset in the affine variety  $X' = \text{Spec } A$ . The system  $(a_1, \dots, a_n)$  extends to a system of generators  $(a_1, \dots, a_n, a_{n+1}, \dots)$  of  $\mathcal{O}_X(X)$ . Let  $A_d = \mathbb{K}[a_1, \dots, a_d] \subset \mathcal{O}_X(X)$ . For any  $g \in \text{Aut } X$  there is  $d = d(g)$  such that  $g(A) \subset A_d$ . Consider the surjection

$$\mathbb{K}[x_1, \dots, x_d] \rightarrow A_d \cong \mathbb{K}[x_1, \dots, x_d]/I_d, \quad x_i \mapsto a_i,$$

where  $I_d$  is the corresponding ideal of relations. Let  $V_{d,m}$  be the image in  $A_d$  of the subspace  $\mathbb{K}[x_1, \dots, x_d]_{\leq m}$  of polynomials of degree at most  $m$ , and let

$$\Sigma_d = \{g \in \text{Aut } X \mid g(a_i), g^{-1}(a_i) \in V_{d,d} \forall i = 1, \dots, n\}.$$

It is easily seen that  $\text{Aut } X = \varinjlim \Sigma_d$  is an ind-group. The last assertion is now immediate.  $\square$

**Remark 2.2.** According to Ramanujam ([88]), a family of automorphisms in  $\text{Aut } X$  parameterized by a variety  $R$  is called a *subvariety of  $\text{Aut } X$*  if the action  $R \times X \rightarrow X$ ,  $(r, x) \mapsto r(x)$ , is a morphism. In this case the natural map  $R \rightarrow \text{Aut } X$  is a morphism of ind-varieties. The *neutral component of  $\text{Aut } X$  in the sense of Ramanujam* is the union of all the irreducible subvarieties in  $\text{Aut } X$  passing through the neutral element

$e \in \text{Aut } X$ . Clearly, it coincides with the neutral component  $\text{Aut}^\circ X$  of the ind-group  $\text{Aut } X$  as defined before.

**Proposition 2.3.** *Let  $X$  be an affine algebraic variety, and let  $I \subset \mathcal{O}_X(X)$  be an ideal. Let  $\text{Aut}(X, I)$  be the subgroup of  $\text{Aut } X$  of all automorphisms of  $X$  leaving  $I$  invariant. Then  $\text{Aut}(X, I)$  is a closed ind-subgroup in both  $\text{Aut } X$  and  $\text{Aut}(X \setminus \text{supp } I)$ .*

*Proof.* We use the notation  $V_d = V_{d,d}$  and  $\Sigma_d$  introduced in the proof of Lemma 2.1. Let  $I = (b_1, \dots, b_k)$ , where  $b_i \in \mathcal{O}_X(X)$ ,  $i = 1, \dots, k$ . Clearly,  $g \in \text{Aut}(X, I)$  if and only if  $b_i \circ g \in I \ \forall i = 1, \dots, k$ . Let us show that, given  $d$  and  $i$ , the latter condition defines a closed subset of  $\Sigma_d$ . Observe that there exists  $m = m(d, i)$  such that  $b_i \circ g \in V_m$  for any  $g \in \Sigma_d$ . Consider the map  $\psi_i: \text{Aut } X \rightarrow \mathcal{O}_X(X)$ ,  $g \mapsto g \circ b_i$ , and let  $\psi_{i,d} = \psi_i|_{\Sigma_d}: \Sigma_d \rightarrow V_m \subset \mathcal{O}_X(X)$ . By virtue of Lemma 2.1,  $\psi_{i,d}$  is a morphism. Since  $I \cap V_m$  is a linear subspace of the finite dimensional vector space  $V_m$ , the inverse image  $\psi_{i,d}^{-1}(I \cap V_m) = \psi_i^{-1}(I) \cap \Sigma_d$  is closed in  $\Sigma_d$ , as stated. It follows that  $\text{Aut}(X, I) \subset \text{Aut } X$  is a closed ind-subgroup.

To show the converse, letting  $Y = X \setminus \text{supp } I$  consider the subgroup  $H = \text{Aut}(X, I)$  of the ind-group  $G = \text{Aut } Y = \varinjlim \Sigma'_d$  (see Lemma 2.1). We must show that  $H$  is a closed ind-subgroup of  $G$ . Suppose first that the ideal  $I = \sqrt{I}$  is radical. Choose a system of generators  $a_1, \dots, a_n$  of  $A = \mathcal{O}_X(X)$ . Under our assumption we have

$$H = \{g \in G \mid a_i \circ g \in A \ \forall i = 1, \dots, n\}.$$

Consider the map  $\phi_i: G \rightarrow \mathcal{O}_Y(Y)$ ,  $g \mapsto a_i \circ g$ . For any  $d \in \mathbb{N}$ ,  $\Sigma'_d \subset G$  is an algebraic variety such that the natural map  $\Sigma'_d \times Y \rightarrow Y$  is a morphism. Let  $\eta_d: \mathcal{O}_Y(Y) \rightarrow \mathcal{O}_Y(Y) \otimes \mathcal{O}_{\Sigma'_d}(\Sigma'_d)$  be the induced morphism, and let

$$\eta_d(a_i) = \sum_{j=1}^m b_{i,j} f_{i,j} \quad \text{where } b_{i,j} \in \mathcal{O}_Y(Y) \quad \text{and} \quad f_{i,j} \in \mathcal{O}_{\Sigma'_d}(\Sigma'_d).$$

Letting  $W_{i,d} = \text{span}(b_{i,j} \mid j = 1, \dots, m) \subset \mathcal{O}_Y(Y)$ , for any  $g \in \Sigma'_d$  we obtain

$$\phi_i(g) = a_i \circ g = \sum_{j=1}^m b_{i,j} f_{i,j}(g) \in \bar{X}_{i,d}.$$

Since  $W_{i,d} \subset \mathcal{O}_Y(Y)$  is a finite dimensional subspace and  $W_{i,d} \cap A$  is as well,  $\phi_i^{-1}(W_{i,d} \cap A) = \phi_i^{-1}(A) \cap \Sigma'_d$  is closed in  $\Sigma'_d$  for any  $i, d \geq 1$ . Hence  $H \subset G$  is closed.

For an arbitrary ideal  $I$  of  $\mathcal{O}_X(X)$ , consider its radical  $\sqrt{I} \supset I$ . By the preceding,  $\text{Aut}(X, \sqrt{I})$  is closed in  $\text{Aut}(Y)$ . It remains to show that  $\text{Aut}(X, I)$  is a closed ind-subgroup of  $\text{Aut}(X, \sqrt{I})$ . However, the latter follows because both these groups are closed in  $\text{Aut } X$  by the first part of the proof.  $\square$

**Definition 2.4.** Let  $X$  be an affine variety. An element  $g \in \text{Aut } X$  is called *semisimple*, if there exists a finite-dimensional  $g$ -stable subspace  $V \subset \mathcal{O}_X(X)$  which contains a system of generators  $a_1, \dots, a_m$  of  $\mathcal{O}_X(X)$  and such that the image of  $g$  in  $\text{GL}(V^*)$  is semisimple.

Recall that an *algebraic quasitorus* is a product of an algebraic torus and a finite Abelian group.

**Lemma 2.5.** *An element  $g \in \text{Aut } X$  is semisimple if and only if  $g$  is contained in an algebraic quasitorus  $T \subset \text{Aut } X$ .<sup>2</sup>*

*Proof.* If  $g$  is semisimple then  $g \in \text{GL}(V^*)$  is contained in an algebraic torus  $T' \subset \text{GL}(V^*)$ . The Zariski closure  $T$  of the cyclic group  $\langle g \rangle \subset T'$  generated by  $g$  is an algebraic quasitorus, say,  $T \subset T'$ . Consider the natural embedding  $\phi: X \hookrightarrow V^*$ . Since

<sup>2</sup>We suppose, of course, that  $T \subset \text{Aut } X$  is a closed subgroup acting morphically on  $X$ .

$g$  leaves invariant the image  $\phi(X) \subset V^*$ , this is also true for  $T$ . Hence there is an algebraic quasitorus in  $\text{Aut } X$  containing  $g$ .

It is well known that an algebraic group  $G$  acting on  $X$  acts *locally finitely* on  $\mathcal{O}_X(X)$ , that is, each finite dimensional subspace of  $\mathcal{O}_X(X)$  extends to a finite dimensional  $G$ -invariant subspace. This implies that any  $g \in \text{Aut } X$  contained in an algebraic quasitorus in  $\text{Aut } X$  is semisimple.  $\square$

## 2.2. Nested ind-groups.

**Definition 2.6.** We say that a group  $G$  is a *nested ind-group* if

$$G = \varinjlim G_i, \quad \text{where } G_1 \subset \dots \subset G_i \subset G_{i+1} \subset \dots$$

is an increasing sequence of algebraic groups and their closed embeddings. The *rank* of a nested ind-group  $G$  is defined as  $\text{rk } G = \lim_{i \rightarrow \infty} \text{rk } G_i$ . If all the  $G_i$  are unipotent, then we say that the nested ind-group  $G$  is unipotent. The *unipotent radical*  $R_u(G)$  is the largest closed normal ind-subgroup of  $G$  such that every element  $g \in R_u(G)$  is unipotent in  $G_i$  for all  $i$  sufficiently large.<sup>3</sup>

**Remarks 2.7.** 1. An algebraic group is a nested ind-group. By contrast, the ind-group  $\mathbb{Z}$  is not a nested ind-group.

2. A closed subgroup of a nested ind-group is nested. If  $G = \varinjlim G_i$  is nested, then also its neutral component  $G^\circ = \varinjlim G_i^\circ$  is.

3. If  $G = \varinjlim G_i$  and  $G_i$  is connected for any  $i \in \mathbb{N}$ , then  $G$  is. Conversely, if  $G = \varinjlim G_i$  is connected as an ind-group, then  $G = \varinjlim G_i^\circ$ .

4. If a nested ind-group  $G = \varinjlim G_i$  is a connected ind-subgroup of the automorphism group  $\text{Aut } X$ , where  $X$  is an algebraic variety, then  $G$  is algebraically generated in the sense of [1, Def. 1.1].

Due to the last remark, the following result is immediate from [1, Prop. 1.7].

**Proposition 2.8.** *If a connected nested ind-group  $G = \varinjlim G_i$  acts morphically on an algebraic variety  $X$ , then the  $G$ -orbits in  $X$  coincide with the  $G_i$ -orbits in  $X$  for some  $i \geq 1$ .*

**Definition 2.9.** Let  $G$  be a nested ind-group, and let  $T$  be a maximal torus in  $G$  of finite rank. The *Cartan subgroup*  $C_G^\circ(T)$  associated to  $T$  is the neutral component of the centralizer  $C_G(T)$  of  $T$  in  $G$ .

Clearly,  $C_G^\circ(T)$  is a closed nested ind-subgroup of  $G$ , cf. Remarks 2.7.2-2.7.3. Note that if  $\text{rk } G = 0$ , i.e.,  $G^\circ$  is unipotent, then  $T = \{1\}$  and  $C_G^\circ(T) = G^\circ$ . In a semisimple algebraic group  $G$  one has  $C_G^\circ(T) = T$ , and in a product  $\tilde{G} = G \times U$ , where  $G$  is semisimple and  $U$  is unipotent, one has  $C_{\tilde{G}}^\circ(T) = T \times U$ .

In the following proposition we provide an analog of the classical fact on the conjugacy classes in algebraic groups. Another analog of this fact is known to hold in the group  $\text{Aut } \mathbb{A}^2$ , which is not a nested ind-group ([43]).

**Proposition 2.10.** *Let  $G = \varinjlim G_i$  be a connected nested ind-group. Then for any element  $g$  of a Cartan subgroup  $C_G^\circ(T)$ , the conjugacy class  $\text{Cl}_G(g)$  of  $g$  is closed in  $G$ .*

*Proof.* Due to Remark 2.7.3 one may assume that all the  $G_i$ ,  $i \geq 1$ , are connected algebraic groups. For any element  $g$  of a connected algebraic group  $H$  the conjugacy class  $\text{Cl}_H(g)$  of  $g$  in  $H$  is an irreducible locally closed subvariety of  $H$ . We will use the fact that an increasing sequence of irreducible locally closed subvarieties in an algebraic variety stabilizes.

<sup>3</sup>See the notions of a locally linear ind-group and of its unipotent radical in [25, I.3].

Let  $g \in C_G^\circ(T)$ . Then for  $i \geq 1$  sufficiently large we have  $T \subset G_i$  and  $g \in C_{G_i}^\circ(T)$ . Hence the conjugacy class  $\text{Cl}_{G_i}(g)$  of  $g$  is closed in  $G_i$  ([98, Prop. 6.14]). On the other hand, for any  $j > 0$ ,

$$\text{Cl}_G(g) \cap G_j = \varinjlim (\text{Cl}_{G_i}(g) \cap G_j)$$

is closed being an increasing union of closed irreducible subsets, which stabilizes. Thus,  $\text{Cl}_G(g)$  is closed in  $G$ .  $\square$

In a nested ind-group of automorphisms, one has the following analog of the Mostow decomposition in a connected algebraic group ([81]; cf. [71, Thm. 4.10]).

**Theorem 2.11.** *Let  $X$  be a quas affine algebraic variety, and let  $G = \varinjlim G_i \subset \text{Aut } X$  be a connected nested subgroup. Then  $G$  admits a Mostow decomposition  $G = R_u(G) \rtimes L$ , where  $L$  is a maximal reductive algebraic subgroup in  $G$  and  $R_u(G)$  is the unipotent radical of  $G$ . Furthermore, any semisimple element  $g \in G$  is contained in a maximal torus of  $G$ , and any two such tori are conjugated in  $G$ .*

*Proof.* Due to Remark 2.7.3 one may assume that  $G_i$  is connected for any  $i \geq 1$ . For each  $i \gg 1$  consider a Mostow decomposition  $G_i = U_i \rtimes L_i$ , where  $U_i = R_u(G_i)$ . Since  $G_i$  is connected,  $L_i$  is connected as well for  $i \gg 1$ . The rank of the Levi factor  $L_i$  is bounded by  $\dim X$  (see Proposition 4.1.a). Hence the dimensions  $\dim L_i$  are uniformly bounded. If  $L = L_k$  is of maximal dimension, then  $L$  is a Levi subgroup of  $G_i$  for any  $i \geq k$ . Thus,  $G_i = U_i \rtimes L \ \forall i \geq k$ .

Let us show that  $U_i = U_{i+1} \cap G_i$  for  $i \geq k$ . Indeed, since  $L \subset G_i \subset U_{i+1} \rtimes L$ , given  $(u, l) \in (U_{i+1} \rtimes L) \cap G_i$ , one has  $(u, 1) = (u, l) \cdot (1, l^{-1}) \in G_i$ . Thus,  $U_i = U_{i+1} \cap G_i$ . Clearly,  $R_u(G) = \varinjlim U_i$ , and so, the first assertion follows. The remaining statements hold because they hold for any connected algebraic subgroup  $G_i$ ,  $i \gg 1$ , see, e.g., [52, Prop. 19.4 and Cor. 21.3.A].  $\square$

**Corollary 2.12.** *Let  $G = R_u(G) \rtimes L$  and  $G_i = U_i \rtimes L$  be as before. If the unipotent radical  $R_u(G)$  is Abelian, then there exists a decomposition  $R_u(G) = \bigoplus_{j=1}^\infty H_j$  such that  $H_i$  is normal in  $G$  for all  $i \geq 1$  and  $U_i = \bigoplus_{j=1}^i H_j$ .*

*Proof.* Since the adjoint representation of  $L$  on the Lie algebra  $\text{Lie } G_i$  is completely reducible, and the subalgebras  $\text{Lie } U_i$  and  $\text{Lie } U_{i-1}$  are  $L$ -stable, there is an  $L$ -stable subspace  $V_i \subset \text{Lie } U_i$  complementary to  $\text{Lie } U_{i-1}$ . By our assumption,  $U_i$  is Abelian. Hence the Lie subalgebra  $V_i$  corresponds to a subgroup  $H_i \subset U_i$  normalized by  $L$  and  $U_i = U_{i-1} \oplus H_i$ . Now the assertions follow.  $\square$

**Example 2.13.** Let  $X$  be a normal affine surface, let  $\mu: X \rightarrow B$  be an  $\mathbb{A}^1$ -fibration over a smooth affine curve  $B$ , and let  $\text{Aut}(X, \mu)$  be the group of all automorphisms of  $X$  preserving  $\mu$ . Then the neutral component  $\text{Aut}^\circ(X, \mu)$  is a nested ind-group with an Abelian unipotent radical, see Corollary 8.25. Hence Corollary 2.12 applies to this case.

**Lemma 2.14.** *Let  $\phi: G \rightarrow H$  be a morphism of nested ind-groups  $G = \varinjlim G_i$  and  $H = \varinjlim H_j$ . Assume that the orders  $|G_i/G_i^\circ|$ ,  $i \in \mathbb{N}$ , are bounded above. Then  $\phi(G)$  is a closed nested ind-subgroup in  $H$ .*

*Proof.* For a fixed  $j$  consider the increasing sequence of algebraic subgroups  $\phi(G_i) \cap H_j$ ,  $i \in \mathbb{N}$ , of the group  $H_j$ . Since their dimensions and the numbers of connected components are bounded, this sequence stabilizes. Hence  $\phi(G) \cap H_j$  is an algebraic subgroup.  $\square$



**2.3. Amalgams.** Recall ([95]) that a *tree of groups*  $(T, \mathcal{G})$  consists in a combinatorial tree  $T$  along with a collection  $\mathcal{G}$  of vertex groups  $(G_P)_{P \in \text{vert } T}$ , edge groups  $(G_\nu)_{\nu \in \text{edge } T}$ , and for each edge  $\nu = [P, Q]$  of  $T$ , monomorphisms  $G_\nu \rightarrow G_P$  and  $G_\nu \rightarrow G_Q$  identifying  $G_\nu$  with (common) subgroups of the vertex groups  $G_P$  and  $G_Q$ . Given such a tree of groups  $(T, \mathcal{G})$ , one can construct a unique group  $G = *_T G_P$  called the *free amalgamated product*, or simply the *amalgam* of  $(T, \mathcal{G})$ , where  $G$  is freely generated by the subgroups  $(G_P)$  and  $(G_\nu)$  with joint subgroups  $G_P \cap G_Q = G_\nu$  for each  $\nu = [P, Q] \in \text{edge } T$ . We refer to [95, Ch. I, §§4,5] for the existence and uniqueness of the amalgam  $G = *_T G_P$ , its presentation and the universal property.

**Serre Theorem 2.15** ([95, Ch. I, §4.3, Thm. 8 and §4.4, Ex. 2]). *Any subgroup of bounded length of an amalgam  $G = *_T G_P$  is contained in a conjugate to one of the factors  $G_P$ , where  $P \in \text{vert } T$ .*

The conclusion of Serre's Theorem holds true for algebraic subgroups of  $\text{Aut } \mathbb{A}^2$ , where  $\text{Aut } \mathbb{A}^2$  is endowed with its usual amalgam structure, see [58, Cor. 4.2] and [101]. More generally, we have the following result.

**Proposition 2.16.** *Let  $F = \varinjlim \Sigma_i$  be an ind-group<sup>4</sup>, and let  $G = *_T G_P \subset F$  be the amalgam of a tree of groups  $(T, \mathcal{G})$ , where the vertex groups  $G_P$  are ind-groups. Then any algebraic subgroup  $H \subset F$  contained in  $G$  is of bounded length, and so, is conjugated to a subgroup of some vertex group  $G_P$ ,  $P \in \text{vert } T$ .*

*Proof.* By Serre's Theorem 2.15, the latter assertion follows from the former. Thus, it suffices to show that  $H$  is of bounded length in  $G$ . Let  $G_P = \varinjlim \Sigma_{P,n}$  be the ind-structure of the group  $G_P$  for  $P \in \text{vert } T$ . Given a finite sequence  $((P_1, n_1), \dots, (P_l, n_l)) \in ((\text{vert } T) \times \mathbb{N})^l$ , consider the morphism  $\Sigma_{P_1, n_1} \times \dots \times \Sigma_{P_l, n_l} \rightarrow F$  induced by the multiplication. Its image is a constructible subset of the variety  $\Sigma_i$  for some  $i \in \mathbb{N}$ . The group  $G$  is covered by these constructible sets. If  $E$  is an algebraic subvariety of  $G$ , then  $E$  is covered by a finite number of them, and so,  $E$  has bounded length. In particular, any algebraic subgroup  $H$  of  $G$  has bounded length, as claimed.  $\square$

**Remark 2.17** (*Pushing forward amalgamated free product structures*). Let  $X' \rightarrow X$  be an étale Galois covering with the Galois group  $\Gamma$ , where  $X$  and  $X'$  are affine varieties. Assume that every automorphism  $\alpha \in \text{Aut } X$  admits a lift to an automorphism  $\tilde{\alpha} \in \text{Aut } X'$ . By the Monodromy Theorem, the latter holds, for instance, if  $\mathbb{K} = \mathbb{C}$  and  $X' \rightarrow X$  is the universal covering. Clearly, under this assumption the subset in  $\text{Aut } X'$  of all lifts of the automorphisms in  $\text{Aut } X$  coincides with the normalizer  $\text{Norm}_{\text{Aut } X'}(\Gamma)$  of  $\Gamma$  in  $\text{Aut } X'$ . Furthermore, we have  $\text{Aut } X \cong \text{Norm}_{\text{Aut } X'}(\Gamma)/\Gamma$ .

Assume that  $\text{Aut } X'$  admits a structure of an amalgamated free product  $\text{Aut } X' \cong A' *_C B'$ , where  $C' = A' \cap B' \supset \Gamma$ . Then we have the inclusion

$$(1) \quad \text{Norm}_{\text{Aut } X'}(\Gamma) \supseteq (\text{Norm}_A(\Gamma)) *_C (\text{Norm}_B(\Gamma)) .$$

This inclusion can be strict, in general. However, in case of equality the following holds (cf. [2, Lemma 4.14]).

**Lemma 2.18.** *In the setup as before, assume that the equality holds in (1). Then  $\text{Aut } X \cong A *_C B$  is an amalgam of  $A = \text{Norm}_A(\Gamma)/\Gamma$  and  $B = \text{Norm}_B(\Gamma)/\Gamma$  along the joint subgroup  $C = A \cap B = \text{Norm}_C(\Gamma)/\Gamma$ .*

**2.4. Bearable automorphism groups.** Let us introduce the following notions.

**Definition 2.19.** A group  $G$  will be called  $\alpha$ -*bearable*, where  $\alpha$  is a cardinal number (e.g., *finitely bearable*, *countably bearable*, etc.) if  $G$  is an amalgam of a tree  $(T, \mathcal{H})$

<sup>4</sup>For instance,  $F$  can be the automorphism group of an affine variety.

of nested ind-groups  $H_P$ ,  $P \in \text{vert } T$ , where  $\text{vert } T$  has cardinal  $\alpha$ . A group will be called *bearable* if it is  $\alpha$ -bearable for some cardinal  $\alpha$ , and *unbearable* otherwise.

**Remarks 2.20.** 1. It is easily seen that a nontrivial bearable group  $G = *_T G_P$  is a nested ind-group if only if  $T$  is reduced to a single vertex.

2. A connected bearable group of automorphisms is algebraically generated in the sense of [1, Def. 1.1]. Hence its orbits are locally closed smooth subvarieties ([1, Prop. 1.3]).

3. For a smooth affine surface  $X$ , the group  $\text{SAut}(X)$  generated by its one-parameter unipotent subgroups (see 1.1) can have an open orbit in  $X$ , which is not closed. The corresponding examples are due to Gizatullin, Danilov, and the first author, see [62] and the references therein.

We need the following fact. <sup>5</sup>

**Theorem 2.21.** *Let  $G$  be a reductive algebraic group. Then the set of conjugacy classes of connected reductive subgroups of  $G$  is at most countable.*

The proof is based on the following result of Richardson ([90, Cor. 11.5(b) and Prop. 12.1], see also [91, Thm. 8.1]). Alternatively, the lemma can be deduced from the classification of semisimple subgroups of reductive groups started in [28].

**Lemma 2.22.** *The set of conjugacy classes of connected semisimple algebraic subgroups of a reductive algebraic group  $G$  is finite.*

*Proof of Theorem 2.21.* Let  $H \subset G$  be a connected reductive subgroup. Consider the Levi decomposition  $H = S \cdot T$ , where  $T = \text{Rad } H$  is a central torus in  $H$  and  $S = [H, H]$  is the commutator subgroup, see [12, Prop. 2.2]. Since  $S$  is semisimple, its conjugacy class in  $G$  is chosen among a finite set of such classes, see the claim. The torus  $T$  is contained in the centralizer  $C_G(S)$ . Together with  $S$ , the centralizer  $C_G(S)$  also runs over a finite set of conjugacy classes in  $G$ , along with a maximal torus in  $C_G(S)$  which contains  $T$ , since any two maximal tori in  $C_G(S)$  are conjugated. Furthermore, the conjugation by elements of  $C_G(S)$  act trivially on  $T$ , hence  $T$  is contained in any maximal torus of  $C_G(S)$ . Fixing one of them, say,  $\mathcal{T}$ , by the rigidity of subtori of  $\mathcal{T}$  there is at most countable number of possibilities to choose  $T \subset \mathcal{T}$ . The conjugacy class of  $H = S \cdot T$  is then also chosen among at most countable number of such classes.  $\square$

The following extension of Theorem 2.21 to the reductive subgroups in countably bearable groups will be used in § 5.2, see Corollary 5.6.

**Proposition 2.23.** *In a countably bearable group  $G$ , the set of conjugacy classes of connected reductive algebraic subgroups is at most countable.*

*Proof.* Any connected reductive algebraic subgroup  $F$  in a nested group  $H = \varinjlim H_i$  is contained in an algebraic subgroup  $H_i$  for some  $i$ , and, moreover, in a maximal connected reductive subgroup of  $H_i$ . By the Mostow decomposition theorem, any two maximal connected reductive subgroups are conjugated in  $H_i$ . By Theorem 2.21, the conjugacy class of  $F$  in  $H_i$  runs over at most countable set of such classes. It follows that the set of conjugacy classes of connected reductive algebraic subgroups of  $H$  is at most countable.

By Proposition 2.16, any algebraic subgroup of the countably bearable group  $G$  is conjugated to an algebraic subgroup of one of the countable collection of generating nested ind-groups. Hence the set of conjugacy classes of connected reductive algebraic subgroups of  $G$  is at most countable.  $\square$

<sup>5</sup>The authors are grateful to V. Arzhantsev, R. Avdeev, M. Borovoi, M. Brion, D. Panushev, G. Soifer, D. Timashev, and E. Vinberg for useful discussions and indications.

**Remark 2.24.** The same argument shows that in an  $\alpha$ -bearable group  $\text{Aut } X$  of rank  $r$  the set of conjugacy classes of  $r$ -tori has cardinality at most  $\alpha$ .

**Example 2.25.** Consider, for instance, the Danielewski surface  $S_n = \{x^n y - (z^2 - 1) = 0\}$  in  $\mathbb{A}^3$ . The group  $\text{Aut } S_n$  is 2-bearable, see Theorem 5.26. However, the group  $\text{Aut}(S_n \times \mathbb{A}^1)$  is not finitely bearable. Indeed ([Da]), this group of rank 2 contains a sequence of pairwise non-conjugated 2-tori. Is this group countably bearable?

Similarly, using Proposition 2.16 we obtain the following result. Recall that two  $\mathbb{A}^1$ -fibrations on an affine variety  $X$  are called *equivalent* if one can be transformed into the other by an automorphism of  $X$ .

**Proposition 2.26.** *Let  $X$  be a normal affine variety. Assume that  $X$  does not admit a unipotent group action with a general orbit of dimension  $\geq 2$ .<sup>6</sup> If the group  $\text{Aut}^\circ X$  is  $\alpha$ -bearable for some cardinal  $\alpha$ , then the set of non-equivalent  $\mathbb{A}^1$ -fibrations on  $X$  over affine bases is of cardinality at most  $\alpha$ .*

*Proof.* Let  $\text{Aut}^\circ X = \ast_T G_P$ , where  $(T, \mathcal{G})$  is a tree of nested ind-groups  $(G_P)_{P \in \text{vert } T}$ , where  $\text{vert } T$  is a set of cardinality  $\alpha$ .

For any  $\mathbb{A}^1$ -fibration  $\mu: X \rightarrow Z$ , where  $Z$  is a normal affine variety, one can find a one-parameter unipotent subgroup  $U_\mu \subset \text{Aut } X$  acting along the fibers of  $\mu$ .

Assume to the contrary that the set of pairwise non-equivalent  $\mathbb{A}^1$ -fibrations on  $X$  with affine bases is of cardinal larger than  $\alpha$ . By Proposition 2.16, any one-parameter group  $U_\mu$  is conjugated to a subgroup  $U'_\mu$  of one of the vertex groups  $G_P$ ,  $P \in \text{vert } T$ . From our assumption, for some vertex  $P \in \text{vert } T$ , the vertex group  $G_P$  contains at least two one-parameter unipotent subgroup  $U'_\mu \subset \text{Aut } X$ , which act along pairwise non-equivalent  $\mathbb{A}^1$ -fibrations on  $X$ .

Since  $G_P = \varinjlim G_{P,n}$  is a nested ind-group, any one-parameter unipotent subgroup in  $G_P$  is contained in some algebraic subgroup  $G_{P,n}$ . It follows that for some  $n \geq 1$ , the algebraic group  $H = G_{P,n}$  contains at least two unipotent one parameter subgroups  $U'_{\mu_1}, U'_{\mu_2}$  acting along two non-equivalent  $\mathbb{A}^1$ -fibrations  $\mu_i: X \rightarrow Z_i$ ,  $i = 1, 2$ .

Let  $U_{\max}$  be a maximal unipotent subgroup of  $H$ .<sup>7</sup> Since any two such subgroups are conjugated, any one-parameter unipotent subgroup  $U \subset H$  is conjugated to a subgroup of  $U_{\max}$ . Hence  $U_{\max}$  contains two one-parameter unipotent subgroups, say,  $U_1, U_2$  acting on  $X$  along two non-equivalent  $\mathbb{A}^1$ -fibrations. It follows that the general orbits of  $U_{\max}$  in  $X$  are at least two-dimensional, contrary to our assumption.  $\square$

### 3. AUTOMORPHISM GROUPS AND AMALGAMS: THE FIRST EXAMPLES

**3.1 (Classical surfaces).** In this section we describe the automorphism groups of the surfaces

$$(2) \quad \mathbb{A}^2, \quad \mathbb{A}^1 \times \mathbb{A}_*^1, \quad (\mathbb{A}_*^1)^2, \quad V_{d,e} = \mathbb{A}^2 / \mu_{d,e}, \quad \mathbb{P}^2 \setminus C, \quad \text{and} \quad (\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta,$$

where  $C \subset \mathbb{P}^2$  is a smooth conic,  $\Delta \subset \mathbb{P}^1 \times \mathbb{P}^1$  the diagonal, and for any  $1 \leq e < d$  with  $\text{gcd}(d, e) = 1$ ,  $\mu_{d,e} \subset \text{GL}(2, \mathbb{K})$  stands for the cyclic group  $\{\text{diag}(\zeta, \zeta^e) \mid \zeta^d = 1\}$ . For any one of these surfaces, its automorphism group carries an amalgam structure.

**3.2 (Toric affine surfaces).** These are the normal affine surfaces  $X$  with the group  $\text{Aut } X$  of rank 2. Any toric affine surface is one from the list

$$(3) \quad \mathbb{A}^2, \quad \mathbb{A}^1 \times \mathbb{A}_*^1, \quad (\mathbb{A}_*^1)^2, \quad \text{and} \quad V_{d,e}.$$

The torus action on  $X$  comes from the action of the diagonal 2-torus  $\mathbb{T}$  on  $\mathbb{A}^2$  (see [16]). The smooth toric surfaces  $\mathbb{A}^2$ ,  $\mathbb{A}^1 \times \mathbb{A}_*^1$ , and  $(\mathbb{A}_*^1)^2$  are the underlying

<sup>6</sup>The latter holds, in particular, for any affine surface different from  $\mathbb{A}^2$ , see Proposition 4.1(b).

<sup>7</sup>This means that  $U_{\max}$  is the set of the unipotent elements in a Borel subgroup of  $H$ .

homogeneous spaces of the solvable algebraic groups  $\mathbb{G}_a^2$ ,  $\text{Aff } \mathbb{A}^1 \cong \mathbb{G}_a \rtimes \mathbb{G}_m$ , and  $\mathbb{T} \cong \mathbb{G}_m^2$ , respectively. The toric affine surfaces fall into 3 classes as follows:

- $(\text{ML}_0, 2) = \{\mathbb{A}^2, V_{d,e}\};$
- $(\text{ML}_1, 2) = \{\mathbb{A}^1 \times \mathbb{A}_*^1\};$
- $(\text{ML}_2, 2) = \{(\mathbb{A}_*^1)^2\}.$

In 3.3–3.6 we describe their automorphism groups.

**3.3** (*The group  $\text{Aut } \mathbb{A}^2$* ). By the Jung–van der Kulk Theorem<sup>8</sup> (see [58, Thm. 2] and the references therein),  $\text{Aut } \mathbb{A}^2$  is the amalgamated free product  $A \star_C B$  of the affine group  $A = \text{Aff}(\mathbb{A}^2)$  and the de Jonquères subgroup  $B = \text{J}(\mathbb{A}^2)$  over their intersection  $C = A \cap B$ . The solvable group  $\text{J}(\mathbb{A}^2)$  of rank 2 can be decomposed as  $\text{J}(\mathbb{A}^2) \cong \mathbb{U} \rtimes \mathbb{T}$ , where the unipotent radical  $\mathbb{U} = \text{R}_u(\text{J}(\mathbb{A}^2))$  is Abelian and consists of the triangular transformations  $(x, y) \mapsto (x + f(y), y)$  with  $f \in \mathbb{K}[y]$ . In particular,  $\text{J}(\mathbb{A}^2)$  is a nested solvable ind-group, see Definition 2.6. More precisely,  $\text{J}(\mathbb{A}^2)$  is an inductive limit of a sequence of solvable, connected affine algebraic groups of rank 2 with Abelian unipotent radicals.

It follows that any algebraic subgroup  $G \subset \text{Aut } \mathbb{A}^2$  is conjugate to a subgroup of one of the factors  $\text{Aff}(\mathbb{A}^2)$  and  $\text{J}(\mathbb{A}^2)$ , see [57, 4.3–4.4], [101]. If  $G$  is conjugate to a subgroup of  $\text{J}(\mathbb{A}^2)$ , then  $G$  is solvable of rank  $\leq 2$ . Hence, if a subgroup  $G \subset \text{Aut } \mathbb{A}^2$  is algebraic and non-solvable, then  $G$  is conjugated to a subgroup of  $\text{Aff}(\mathbb{A}^2)$  (cf. [58, Cor. 4.4.]). Any reductive subgroup  $G \subset \text{Aut } \mathbb{A}^2$  is linearizable, i.e., is conjugated to a subgroup of  $\text{GL}(2, \mathbb{K})$  ([66, Thm. 2.3]). Therefore, it is conjugated either to a subgroup of  $\mathbb{T}$ , or to  $\text{SL}(2, \mathbb{K})$ , or finally to  $\text{GL}(2, \mathbb{K})$ .

**3.4** (*Automorphism groups of singular toric affine surfaces*). For the toric affine surfaces  $X = \mathbb{A}^2/\mu_{d,e}$  of class  $(\text{ML}_0)$ , there are analogs of the Jung–van der Kulk and Kambayashi–Wright Theorems, see [2, Thms. 4.2, 4.15, 4.17]. Once again,  $\text{Aut } X$  is an amalgamated free product  $A^+ \star_C A^-$ , where  $C = A^+ \cap A^-$ . To describe this decomposition in more detail, one has to distinguish between the following cases:

- (i)  $e = 1$ ;
- (ii)  $e^2 \equiv 1 \pmod{d}$  and  $e \neq 1$ ;
- (iii)  $e^2 \not\equiv 1 \pmod{d}$ .

Note that the involution  $\tau : (x, y) \mapsto (y, x)$  acting on  $\mathbb{A}^2$  normalizes the cyclic subgroup  $\mu_{d,e}$  in  $\text{GL}(2, \mathbb{K})$  in cases (i) and (ii), and does not normalize it in case (iii).

In case (i)  $X = \mathbb{A}^2/\mu_{d,1}$  is the *Veronese cone*  $V_d$ , i.e. the affine cone over a rational normal curve in  $\mathbb{P}^d$ . Since  $\mu_{d,1} \subset \mathbb{T}$  is central, the standard  $\text{SL}(2, \mathbb{K})$ -action on  $\mathbb{A}^2$  descends to  $V_d$ . The complement to the vertex of  $V_d$  is the open  $\text{SL}(2, \mathbb{K})$ -orbit; the same is true for the natural  $\text{GL}(2, \mathbb{K})$ -action on  $V_d$ . Thus  $V_d$  is a quasihomogeneous  $\text{SL}(2, \mathbb{K})$ -variety.

The amalgam structure on  $\text{Aut } X$  is naturally related to that on  $\text{Aut } \mathbb{A}^2$ , see 3.3. Consider the normalizer

$$\mathcal{N}_{d,e}^+ = \text{Norm}_{\text{J}(\mathbb{A}^2)}(\mu_{d,e}) = \mathbb{U}_{d,e} \rtimes \mathbb{T} \subset \text{J}(\mathbb{A}^2),$$

where

$$\mathbb{U}_{d,e} = \{(x, y) \mapsto (x + f(y), y) \mid f \in y^e \mathbb{K}[y^d]\} \subset \mathbb{U}.$$

In case (i) we have  $\text{Aut } V_d \cong A^+ \star_C A^-$ , where

$$A^+ = \mathcal{N}_{d,1}^+/\mu_{d,1}, \quad A^- = \text{GL}(2, \mathbb{K})/\mu_{d,1}, \quad \text{and} \quad C = A^+ \cap A^-$$

(cf. [23, §11]).

---

<sup>8</sup>Valid over an arbitrary field.

Similarly, in case (ii) we have  $\text{Aut } X \cong A^+ *_C A^-$ , where

$$A^+ = \mathcal{N}_{d,e}^+ / \mu_{d,e}, \quad A^- = \langle \mathbb{T}, \tau \rangle / \mu_{d,e}, \quad \text{and} \quad C = A^+ \cap A^-.$$

Finally, in case (iii) we have  $\text{Aut } X \cong A^+ *_C A^-$ , where

$$A^\pm = \mathcal{N}_{d,e}^\pm / \mu_{d,e} \quad \text{with} \quad \mathcal{N}_{d,e}^- = \tau \mathcal{N}_{d,e}^+ \tau \quad \text{and} \quad C = A^+ \cap A^-.$$

**3.5** (*The group  $\text{Aut}(\mathbb{A}^1 \times \mathbb{A}_*^1)$* ). For the surface  $X = \mathbb{A}^1 \times \mathbb{A}_*^1$  of class  $(\text{ML}_1)$  we have

$$\text{Aut } X \cong \text{Aut}^\circ X \rtimes (\mathbb{Z} \rtimes (\mathbb{Z}/2\mathbb{Z})) = (U \rtimes \mathbb{T}) \rtimes (\mathbb{Z} \rtimes (\mathbb{Z}/2\mathbb{Z})),$$

where the factor  $\mathbb{Z}$  is generated by the transformation  $(x, y) \mapsto (xy, y)$ , the factor  $\mathbb{Z}/2\mathbb{Z}$  by the involution  $(x, y) \mapsto (x, y^{-1})$ , and the Abelian unipotent radical  $U$  of  $\text{Aut } X$  is

$$U = \left\{ \exp\left(p(y) \frac{d}{dx}\right) : (x, y) \mapsto (x + p(y), y) \mid p(y) \in \mathbb{K}[y, y^{-1}] \right\}.$$

The group  $\text{Aut } X$  is solvable, and so, any algebraic group acting effectively on  $\mathbb{A}^1 \times \mathbb{A}_*^1$  is as well. In fact, for any affine surface of class  $(\text{ML}_1)$  the automorphism group has similar properties, see Section 6.

**3.6** (*The group  $\text{Aut}(\mathbb{A}_*^1)^2$* ). If  $X = (\mathbb{A}_*^1)^2$ , then

$$\text{Aut } X = \left\{ (x, y) \mapsto (t_1 x^a y^b, t_2 x^c y^d) \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{Z}), (t_1, t_2) \in \mathbb{T} \right\} \cong \mathbb{T} \rtimes \text{GL}_2(\mathbb{Z}).$$

Indeed,  $\text{Aut } X$  surjects onto the automorphism group of the multiplicative group

$$\mathcal{O}_X^\times(X) = \{t x^a y^b \mid t \in \mathbb{K}^*, a, b \in \mathbb{Z}\}$$

of the ring  $\mathcal{O}_X(X)$ .

**3.7** (*The group  $\text{Aut}((\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta)$* ). The surface  $X' = (\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta$  is isomorphic to the smooth quadric in  $\mathbb{A}^3$  with equation  $xy - z^2 = -1$ . The group  $\text{Aut } X'$  was studied in [23, 10.1], [69, §2.1, Thm. 4], and [72]. The result in [69] can be interpreted as follows. There is an amalgam

$$\text{Aut } X' \cong A' *_C B' \quad \text{with} \quad C' = A' \cap B' \cong \mathbb{G}_m \times \langle \tau \rangle,$$

where  $\tau : (u, v) \mapsto (v, u)$  is the involution interchanging the factors of  $\mathbb{P}^1 \times \mathbb{P}^1$ ,

$$(4) \quad A' = \text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1, \Delta) \cong \text{PSL}(2, \mathbb{K}) \times \langle \tau \rangle,$$

and

$$(5) \quad B' = (U'_\infty \rtimes \mathbb{G}_m) \rtimes \langle \tau \rangle \quad \text{with} \quad U'_\infty = R_u(B') \cong \mathbb{K}[t']$$

being the unipotent radical of  $B'$ . In particular,  $A'$  is semisimple and  $B'$  is solvable of rank 1. In the affine coordinates  $(u, v)$  in  $\mathbb{P}^1 \times \mathbb{P}^1$ , where  $u = u_0/u_1$  and  $v = v_0/v_1$ , we have

$$U'_\infty \rtimes \mathbb{G}_m = \left\{ (u, v) \mapsto (cu + P, cv + P) \mid c \in \mathbb{G}_m, P \in \mathbb{K} \left[ \frac{1}{u-v} \right] \right\}.$$

**3.8** (*The group  $\text{Aut}(\mathbb{P}^2 \setminus C)$* ). For  $X = \mathbb{P}^2 \setminus C$  the group  $\text{Aut } X$  was studied in [23, §2]. By *loc.cit.*, there is an amalgam  $\text{Aut } X \cong A *_C B$  with

$$A = \text{Aut}(\mathbb{P}^2, C) \cong \text{SO}(3, \mathbb{K}) \cong \text{PSL}(2, \mathbb{K}) \quad \text{and} \quad B = U_\infty \rtimes \mathbb{G}_m,$$

where  $U_\infty \cong \mathbb{K}[t]$  and  $C = A \cap B \cong \text{Aff}(\mathbb{A}^1)$ .

**Remark 3.9.** The amalgam in 3.8 is pushforward of that in 3.7 via the construction of Remark 2.17. Indeed, the surface  $X' = (\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta$  is the Galois covering of  $X = \mathbb{P}^2 \setminus C$ , and more precisely,  $X = X'/\mathbb{Z}_2$ , where  $\mathbb{Z}_2 = \langle \tau \rangle$ . In the affine coordinates  $(u, v)$ , the quotient morphism  $\mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^2$  is given by the elementary symmetric polynomials in two variables via the classical Vieta formulas; cf. [33, Ex. 5.1]. The Galois  $\mathbb{Z}_2$ -covering  $X' \rightarrow X$  being the universal covering, we have

$\text{Aut } X \cong \text{Norm}_{\text{Aut } X'}(\tau)/\langle \tau \rangle$ . A comparison of the explicit formulas in [23, (2.4.3;1)] and [69, §2] yields the isomorphism

$$\text{Norm}_{\text{Aut } X'}(\tau) \cong \text{Norm}_{A'}(\tau) *_{\text{Norm}_{C'}(\tau)} \text{Norm}_{B'}(\tau),$$

where

$$\text{Norm}_{A'}(\tau) = A' \cong \text{PSL}(2, \mathbb{K}) \times \langle \tau \rangle \quad \text{and} \quad \text{Norm}_{B'}(\tau) = B_0 \times \langle \tau \rangle \subset B',$$

and where

$$B_0 = \left\{ (u, v) \mapsto (cu + P, cv + P) \mid c \in \mathbb{K}^\times, P \in \mathbb{K} \left[ \frac{1}{(u-v)^2} \right] \right\} = U_\infty \times \mathbb{G}_m$$

and  $U_\infty = \mathbb{K}[t^2] \subset U'_\infty = \mathbb{K}[t']$  with  $t' = \frac{1}{u-v}$ . Finally,

$$A \cong \text{Norm}_{A'}(\tau)/\langle \tau \rangle = A'/\langle \tau \rangle \cong \text{PSL}(2, \mathbb{K})$$

and

$$B \cong \text{Norm}_{B'}(\tau)/\langle \tau \rangle = B_0 \cong U_\infty \times \mathbb{G}_m.$$

Summarizing the results in 3.3-3.8 we arrive at the following conclusion.

**Theorem 3.10.** *For any surface  $X$  in (2) the neutral component  $\text{Aut}^\circ X \cong A^\circ *_{C^\circ} B^\circ$  is finitely bearable, where  $A^\circ$ ,  $B^\circ$ , and  $C^\circ = A^\circ \cap B^\circ$  are connected, and either  $A^\circ$  and  $B^\circ$  are both solvable nested ind-groups, or  $A^\circ$  is such a group and  $B^\circ$  is a reductive affine algebraic group.*

**Corollary 3.11.** <sup>9</sup> *Let  $X$  in (2) be one of the surfaces  $\mathbb{A}^1 \times \mathbb{A}_*^1$ ,  $(\mathbb{A}_*^1)^2$ , or  $V_{d,e}$  with  $e > 1$ . Then any connected algebraic group  $G$  acting effectively on  $X$  is solvable.*

*Proof.* Indeed, by Proposition 2.16,  $G$  is conjugated to a subgroup of one of the factors  $A^\circ$  and  $B^\circ$ , which are both solvable in these cases.  $\square$

#### 4. ALGEBRAIC GROUP ACTIONS ON AFFINE SURFACES

**4.1. Generalities.** In this section we recall some general facts about algebraic group actions on affine varieties and their specialization to the case of affine surfaces. By a  $G$ -variety we mean a variety with an effective (regular) action of an (algebraic) group  $G$ . The next proposition is well known (see, e.g., [86, 87] and [33, Lem. 2.7 and 2.9]); for the reader's convenience we provide either a short argument, or a reference.

**Proposition 4.1.** *Let  $G$  be a connected affine algebraic group, and let  $X$  be a normal affine  $G$ -variety. Then the following hold.*

- (a) *We have  $\text{rk } G \leq \dim X$ , and  $\text{rk } G = \dim X$  if and only if  $X$  is toric.*
- (b) *If  $G$  is solvable and acts transitively on  $X$ , then  $X \cong \mathbb{A}^k \times (\mathbb{A}_*^1)^l$  for some  $k, l \geq 0$ .*
- (c) *If  $G$  is solvable and acts on  $X$  with an open orbit  $O$ , then either  $O = X$ , or  $X \setminus O$  is a divisor.*
- (d) *If  $G$  is unipotent and has an open orbit in  $X$ , then  $X \cong \mathbb{A}^n$ .*
- (e) *If  $G$  is reductive and acts with an open orbit, then it has a unique closed orbit, and this orbit lies in the closure of any other orbit.*
- (f) *If  $G$  is semisimple, then  $G$  has no one-dimensional orbit in  $X$ .*

*Proof.* (a) Let  $T \subset G$  be a maximal torus. By the rigidity of algebraic subtori, algebraic subgroups of  $T$  form a countable set. Since  $T$  acts effectively on  $X$ , the isotropy subgroup of  $T$  in a generic point of  $X$  is trivial due to the aforementioned rigidity. Hence  $\text{rk } G = \dim T \leq \dim X$ . The second assertion in (a) follows by definition of a toric variety.

<sup>9</sup>Cf. Proposition 4.7.

For (b) see [87, Thm. 2].

(c) follows from (b) since the open orbit is affine in this case.

(d) Let  $O$  be the open orbit of  $U$ . Since any orbit of a unipotent group acting on an affine variety is closed,  $O = X$ . Now the result follows from the corollary of Theorem 2 in *loc.cit.*

(e) is proven in [86, Prop. 2].

(f) Since  $G$  is semisimple, it admits no non-trivial homomorphism to  $\text{Aff}(\mathbb{A}^1)$ . Indeed, otherwise, it would act non-trivially on  $\mathbb{P}^1$  with a fixed point. Such an action can be lifted to a non-trivial linear representation of the universal covering group  $\tilde{G}$  in  $\text{GL}(2, \mathbb{K})$  with a trivial one-dimensional subrepresentation, which is impossible.

It follows that  $G$  cannot act non-trivially on a curve. Now (f) follows.  $\square$

**Corollary 4.2.** *Let  $X$  be a normal affine  $G$ -surface. Then the following hold.*

(g) *If  $G$  is nonabelian and unipotent, then  $X \cong \mathbb{A}^2$ .*

(h) *If  $G$  semisimple, then it has an open orbit in  $X$  with a finite complement.*

*Proof.* (g) If  $G$  acts with an open orbit on  $X$ , then the result follows from Proposition 4.1(d). Otherwise,  $G$  acts with one-dimensional general orbits, and so, any one-parameter subgroup  $H \subset G$  has the same algebra of invariants:  $\mathcal{O}(X)^H = \mathcal{O}(X)^G$ . This algebra is affine and its spectrum  $Z$  is a smooth affine curve (see [30, Lem. 1.1]). Any fiber of the induced  $\mathbb{A}^1$ -fibration  $\pi: X \rightarrow Z$  is stable under the  $G$ -action.

If  $H'$  is another one-parameter subgroup of  $G$ , then the actions of  $H$  and  $H'$  commute. Hence these subgroups commute, and so,  $G$  is Abelian, contrary to our assumption.

Finally, (h) is immediate from (f).  $\square$

**Remark 4.3.** The affine plane is exceptional with respect to the property in (g). Indeed, the Heisenberg group

$$H = \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}, \quad a, b, c \in \mathbb{K} \right\}$$

is a nonabelian unipotent group acting effectively on  $\mathbb{A}^2$  via  $(x, y) \mapsto (x+ay+b, y+c)$ .

Let us also mention the following result.

**Proposition 4.4.** *If the neutral component  $\text{Aut}^\circ X$  of an affine algebraic variety  $X$  with  $\dim X \geq 2$  is an algebraic group, then  $\text{Aut}^\circ X \cong \mathbb{G}_m^k$  is an algebraic torus. This is the case, in particular, for the surfaces of class  $(\text{ML}_2)$ .*

*Proof.* The first assertion follows easily by a lemma of Iitaka [53, Lem. 3], see, e.g., [65, Thm. 1.3] and [71, Thm. 4.10(a)]. The second will be proven in a forthcoming paper ([84]).  $\square$

## 4.2. Quasihomogeneous affine surfaces.

**Theorem 4.5** (Gizatullin–Popov, [48], [87]). *A normal affine surface  $X$  admitting an action of an algebraic group with an open orbit whose complement is finite, is one of the surfaces*

$$(6) \quad \mathbb{A}^2, \quad \mathbb{A}^1 \times \mathbb{A}_*^1, \quad (\mathbb{A}_*^1)^2, \quad V_d, \quad d \geq 2, \quad \mathbb{P}^2 \setminus C, \quad \text{and} \quad (\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta,$$

where  $V_d = V_{d,1}$  is a Veronese cone (see 3.1),  $C$  is a smooth conic in  $\mathbb{P}^2$ , and  $\Delta$  is the diagonal in  $\mathbb{P}^1 \times \mathbb{P}^1$ .

Using Proposition 4.1(f) we deduce the following corollary.

**Corollary 4.6.** *Let  $X$  be a normal affine surface. Then the following conditions are equivalent:*

- (i)  $X$  admits a nontrivial action of a connected semisimple group;
- (ii)  $X$  is spherical, that is, it admits a semisimple group action, such that a Borel subgroup has an open orbit;
- (iii)  $X$  is one of the surfaces

$$(7) \quad \mathbb{A}^2, \quad V_d, \quad d \geq 2, \quad \mathbb{P}^2 \setminus C, \quad (\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta.$$

The following proposition is a version of Proposition 4.14 in [33]. We provide a new proof.

**Proposition 4.7.** *Let  $X$  be a normal affine  $G$ -surface, where  $G \subset \text{Aut } X$  is a connected reductive algebraic group different from a torus. Then the pair  $(X, G)$  is one of the following:*

- $(\mathbb{A}^2, \text{GL}(2, \mathbb{K}))$  and  $(\mathbb{A}^2, \text{SL}(2, \mathbb{K}))$ ;
- $(V_d, \text{GL}(2, \mathbb{K})/\mu_d)$ ,  $d \geq 2$ ,  $(V_d, \text{SL}(2, \mathbb{K}))$ ,  $d \geq 3$  odd, and  $(V_d, \text{PSL}(2, \mathbb{K}))$ ,  $d \geq 2$  even;
- $(\mathbb{P}^2 \setminus C, \text{PSL}(2, \mathbb{K}))$  and  $((\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta, \text{PSL}(2, \mathbb{K}))$ .

Furthermore, the action of  $G$  on  $X$  is unique up to a conjugation in the group  $\text{Aut } X$ .

*Proof.* Under our assumptions  $G$  contains a nontrivial semisimple subgroup. Hence  $G$  acts with an open orbit, which has a finite complement in  $X$ , see Corollary 4.2(h). By Gizatullin-Popov Theorem 4.5  $X$  is one of the list (7). Due to the results in Section 3, the group  $\text{Aut } X$  admits a structure of an amalgam of two subgroups. By Proposition 2.16,  $G$  is conjugated to a subgroup of the non-solvable factor in the amalgam decomposition of  $\text{Aut } X$ .

If  $X$  is one of the surfaces  $\mathbb{P}^2 \setminus C$  and  $(\mathbb{P}^1 \times \mathbb{P}^1) \setminus \Delta$ , then by 3.7 and 3.8,  $G = \text{PSL}(2, \mathbb{K})$ , and the  $G$ -action on  $X$  is unique up to conjugation; see also [86], [23], and [33, Prop. 4.14] for alternative proofs.

Similarly, for  $X = \mathbb{A}^2$  the assertion follows from 3.3. If  $X = V_d$ ,  $d \geq 2$ , then by 3.4, case (i),  $G$  is conjugated in  $\text{Aut } X$  to a subgroup of the non-solvable factor  $\text{GL}(2, \mathbb{K})/\mu_{d,1}$ , where  $\mu_{d,1} \cong \mathbb{Z}/d\mathbb{Z}$  is contained in the center of  $\text{GL}(2, \mathbb{K})$ . Hence  $G$  is conjugated either to  $\text{GL}(2, \mathbb{K})/\mu_{d,1}$  itself, or to  $\text{SL}(2, \mathbb{K})$ , or to  $\text{PSL}(2, \mathbb{K})$  canonically embedded in  $\text{GL}(2, \mathbb{K})/\mu_{d,1}$ , depending on the parity of  $d$ .  $\square$

**4.3. Actions with an open orbit.** Recall that an effective  $\mathbb{G}_m$ -action on a normal affine variety defines a grading  $A = \bigoplus_{j \in \mathbb{Z}} A_j$  on the algebra  $A = \mathcal{O}(X)$ . For  $\dim X = 2$  the  $\mathbb{G}_m$ -action is called *elliptic* if  $A_j = 0 \ \forall j < 0$  and  $A_0 = \mathbb{K}$ , *parabolic* if  $A_j = 0 \ \forall j < 0$  and  $A_0 \neq \mathbb{K}$ , and *hyperbolic* if  $A_{-1} \neq 0 \neq A_1$ .

The following result is essentially Proposition 2.10 in [33]; cf. also [8, Prop. 2.5]<sup>10</sup>. For the reader's convenience, we sketch a proof.

**Proposition 4.8.** *Let  $X$  be a normal affine surface different from the surfaces in (2). Then the following are equivalent:*

- (i)  $X$  admits an effective action of a connected affine algebraic group  $G$  with an open orbit;
- (ii)  $X$  admits an action of a semi-direct product  $\mathbb{G}_a \rtimes \mathbb{G}_m$  with an open orbit;
- (iii)  $X$  admits effective  $\mathbb{G}_a$ - and  $\mathbb{G}_m$ -actions such that  $\mathcal{O}(X)^{\mathbb{G}_m} \neq \mathcal{O}(X)^{\mathbb{G}_a}$ .

Moreover, any  $\mathbb{G}_m$ -action on  $X$  as in (iii) is hyperbolic,  $X$  is of class  $(\text{ML}_0 \cup \text{ML}_1, 1)$ , and  $X$  is a cyclic quotient of a normalization of a Danielewski surface  $\{xy - P(z) = 0\}$  in  $\mathbb{A}^3$  for some  $P \in \mathbb{K}[z]$ .

<sup>10</sup>In [8, Prop. 2.5] the condition  $\mathcal{O}(X)^{\mathbb{G}_m} \neq \mathcal{O}(X)^{\mathbb{G}_a}$  is lacking; furthermore, the proof in [8] assumes implicitly smoothness of  $X$ .



*Proof.* (i) $\Leftrightarrow$ (ii). As follows from Proposition 4.7 and Theorem 4.5, under our assumption the group  $G$  as in (i) does not contain any semisimple subgroup, and so, is solvable. Since by assumption  $X$  is non-toric,  $\text{rk } G \leq 1$ . In fact,  $\text{rk } G = 1$ . Indeed, otherwise  $G$  is unipotent and acts with an open orbit. Since the orbits of a unipotent group acting on an affine variety are closed,  $G$  is transitive in  $X$ . By Proposition 4.1(b),  $X \cong \mathbb{A}^2$ , which is excluded by our assumption.

The open orbit  $O$  of  $G$  in  $X$  coincides with an open orbit (isomorphic to one of  $\mathbb{A}^2$ ,  $\mathbb{A}^1 \times \mathbb{A}_*^1$ ,  $(\mathbb{A}_*^1)^2$ ) of a two-dimensional subgroup  $H \cong \mathbb{G}_a \rtimes \mathbb{G}_m$  of  $G$  ([33, Lem. 2.9(b)]). This gives (i) $\Rightarrow$ (ii). The implication (ii) $\Rightarrow$ (i) is immediate, hence we have (i) $\Leftrightarrow$ (ii).

(ii) $\Rightarrow$ (iii). By (ii)  $X$  is a  $\mathbb{G}_m$ -surface admitting a *horizontal*  $\mathbb{G}_a$ -action. The latter means that the general  $\mathbb{G}_a$ -orbits are not the closures of general  $\mathbb{G}_m$ -orbits, which implies (iii).

(iii) $\Rightarrow$ (ii). We claim that any affine variety with effective  $\mathbb{G}_a$ - and  $\mathbb{G}_m$ -actions as in (iii) possesses an effective action of a semi-direct product  $\mathbb{G}_a \rtimes \mathbb{G}_m$ . Indeed, let  $A = \bigoplus_{i \in \mathbb{Z}} A_i$  be the grading of the algebra  $A = \mathcal{O}(X)$  induced by the  $\mathbb{G}_m$ -action, and let  $\partial \in \text{Der } A$  be the locally nilpotent derivation corresponding to the  $\mathbb{G}_a$ -action. Write  $\partial = \sum_{i=k}^l \partial_i$ , where  $k \leq l$  and  $\partial_i \in \text{Der } A$  is a homogeneous derivation of degree  $i$  with  $\partial_k \neq 0 \neq \partial_l$ . Then  $\partial_k, \partial_l$  are again locally nilpotent ([33, Lem. 2.1], [89]). Then the  $\mathbb{G}_a$ -actions on  $X$  generated by  $\partial_k$  and  $\partial_l$  are normalized by the  $\mathbb{G}_m$ -action. This yields the existence of an  $(\mathbb{G}_a \rtimes \mathbb{G}_m)$ -action on  $X$ ; see [33, Lem. 2.2]. Notice that, if  $k = l$ , then  $\partial = \partial_k = \partial_l$ , and the induced  $\mathbb{G}_a$ -action is horizontal due to our assumption that  $\mathcal{O}(X)^{\mathbb{G}_m} \neq \mathcal{O}(X)^{\mathbb{G}_a}$ . Otherwise, at least one of the indices  $k$  and  $l$  is different from  $-1$ , and again the induced  $\mathbb{G}_a$ -action is horizontal, since otherwise the degree of the corresponding locally nilpotent derivation equals  $-1$ , see [33, Thm. 3.12]. In any case, the associate  $(\mathbb{G}_a \rtimes \mathbb{G}_m)$ -action on  $X$  has an open orbit, as required in (ii).

Finally we have the equivalences (i) $\Leftrightarrow$ (ii) $\Leftrightarrow$ (iii).

To show the last assertions, note that under condition (iii) the horizontal  $\mathbb{G}_a$ -action on  $X$  is normalized by the given  $\mathbb{G}_m$ -action. If the latter action were elliptic or parabolic, then  $X$  would be a toric surface  $\mathbb{A}^2$  or  $V_{d,e}$ , contrary to our assumption, see [33, Thms. 3.3 and 3.16]. Hence the  $\mathbb{G}_m$ -action on  $X$  is hyperbolic, as claimed.

By exclusion  $X$  belongs to one of the classes  $(\text{ML}_i, 1)$ ,  $i = 0, 1$ . Furthermore, due to [33, Cor. 3.27 and 3.30], any hyperbolic  $\mathbb{G}_m$ -surface  $X \in (\text{ML}_0) \cup (\text{ML}_1)$  is a cyclic quotient of the normalization of some Danielewski surface  $x^n y - P(z) = 0$  in  $\mathbb{A}^3$ , where  $P \in \mathbb{K}[z]$  and  $n \geq 1$ .  $\square$

**Remark 4.9.** If a surface  $X \in (\text{ML}_0)$  is a complete intersection, then it can be realized as a hypersurface  $xy - P(z) = 0$  in  $\mathbb{A}^3$ , where  $P \in \mathbb{K}[z]$  is nonconstant, see [4], [20], [21]. In particular,  $X$  is a hyperbolic  $\mathbb{G}_m$ -surface.

**4.4.  $\mathbb{G}_m$ -surfaces: Dolgachev-Pinkham-Demazure presentation.** The  $\mathbb{G}_m$ -surfaces can be described in terms of their *Dolgachev-Pinkham-Demazure presentation*, or *DPD presentation*, for short. Let us recall this description, see [32].

**Definition 4.10.** In the elliptic and the parabolic cases, the DPD construction associates to any pair  $(C, D)$ , where  $C$  is a smooth curve and  $D$  is an ample  $\mathbb{Q}$ -divisor on  $C$ , the graded  $\mathbb{K}$ -algebra

$$A = \bigoplus_{j \geq 0} A_j, \quad \text{where } A_j = H^0(C, \mathcal{O}_C([jD])).$$

The induced effective  $\mathbb{G}_m$ -action on the normal affine surface  $X = \text{Spec } A$  is elliptic if  $C$  is projective and parabolic otherwise. In the hyperbolic case, the DPD construction associates to any triple  $(C, D_+, D_-)$ , where  $C$  is a smooth affine curve and

$D_{\pm}$  are  $\mathbb{Q}$ -divisors on  $C$  with  $D_+ + D_- \leq 0$ , the graded  $\mathbb{K}$ -algebra

$$A = \bigoplus_{j \in \mathbb{Z}} A_j, \quad \text{where } A_{\pm j} = H^0(C, \mathcal{O}_C([jD_{\pm}])) \text{ for } j \geq 0.$$

The resulting effective  $\mathbb{G}_m$ -action on  $X = \text{Spec } A$  is hyperbolic. In fact, any normal affine  $\mathbb{G}_m$ -surface  $X$  arises in this way, and the corresponding DPD presentation is unique up to isomorphisms of pairs  $(C, D)$  and of triples  $(C, D_+, D_-)$  and up to replacing  $D$  (the pair  $(D_+, D_-)$ , respectively) by a linearly equivalent divisor  $\hat{D}$  (by a pair  $(D_+ + D', D_- - D')$ , where  $D'$  is a principal divisor on  $C$ , respectively), see [32, Thms. 2.2, 3.2, and 4.3].

The classification of the normal affine  $\mathbb{G}_m$ -surfaces according to the Makar-Limanov complexity is as follows, see [33] and [70, Cor. 3.30]. We let  $\{D\}$  be the fractional part of a  $\mathbb{Q}$ -divisor  $D$ .

**Proposition 4.11.** *Let  $X$  be a normal affine  $\mathbb{G}_m$ -surface with an associate DPD presentation  $(C, D)$  for an elliptic or a parabolic  $\mathbb{G}_m$ -action, and  $(C, D_+, D_-)$  for a hyperbolic one. Then*

- $X \in (\text{ML}_0)$  if and only if one of the following holds:
  - $X$  is elliptic,  $C = \mathbb{P}^1$ , and  $\{D\}$  is supported in at most two points;
  - $X$  is parabolic,  $C = \mathbb{A}^1$ , and  $\{D\}$  is supported in at most one point;
  - $X$  is hyperbolic,  $C = \mathbb{A}^1$ , and  $\{D_{\pm}\}$  is supported in at most one point  $p_{\pm}$ .
- $X \in (\text{ML}_1)$  if and only if one of the following holds:
  - $X$  is parabolic, and either  $C$  is non-rational, or  $\{D\}$  is supported in at least two points;
  - $X$  is hyperbolic,  $C = \mathbb{A}^1$ , and exactly one of the  $\mathbb{Q}$ -divisors  $\{D_+\}$ ,  $\{D_-\}$  is supported in at most one point.
- $X \in (\text{ML}_2)$  otherwise.

**Remarks 4.12.** 1. The elliptic and the parabolic  $\mathbb{G}_m$ -surfaces of class  $(\text{ML}_0)$  are exactly the nondegenerate toric surfaces<sup>11</sup>  $\mathbb{A}^2$  and  $V_{d,e}$ , see [33, Thms. 3.3 and 3.16]. A hyperbolic  $\mathbb{G}_m$ -surface  $X$  of class  $(\text{ML}_0)$  is a nondegenerate toric surface if and only if  $\text{supp}\{D_+\} = \text{supp}\{D_-\} = \{p_0\}$  for some point  $p_0 \in C = \mathbb{A}^1$ , and  $(D_+, D_-) = (D_0 + \{D_+\}, -D_0 + \{D_-\})$  for some integral divisor  $D_0$  on  $\mathbb{A}^1$  ([37, Lem. 4.2(b)]).

2. There exist smooth surfaces of class  $(\text{ML}_0, 0)$ , that is, smooth Gizatullin surfaces which do not admit any nontrivial  $\mathbb{G}_m$ -action, see [37, Cor. 4.9]. The subgroup  $\text{SAut } X \subset \text{Aut } X$  of such a surface acts on  $X$  with an open orbit, while there is no algebraic group action on  $X$  with an open orbit.

3. If  $X \in (\text{ML}_1)$  is a parabolic  $\mathbb{G}_m$ -surface, then the  $\mathbb{G}_a$ -action on  $X$  is *vertical* (or *fiberwise*), that is,  $\mathcal{O}(X)^{\mathbb{G}_m} = \mathcal{O}(X)^{\mathbb{G}_a}$ . This follows from [33, Thm. 3.16], cf. Remark 1 above.

4. If  $X \in (\text{ML}_2)$  is a  $\mathbb{G}_m$ -surface, then  $\text{Aut}^\circ X \cong \mathbb{G}_m$  or  $\mathbb{G}_m^2$ , see Proposition 4.4.

In terms of the DPD presentation, the criterion of Proposition 4.8 becomes more concrete.

**Corollary 4.13.** *Let  $X$  be a normal affine surface. The group  $\text{Aut } X$  acts on  $X$  with an open orbit if and only if either  $X$  is one of the surfaces in (2), or  $X$  is of class  $(\text{ML}_0)$ ,<sup>12</sup> or  $X$  is a hyperbolic  $\mathbb{G}_m$ -surface of class  $(\text{ML}_1)$ .*

*Proof.* Suppose that  $X$  does not appear in (2). By Proposition 4.8 the group  $\text{Aut } X$  acts on  $X$  with an open orbit if and only if a semi-direct product  $\mathbb{G}_a \rtimes \mathbb{G}_m$  does, and

<sup>11</sup>Recall that a toric affine variety  $X$  is *nondegenerate* if any invertible function on  $X$  is constant.

<sup>12</sup>That is,  $X$  is a Gizatullin surface.

so,  $X$  is a  $\mathbb{G}_m$ -surface of class  $(\text{ML}_0) \cup (\text{ML}_1)$ . It remains to note that the  $(\text{Aut } X)$ -action on a parabolic  $\mathbb{G}_m$ -surface of class  $(\text{ML}_1)$  has one-dimensional orbits, see Remark 4.12.2.  $\square$

## 5. AUTOMORPHISM GROUPS OF GIZATULLIN SURFACES

**5.1. Definition, characterizations, examples.** We adopt the following definition.

**Definition 5.1.** A *Gizatullin surface* is a normal affine surface  $X$  non-isomorphic to  $\mathbb{A}^1 \times \mathbb{A}_*^1$  that can be completed by a chain of smooth rational curves (a *zigzag*) into an SNC-pair  $(\bar{X}, D)$ .

The following characterization goes back to Gizatullin [49]; see also [8, Thm. 1.8], [26].

**Theorem 5.2.** *Given a normal affine surface  $X$ , the following are equivalent:*

- $X$  is a Gizatullin surface;
- $X$  is of class  $(\text{ML}_0)$ ;
- $X$  admits two distinct  $\mathbb{A}^1$ -fibrations  $X \rightarrow \mathbb{A}^1$ ;
- the group  $\text{SAut } X$  acts on  $X$  with an open orbit.

The affine plane  $\mathbb{A}^2$  and the toric affine surfaces  $\mathbb{A}^2/\mu_{d,e}$  are examples of Gizatullin surfaces. Another important examples are the *Danilov-Gizatullin surfaces* and the *special Gizatullin surfaces*. Let us consider these classes along with their DPD presentations.

**Example 5.3** (*Danilov-Gizatullin surfaces*). Such a surface is the complement  $X = \mathbb{F}_n \setminus S$  to an ample section  $S$  in a Hirzebruch surface  $\mathbb{F}_n = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(n)) \rightarrow \mathbb{P}^1$ . A section  $S$  is ample if and only if  $d := S^2 > n$ . Two Danilov-Gizatullin surfaces are isomorphic if and only if they share the same invariant  $d = S^2$ , see [23, Thm. 5.8.1] (see also [14, Cor. 4.8], [40]). We let  $X_d$  denote the Danilov-Gizatullin surface with invariant  $d = S^2$ . The surface  $X_d$  possesses exactly  $d - 1$  pairwise non-conjugated  $\mathbb{G}_m$ -actions with the DPD presentations

$$(C, D_+, D_-) = \left( \mathbb{A}^1, -\frac{1}{r}[p_0], -\frac{1}{d-r}[p_1] \right), \quad r = 1, \dots, d-1,$$

where  $p_0, p_1 \in \mathbb{A}^1$ ,  $p_0 \neq p_1$ . see [33, 5.3] and [37, Prop. 5.15]. The automorphism group of a Danilov-Gizatullin surface  $X_d$  with  $2 \leq d \leq 5$  is an amalgam, see [23, §§5-8].

**Example 5.4** (*Special Gizatullin surfaces*). A smooth Gizatullin surface  $X$  equipped with a hyperbolic  $\mathbb{G}_m$ -action is called *special* if the associate DPD presentation is

$$(C, D_+, D_-) = \left( \mathbb{A}^1, -\frac{1}{r}[p_+], -\frac{1}{d-r}[p_-] - D_0 \right)$$

with  $d \geq 3$ ,  $1 \leq r \leq d-1$ , under the convention that  $D_+ = 0$  if  $r = 1$  and  $D_- = 0$  if  $r = d-1$ , and otherwise  $p_+ \neq p_-$ , and with a reduced divisor  $D_0 = \sum_{i=1}^s [p_i]$  on  $\mathbb{A}^1$ , where  $s > 0$  and  $p_i \neq p_{\pm} \forall i$ .

**5.2. One-parameter subgroups and bearability on Gizatullin surfaces.** For Gizatullin surfaces the following theorem is proven in [39, Thms. 1.0.1, 1.0.5, and Ex. 6.3.21], see also [37, §5.3] and [38, Cor. 5.15].

**Theorem 5.5.** *Smooth affine surfaces  $X$  admitting an effective  $\mathbb{G}_m$ -action can be divided into the following 4 classes:*

- 1) toric affine surfaces;
- 2) Danilov-Gizatullin surfaces  $X_d$ ,  $d \geq 4$ ;

- 3) special Gizatullin surfaces;
- 4) all the others,

so that the set of conjugacy classes of 1-tori in  $\text{Aut } X$

- is infinite countable in case 1)<sup>13</sup>;
- is finite of cardinality  $\lfloor d/2 \rfloor$  for  $X_d$  in case 2);
- forms a 1- or 2-parameter family in case 3);
- is finite of cardinality at most 2 in case 4).

Furthermore, the set of equivalence classes of  $\mathbb{A}^1$ -fibrations  $X \rightarrow \mathbb{A}^1$

- is finite of cardinality at most 2 in cases 1) and 4);
- forms an  $m$ -parameter family in case 2), where  $m \geq 1$  if  $d \geq 7$ , and  $m = m(d) \rightarrow +\infty$  as  $d \rightarrow +\infty$ ;
- forms an  $m$ -parameter family in case 3), where  $m \geq 1$ .

*Proof.* For surfaces of class  $\text{ML}_0$ , that is, for Gizatullin surfaces, the assertions follow due to the references preceding the theorem. Thus we need to consider just the  $\text{ML}_1$ - and  $\text{ML}_2$ -surfaces. By definition, such a surface  $X$  admits at most one  $\mathbb{A}^1$ -fibration over an affine curve. By Corollary 6.11 the group  $\text{Aut}^\circ X$  is nested, that is, 1-bearable, and has at most one conjugacy class of maximal tori. Hence the assertions follow also in this case.  $\square$

**Corollary 5.6.** *If  $X$  is either a special Gizatullin surface, or a Danilov-Gizatullin surface  $X_d$  with  $d \geq 7$ , then the group  $\text{Aut}^\circ X$  is not countably bearable. Furthermore,  $\text{Aut } X_d$  ( $\text{Aut}^\circ X_d$ , respectively) is not a nested ind-group for  $d = 4, 5$ , and cannot be a nontrivial amalgam of two nested ind-groups for  $d = 6$ .*

*Proof.* By Theorem 5.5, for a special Gizatullin surface  $X$  (for a Danilov-Gizatullin surface  $X_d$  with  $d \geq 7$ , respectively) the set of conjugacy classes of  $\mathbb{G}_m$ -subgroups (of  $\mathbb{A}^1$ -fibrations over  $\mathbb{A}^1$ , respectively) in  $\text{Aut}^\circ X$  is uncountable. In these cases the assertion follows from Propositions 2.23 and 2.26, respectively. For  $d \leq 6$ , the groups  $\text{Aut } X_d$  and  $\text{Aut}^\circ X_d$  have both  $\lfloor d/2 \rfloor$  conjugacy classes of  $\mathbb{G}_m$ -subgroups, that is, of maximal tori. Indeed,  $\text{rk } \text{Aut } X_d = 1$ , since this surface is not toric. Hence for  $d \leq 6$  the assertion follows from Remark 2.24.  $\square$

**Remarks 5.7.** 1. The presentation of the group  $\text{Aut } X_d$  as an amalgam in [23, §§6-8] involves 2 factors if  $d = 3$  and 3 factors if  $d = 4, 5$ . It seems that for  $d \geq 6$ , no explicit amalgamated structure on  $\text{Aut } X_d$  is known. However, the authors of [23] mention that their methods allow in principle to compute such a structure for any  $d$ .

2. See [10] and [63] for spectacular examples of Gizatullin surfaces  $X$  such that, if  $N(X) \subset \text{Aut } X$  is the normal subgroup generated by all algebraic subgroups of  $\text{Aut } X$ , then the quotient  $(\text{Aut } X)/N(X)$  contains a free group on an uncountable set of generators.

**5.3. Standard completions and extended divisors of Gizatullin surfaces.** These combinatorial invariants are indispensable in studies on Gizatullin surfaces.

**Notation 5.8.** Let  $X$  be a Gizatullin surface, and let  $(\bar{X}, D)$  be an SNC completion of the minimal resolution of singularities of  $X$  by a zigzag  $D$ , where

$$D = C_0 + \cdots + C_n \quad \text{with} \quad C_i \cdot C_j = 1 \quad \text{if} \quad |i - j| = 1 \quad \text{and} \quad C_i \cdot C_j = 0 \quad \text{otherwise.}$$

The boundary components  $C_i$ ,  $0 \leq i \leq n$ , serve as the vertices  $v_i$  of the dual linear graph  $\Gamma_D$  of  $D$ . Each vertex  $v_i$  is weighted by the corresponding self-intersection

<sup>13</sup>The latter holds for any toric affine surface  $X$ , not necessarily smooth.

number  $w_i = C_i^2$ . Thus  $\Gamma_D$  is of the form

$$\Gamma_D : \begin{array}{ccccccc} & v_0 & v_1 & & \cdots & & v_n \\ & \circ & \circ & \text{---} & \cdots & \text{---} & \circ \\ & w_0 & w_1 & & & & w_n \end{array} .$$

The string of weights  $[[w_0, w_1, \dots, w_n]]$  can be putted into a standard form by means of elementary transformations of weighted graphs.

**Definition 5.9.** Given an at most linear vertex  $v$  of a weighted graph  $\Gamma$  with weight 0 one can perform the following transformations. If  $v$  is linear with neighbors  $v_1, v_2$  then we blow up the edge connecting  $v$  and  $v_1$  in  $\Gamma$  and blow down the proper transform of  $v$ :

$$(8) \quad \begin{array}{ccccccc} & v_1 & v' & v_2 & & & \\ & \circ & \circ & \circ & \cdots & & \\ & w_1 - 1 & 0 & w_2 + 1 & & & \end{array} \rightarrow \begin{array}{ccccccc} & v_1 & v' & v & v_2 & & \\ & \circ & \circ & \circ & \circ & \cdots & \\ & w_1 - 1 & -1 & -1 & w_2 & & \end{array} \rightarrow \begin{array}{ccccccc} & v_1 & v & v_2 & & & \\ & \circ & \circ & \circ & \cdots & & \\ & w_1 & 0 & w_2 & & & \end{array} .$$

Similarly, if  $v$  is an end vertex of  $\Gamma$  connected to the vertex  $v_1$  then one proceeds as follows:

$$(9) \quad \begin{array}{ccccccc} & v_1 & v' & & & & \\ & \circ & \circ & & & & \\ & w_1 - 1 & 0 & & & & \end{array} \rightarrow \begin{array}{ccccccc} & v_1 & v' & v & & & \\ & \circ & \circ & \circ & & & \\ & w_1 - 1 & -1 & -1 & & & \end{array} \rightarrow \begin{array}{ccccccc} & v_1 & v & & & & \\ & \circ & \circ & & & & \\ & w_1 & 0 & & & & \end{array} .$$

These operations (8) and (9) and their inverses are called *elementary transformations* of  $\Gamma$ . If such an elementary transformation involves only an inner blowup then we call it *inner*. Thus (8) and (9) are inner whereas the inverse of (9) is not as it involves an outer blowup.

Consider a Gizatullin surface  $X$  along with a resolved SNC completion  $(\bar{X}, D)$ , where  $\bar{X}$  is a smooth projective surface and  $D \subset \bar{X}$  is a zigzag. By a sequence of blowups and blowdowns one can transform the dual graph  $\Gamma_D$  into a *standard form*, where  $C_0^2 = C_1^2 = 0$  and  $C_i^2 \leq -2$  for all  $i \geq 2$  if  $n \geq 4$  or  $C_i^2 = 0$  for all  $i$  if  $n \leq 3$  (cf. [19], [22], [36]). Moreover, this representation is unique up to reversion. The latter means that for two standard forms  $[[0, 0, w_2, \dots, w_n]]$  and  $[[0, 0, w'_2, \dots, w'_n]]$  of  $\Gamma_D$ , either  $w_i = w'_i$  or  $w_i = w'_{n+2-i}$  holds ([41]).

The reversion process can be described as follows. Start with a boundary divisor of type  $[[0, 0, w_2, \dots, w_n]]$ . Performing the elementary transformation (8) at the vertex corresponding to  $C_1$  one gets a boundary divisor of type  $[[ -1, 0, w_2 + 1, w_3, \dots, w_n ]]$ . After  $|w_2|$  steps one arrives at a boundary divisor of type  $[[w_2, 0, 0, w_3, \dots, w_n]]$ . Thus, one can move pairs of zeros to the right. Repeating this, one obtains finally a boundary divisor of type

$$[[w_2, \dots, w_n, 0, 0]] = [[0, 0, w_n, \dots, w_2]].$$

Notice that all the birational transformations involved are centered at the boundary (so to say, they yield the identity on the affine parts).

Let us recall the notion of an  $m$ -standard zigzag (see [22, (1.2)]).

**Definition 5.10.** A zigzag  $D$  of type  $[[0, -m, w_2, \dots, w_n]]$  with  $n \geq 1$  and  $w_i \leq -2$  is called  *$m$ -standard* (in the case  $n = 1$  there is no weight  $w_i$ ).

An  *$m$ -standard pair* is a pair  $(\bar{X}, D)$  consisting of a smooth projective surface  $\bar{X}$  and an  $m$ -standard zigzag  $D$  on  $\bar{X}$ . If  $m = 0$ , then  $(\bar{X}, D)$  is called a *standard pair*. A *birational map*  $\varphi : (\bar{X}, D) \rightarrow (\bar{X}', D')$  between  $m$ -standard pairs is a birational map  $\varphi : \bar{X} \rightarrow \bar{X}'$  which restricts to an isomorphism  $\varphi|_{\bar{X} \setminus D} : \bar{X} \setminus D \xrightarrow{\sim} \bar{X}' \setminus D'$ .

**Examples 5.11.** 1. The Danilov-Gizatullin surface  $X_d$  (see 5.3) has a boundary zigzag of type  $[[d]]$  with the standard form  $[[0, 0, (-2)_{d-1}]]$  (the index  $d-1$  means that there are  $d-1$  consecutive components with self-intersection index  $-2$ ). Any smooth affine surface  $X$  completable by a standard zigzag  $[[0, 0, (-2)_{d-1}]]$ ,  $d \geq 2$ , and non-isomorphic to  $\mathbb{P}^2 \setminus C$ , where  $C$  is a smooth conic, is isomorphic to the Danilov-Gizatullin surface  $X_d$ .

2. For a special Gizatullin surface (see 5.4) the standard zigzag is

$$[[0, 0, -2, \dots, -2, -w_s, -2, \dots, -2]], \quad \text{where } w_s < -2.$$

However, a Gizatullin surface with such a sequence of weights does not need to be special.

**Definition 5.12** (*Extended divisor*). Since the underlying smooth projective surface  $\bar{X}$  of a 0-standard pair is rational and  $C_0^2 = C_1^2 = 0$ , it is equipped with rational fibrations  $\Phi_i = \Phi_{|C_i|} : \bar{X} \rightarrow \mathbb{P}^1$  defined by the complete linear systems  $|C_i|$  on  $\bar{X}$ ,  $i = 0, 1$ , respectively. This defines a birational morphism ([37, Lem. 2.19])

$$\Phi = \Phi_0 \times \Phi_1 : \bar{X} \rightarrow \mathbb{P}^1 \times \mathbb{P}^1.$$

After a suitable coordinate change one may suppose that  $C_0 = \Phi_0^{-1}(\infty)$ ,  $\Phi(C_1) = \mathbb{P}^1 \times \{\infty\}$ , and  $C_2 \cup \dots \cup C_n \subseteq \Phi_0^{-1}(0)$ . The divisor  $D_{\text{ext}} := C_0 \cup C_1 \cup \Phi_0^{-1}(0)$  is called the *extended divisor*.

In order to determine the structure of the extended divisor, let us recall the notion of a *feather* ([37, Def. 5.5]).

**Definition 5.13** (*feathers*). (1) A *feather* is a linear chain

$$F : \begin{array}{ccccccc} & B & F_1 & & \dots & & F_s \\ & \circ & \circ & \text{---} & \dots & \text{---} & \circ \end{array}$$

of smooth rational curves such that  $B^2 \leq -1$  and  $F_i^2 \leq -2$  for all  $i \geq 1$ . The curve  $B$  is called the *bridge curve*.

(2) A *collection of feathers*  $\{F_\rho\}$  consists of pairwise disjoint feathers  $F_\rho$ ,  $\rho = 1, \dots, r$ . Such a collection will be denoted by a plus box

$$\begin{array}{c} \{F_\rho\} \\ \boxplus \end{array}.$$

(3) Let  $D = C_0 + \dots + C_n$  be a zigzag. A collection  $\{F_\rho\}$  is *attached to a curve*  $C_i$  if the bridge curves  $B_\rho$  of the feathers  $F_\rho$  meet  $C_i$  in pairwise distinct points and the feathers  $F_\rho$  are disjoint with the curves  $C_j$  for  $j \neq i$ .

**Lemma 5.14.** ([38, Prop. 1.11]) *Let  $(\tilde{X}, D)$  be a minimal SNC completion of the minimal resolution of singularities of a Gizatullin surface  $X$ . Furthermore, let  $D = C_0 + \dots + C_n$  be the boundary divisor in standard form. Then the extended divisor  $D_{\text{ext}}$  has the dual graph*

$$(10) \quad \Gamma_{\text{ext}} : \begin{array}{ccccccc} & & \{F_{2,j}\} & & \{F_{i,j}\} & & \{F_{n,j}\} \\ & & \boxplus & & \boxplus & & \boxplus \\ 0 & 0 & & \dots & & \dots & \\ \circ & \circ & \circ & \text{---} & \circ & \text{---} & \circ \\ C_0 & C_1 & C_2 & & C_i & & C_n \end{array},$$

where  $\{F_{i,j}\}$ ,  $j \in \{1, \dots, r_i\}$ , are feathers attached to the curve  $C_i$ . Moreover,  $\tilde{X}$  is obtained from  $\mathbb{P}^1 \times \mathbb{P}^1$  by a sequence of blowups with centers in the images of the components  $C_i$ ,  $i \geq 2$ .

**Remark 5.15.** Consider the feathers  $F_{i,j} := B_{i,j} + F_{i,j,1} + \cdots + F_{i,j,k_{i,j}}$  mentioned in Lemma 5.14. The collection of linear chains  $R_{i,j} := F_{i,j,1} + \cdots + F_{i,j,k_{i,j}}$  corresponds to the minimal resolution of singularities of  $X$ . Thus, if  $(\bar{X}, D)$  is a standard completion of  $X$  and  $(\tilde{X}, D)$  is the minimal resolution of singularities of  $(\bar{X}, D)$ , then the chain  $R_{i,j}$  contracts via  $\mu : (\tilde{X}, D) \rightarrow (\bar{X}, D)$  to a singular point of  $X$ , which is a cyclic quotient singularity. In particular,  $X$  has at most cyclic quotient singularities ([78, §3, Lem. 1.4.4(1)] and [38, Rem. 1.12]).

Hence  $X$  is smooth if and only if every  $R_{i,j}$  is empty, i. e. , if every feather  $F_{i,j}$  is irreducible and reduces to a single bridge curve  $B_{i,j}$  ([38, 1.8, 1.9 and Rem. 1.12]).

Let us introduce the notions of a  $*$ -component and a  $+$ -component.

**Definition 5.16.** (1) For a general feather  $F$  with dual graph

$$\Gamma_F : \begin{array}{ccccccc} \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ \\ B & & D_1 & & & & D_k \end{array}$$

and bridge curve  $B$  we call  $D_k$  the *tip component* of  $F$ .

(2) The component  $C_i$  is called a  $*$ -component if

- (i)  $D_{\text{ext}}^{\geq i+1}$  is not contractible and
- (ii)  $D_{\text{ext}}^{\geq i+1} - F_{j,k}$  is not contractible for every feather  $F_{j,k}$  of  $D_{\text{ext}}^{\geq i+1}$  such that the tip component of  $F_{j,k}$  has *mother component*  $C_\tau$ , that is, the component  $C_\tau$  with  $\tau < i$  carrying the center of blowup in which the tip component of  $F_{j,k}$  is born.

Otherwise  $C_i$  is called a  $+$ -component.

**Lemma 5.17.** Let  $D_{\text{ext}}$  be the extended divisor of the minimal resolution of singularities of a 1-standard completion of a Gizatullin surface  $X$ . Suppose that every  $C_i$ ,  $3 \leq i \leq n-1$ , is a  $*$ -component and that there is no feather attached to the component  $C_n$ . Then every feather  $F_{i,j}$  is an  $A_k$ -feather, that is, every  $F_{i,j}$  is contractible and therefore has the dual graph

$$\Gamma_{F_{i,j}} : \begin{array}{ccccccc} -1 & & -2 & & & & -2 \\ \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ \\ B & & D_1 & & & & D_k \end{array} ,$$

with  $k$  depending on  $i$  and  $j$ .


Note that for an  $A_k$ -feather the mother components of all curves  $D_1, \dots, D_k$  coincide, since any  $A_k$ -feather is born by successive blowups of a point on the boundary component it is attached to.

**Examples 5.18.** 1. A Gizatullin surface  $V$  is isomorphic to a nondegenerate toric surface  $V_{d,e} = \mathbb{A}^2/\mu_{d,e}$  if and only if for some (and then also for any) resolved standard completion  $(\tilde{X}, D)$  of  $X$  the dual graph  $\Gamma_{\text{ext}}$  of the associated extended divisor  $D_{\text{ext}}$  is a linear chain ([37, Lem. 2.20]).

2. Given a Danilov-Gizatullin surface  $X_d$  with  $d \neq 4$ , there are only  $d-1$  possible associated extended divisors (which do not depend, up to an isomorphism, on any further continuous parameter), with the dual graphs

$$\Gamma_{\text{ext}} : \begin{array}{ccccccccccccccc} & & & & & & & & & & 1-r & & & & & & & & & & & -1 \\ & & & & & & & & & & \circ & & & & & & & & & & & \circ \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \circ \\ C_0 & & C_1 & & C_2 & & C_3 & & & & C_{r-1} & & C_r & & C_{r+1} & & & & & & C_{d-2} & & C_{d-1} \end{array} ,$$

where  $2 \leq r \leq d-1$ . In addition, for  $d=4$ , there is another extended divisor possible; the corresponding affine surface is called the *affine pseudo-plane*. Its dual graph is



**5.4. Associated graph of groups.** Following [22] and [9], for an  $\mathbb{A}^1$ -fibered surface  $X$  we introduce a (not necessarily finite) graph  $\mathcal{F}_X$ , which reflects the structure of the group  $\text{Aut } X$ .

**Definition 5.19.** To any  $\mathbb{A}^1$ -fibered smooth affine surface  $\mu: X \rightarrow \mathbb{A}^1$  one associates the oriented graph  $\mathcal{F}_X$  as follows:

- A vertex of  $\mathcal{F}_X$  is an equivalence class of a 1-standard pair  $(\bar{X}, D)$  such that  $\bar{X} \setminus D \cong X$ , where two 1-standard pairs  $(\bar{X}_1, D_1, \bar{\mu}_1)$  and  $(\bar{X}_2, D_2, \bar{\mu}_2)$  define the same vertex if and only if  $(\bar{X}_1 \setminus D_1, \mu_1) \cong (\bar{X}_2 \setminus D_2, \mu_2)$ .
- An arrow of  $\mathcal{F}_X$  is an equivalence class of reversions. If  $\varphi: (\bar{X}, D) \rightarrow (\bar{X}', D')$  is a reversion, then the class  $[\varphi]$  of  $\varphi$  is an arrow starting from  $[(\bar{X}, D)]$  and ending at  $[(\bar{X}', D')]$ . Two reversions  $\varphi_1: (\bar{X}_1, D_1) \rightarrow (\bar{X}'_1, D'_1)$  and  $\varphi_2: (\bar{X}_2, D_2) \rightarrow (\bar{X}'_2, D'_2)$  define the same arrow if and only if there exist isomorphisms  $\theta: (\bar{X}_1, D_1) \rightarrow (\bar{X}_2, D_2)$  and  $\theta': (\bar{X}'_1, D'_1) \rightarrow (\bar{X}'_2, D'_2)$ , such that  $\varphi_2 \circ \theta = \theta' \circ \varphi_1$ .

**Remark 5.20.** It follows from the definition that for a 1-standard pair  $(\bar{X}, D)$  two reversions  $\varphi_1: (\bar{X}, D) \rightarrow (\bar{X}_1, D_1)$  and  $\varphi_2: (\bar{X}, D) \rightarrow (\bar{X}_2, D_2)$  centered at the points  $p_1$  and  $p_2$  define the same arrow if and only if there exists an automorphism  $\psi \in \text{Aut}(\bar{X}, D)$  such that  $\psi(p_1) = p_2$ .

The structure of the graph  $\mathcal{F}_X$  allows to decide, whether the automorphism group  $\text{Aut } X$  is generated by automorphisms of  $\mathbb{A}^1$ -fibrations. One says that  $\varphi \in \text{Aut } X$  is an *automorphism of  $\mathbb{A}^1$ -fibrations* if there exists an  $\mathbb{A}^1$ -fibration  $\mu: X \rightarrow \mathbb{A}^1$  such that  $\varphi$  induces an isomorphism  $\varphi: (X, \mu) \xrightarrow{\cong} (X, \mu)$ . Indeed, we have the following important fact.

**Theorem 5.21.** ([9, Prop. 4.0.7]) *Suppose that  $D$  has a component  $C_i$  with  $C_i^2 \leq -3$ . Then  $\text{Aut } X$  is generated by automorphisms of  $\mathbb{A}^1$ -fibrations if and only if  $\mathcal{F}_X$  is a tree. Furthermore, there is an exact sequence*

$$1 \longrightarrow H \longrightarrow \text{Aut } X \longrightarrow \pi_1(\mathcal{F}_X) \longrightarrow 1,$$

where  $H$  is the normal subgroup of  $\text{Aut } X$  generated by the automorphisms of  $\mathbb{A}^1$ -fibrations and  $\pi_1(\mathcal{F}_X)$  is the fundamental group of the graph  $\mathcal{F}_X$ .

**Remark 5.22.** Due to Corollary 8.25 below, each of the automorphism groups of  $\mathbb{A}^1$ -fibrations which generate  $H$  is an extension of a metabelian connected nested ind-group of rank  $\leq 2$  by an at most countable group. The same concerns the factors of the amalgams considered in the next subsection.

One can equip  $\mathcal{F}_X$  with a structure of a graph of groups as follows.

**Definition 5.23.** Let  $X$  be a normal quasi-projective surface, and let  $\mathcal{F}_X$  be its associated graph. Then  $\mathcal{F}_X$  admits a structure of a graph of groups by the following choice:

- For any vertex  $v$  of  $\mathcal{F}_X$ , fix a 1-standard pair  $(\bar{X}_v, D_v, \bar{\mu}_v)$  in the class  $v$ . The group  $G_v$  is equal to  $\text{Aut}(\bar{X}_v \setminus D_v, \mu_v)$ .
- For any arrow  $\sigma$  of  $\mathcal{F}_X$ , fix a reversion  $r_\sigma: (\bar{X}_\sigma, D_\sigma, \bar{\mu}_\sigma) \rightarrow (\bar{X}'_\sigma, D'_\sigma, \bar{\mu}'_\sigma)$  in the class of  $\sigma$  and also an isomorphism  $\phi_\sigma: (\bar{X}'_\sigma \setminus D'_\sigma, \mu'_\sigma) \rightarrow (\bar{X}_{t(\sigma)} \setminus D_{t(\sigma)}, \mu_{t(\sigma)})$ . Then the group  $G_\sigma$  is equal to

$$\{(\varphi, \varphi') \in \text{Aut}(\bar{X}_\sigma, D_\sigma) \times \text{Aut}(\bar{X}'_\sigma, D'_\sigma) \mid r_\sigma \circ \varphi = \varphi' \circ r_\sigma\}$$



and the monomorphisms  $\kappa_\sigma : G_\sigma \rightarrow G_{s(\sigma)}$  and  $\lambda_\sigma : G_\sigma \rightarrow G_{t(\sigma)}$  are given by  $\kappa_\sigma((\varphi, \varphi')) = \phi_{\sigma^{-1}} \circ \varphi \circ \phi_{\sigma^{-1}}^{-1}$  and  $\lambda_\sigma((\varphi, \varphi')) = \phi_\sigma \circ \varphi' \circ \phi_\sigma^{-1}$ .

The first version of the following theorem was established by Danilov and Gizatullin ([22, Thm. 5]). It connects the structure of the graph of groups on  $\mathcal{F}_X$  with the group  $\text{Aut } X$ .

**Theorem 5.24.** ([22, Thm. 5], see also [9, Thm. 4.0.11]) *Let  $(\bar{X}, D)$  be a 1-standard pair such that  $D$  has a component  $C_i$  with  $C_i^2 \leq -3$ . Let  $X = \bar{X} \setminus D$ . If  $\mathcal{F}_X$  is equipped with a structure of a graph of groups as in Definition 5.23, then  $\text{Aut } X \cong \pi_1(\mathcal{F}_X)$ .*

There is the following important consequence concerning the structure of the automorphism groups of Gizatullin surfaces.

**Corollary 5.25.** *Under the assumptions of Theorem 5.24 suppose in addition that  $\mathcal{F}_X$  is a tree with vertices  $[(\bar{X}_i, D_i)]$ ,  $i \in I$ . Then  $\text{Aut } X$  is an amalgam of the automorphism groups  $\text{Aut}(\bar{X}_i \setminus D_i, \mu_i)$  of  $\mathbb{A}^1$ -fibrations over  $\mathbb{A}^1$ .*

**5.5. Amalgam structures for Gizatullin surfaces.** In this section we list all the Gizatullin surfaces known to the authors, where the automorphism group is an amalgam (however, see [27] for further potential examples). The easiest way to present such surfaces is to describe various 1-standard completions of them in terms of the dual graphs of their extended divisors.

Although the following theorem is a special case of Theorem 5.27, it is worth to be mentioned independently.

**Theorem 5.26.** ([9, Thm. 5.4.5]) *Consider a Danielewski surface*

$$X = \{xy - P(z) = 0\} \subseteq \mathbb{A}^3, \quad \text{where } P(z) \in \mathbb{K}[z]$$

*has degree  $n \geq 1$ . Then  $X$  has a standard completion  $(\bar{X}, D)$  of type  $[[0, 0, -n]]$ . Letting  $\tau \in \text{Aut } X$  be the involution  $(x, y, z) \mapsto (y, x, z)$  and  $\mu : X \rightarrow \mathbb{A}^1$  be the  $\mathbb{A}^1$ -fibration  $(x, y, z) \mapsto x$ , we let  $A = \langle \text{Aut}(\bar{X}, D), \tau \rangle \subseteq \text{Aut } X$  and  $J = \text{Aut}(X, \mu)$ . Then  $A \cap J = \text{Aut}(\bar{X}, D)$  and*

$$\text{Aut } X = A \star_{A \cap J} J.$$

This result can be generalized as follows.

**Theorem 5.27.** ([62, Cor. 3.19, cf. Thm. 4.4]) *Let  $X$  be a smooth Gizatullin surface satisfying the following condition (see diagram (10)):*

- (\*)  *$X$  admits a 1-standard completion  $(\bar{X}, D)$  such that  $C_3, \dots, C_{n-1}$  are  $*$ -components and there is no feather attached to  $C_2$  and to  $C_n$ .*

*Fix an  $\mathbb{A}^1$ -fibration  $\mu : X \rightarrow \mathbb{A}^1$ , and let  $\mu^\vee : X \rightarrow \mathbb{A}^1$  be the  $\mathbb{A}^1$ -fibration induced by the reversion  $\psi : (\bar{X}, D) \rightarrow (\bar{X}^\vee, D^\vee)$  with center  $p \in C_0 \setminus C_1$ . Then  $\mathcal{F}_X$  has one of the following structures:*

$$\mathcal{F}_X : [(\bar{X}, D)] \bullet \longleftrightarrow \bullet [(\bar{X}^\vee, D^\vee)] \quad \text{or} \quad \mathcal{F}_X : [(\bar{X}, D)] \bullet \circlearrowleft .$$

*If  $\mathcal{F}_X$  is of the form  $\bullet \circlearrowleft$ , then  $D^{\geq 2}$  is a palindrome.*

- (a) *Let  $\mathcal{F}_X$  be of the form  $\bullet \circlearrowleft$ , that is,  $(\bar{X}, D) \cong (\bar{X}^\vee, D^\vee)$ . Then*

$$\text{Aut } X = A \star_{A \cap J} J,$$


*where  $A = \langle \text{Aut}(\bar{X}, D), \psi \rangle$ ,  $J = \text{Aut}(X, \mu)$ , and  $A \cap J = \text{Aut}(\bar{X}, D)$ .*

- (b) *Let  $\mathcal{F}_X$  be of the form  $[(\bar{X}, D)] \bullet \longleftrightarrow \bullet [(\bar{X}^\vee, D^\vee)]$ . Denote by  $A$  the subgroup corresponding to the edge and by  $J$  and  $J^\vee$  the subgroups  $J = \text{Aut}(X, \mu)$  and  $J^\vee = \text{Aut}(X, \mu^\vee)$ . Identifying  $J \leftrightarrow A \leftrightarrow J^\vee$  we have*


$$\text{Aut } X = J \star_A J^\vee.$$

An important particular case of Theorem 5.27 is that of the toric affine surfaces with the amalgam structures on their automorphism groups as exposed in 3.3-3.6.


Another interesting example of a family of smooth Gizatullin surfaces, for which the automorphism groups are amalgams, is the following one. Consider any smooth 1-standard pair  $(\bar{X}, D)$  such that the dual graph of  $D_{\text{ext}}$  has the following form:



where  $C_3, \dots, C_{n-1}$  are  $\star$ -components. Hence any feather  $F_j$  has self-intersection index  $F_j^2 = -1$ . Reversion of  $(\bar{X}, D)$  may lead to two different completions, namely those with the dual graphs of the extended divisors



and



respectively, depending on the choice of the center of reversion  $\lambda \in C_0 \setminus C_1$ . It is not difficult to see that the action of  $\text{Aut}(\bar{X}, D)$  on  $C_0 \setminus C_1$  admits two orbits, namely an open orbit  $(C_0 \setminus C_1) \setminus \{p\}$  and a point  $\{p\}$  (depending on the position of the feather  $G$  attached to  $C_n$ ). Moreover, the actions of  $\text{Aut}(\bar{X}', D')$  and  $\text{Aut}(\bar{X}'', D'')$ , respectively, on  $C'_0 \setminus C'_1$  and  $C''_0 \setminus C''_1$ , respectively, are transitive. These observations lead to the following proposition.

**Proposition 5.28.** *The graph  $\mathcal{F}_X$  associated to  $X = \bar{X} \setminus D$  is*

$$\mathcal{F}_X : [(\bar{X}', D')] \xleftrightarrow{\sigma'} [(\bar{X}, D)] \xleftrightarrow{\sigma''} [(\bar{X}'', D'')].$$

Fixing arbitrary reversions  $\alpha : (\bar{X}, D) \rightarrow (\bar{X}', D')$  and  $\beta : (\bar{X}, D) \rightarrow (\bar{X}'', D'')$ , it follows that  $\text{Aut} X$  is an amalgam of the groups  $\text{Aut}(X, \mu)$ ,  $\text{Aut}(X, \mu' \circ \alpha)$ , and  $\text{Aut}(X, \mu'' \circ \beta)$ , amalgamated over their pairwise intersections.

**5.29.** The last statement requires an explanation. By [9, Thm. 3.0.2], every automorphism of a Gizatullin surface admits (an essentially unique) decomposition in fibered modifications and reversions. Let us fix two reversions  $\alpha : (\bar{X}, D) \rightarrow (\bar{X}', D')$  and  $\beta : (\bar{X}, D) \rightarrow (\bar{X}'', D'')$  (which are, as we have seen, unique up to equivalence). Then every automorphism of  $X$  has an (essentially unique) decomposition into maps of the form  $\phi \in \text{Aut}(X, \mu)$ ,  $\alpha^{-1}\varphi'\alpha \in \text{Aut}(X, \mu' \circ \alpha)$  with  $\varphi' \in \text{Aut}(X, \mu')$ , and  $\beta^{-1}\varphi''\beta \in \text{Aut}(X, \mu'' \circ \beta)$  with  $\varphi'' \in \text{Aut}(X, \mu'')$ .

**Examples 5.30.** 1. Given a Danilov-Gizatullin surface  $V_d$  (see Example 5.3), for  $d = 2, 3, 4, 5$  the group  $\text{Aut}(V_d)$  is an amalgam of a finite set of nested subgroups, hence is finitely bearable, see [23], §§6 - 10 for details. Whereas for  $d \geq 7$  this group is not countably bearable, see Corollary 5.6.

2. An interesting example of a smooth Gizatullin surface  $X$  with an amalgam structure of  $\text{Aut } X$  is given by the following construction, see [9, 5.5]. For  $a, b \in \mathbb{K}^*$ ,  $c \in \mathbb{K}$ , and  $a \neq b$ , consider the smooth Gizatullin surface  $X_{a,b,c}$  in  $\mathbb{A}^4$  given by the equations

$$\begin{aligned}xz &= y(y-a)(y-b), \\yw &= z(z-c), \\xw &= (y-a)(y-b)(z-c).\end{aligned}$$

The (abstract) isomorphism type of  $X_{a,b,c} =: X$  does not depend on the parameters  $a, b, c$ , see [9, 5.5.6]. Furthermore,  $X$  possesses a 1-standard completion of type  $[[0, -1, -2, -3]]$ . It is an easy exercise to show that  $X$  admits 4 different families of 1-standard completions  $(\bar{X}_1, D_1)$ ,  $(\bar{X}_{2,t}, D_{2,t})$ ,  $(\bar{X}_{3,t}, D_{3,t})$ , and  $(\bar{X}_4, D_4)$ , two of them depending on a parameter  $t \in \mathbb{K} \setminus \{0, 1\}$  (and these are isomorphic if and only if the parameters  $t, t'$  are equivalent under the relation  $\sim$  generated by  $t \sim t^{-1}$ ) and the other two are independent on any parameter. It is shown in [9, 5.5.4] that the associated graph  $\mathcal{F}_X$  has the following structure:

$$\begin{array}{ccccc} & & & & [(X_{2,s}, D_{2,s})] \longleftrightarrow [(X_{3,s}, D_{3,s})] \\ & & & \nearrow & \\ & & & & \vdots \\ [(X_4, D_4)] \longleftrightarrow [(X_1, D_1)] & & & & \vdots \\ & & & \searrow & \\ & & & & [(X_{2,t}, D_{2,t})] \longleftrightarrow [(X_{3,t}, D_{3,t})]\end{array}$$

where  $(s, t) \in (\mathbb{K} \setminus \{0, 1\})^2$  factorized by the equivalence relation  $s \sim s^{-1}$ . The group  $\text{Aut } X$  is an amalgam of the (uncountable set of) groups of automorphisms of  $\mathbb{A}^1$ -fibrations, see [9, 5.5.5]. In particular, this group is uncountably bearable.

Let us conclude this section with the following problem.

**Problem.** *Describe all the Gizatullin surfaces  $X$  such that the neutral component  $\text{Aut}^\circ X$  is a (finitely or countably) bearable group.*

## 6. AUTOMORPHISM GROUPS OF $\mathbb{A}^1$ -FIBRATIONS

As we have seen in Section 5, the automorphism groups of  $\mathbb{A}^1$ -fibrations play an essential role in studying the full automorphism group. For an  $\mathbb{A}^1$ -fibered variety of arbitrary dimension, we describe in Subsection 6.1 the unipotent radical of such a group as a nested ind-group. In Subsection 6.2 we give some immediate applications to the neutral component of the automorphism group of a given  $\mathbb{A}^1$ -fibration  $\mu: X \rightarrow B$  on a normal affine surface  $X$  over an affine curve  $B$ . Then in Sections 7 and 8 we dwell to a description of the full group of automorphisms  $\text{Aut}(X, \mu)$  in the surface case. For an  $\text{ML}_1$ -surface  $X$ , this group coincides with the full automorphism group  $\text{Aut } X$ .

### 6.1. Generalized de Jonquières groups.

**Definition 6.1.** Let  $X$  be a normal affine variety, and let  $\mu: X \rightarrow Z$  be an  $\mathbb{A}^1$ -fibration over a normal affine variety  $Z$ , that is, a morphism with general scheme theoretical fibers isomorphic to the affine line. We assume that  $\text{codim}_Z(Z \setminus \mu(X)) \geq 2$ . Consider the subgroups

- $\text{Aut}(X, \mu) \subset \text{Aut } X$  of all automorphisms of  $X$  preserving the fibration  $\mu$ ;
- $\text{Aut}_Z(X, \mu) \subset \text{Aut}(X, \mu)$  of those automorphisms which preserve each  $\mu$ -fiber;
- $\mathbb{U}_\mu \subset \text{Aut}_Z(X, \mu)$  of those automorphisms which restrict to translations on general  $\mu$ -fibers.

Clearly,  $\mathbb{U}_\mu$  is an infinite dimensional Abelian group. If  $X = \mathbb{A}^2$  and  $\mu: (x, y) \mapsto x$ , then  $\mathbb{U}_\mu$  is the maximal unipotent subgroup of the de Jonquières group, see 3.3. In the general case we call  $\text{Aut}(X, \mu)$  a *generalized de Jonquières group*, and  $\mathbb{U}_\mu$  a *generalized unipotent de Jonquières group*.

**Remarks 6.2.** 1. Recall that two  $\mathbb{A}^1$ -fibrations  $\mu_i : X \rightarrow Z_i$  on a normal affine variety  $X$  over normal affine varieties  $Z_i$ ,  $i = 1, 2$ , are said to be equivalent if one can be sent into another by an automorphism of  $X$  which induces an isomorphism of  $Z_1$  and  $Z_2$ . Clearly,  $\mu_1$  and  $\mu_2$  are equivalent if and only if the corresponding subgroups  $\mathbb{U}_{\mu_1}$  and  $\mathbb{U}_{\mu_2}$  are conjugated in  $\text{Aut } X$ .

2. If  $\mu$  is locally trivial then  $\mathbb{U}_\mu$  is the union of its unipotent one-parameter subgroups. In fact, for any  $\alpha \in \mathbb{U}_\mu$  there is a locally nilpotent regular vertical vector field  $\partial$  on  $X$  such that  $\alpha = \exp \partial$ , and so,  $\alpha$  belongs to the unipotent one-parameter subgroup  $H = \{\exp(t\partial)\}_{t \in \mathbb{K}} \subset \mathbb{U}_\mu$ . The latter holds as well if  $X$  can be covered by affine charts  $(U_i)_{i \in I}$  such that the restriction  $\mu|_{U_i}$  is locally trivial for each  $i \in I$  (such charts are automatically  $\alpha$ -stable). Moreover, the same conclusion remains true under a weaker assumption that in each chart  $U_i$  of the covering the  $\mathbb{A}^1$ -fibration  $\mu$  becomes locally trivial after a cyclic base change. This is the case, for instance, for any normal  $\mathbb{A}^1$ -fibered affine surface.

3. Let  $X$  be a normal affine surface. Then any  $\mathbb{G}_a$ -action on  $X$  acts along the fibers of an  $\mathbb{A}^1$ -fibration  $\mu : X \rightarrow Z$  over a smooth affine curve  $Z$ , see, e.g., [30, Lem. 1.1]. Thus, the group  $\text{SAut } X$  is generated by the unipotent de Jonquières subgroups  $\mathbb{U}_\mu$ , where  $\mu$  runs over the set of all the  $\mathbb{A}^1$ -fibrations on  $X$  with affine bases. If  $X$  is an  $\text{ML}_1$ -surface, then  $\mu$  is unique, and so,  $\text{SAut } X = \mathbb{U}_\mu$ .

The group  $\mathbb{U}_\mu$  admits the following presentation. We let  $\text{Frac}(A)$  denote the quotient field of an integral domain  $A$ , and  $\mathbb{K}(Y)$  the function field of an algebraic variety  $Y$  over  $\mathbb{K}$ .

**Theorem 6.3.** *Let  $\mu : X \rightarrow Z$  be an  $\mathbb{A}^1$ -fibration as in Definition 6.1. Suppose that  $\mathbb{U}_\mu$  is the union of its unipotent one-parameter subgroups (see Remark 6.2.2). Then*

$$\mathbb{U}_\mu \cong H^0(Z, \mathcal{O}(D))$$

for a divisor  $D$  on  $Z$ , where the class  $[D] \in \text{Pic } Z$  is uniquely defined by  $\mu$ . If  $\text{Pic } Z = 0$ , then there exists a locally nilpotent derivation  $\partial_0 \in \text{Der } \mathcal{O}(X)$  such that  $\mathbb{U}_\mu = \exp((\ker \partial_0) \cdot \partial_0)$ .

*Proof.* Let  $A = \mathcal{O}(X)$ . Shrinking  $Z$  appropriately one can obtain an affine ruling, and even a locally trivial  $\mathbb{A}^1$ -bundle  $X_\omega \rightarrow \omega$  on the normal affine variety  $X_\omega = \mu^{-1}(\omega)$  over a normal affine base  $\omega$ , where  $\omega$  is a principal open subset of  $Z$  ([60]; see also [59]). Shrinking the base further, one may assume that  $X_\omega \cong \omega \times \mathbb{A}^1 \rightarrow \omega$  is a principal cylinder in  $X$ . There exists a  $\mathbb{G}_a$ -action  $U = \exp(\mathbb{K} \cdot \partial)$  along the fibers of  $\mu$ , where  $\partial \in \text{Der}(A)$  is locally nilpotent and  $\ker \partial = \mu^*(\mathcal{O}_Z(Z))$ , see, e.g., [61, Prop. 3.1.5].

If  $U' = \exp(\mathbb{K} \cdot \partial')$  is a second  $\mathbb{G}_a$ -action on  $X$  along the fibers of  $\mu$ , then  $\partial' = f\partial$  for some  $f \in \text{Frac}(\ker \partial) = \mu^*(\mathcal{O}_Z(Z))$  such that  $f \cdot \partial(A) \subset A$ . Conversely, for any rational function  $f \in \mu^*(\mathcal{O}_Z(Z))$  such that  $f \cdot \partial(A) \subset A$ , the derivation  $\partial' = f\partial$  is locally nilpotent on  $A$ , and so  $U' = \exp(\mathbb{K} \cdot \partial') \subset \mathbb{U}_\mu$ , see, e.g., [33, Prop. 1.1(b)].

It follows that  $\mathbb{U}_\mu = \exp(\mu^* \mathcal{H} \cdot \partial)$ , where

$$\mathcal{H} = \{u \in \mathcal{O}_Z(Z) \mid \partial(A) \subset \mu^*(u^{-1})A\}.$$

Note that  $\mathcal{H}$  is an  $\mathcal{O}_Z(Z)$ -module. There is an isomorphism  $\mathcal{H} \xrightarrow{\cong} \mathbb{U}_\mu$ ,  $u \mapsto \exp(\mu^*(u) \cdot \partial)$ . Assume that the set  $\mathfrak{D} = \{-\text{div}(u) \mid u \in \mathcal{H}\}$  is bounded above, and consider the divisor  $D = \sup \mathfrak{D}$  on  $Z$ . We claim that  $\mathcal{H} = H^0(Z, \mathcal{O}_Z(D))$ , that is,  $u \in \mathcal{H}$  if and only if  $-\text{div}(u) \leq D$ . Since  $\mathcal{H} \cong \mathbb{U}_\mu$ , this yields the required isomorphism  $\mathbb{U}_\mu \cong H^0(Z, \mathcal{O}_Z(D))$ .

To show the claim, it suffices to establish the inclusion  $H^0(Z, \mathcal{O}_Z(D)) \subset \mathcal{H}$ , the converse inclusion being clear from the definition of  $D$ . Let  $u \in \mathcal{O}_Z(Z)$  be such that  $-\operatorname{div}(u) \leq D$ . Then there exists a cortège  $(u_1, \dots, u_n) \in \mathcal{H}^n$  such that  $-\operatorname{div}(u) \leq \max_{1 \leq i \leq n} \{-\operatorname{div}(u_i)\}$ . We claim that  $u \in \mathcal{H}$ , i.e.,  $\mu^*(u)\partial A \subset A$ , or, which is equivalent, that  $-\operatorname{div}(\mu^*(u)\partial a) \leq 0$  for any  $a \in A$ . Indeed, one has

$$-\operatorname{div}(\mu^*(u)\partial a) \leq \max_{1 \leq i \leq n} \{-\operatorname{div}(\mu^*(u_i)\partial a)\} \leq 0,$$

since  $-\operatorname{div}(\mu^*(u_i)\partial a) \leq 0$  for  $i = 1, \dots, n$ . This proves our claim.

To finish the proof of the first assertion, it remains to show that  $\mathfrak{D}$  is bounded above. Choose an element  $a \in \ker \partial^2 \setminus \ker \partial \subset A$ . Then  $\partial a = \mu^*(h) \in \ker \partial = \mu^*(\mathcal{O}_Z(Z))$ , where  $h \in \mathcal{O}_Z(Z)$ . For  $u \in \mathcal{H}$  we have  $\mu^*(h) \in \partial(A) \subset \mu^*(u^{-1})A$ . Hence there exists  $b \in \mathcal{O}(Z)$  such that  $h = u^{-1}b$ . Thus,  $-\operatorname{div} u \leq \operatorname{div} h$  and so,  $\operatorname{div} h$  is an upper bound for  $\mathfrak{D}$ . Actually our argument shows that  $D \leq D_0$ , where the effective divisor  $D_0 = \inf\{\operatorname{div}(h) \mid \mu^*h \in \partial(\ker \partial^2)\}$  on  $Z$  is given by the zero locus of the ideal  $I = \mu_*(\partial(\ker \partial^2)) \subset \mathcal{O}_Z(Z)$ .

Replacing in our construction  $\partial$  by  $\partial'$  results in replacing the divisor  $D$  by a linearly equivalent one  $D'$ . Starting with a suitable derivation  $\partial'$  of the form  $f\partial$ , where  $f \in \mu^*(\mathcal{O}_Z(Z))$ , one can get as  $D'$  an arbitrary representative of the class  $[D] \in \operatorname{Pic} Z$ . Thus our construction associates canonically the class  $[D]$  to the  $\mathbb{A}^1$ -fibration  $\mu: X \rightarrow Z$ .

Now the second assertion follows easily. Indeed, if  $\operatorname{Pic} Z = 0$ , then  $D = -\operatorname{div}(u) = \operatorname{div}_\infty(u)$  for a rational function  $u \in \mathbb{K}(Z)$ . Then the locally nilpotent derivation  $\partial_0 = \mu^*(u)\partial \in \operatorname{Der} A$  satisfies  $\mathbb{U}_\mu = \exp((\ker \partial_0) \cdot \partial_0)$ . We leave the details to the reader.  $\square$

**Corollary 6.4.**  $\mathbb{U}_\mu$  is a unipotent Abelian nested ind-group.

*Proof.* Let  $(\bar{Z}, D')$  be a completion of  $Z$  by an SNC divisor  $D' = \bar{Z} \setminus Z$ . Then

$$H^0(Z, \mathcal{O}_Z(D)) = \varinjlim_n H^0(\bar{Z}, \mathcal{O}_{\bar{Z}}(D - nD')),$$

where  $H^0(\bar{Z}, \mathcal{O}_{\bar{Z}}(D - nD'))$  is a finite dimensional vector group for each  $n$ . Therefore, the vector group  $\mathbb{U}_\mu \cong H^0(Z, \mathcal{O}_Z(D))$  is a nested ind-group.  $\square$

**Remarks 6.5.** 1. Suppose that the affine variety  $X$  as in Definition 6.1 admits a free  $\mathbb{G}_a$ -action along the  $\mu$ -fibers. Then the corresponding locally nilpotent  $\mu$ -vertical vector field  $\partial_0$  (that is,  $\partial_0$  is tangent to the  $\mu$ -fibers) has no zero, and so, divides any other locally nilpotent  $\mu$ -vertical vector field on  $X$ . Then the equality  $\mathbb{U}_\mu = \exp((\ker \partial_0) \cdot \partial_0)$  holds. This is the case, for instance, for any smooth Danielewski surface  $xy - p(z) = 0$  in  $\mathbb{A}^3$ .

2. Consider a line bundle  $L = (\mu: X \rightarrow Z)$ , that is, a locally trivial  $\mathbb{A}^1$ -bundle on  $Z$  with a fixed (zero) section. Then any  $\mathbb{G}_a$ -action on  $X$  along the  $\mu$ -fibers is uniquely defined by the image of the zero section. Vice versa, given a section  $S$  of  $\mu$ , there is a unique  $\mathbb{G}_a$ -action on  $X$  along the  $\mu$ -fibers which sends the zero section to  $S$ . Hence  $\mathbb{U}_\mu \cong H^0(Z, \mathcal{O}_Z(L)) \cong H^0(Z, \mathcal{O}_Z(D))$  for any divisor  $D$ , which represents the class of  $L$  in  $\operatorname{Pic} Z$ .

**Proposition 6.6.** Under the assumptions of Theorem 6.3 there is an exact sequence

$$(11) \quad 1 \rightarrow \mathbb{U}_\mu \rightarrow \operatorname{Aut}_Z(X, \mu) \rightarrow \Delta_\mu \rightarrow 1,$$

where  $\Delta_\mu \cong \Gamma_\mu \times \mathbb{Z}^l$  for some  $l \geq 0$  and some subgroup  $\Gamma_\mu \subset \mathbb{G}_m$ .

*Proof.* We use the notation from the proof of Theorem 6.3. Since  $X_\omega \cong_Z \omega \times \mathbb{A}^1$  we have  $\operatorname{Aut}_\omega(X_\omega, \mu) \cong \mathcal{O}_\omega^+(A) \rtimes \mathcal{O}_\omega^\times(A)$ , where  $A^+$  stands for the additive group of an

algebra  $A$  and  $A^\times$  for its multiplicative group. The natural embedding  $\text{Aut}_Z(X, \mu) \hookrightarrow \text{Aut}_\omega(X_\omega, \mu)$  induces the commutative diagram

$$(12) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \mathcal{O}_\omega^+(\omega) & \longrightarrow & \text{Aut}_\omega(X_\omega, \mu) & \longrightarrow & \mathcal{O}_\omega^\times(\omega) \longrightarrow 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ 1 & \longrightarrow & \mathbb{U}_\mu & \longrightarrow & \text{Aut}_Z(X, \mu) & \longrightarrow & \Delta_\mu \longrightarrow 1 \end{array}$$

where  $\Delta_\mu$  is the image of  $\text{Aut}_Z(X, \mu)$  in  $\mathcal{O}_\omega^\times(\omega)$ . By Samuel's Units Lemma ([93, Lem. 1], see also [71, Lem. 4.3]) we have  $\mathcal{O}_\omega^\times(\omega) \cong \mathbb{G}_m \times \mathbb{Z}^N$  for some  $N \geq 0$ . It is easily seen that any subgroup of the product  $\mathbb{G}_m \times \mathbb{N}^N$  is a product of subgroups of the factors. Hence  $\Delta_\mu \cong \Gamma_\mu \times \mathbb{Z}^l$  for some  $l \geq 0$  and some subgroup  $\Gamma_\mu \subset \mathbb{G}_m$ .  $\square$

**6.2. The surface case.** Here we give some immediate applications of Theorem 6.3 to the neutral component and the unipotent radical of the group  $\text{Aut}(X, \mu)$  in the case of a normal affine surface  $X$ . In the next two sections we enterprise a more thorough study of this group.

**Remarks 6.7.** 1. In the surface case, sequence (11) splits and the subgroup  $\Gamma_\mu \subset \mathbb{G}_m$  is closed, see Proposition 8.23 and Theorem 8.24. Plausibly, the latter holds in the general case as well.

2. Rentschler's Theorem ([89]) says that any locally nilpotent derivation of the polynomial ring  $\mathbb{K}[x, y]$  is conjugated to the derivation  $\partial_0 = f(x)\partial/\partial y$  for some  $f \in \mathbb{K}[x]$ . This can be generalized to the surfaces of class  $(\text{ML}_0)$  as follows.

**Proposition 6.8.** *Let  $X$  be a smooth Gizatullin  $\mathbb{G}_m$ -surface, which is neither Danilov-Gizatullin, nor special (see Examples 5.3 and 5.4). Then there exists two locally nilpotent derivations  $\partial_0, \partial_1 \in \text{Der } \mathcal{O}_X(X)$  such that any other locally nilpotent derivation of  $\mathcal{O}_X(X)$  is conjugated to a one of the form  $f_i \partial_i$  for some  $i \in \{0, 1\}$  and  $f_i \in \ker \partial_i$ .*

*Proof.* This is an immediate consequence of Theorems 5.5 and 6.3.  $\square$

For the next theorem we address the reader to [8], [84]; cf. Proposition 4.4.

**Theorem 6.9.** *Let  $X$  be a normal affine surface of class  $(\text{ML}_1)$  or  $(\text{ML}_2)$ . Then the neutral component  $\text{Aut}^\circ X \subset \text{Aut } X$  is isomorphic to*

- $\mathbb{G}_m^r$  if  $X$  is of type  $(\text{ML}_2, r)$ ,  $r \in \{0, 1, 2\}$ , and
- $\mathbb{U}_\mu \rtimes \mathbb{G}_m^r$  if  $X$  is of type  $(\text{ML}_1, r)$ ,  $r \in \{0, 1, 2\}$ , where  $\mu: X \rightarrow B$  is a unique  $\mathbb{A}^1$ -fibration on  $X$  over an affine curve  $B$ .

**Remark 6.10.** Recall (see 3.2) that the class  $(\text{ML}_2, 2)$  consists of a single surface  $(\mathbb{A}_x^1)^2$ , and the class  $(\text{ML}_1, 2)$  of a single surface  $\mathbb{A}^1 \times \mathbb{A}_x^1$ ; see 3.5 and 3.6 for a description of the corresponding automorphism groups.

From Corollary 6.4 and Theorem 6.9 we deduce the following result; see [8, Cor. 2.3] for an alternative proof in the case of a rational surface.

**Corollary 6.11.** *If  $X$  is a surface of class  $(\text{ML}_1)$  or  $(\text{ML}_2)$ , then  $\text{Aut}^\circ X$  is a solvable nested ind-group, and any two maximal tori in  $\text{Aut}^\circ X$  are conjugated.*

## 7. FORMAL NEIGHBORHOOD OF A FIBER IN AN $\mathbb{A}^1$ -FIBRATION

In this section we consider a normal affine surface  $X$  equipped with an  $\mathbb{A}^1$ -fibration  $\mu: X \rightarrow B$  over a smooth affine curve  $B$ . We study formal neighborhoods of fibers, the corresponding arc spaces, and their stabilizers. These technical tools are used in the next section in the proofs of our main Theorems 8.12 and 8.24 on the structure of the automorphism group of an  $\mathbb{A}^1$ -fibration.

## 7.1. Chain of contractions.

**Notation 7.1.** Fix a minimal resolved SNC completion  $(\bar{X}, D)$  of  $X$  such that  $\mu$  extends to a  $\mathbb{P}^1$ -fibration  $\bar{\mu}: \bar{X} \rightarrow \bar{B}$ . Recall that  $\bar{X}$  is smooth and contains the minimal resolution of singularities of  $X$  (see Notation 5.8).

We call the degenerate  $\bar{\mu}$ -fibers *special* and denote their union by  $\mathcal{T} \subset \bar{X}$ . The components of special fibers are smooth rational curves, and their dual graphs are trees.

Let  $S$  be the unique component of the boundary divisor  $D = \bar{X} \setminus X$ , which is a section of  $\bar{\mu}$ . The dual graph  $\Gamma_D$  of  $D$  is a rooted tree with a root vertex  $S$ . The irreducible components of  $F := \bar{\mu}^{-1}(\bar{B} \setminus B)$  correspond to the 0-vertices of degree 1 that are neighbors of  $S$ . The other branches of  $\Gamma_D$  at  $S$  are nonempty connected subgraphs of the dual graphs of the special fibers over  $B$ ; their intersection bilinear forms are negative definite. Recall that  $D \subset D_{\text{ext}}$ , where  $D_{\text{ext}} = S + F + \mathcal{T}$  is the extended divisor of  $(\bar{X}, \bar{\mu})$ .

**Remark 7.2.** By Miyanishi's Theorem [77, Ch. 3, Lem. 1.4.4(1)],  $X$  has only cyclic quotient singularities. The weighted dual graph of the minimal resolution of such a singular point is a Hirzebruch-Jung string, see [6].

The following lemma is well known. For the reader's convenience, we recall the proof.

**Lemma 7.3.** *Let  $V$  be a smooth projective surface,  $Z$  be a smooth projective curve, and let  $\pi: V \rightarrow Z$  be a  $\mathbb{P}^1$ -fibration, which admits a section  $s: Z \rightarrow V$  with image  $S$ . Then there is a sequence of contractions*

$$V = V_n \rightarrow V_{n-1} \rightarrow \dots \rightarrow V_0$$

*of  $(-1)$ -components of degenerate fibers disjoint from  $S$  and from the subsequent images of  $S$  that terminates by a ruled surface  $V_0$  with an induced ruling  $\pi_0: V_0 \rightarrow Z$ .*

*Proof.* We have the following

*Claim.* *If a fiber  $F$  of  $\pi$  is degenerate, then either  $F$  contains at least two  $(-1)$ -components, or such a component is unique and multiple. In any case, at least one of the  $(-1)$ -components of  $F$  is disjoint from  $S$ .*

Since  $F \cdot S = 1$ , the second assertion follows from the first. Then also the lemma follows by induction on the total number of components of degenerate fibers. Indeed, while contracting a  $(-1)$ -component of the fiber as an induction step, we reproduce the setting of the lemma with a smaller total number of fiber components.

To fix the first assertion of the claim, we let  $F = \pi^*(z) = \sum_{i=1}^n m_i F_i$ ,  $z \in Z$ . Since  $F$  is degenerate and non-multiple we have  $n \geq 2$ . Since  $F^2 = 0$ , by the adjunction formula we obtain

$$-K_V \cdot F = 2 - 2\pi_a(F) = 2 \quad \text{and} \quad -K_V \cdot F_i = F_i^2 + 2 - 2\pi_a(F_i) = F_i^2 + 2, \quad i = 1, \dots, n.$$

Hence

$$\sum_{i=1}^n m_i (F_i^2 + 2) = 2.$$

Since  $F_i^2 \leq -1 \forall i$ , the positive summands correspond exactly to the  $(-1)$ -components of  $F$ . If  $F_i$  is a unique such component, then necessarily  $m_i > 1$ . Now the assertion follows.  $\square$

**Notation 7.4.** Let  $\Phi: \bar{X} \rightarrow \bar{X}_0$  be a birational morphism, which contracts all degenerate  $\bar{\mu}$ -fibers to non-degenerate ones yielding a  $\mathbb{P}^1$ -fiber bundle  $\bar{\mu}_0: \bar{X}_0 \rightarrow \bar{B}$ . Then  $\Phi$  can be decomposed into a sequence of blowups of smooth points

$$(13) \quad \Phi: \bar{X} = \bar{X}_m \xrightarrow{\sigma_m} \bar{X}_{m-1} \xrightarrow{\sigma_{m-1}} \dots \xrightarrow{\sigma_1} \bar{X}_0,$$

where  $\sigma_i$  contracts the component  $T_i \subset \bar{X}_i$  of the image  $\sigma_{i+1} \circ \dots \circ \sigma_m(\mathcal{T}) \subset \bar{X}_i$  to a point  $p_i \in \bar{X}_{i-1}$ . The proper transforms of the curves  $T_i$  ( $i < j$ ),  $S$ , and  $F = \bar{\mu}^{-1}(\bar{B} \setminus B)$  on the surfaces  $\bar{X}_j$  will be denoted by the same letters. The  $\mathbb{P}^1$ -fibration  $\bar{\mu}: \bar{X} \rightarrow \bar{B}$  induces  $\mathbb{P}^1$ -fibrations  $\bar{\mu}_i: \bar{X}_i \rightarrow \bar{B}$  so that  $\sigma_i: \bar{X}_i \rightarrow \bar{X}_{i-1}$  becomes a morphism of  $\mathbb{P}^1$ -fibrations identical on  $\bar{B}$ ,  $i = 1, \dots, m$ . By Lemma 7.3 we may assume that  $T_1, \dots, T_m$  do not meet  $S$ , and so, the centers of blowups  $p_1, \dots, p_m$  do not belong to  $S$  or its images. Thus,  $\Phi(T_1 \cup \dots \cup T_m)$  is a finite subset of the smooth affine surface  $X_0 := \bar{X}_0 \setminus (S \cup F)$ . The induced  $\mathbb{A}^1$ -fibration  $\mu_0 = \bar{\mu}_0|_{X_0}: X_0 \rightarrow B$  is a locally trivial bundle with fiber  $\mathbb{A}^1$ .

**Lemma 7.5.**  $\mu_0: X_0 \rightarrow B$  admits a structure of a line bundle.

*Proof.* The ruling  $\bar{\mu}_0: \bar{X}_0 \rightarrow \bar{B}$  is the projectivization of a rank 2 vector bundle  $V \rightarrow \bar{B}$ . The section  $S$  of  $\bar{\pi}$  corresponds to a line subbundle  $L \subset V$ . The exact sequence of vector bundles over the affine curve  $B$ ,

$$0 \rightarrow L|_B \rightarrow V|_B \rightarrow (V/L)|_B \rightarrow 0$$

splits. Indeed, the obstacle to splitting sits in the group

$$\text{Ext}^1(L, V/L) \cong H^1(B, \mathcal{H}om_{\mathcal{O}_B}(\mathcal{O}_B(L), \mathcal{O}_B(V/L))) \cong H^1(B, \mathcal{O}_B(L)^\vee \otimes_{\mathcal{O}_B} \mathcal{O}_B(V/L))$$

that vanishes due to Serre's Vanishing Theorem. This provides a section of  $V|_B \rightarrow B$  disjoint with  $S$ , which can be taken for the zero section of a line bundle  $\mu_0: X_0 \rightarrow B$ .  $\square$

**Definition 7.6.** The blowup  $\sigma_i$  is called *inner* if  $p_i$  is a node of  $\mathcal{T}^{(i-1)}$ , and *outer* otherwise. The corresponding component  $T_i$  is also called *inner* or *outer*, respectively.

**Notation 7.7.** Fix a special fiber, say,  $\mathcal{T}' = \bar{\mu}^{-1}(\beta')$ , where  $\beta' \in B$ . Since the blowups with centers in different fibers commute, with a suitable enumeration we may assume that

$$\mathcal{T}' = T_0 \cup T_1 \cup \dots \cup T_{m'},$$

where  $m' \leq m$  and  $T_0$  is the proper transform of  $\bar{\mu}_0^{-1}(\beta') \subset \bar{X}_0$  and the only component of  $\mathcal{T}'$  meeting  $S$ . We denote by  $\mathcal{T}^{(i)}$  the image of  $\mathcal{T}'$  in  $\bar{X}_i$ .

**Definition 7.8.** Let  $T_i, T_k$  ( $k < i$ ) be two components of  $\mathcal{T}'$  such that  $p_i = \sigma_i(T_i) \in T_k$ , see 7.4. We say that  $T_k$  is the *parent* of  $T_i$  if either  $T_i$  is outer, or  $T_i$  is inner and  $p_i \in T_k \cap T_j$  on  $\bar{X}_{i-1}$  for some  $j < k$ . Every component  $T_i$ , where  $i > 0$ , has exactly one parent. <sup>14</sup>

## 7.2. Formal neighborhoods and coordinate charts.

**Definition 7.9.** Given an algebraic variety  $Y$  and a closed subset  $Z \subset Y$ , we denote by  $\hat{\mathcal{O}}_{Y,Z}$  the completion of the local sheaf  $\mathcal{O}_{Y,Z}$  with respect to the filtration by powers of the ideal sheaf  $\mathcal{I}$  of  $Z$ . The corresponding formal scheme  $\text{Spf } \hat{\mathcal{O}}_{Y,Z}$  is called a *formal neighborhood* of  $Z$  in  $Y$ , see, e.g., [3].

**Notation 7.10.** Given a surface  $V$  and a local coordinate chart  $(x_p, y_p)$  on  $V$  centered at a smooth point  $p \in V$  <sup>15</sup>, we identify the completion  $\hat{\mathcal{O}}_{V,p}$  of the local ring  $\mathcal{O}_{V,p}$  with the ring  $\mathbb{K}[[x_p, y_p]]$ . Let  $\sigma: \tilde{V} \rightarrow V$  be the blowup of  $p$  with exceptional  $(-1)$ -curve  $E$ . The rational function  $y_E = y_p/x_p$  defines an isomorphism  $y_E: E \xrightarrow{\cong} \mathbb{P}^1$ . For

<sup>14</sup>The notion of *parent* differs from the notion of *mother component* in 5.16.

<sup>15</sup>Such a coordinate chart  $(x_p, y_p)$  on  $V$  can be defined as follows. Consider an affine neighborhood  $X$  of  $p$  in  $V$ , a closed embedding  $X \hookrightarrow \mathbb{A}^N$ , and a linear projection  $\pi: \mathbb{A}^N \rightarrow T_p X$ . Let  $U, V$  be linear functions on  $\mathbb{A}^N$  that restrict to coordinates in the tangent plane  $T_p X$ . Then we let  $x_p = U|_X$  and  $y_p = V|_X$ .



each  $q \in \mathbb{P}^1 \cong \mathbb{A}^1 \cup \{\infty\}$  we let  $E(q) = y_E^{-1}(q) \in E$ . The sheaf  $\hat{\mathcal{O}}_{\tilde{V}, E}$  inherits coordinates of  $\hat{\mathcal{O}}_{V, p}$  as follows:

$$\begin{aligned}\hat{\mathcal{O}}_{\tilde{V}, E}(\{y_E \neq 0\}) &= \hat{\mathcal{O}}_{V, p} \left[ \frac{1}{y_E} \right] = \mathbb{K}[1/y_E][[y_p]], \\ \hat{\mathcal{O}}_{\tilde{V}, E}(\{y_E \neq \infty\}) &= \hat{\mathcal{O}}_{V, p}[y_E] = \mathbb{K}[y_E][[x_p]].\end{aligned}$$

For a point  $E(q) \in E$  the local coordinate chart centered at  $E(q)$  is given by

$$(14) \quad (x_{E(q)}, y_{E(q)}) = \begin{cases} (x_p, y_E - q), & q \neq \infty, \\ \left( \frac{1}{y_E}, y_p \right), & q = \infty, \end{cases}$$

thus in the other sense we have

$$(15) \quad (x_p, y_p) = \begin{cases} (x_{E(q)}, x_{E(q)}(y_{E(q)} + q)), & q \neq \infty, \\ (x_{E(q)}y_{E(q)}, y_{E(q)}), & q = \infty. \end{cases}$$

Letting  $x_E = x_p$  we call  $(x_E, y_E)$  the *local coordinates near E*.

**Notation 7.11.** We write  $p' \geq p$  if  $p' \in \mathcal{T}^{(i+k)}$  and  $\sigma_{i+1} \circ \dots \circ \sigma_{i+k}(p') = p \in \mathcal{T}^{(i)}$  for some  $k \geq 0$  (that is,  $p'$  is an infinitesimally near point of  $p$ ). If  $p$  is the center of a blowup  $\sigma_{i+j}$ ,  $1 \leq j \leq k$ , then we write  $p' > p$ . Otherwise, by abuse of notation, we write  $p = p'$ .

**Notation 7.12.** Tensoring with the ring  $\hat{\mathcal{O}}_{B, \beta'} = \mathbb{K}[[t]]$ , where  $t$  is a local coordinate on  $B$  centered at  $\beta'$ , we restrict  $\mu_0$  to the formal neighborhood of the fiber  $T_0^* := T_0 \setminus S = \mu_0^{-1}(\beta') \cong \mathbb{A}^1$  in  $X_0$ , namely, to

$$\hat{\mathcal{O}}_{X_0, T_0^*}(T_0^*) = \mathbb{K}[y_0][[x_0]],$$

where the coordinates  $(x_0, y_0)$  near the fiber  $T_0^* \subset X_0$  are chosen so that  $x_0 = \mu_0^*(t)$ , and  $y_0 = 0$  defines the zero section of  $\mu_0: X_0 \rightarrow B$ . Regarding  $y_0$  as a  $\mathbb{P}^1$ -coordinate on  $T_0 \subset \bar{X}_0$ , we define local coordinates at each point  $T_0(q)$  of  $T_0^* = T_0 \cap X_0$  via (14).

For every  $i = 1, \dots, m'$  we define recursively local coordinates at the points of the fiber  $\mathcal{T}^{(i)} \subset \bar{X}_i$  as follows. Assume that the local coordinates  $(x_{p_i}, y_{p_i})$  centered at a point  $p_i \in \bar{X}_{i-1}$  are already defined. As in 7.10 we infer first a  $\mathbb{P}^1$ -coordinate  $y_i = y_{T_i}$  on  $T_i$  and then local coordinates near  $T_i(q)$  for each  $q \in \mathbb{P}^1$ . For any point  $p \in \mathcal{T}^{(i)} \setminus (T_i \cup S)$  we keep the same local chart  $(x_p, y_p) = (x_{\sigma_i(p)}, y_{\sigma_i(p)})$  as on the surface  $X_{i-1}$ . In more detail, each point  $p \in \mathcal{T}^{(i)}$  admits a unique representation of the form  $p = T_j(q)$  as in Notation 7.11, where  $j \leq i$  and  $q \in \mathbb{P}^1$ . In particular, if  $p = T_j \cap T_k$  with  $j > k$ , then  $p = T_j(q)$ , where  $q \in \mathbb{P}^1$ , and  $p$  cannot be represented as  $T_k(q')$  for  $q' \in \mathbb{P}^1$ . Then we let  $(x_p, y_p) = (x_{T_j(q)}, y_{T_j(q)})$ .

**Remark 7.13.** If  $T_j \cap T_k = \{p\} \subset \bar{X}_i$ , then  $T_j$  and  $T_k$  are the coordinate lines in the local coordinates  $(x_p, y_p)$  near  $p$ . Up to permuting  $j$  and  $k$  one has  $p \geq T_j(\infty)$ ,  $p \geq T_k(q)$ , where  $q \neq \infty$ , and

$$(x_p, y_p) = \left( \frac{1}{y_j}, y_k - q \right).$$

Note that  $T_j(\infty)$  is the point of  $T_j$  closest to  $S$ , and the component  $T_k$  separates  $T_j$  and  $S$  in  $\mathcal{T}^{(i)}$ .

**7.3. Arcs and multiplicities.** In this subsection we introduce the arc spaces of an  $\mathbb{A}^1$ -fibration (see 7.14, 7.10) and the multiplicities of arcs (see 7.18 and 7.19).

**Definition 7.14** (*arc space*; see, e.g., [29, 54]). Given a variety  $V$ , an *arc* in  $V$  is a parameterized formal curve germ  $\text{Spec } \mathbb{K}[[t]] \rightarrow V$ . The *arc space*  $\text{Arc}(V)$  is the  $\mathbb{K}[[t]]$ -scheme consisting of all the  $\mathbb{K}[[t]]$ -rational points of  $V$ . Given a closed subvariety  $Z \subset V$ , we let  $\text{Arc}(V)_Z$  denote the subscheme of arcs centered in  $Z$ .

**Example 7.15.** Let  $V$  be a surface and  $p \in V$  be a smooth point with a local coordinate chart  $(x_p, y_p)$  centered at  $p$ . The corresponding arc space is

$$\begin{aligned} \text{Arc}(V)_p &= \{h: \hat{\mathcal{O}}_{V,p} \rightarrow \mathbb{K}[[t]] \mid h(\mathfrak{m}_{V,p}) \subseteq t\mathbb{K}[[t]]\} \\ &= \{(x_p, y_p) \mapsto (x(t), y(t)) \mid x, y \in t\mathbb{K}[[t]]\}. \end{aligned}$$

If  $\sigma: \tilde{V} \rightarrow V$  is the blowup of  $p$  with exceptional  $(-1)$ -curve  $E$ , then  $\sigma$  induces an isomorphism  $\text{Arc}(\tilde{V})_E \setminus \text{Arc}(E) \cong \text{Arc}(V)_p \setminus \{0\}$ .

**Notation 7.16.** For a point  $p \in \mathcal{T}^{(i)} \setminus S \subset \bar{X}_i \setminus S$  ( $i \in \{0, \dots, m'\}$ ) we consider the subset  $\text{Arc}(\bar{X}_i)_p^* \subset \text{Arc}(\bar{X}_i)_p$  of all arcs in  $\bar{X}_i$  centered at  $p$  whose generic point does not belong to  $\mathcal{T}'$ . The map  $\text{Arc}(\bar{X}_i)_p^* \rightarrow \text{Arc}(X_0)_{T_0}^* \subset \text{Arc}(\bar{X}_0)_{T_0}^*$  induced by  $\sigma_1 \circ \dots \circ \sigma_i: \bar{X}_i \rightarrow \bar{X}_0$  is injective. This allows to identify  $\text{Arc}(\bar{X}_i)_p^*$  with its image in  $\text{Arc}(X_0)_{T_0}^*$ , where

$$\begin{aligned} \text{Arc}(X_0)_{T_0}^* &= \text{Arc}(X_0)_{T_0} \setminus \text{Arc}(T_0) \\ &= \{(x_0, y_0) \mapsto (x(t), y(t)) \mid x \in t\mathbb{K}[[t]] \setminus \{0\}, y \in \mathbb{K}[[t]]\}. \end{aligned}$$

**Notation 7.17.** Let  $\text{mult } T_i = m_i$  be the multiplicity of  $T_i$  in the special fiber  $\bar{\mu}_{m'}^*(\beta') = \sum_{i=0}^{m'} m_i T_i$ . Thus,  $\text{mult } T_i = \text{mult } T_j$  if  $T_i$  is outer and  $\sigma_i(T_i) = p_i \in T_j$ , and  $\text{mult } T_i = \text{mult } T_j + \text{mult } T_k$  if  $T_i$  is inner and  $p_i \in T_j \cap T_k$  (see 7.7). In particular,  $\text{mult } T_i$  is the same on every surface  $\bar{X}_j$ ,  $j \geq i$ .

**Definition 7.18** (*multiplicity of an arc*). For an arc  $h \in \text{Arc}(\bar{X}_0)_{T_0}^*$ ,  $(x_0, y_0) \mapsto (x(t), y(t))$ , we define its multiplicity by  $\text{mult } h = \text{ord}_t x$  ( $= \text{ord}_t(\bar{\mu}_0)_* h$ ). The multiplicity of an arc  $h \in \text{Arc}(\bar{X}_i)_{\mathcal{T}^{(i)}}^*$  is defined as the multiplicity of the image of  $h$  in  $\text{Arc}(\bar{X}_0)_{T_0}^*$ .

**Proposition 7.19.** Let  $p \in \mathcal{T}^{(i)}$  and  $h \in \text{Arc}(\bar{X}_i)_p^*$ ,  $h: (x_p, y_p) \mapsto (x(t), y(t))$ .<sup>16</sup> If  $p \notin \text{Sing } \mathcal{T}^{(i)}$  is a point of  $T_j$ , then

$$(16) \quad \text{mult } h = \text{mult}(T_j) \text{ord}_t x.$$

If  $p \in \text{Sing } \mathcal{T}^{(i)}$  is an intersection of components  $T_j$  and  $T_k$  with  $\sigma_{k+1} \circ \dots \circ \sigma_i(p) = T_k(\infty)$  (cf. 7.13), then

$$(17) \quad \text{mult } h = \text{mult}(T_j) \text{ord}_t x + \text{mult}(T_k) \text{ord}_t y.$$

*Proof.* We proceed by induction on  $i$ . The case  $i = 0$  is trivial. Assume that the assertion holds on  $\bar{X}_{i-1}$ . Then it also holds on  $\mathcal{T}^{(i)} \setminus T_i$ . Let then  $p = T_i(q)$ ,  $q \in \mathbb{P}^1$ , so  $\sigma_i(p) = p_i$ . We distinguish the following five cases:

- $T_i$  is outer and  $q = \infty$ ;
- $T_i$  is outer and  $q \neq \infty$ ;
- $T_i$  is inner and  $q = \infty$ ;
- $T_i$  is inner and  $q = 0$ ;
- $T_i$  is inner and  $q \neq 0, \infty$ .

These are depicted below in Figure 1 for  $T_i$  outer and in Figure 2 for  $T_i$  inner. All these cases are treated similarly, so we consider just two of them and leave the others to the reader.

Let first  $T_i$  be outer with a parent  $T_j$ , where  $p = T_i(q)$ ,  $q \neq \infty$ . Then  $(x_{p_i}, y_{p_i}) = (x_p, x_p(y_p + q))$  by (15), so  $h: (x_{p_i}, y_{p_i}) \mapsto (x(t), x(t)(q + y(t)))$  on  $X_{i-1}$ , and by the induction conjecture

$$\text{mult } h = \text{mult}(T_j) \text{ord}_t x = \text{mult}(T_i) \text{ord}_t x.$$

<sup>16</sup>Since  $h \in \text{Arc}(\bar{X}_i)_p^*$ ,  $x(t) \neq 0$  in the former case and  $x(t), y(t) \neq 0$  in the latter one, thus the formulas (16)–(17) are well defined.




FIGURE 1.  $T_i$  is outer.




FIGURE 2.  $T_i$  is inner.

Let further  $T_i$  be inner and  $p = T_i(\infty)$ . Then  $h: (x_{p_i}, y_{p_i}) \mapsto (x(t)y(t), y(t))$  on  $X_{i-1}$  by (15), and so, by the inductive conjecture,

$$\begin{aligned} \text{mult } h &= \text{mult}(T_j)(\text{ord}_t x + \text{ord}_t y) + \text{mult}(T_s) \text{ord}_t y = \text{mult}(T_j) \text{ord}_t x + \\ &(\text{mult}(T_j) + \text{mult}(T_s)) \text{ord}_t y = \text{mult}(T_j) \text{ord}_t x + \text{mult}(T_i) \text{ord}_t y. \end{aligned}$$

□

**7.4. Puiseux arc spaces.** In this subsection we introduce and study Puiseux arc spaces.

**7.20 (Puiseux arcs).** Fix a point<sup>17</sup>  $p \in \mathcal{T}^{(i)} \setminus S$ .

1. An *invertible substitution* is a change of variable  $t \mapsto A(t)$ , where  $\text{ord}_t A = 1$ . These form a group acting on the arc space  $\text{Arc}(\bar{X}_i)_p^*$  via

$$A(t).h: (x_p, y_p) \mapsto (x_h(A(t)), y_h(A(t))),$$

where  $h: (x_p, y_p) \mapsto (x_h(t), y_h(t))$ . We call two arcs *equivalent* if they belong to the same orbit of the action.

2. The coordinate line  $x_p = 0$  in a local chart  $(x_p, y_p)$  is a part of a component of  $\mathcal{T}^{(i)}$ . Hence  $x(t) \neq 0$  for any arc  $h \in \text{Arc}(\bar{X}_i)_p^*$ ,  $h: (x_p, y_p) \mapsto (x(t), y(t))$ . So, a suitable invertible substitution  $A(t)$  sends  $h$  to an arc  $\tilde{h} = A(t).h: (x_p, y_p) \mapsto (t^n, \tilde{y}(t))$ , where  $n = \text{ord}_t x$  and  $\text{ord}_t \tilde{y} = \text{ord}_t y$ . Such a change of variable  $t \mapsto A(t)$  and an arc  $\tilde{h}$  are defined uniquely up to a composition with  $t \mapsto \alpha \cdot t$ , where  $\alpha \in \mathbb{K}$ ,  $\alpha^n = 1$ . All the arcs equivalent to  $\tilde{h}$  enjoy the same Puiseux expansion  $y = \tilde{y}(x^{1/n})$ .
3. An arc  $\tilde{h}: (x_p, y_p) \mapsto (t^n, \tilde{y}(t))$  with  $\tilde{y} = \sum_{i=\text{ord}_t y}^{\infty} a_i t^i$  is called a *Puiseux arc* if

$$\gcd(\{i \mid a_i \neq 0\} \cup \{n\}) = 1.$$

**Definition 7.21 (Puiseux arc space).** Consider a point  $p \in \mathcal{T}^{(i)} \setminus S$ , a pair of positive integers  $n, d \in \mathbb{Z}_{>0}$ , and a polynomial  $\psi = \sum_{i=1}^{d-1} \psi_i t^i \in t\mathbb{K}[t]$  such that

$$(18) \quad \gcd(\{i \mid \psi_i \neq 0\} \cup \{n\}) = 1,$$

that is, such that  $(x_p, y_p) \mapsto (t^n, \psi(t))$  is a Puiseux arc. Assume also that  $\psi \neq 0$  if  $p$  is a node. The *Puiseux arc space*  $W = \text{Pui}_{\bar{X}_i, p}(\psi, n, d)$  on  $\bar{X}_i$  relative to the coordinate system  $(x_p, y_p)$  centered at  $p$  consists of all arcs equivalent to Puiseux arcs of the form  $h: (x_p, y_p) \mapsto (t^n, y(t))$  with  $y \in \psi + t^d \mathbb{K}[[t]]$ . In other words,

$$W = \{(x_p, y_p) \mapsto (A(t)^n, y(A(t))) \mid \text{ord}_t A = 1, y \in \psi + t^d \mathbb{K}[[t]]\}.$$

Thus, the elements of  $W$  share the same starting piece  $\psi$  of the Puiseux expansion. The condition that  $\psi \neq 0$  if  $p$  is a node ensures that  $W \subset \text{Arc}(\bar{X}_i)_p^*$ . Furthermore, the elements of  $W$  share the same multiplicity (see 7.18), which we denote by  $\text{mult } W$ .

<sup>17</sup>Warning: in what follows we never consider the points ‘at infinity’  $p \in \mathcal{T}^{(i)} \cap S$ .

Any Puiseux arc space can be expressed in the coordinates  $(x_0, y_0)$  on  $X_0$ , see 7.16. To this end, given a pair of positive integers  $n, d \in \mathbb{Z}_{>0}$  and a polynomial  $\psi = \sum_{i=0}^{d-1} \psi_i t^i \in \mathbb{K}[t]$  with a possibly nonzero constant term, we let

$$\text{Pui}(\psi, n, d) = \text{Pui}_{\bar{X}_0, T_0(\psi_0)}(\psi - \psi_0, n, d).$$

Thus, the constant term  $\psi_0$  is responsible for the choice of the center  $T_0(\psi_0) \in T_0 \setminus S$ .

**Lemma 7.22.** *For any point  $p = T_i(q) \in \mathcal{T}^{(i)} \setminus S$ ,  $i \in \{0, \dots, m'\}$ , and any Puiseux arc space  $W = \text{Pui}_{\bar{X}_i, p}(\psi, n, d) \subset \text{Arc}(\bar{X}_i)_p^*$ , the image  $W'$  of  $W$  under the embedding  $\text{Arc}(\bar{X}_i)_p^* \hookrightarrow \text{Arc}(\bar{X}_{i-1})_{p_i}^*$  induced by  $\sigma_i: \bar{X}_i \rightarrow \bar{X}_{i-1}$  is a Puiseux arc space in  $\text{Arc}(\bar{X}_{i-1})_{p_i}^*$ . More precisely,*

$$(19) \quad W' = \text{Pui}_{\bar{X}_{i-1}, p_i}(t^n(q + \psi), n, d + n) \quad \text{if } q \neq \infty,$$

$$(20) \quad W' = \text{Pui}_{\bar{X}_{i-1}, p_i}(\tilde{\psi}, n + \text{ord } \psi, d), \quad \text{if } q = \infty,$$

where  $\text{ord } \tilde{\psi} = \text{ord } \psi$ .

*Proof.* According to Definition 7.21 one has

$$W = \{(x_p, y_p) \mapsto (A(t)^n, \psi(A(t)) + \eta(A(t))) \mid \text{ord}_t A = 1, \eta \in t^d \mathbb{K}[[t]]\}.$$

Suppose first that  $p = T_i(q)$ , where  $q \neq \infty$ . By (14) we have  $(x_{p_i}, y_{p_i}) = (x_p, x_p(y_p + q))$ . Hence

$$W' = \{(x_{p_i}, y_{p_i}) \mapsto (A(t)^n, y(A(t))) \mid \text{ord}_t A = 1, y \in t^n(q + \psi) + t^{d+n} \mathbb{K}[[t]]\}$$

is again a Puiseux arc space relative to the coordinate system  $(x_{p_i}, y_{p_i})$  in  $\bar{X}_{i-1}$ . More precisely,  $W' = \text{Pui}_{\bar{X}_{i-1}, p_i}(\tilde{\psi}, n, d + n)$ , where  $\tilde{\psi}(t) = t^n(q + \psi(t)) \in \mathbb{K}[t]$  is a polynomial of degree  $< d + n$  satisfying (18).

Let further  $p = T_i(\infty)$ . Then one can write  $t = \sum_{i=1}^{\infty} \alpha_i s^i \in s \mathbb{K}[[s]]$ , where  $\alpha_1 \neq 0$  and  $t^n \psi(t) = s^{n+\text{ord } \psi}$ . Plugging in  $t = t(s)$  sends the set  $\psi(t) + t^d \mathbb{K}[[t]]$  into  $\tilde{\psi}(s) + s^d \mathbb{K}[[s]]$  for some polynomial  $\tilde{\psi} \in s \mathbb{K}[s]$  of degree  $< d$  and of order  $\text{ord } \tilde{\psi} = \text{ord } \psi$ . By (14) we have  $(x_{p_i}, y_{p_i}) = (x_p y_p, y_p)$ . Hence

$$\begin{aligned} W' &= \{(x_{p_i}, y_{p_i}) \mapsto (A(t)^n y(A(t)), y(A(t))) \mid \text{ord}_t A = 1, y \in \psi + t^d \mathbb{K}[[t]]\} \\ &= \{(x_{p_i}, y_{p_i}) \mapsto (\tilde{A}(s)^{n+\text{ord } \psi}, y(\tilde{A}(s))) \mid \text{ord}_s \tilde{A} = 1, y \in \tilde{\psi} + s^d \mathbb{K}[[s]]\} \end{aligned}$$

coincides with the Puiseux arc space  $\text{Pui}_{\bar{X}_{i-1}, p_i}(\tilde{\psi}, n + \text{ord } \psi, d)$  relative to the coordinate system  $(x_{p_i}, y_{p_i})$  in  $\bar{X}_{i-1}$ , where the polynomial  $\tilde{\psi} \in s \mathbb{K}[s]$  still satisfies (18). Indeed, otherwise one can write  $\tilde{\psi}(s) = \tilde{\varphi}(s^k)$ , where  $\tilde{\varphi} \in \mathbb{K}[s]$ ,  $k > 1$ , and  $k|n$ . However, plugging in the expression  $s = s(t) \in t \mathbb{K}[[t]]$  yields  $\psi(t) = \varphi(t^k)$  for some polynomial  $\varphi \in \mathbb{K}[t]$ . The latter contradicts condition (18) for  $\psi$ .  $\square$

The following corollary is straightforward.

**Corollary 7.23.** *Given a Puiseux arc space  $W = \text{Pui}_{\bar{X}_i, p}(\psi, n, d)$ , its image under the embedding  $\text{Arc}(\bar{X}_i)_p^* \hookrightarrow \text{Arc}(\bar{X}_0)_{T_0}^*$  is a Puiseux arc space, say  $\text{Pui}(\tilde{\psi}, \tilde{n}, \tilde{d})$ . In particular,  $\text{mult } W = \tilde{n}$ .*

**Notation 7.24.** Given a point  $p \in \mathcal{T}^{(i)} \setminus \text{Sing}(\mathcal{T}^{(i)} \cup S)$ , we let  $\text{Pui}(p)$  be the image of  $\text{Pui}_{\bar{X}_i, p}(0, 1, 1)$  in  $\text{Arc}(\bar{X}_0)_{T_0}^*$ . By the preceding corollary,  $\text{Pui}(p)$  is a Puiseux arc space.

**Corollary 7.25.** *Let  $\text{Pui}(p) = \text{Pui}(\psi, n, d)$ , where  $p \in T_j \setminus S$  is not a node of  $\mathcal{T}^{(i)}$ . Then  $\text{mult } \text{Pui}(p) = n = \text{mult } T_j$ . Furthermore,  $n = 1$  if and only if  $T_j$  is obtained via a sequence of outer blowups.*

*Proof.* The first assertion follows from Corollary 7.23 and (16). The second follows from the first due to the fact that  $\text{mult}(T_j) = 1$  if and only if the component  $T_j$  is obtained via a sequence of outer blowups.  $\square$

**Remark 7.26.** For each Puiseux space  $\text{Pui}(\psi, n, d) \subset \text{Arc}(X_0)$  centered at a point  $p \in T_0^*$  there exists a surface  $X'$  and a sequence of blowups  $X' \rightarrow \bar{X}_0$  with centers at infinitesimally near points of  $p$  such that  $\text{Pui}(\psi, n, d) = \text{Pui}(p')$  for some point  $p' \in X'$ ,  $p' \geq p$ . So, there is a one-to-one correspondence between infinitesimally near points of  $T_0^* \subset X_0$ , and the Puiseux arc spaces.

**7.5. Stabilizer of a special fiber.** In this subsection we study the action of the automorphism group of an  $\mathbb{A}^1$ -fibration on the Puiseux arc spaces of a special fiber.

**Notation 7.27.** Consider the following groups of automorphisms of the arc space  $\text{Arc}(\bar{X}_0)_{T_0}^*$ :

$$\begin{aligned} \mathcal{H} &= \left\{ (x, y) \mapsto (ax, Q(x)y) \mid a \in \mathbb{K}^\times, Q = \sum_{i=0}^{\infty} b_i x^i \in \mathbb{K}[[x]]^\times \right\}, \\ \mathbb{T} &= \left\{ (x, y) \mapsto (ax, by) \mid (a, b) \in (\mathbb{K}^\times)^2 \right\} \cong \mathbb{G}_m^2, \\ \mathbb{G}_a(i) &= \left\{ (x, y) \mapsto (x, y + c_i x^i) \mid c_i \in \mathbb{K} \right\} \cong \mathbb{G}_a. \end{aligned}$$

We have

$$\begin{aligned} \text{Aut Arc}(\bar{X}_0)_{T_0}^* &= \\ &= \left\{ (x, y) \mapsto (ax, Q(x)y + P(x)) \mid a \in \mathbb{K}^\times, P = \sum_{i=0}^{\infty} c_i x^i \in \mathbb{K}[[x]], Q = \sum_{i=0}^{\infty} b_i x^i \in \mathbb{K}[[x]]^\times \right\} \\ &\cong \left( \prod_{i=0}^{\infty} \mathbb{G}_a(i) \right) \rtimes \mathcal{H}. \end{aligned}$$

We let  $\text{Stab}_{\text{fps}} W \subset \text{Aut Arc}(\bar{X}_0)_{T_0}^*$  be the stabilizer of a subset  $W \subset \text{Arc}(\bar{X}_0)_{T_0}^*$ , where ‘fps’ stands for ‘formal power series’. Attributing the lower index  $B$  to a group of automorphisms means passing to the subgroup of automorphisms that act trivially on the first coordinate, that is, verify  $a = 1$  in the notation as above. In particular, we consider the subgroups  $\text{Aut}_B \text{Arc}(\bar{X}_0)_{T_0}^* \subset \text{Aut Arc}(\bar{X}_0)_{T_0}^*$ ,  $\mathcal{H}_B \subset \mathcal{H}$ , and the one-dimensional torus  $\mathbb{T}_B \subset \mathbb{T}$ .

**Notation 7.28.** Given a Puiseux arc space  $W = \text{Pui}(\psi, n, d)$ , we can decompose

$$(21) \quad \psi(t) = \psi^{\text{reg}}(t^n) + \psi^{\text{sing}}(t),$$

where  $\psi^{\text{sing}} \in \mathbb{K}[t]$  is the sum of all monomials in  $\psi$  with exponents not divided by  $n$ . We let also  $\lceil \frac{d}{n} \rceil$  stand for the smallest integer  $\geq \frac{d}{n}$ .

**Lemma 7.29.** Consider a Puiseux arc space  $W = \text{Pui}(\psi, n, d) \subset \text{Arc}(\bar{X}_0)_{T_0}^*$ . An automorphism  $g \in \text{Aut Arc}(\bar{X}_0)_{T_0}^*$ ,  $g: (x, y) \mapsto (ax, Q(x)y + P(x))$ , stabilizes  $W$  if and only if

$$(22) \quad P(x) = \psi^{\text{reg}}(ax) - \psi^{\text{reg}}(x)Q(x) \quad \text{mod } x^{\lceil \frac{d}{n} \rceil} \mathbb{K}[[x]],$$

and in the case  $n > 1$  also

$$(23) \quad \begin{cases} Q(s^n) \psi^{\text{sing}}(s) = \psi^{\text{sing}}(\alpha s) & \text{mod } s^d \mathbb{K}[[s]] \\ \alpha^n = a \end{cases}$$

for some  $\alpha \in \mathbb{K}$ .

*Proof.* Consider an arc  $h \in W$ ,

$$h: (x_0, y_0) \mapsto (A(t)^n, y(A(t))) \quad \text{with} \quad A \in t\mathbb{K}[[t]] \setminus \{0\} \quad \text{and} \quad y \in \psi + t^d\mathbb{K}[[t]].$$

The automorphism  $g$  sends  $h$  to the arc

$$g.h: (x_0, y_0) \mapsto (a \cdot A(t)^n, Q(A(t)^n) \cdot y(A(t)) + P(A(t)^n)).$$

Hence  $g.h \in W$  if and only if

$$\begin{cases} a \cdot A(t)^n = \tilde{A}(t)^n \\ Q(A(t)^n) \cdot y(A(t)) + P(A(t)^n) = \tilde{y}(\tilde{A}(t)) \end{cases}$$

for some  $\tilde{A}(t) \in t\mathbb{K}[[t]]$  and  $\tilde{y} \in \psi + t^d\mathbb{K}[[t]]$ . The first equation means that  $\tilde{A}(t) = \alpha \cdot A(t)$  for an  $n$ th root  $\alpha$  of  $a$ . Letting  $s = A(t)$ , the second equation holds for some  $\tilde{y} \in \psi + t^d\mathbb{K}[[t]]$  if and only if, with this root  $\alpha$ ,

$$(24) \quad Q(s^n) \cdot \psi(s) + P(s^n) - \psi(\alpha s) \in s^d\mathbb{K}[[s]].$$

Splitting  $\psi$  as in (21) leads to equations (22) and (23).  $\square$

**Proposition 7.30.** *Consider a finite collection of Puiseux arc spaces  $W_k = \text{Pui}(\psi_k, n_k, d_k)$ ,  $k = 1, \dots, r$ , ordered so that  $\frac{d_1}{n_1} \geq \dots \geq \frac{d_r}{n_r}$ . Let  $N = \lceil \frac{d_1}{n_1} \rceil$ . Then*

$$(25) \quad \bigcap_{k=1}^r \text{Stab}_{\text{fps}} \text{Pui}(\psi_k, n_k, d_k) = \left( \prod_{i=N}^{\infty} \mathbb{G}_a(i) \right) \rtimes \bar{\mathcal{H}}^h \subset \text{Aut Arc}(\bar{X}_0)_{T_0}^*$$

for a subgroup  $\bar{\mathcal{H}} \subset \mathcal{H}$ , where  $\bar{\mathcal{H}}^h = h \circ \bar{\mathcal{H}} \circ h^{-1}$  with  $h = \psi_1^{\text{reg}} \in \bigoplus_{i=0}^{N-1} \mathbb{G}_a(i)$ .

*Proof.* In the system of equations (22)–(23) for all  $W_k$ ,  $k = 1, \dots, r$ , we can eliminate  $P(x)$  for  $k = 2, \dots, r$ . This yields the system

$$(26) \quad \begin{cases} P(x) = \psi_1^{\text{reg}}(ax) - \psi_1^{\text{reg}}(x)Q(x) & \text{mod } x^N\mathbb{K}[[x]], \\ \psi_k^{\text{reg}}(ax) - \psi_1^{\text{reg}}(ax) = (\psi_k^{\text{reg}}(x) - \psi_1^{\text{reg}}(x))Q(x) & \text{mod } x^{\lceil \frac{d_k}{n_k} \rceil}\mathbb{K}[[x]], \quad k = 2, \dots, r, \\ Q(s^n)\psi_k^{\text{sing}}(s) = \psi_k^{\text{sing}}(\alpha_k s) & \text{mod } s^{d_k}\mathbb{K}[[s]], \quad k = 1, \dots, r, \\ \alpha_k^{n_k} = a & k = 1, \dots, r. \end{cases}$$

The first equation expresses  $P(x)$  in terms of  $a$  and  $Q(x)$ . This defines the subgroup

$$(27) \quad \left\{ (x, y) \mapsto (ax, Q(x)y + \psi_1^{\text{reg}}(ax) - \psi_1^{\text{reg}}(x)Q(x) + \sum_{i=N}^{\infty} c_i x_i) \right\} \\ = \left( \prod_{i=N}^{\infty} \mathbb{G}_a(i) \right) \rtimes (h \circ \mathcal{H} \circ h^{-1}),$$

where  $h = \psi_1^{\text{reg}}$ . The remaining equations define a subgroup  $\bar{\mathcal{H}} \subset \mathcal{H}$ .  $\square$

**Notation 7.31.** Given a special fiber  $\mathcal{T}' = \bar{\mu}^{-1}(\beta') \subset \bar{X}$ , we let

$$(28) \quad \text{Stab}_{\text{fps}} \mathcal{T}' = \bigcap_{k: \sigma_k \text{ is outer}} \text{Stab}_{\text{fps}}(\text{Pui}(p_k)) \subset \text{Aut Arc}(\bar{X}_0)_{T_0}^*,$$

where  $p_k = \sigma_k(T_k) \in \bar{X}_{k-1} \setminus S$ . Taking in Proposition 7.30 a suitably reordered collection

$$(W_k = \text{Pui}(p_k) \mid \sigma_k \text{ is outer}),$$

we let  $N = N(\mathcal{T}')$ ,  $\bar{\mathcal{H}} = \mathcal{H}(\mathcal{T}')$ , and  $h = h(\mathcal{T}')$  denote the corresponding objects provided by this proposition, so that

$$(29) \quad \text{Stab}_{\text{fps}} \mathcal{T}' = \left( \prod_{i=N}^{\infty} \mathbb{G}_a(i) \right) \rtimes \bar{\mathcal{H}}^h.$$

We let also  $\mathbb{T}(\mathcal{T}') = \mathbb{T} \cap \mathcal{H}(\mathcal{T}')$ .

**Corollary 7.32.** *Given a special fiber  $\mathcal{T}' = \mu^{-1}(\beta') \subset \bar{X}$ , the following conditions are equivalent:*

- $\mathcal{H}(\mathcal{T}') = \mathcal{H}$ ,
- $\mathcal{H}(\mathcal{T}') \supset \mathbb{T}_B$ ,
- the dual graph  $\Gamma_{\mathcal{T}'}$  is a linear chain.

*Proof.* We start with the following observation. Clearly, an inner component  $T_i$  of  $\mathcal{T}'$  which is not a parent is a  $(-1)$ -curve. After contraction of  $T_i$  we obtain a new special fiber, say  $\mathcal{T}''$ , where  $\mathcal{T}''$  and  $\mathcal{T}'$  are both linear or non-linear simultaneously, and  $\text{Stab}_{\text{fps}} \mathcal{T}'' = \text{Stab}_{\text{fps}} \mathcal{T}'$ , hence also  $\mathcal{H}(\mathcal{T}'') = \mathcal{H}(\mathcal{T}')$ . Thus, we may assume in the sequel that each inner component of  $\mathcal{T}'$  is a parent.

Assume first that there exists an inner component  $T_i$  of  $\mathcal{T}'$ . By the previous observation such a component  $T_i$  with a maximal value of  $i$  is a parent of an outer component. Hence in this case  $\Gamma_{\mathcal{T}'}$  is non-linear. Furthermore, being inner,  $T_i$  belongs to the preimage of  $T_k(\infty)$  for some  $T_k$ . Then by (20) the corresponding Puiseux arc space is of form  $\text{Pui}(\psi, n, d)$  with  $n > 1$ . So, the corresponding equations (23) are nontrivial. It follows that  $\dim \mathcal{H}(\mathcal{T}') \cap \mathbb{T} \leq 1$  and  $\mathcal{H}(\mathcal{T}') \cap \mathbb{T}_B$  is finite. Thus, all three conditions of the lemma fail.

Assume further that all components of  $\mathcal{T}'$  are outer. If  $\mathcal{T}'$  is linear, then  $T_{m'}$  is the only non-parent, all the other components of  $\mathcal{T}'$  being its successive parents. By (19) the corresponding Puiseux arc space is of form  $\text{Pui}(\psi, 1, d)$ , and  $\text{Stab}_{\text{fps}} \mathcal{T}' = \text{Stab}_{\text{fps}} \text{Pui}(\psi, 1, d)$ . In this case (26) contains just one equation (of form (22)), and so,  $\mathcal{H}(\mathcal{T}') = \mathcal{H} \supset \mathbb{T}_B$ . Thus, under this setup all three conditions of the lemma are fulfilled.

Finally, assume that  $\mathcal{T}'$  is non-linear. Then there is a component  $T_k \subset \mathcal{T}'$  which is a parent for at least two other components with centers, say,  $T_k(q_1)$  and  $T_k(q_2)$ . The corresponding Puiseux arc spaces are of form  $\text{Pui}(\psi + q_1 t^{d-1}, 1, d)$  and  $\text{Pui}(\psi + q_2 t^{d-1}, 1, d)$  for some  $q_1 \neq q_2$ , some  $d \in \mathbb{Z}_{>0}$ , and some polynomial  $\psi \in \mathbb{K}[t]$  of degree  $\leq d-2$ . Inspecting system (26), for  $i = d-1$  we obtain the equalities  $c_{d-1} = q_1(a^{d-1}-b) = q_2(a^{d-1}-b)$ . This implies that  $b = a^{d-1}$ . Hence we can conclude that  $\dim \mathcal{H}(\mathcal{T}') \cap \mathbb{T} \leq 1$  and  $\mathcal{H}(\mathcal{T}') \cap \mathbb{T}_B$  is finite. So, once again, all three conditions of the lemma fail.  $\square$

**Example 7.33.** Consider a sequence of blowups

$$\bar{X}_3 \xrightarrow{\sigma_3} \bar{X}_2 \xrightarrow{\sigma_2} \bar{X}_1 \xrightarrow{\sigma_1} \bar{X}_0 = \mathbb{P}^1 \times \mathbb{P}^1,$$

where

$$\begin{aligned} \sigma_1: T_1 &\mapsto T_0(0); \\ \sigma_2: T_2 &\mapsto T_1(\infty) = T_0 \cap T_1; \\ \sigma_3: T_3 &\mapsto T_2(1) \end{aligned}$$

The stabilizer of the Puiseux arc space of the point  $\sigma_3(T_3) = T_2(1)$  is contained in the stabilizer of  $\sigma_1(T_1) = T_0(0)$ , hence it coincides with  $\text{Stab}_{\text{fps}} \mathcal{T}'$ . Using Lemma 7.22 we obtain

$$\text{Pui}(\sigma_3(T_3)) = \text{Pui}_{\bar{X}_2, T_2(1)}(0, 1, 1) = \text{Pui}_{\bar{X}_1, T_1(\infty)}(t, 1, 2) = \text{Pui}(t, 2, 2).$$

By Lemma 7.29,  $\text{Stab}_{\text{fps}}(\text{Pui}_{\bar{X}_0, T_0(0)}(t, 2, 2))$  is defined by equations  $c_0 = 0$ ,  $b_0^2 = a$ . It follows that

$$\text{Aut}_{\text{fps}} \mathcal{T}' = \left( \bigoplus_{i=1}^{\infty} \mathbb{G}_a(i) \right) \rtimes \mathbb{T}(\mathcal{T}'),$$

where  $\mathbb{T}(\mathcal{T}') \subset \mathbb{T}$  is the one-parameter subgroup defined by  $b^2 = a$ . The intersection  $\mathbb{T}(\mathcal{T}') \cap \mathbb{T}_B$  has order two and is generated by the involution  $(x, y) \mapsto (x, -y)$ .

## 8. AUTOMORPHISM GROUPS OF $\mathbb{A}^1$ -FIBRATIONS ON SURFACES

For an  $\mathbb{A}^1$ -fibration  $\mu: X \rightarrow B$  on a normal affine surface  $X$  over a smooth affine curve  $B$ , we describe in Subsection 8.3 the automorphism group  $\text{Aut}(X, \mu)$  up to passing to a finite index normal subgroup, see Theorems 8.12 and 8.24. In particular, this applies to the full automorphism group of an  $\text{ML}_1$ -surface.

### 8.1. Notation.

**8.1.** We keep the notation of Section 7. In particular, we consider the induced  $\mathbb{P}^1$ -fibration  $\bar{\mu}: \bar{X} \rightarrow \bar{B}$  on a minimal resolved completion  $\bar{X}$  of  $X$ , and a sequence of blowdowns

$$(30) \quad \Phi: \bar{X} = \bar{X}_m \xrightarrow{\sigma_m} \bar{X}_{m-1} \xrightarrow{\sigma_{m-1}} \dots \xrightarrow{\sigma_1} \bar{X}_0$$

of  $(-1)$ -components of the special fibers  $\mathcal{T}_i = \bar{\mu}^{-1}(\beta_i)$  of  $\bar{\mu}$  with  $\beta_1, \dots, \beta_{n_s} \in B$ , which terminates by a smooth ruling  $\bar{\mu}_0: \bar{X}_0 \rightarrow \bar{B}$ . We assume as before that  $\sigma_i$  contracts the component  $T_i \subset \bar{X}_i \setminus S$  to a point  $p_i \in \bar{X}_{i-1}$ , where  $S$  is the unique horizontal component of  $D = \bar{X} \setminus X$  and a section of  $\bar{\mu}$ . Let also  $F = \bar{\mu}^{-1}(\bar{B} \setminus B)$ . Since  $B$  is affine,  $F \neq \emptyset$ , and we may suppose that  $F$  is a union of irreducible fibers of  $\bar{\mu}$ . We let

$$\mathcal{T} = \bigcup_{i=1}^{n_s} \mathcal{T}_i = \bigcup_{j=1}^m T_j \cup \bigcup T_{0, \beta_i},$$

where  $T_{0, \beta_i} = \bar{\mu}_0^{-1}(\beta_i) \subset \bar{X}_0$ ,  $i = 1, \dots, n_s$ . Thus,  $D \subset D_{\text{ext}} := S \cup F \cup \mathcal{T}$ , and the dual graphs of both  $D$  and  $D_{\text{ext}}$  are trees. We let  $\mathcal{T}_i^{(j)}$  be the image of  $\mathcal{T}_i$  in  $X_j$ .

**Notation 8.2.** Let  $\text{Aut}_B(X, \mu) \subset \text{Aut}(X, \mu)$  be the subgroup of all automorphisms of  $X$  that send each fiber of  $\mu$  into itself, and  $\text{Aut}_\mu B \subset \text{Aut} B$  be the subgroup of all automorphisms of  $B$  induced by the elements of  $\text{Aut}(X, \mu)$ . There is an exact sequence

$$(31) \quad 1 \rightarrow \text{Aut}_B(X, \mu) \rightarrow \text{Aut}(X, \mu) \rightarrow \text{Aut}_\mu B \rightarrow 1.$$

**8.2. Stabilizers of arc spaces.** We need the following fact.

**Lemma 8.3.** *Any automorphism  $\alpha \in \text{Aut}(X, \mu)$  lifts to the minimal resolution of singularities of  $X$  and extends to an automorphism of  $\bar{X} \setminus F$ .*

*Proof.* The first assertion is well known (cf. e.g., [37, Lem. 2.2]) and follows, for instance, from the uniqueness of the minimal resolution of singularities of surfaces. Thus, any automorphism  $g \in \text{Aut}(X, \mu)$  induces a birational automorphism of  $\bar{X}$  regular in  $\bar{X} \setminus D$ . Since  $g$  preserves the  $\mathbb{A}^1$ -fibration  $\mu$ , it extends regularly to the section  $S$ . Furthermore,  $g$  induces a birational transformation  $g_*$  of the dual graphs  $\Gamma_D$  and  $\Gamma_{D_{\text{ext}}}$  fixing the vertex  $S$ , which transforms the dual graphs  $\Gamma_F$  and  $\Gamma_{\mathcal{T}}$  into themselves. By our convention in 7.1 and 8.1  $\Gamma_D$  is minimal. So, the section  $S$  and the components of  $F$  are the only possible zero vertices of  $\Gamma_D$ . All the maximal linear chains in  $\Gamma_{D \cup (S+F)}$  are admissible, that is, with all weights  $\leq -2$ . According to Theorem 3.1 in [36],  $g_*$  can be decomposed into a sequence of elementary transformations in zero vertices in  $F$  followed by an automorphism, see Definition 5.9. Indeed, since  $S$  is fixed by  $g_*$ , also these elementary transformations and the automorphism fix  $S$ . Hence the elementary transformations in the decomposition of  $g_*$  are performed only near components of  $F$ . Now the second assertion follows.  $\square$

**Notation 8.4.** In the notation of 7.1, we let  $\text{Aut}^\bullet(\bar{X}, F) \subset \text{Aut}(X, \mu)$  stand for the subgroup of all automorphisms of  $X$  preserving  $\mu$  and admitting an extension to automorphisms of  $\bar{X} \setminus F$ , which send each component of  $\mathcal{T}$  into itself. Similarly, given  $i \in \{0, \dots, m\}$ , we let  $\text{Aut}^\bullet(\bar{X}_i, F)$  be the group of all birational automorphisms of  $\bar{X}_i$  which preserve  $\bar{\mu}_i$ , send the section  $S$  and each component  $T_{0, \beta_1}, \dots, T_{0, \beta_{n_s}}, T_1, \dots, T_i$



of  $\mathcal{T}^{(i)}$  into itself, and induce automorphisms of  $\bar{X}_i \setminus F$  (see 7.4 and 8.1). Thus,  $\text{Aut}^\bullet(\bar{X}, F) = \text{Aut}^\bullet(\bar{X}_m, F)$ .

**Lemma 8.5.**  $\text{Aut}^\bullet(\bar{X}, F) \subset \text{Aut}(X, \mu)$  is a normal subgroup of finite index.

*Proof.* Let  $\mathbb{S}(n)$  stand for the symmetric group on  $n$  symbols. By Lemma 8.3 there is a natural embedding  $\text{Aut}(X, \mu) \hookrightarrow \text{Aut}(\bar{X} \setminus F, \bar{\mu})$ . Clearly, any  $\alpha \in \text{Aut}(\bar{X} \setminus F, \bar{\mu})$  permutes the special fibers  $\mathcal{T}_j$  of  $\bar{\mu}$  and the components of  $\mathcal{T}$ . Hence  $\alpha$  defines a permutation  $\rho(\alpha) \in \mathbb{S}(m + n_s)$ , where  $\rho: \text{Aut}(X, \mu) \rightarrow \mathbb{S}(m + n_s)$  is a homomorphism. Clearly,  $\text{Aut}^\bullet(X, \mu) = \ker \rho$ . Now the assertion follows.  $\square$

**Notation 8.6** (*stabilizers of arc spaces*). The group  $\text{Aut}^\bullet(\bar{X}_i, F)$  acts naturally on the arc space  $\text{Arc}(\bar{X}_i \setminus F)$ . Given a subset  $W \subset \text{Arc}(\bar{X}_i \setminus F)$ , we let  $\text{Stab}_i(W)$  be the stabilizer of  $W$  in  $\text{Aut}^\bullet(\bar{X}_i, F)$ .

In the next proposition we identify the groups  $\text{Aut}^\bullet(\bar{X}_i, F)$  and  $\text{Stab}_i(W)$  with their images in  $\text{Aut}^\bullet(\bar{X}_0, F)$ .

**Proposition 8.7.** *There is a natural embedding  $\text{Aut}^\bullet(\bar{X}, F) \hookrightarrow \text{Aut}^\bullet(\bar{X}_0, F)$  such that*

$$(32) \quad \text{Aut}^\bullet(\bar{X}, F) = \bigcap_{i: \sigma_i \text{ is outer}} \text{Stab}_0(\text{Pui}(p_i)) \subset \text{Aut}^\bullet(\bar{X}_0, F).$$

*Proof.* We proceed by induction on  $i$ . Assume that our assertion holds for  $\bar{X}_{i-1}$ . Since  $\text{Aut}^\bullet(\bar{X}_i, F)$  stabilizes  $T_i$ , there is a natural homomorphism  $\text{Aut}^\bullet(\bar{X}_i, F) \rightarrow \text{Aut}^\bullet(\bar{X}_{i-1}, F)$ , which embeds  $\text{Aut}^\bullet(\bar{X}_i, F)$  onto the stabilizer of the point  $p_i = \sigma_i(T_i)$  in  $\text{Aut}^\bullet(\bar{X}_{i-1}, F)$ . The latter stabilizer coincides with  $\text{Stab}_{i-1}(\text{Arc}(\bar{X}_{i-1})_{p_i})$ .

If  $T_i$  is inner, then  $p_i$  is already stabilized by  $\text{Aut}^\bullet(\bar{X}_{i-1}, F)$ , thus we have  $\text{Aut}^\bullet(\bar{X}_i, F) \cong \text{Aut}^\bullet(\bar{X}_{i-1}, F)$ . Assume now that  $T_i$  is outer. Then

$$\text{Stab}_{i-1}(\text{Arc}(\bar{X}_{i-1})_{p_i}) = \text{Stab}_{i-1}(\text{Pui}(p_i)),$$

where by abuse of notation we write  $\text{Pui}(p_i)$  for  $\text{Pui}_{\bar{X}_{i-1}, p_i}(0, 1, 1)$  (cf. 7.24). Indeed,  $\text{Pui}(p_i)$  is the subset of arcs of minimal multiplicity (equal to  $\text{mult}(T_i)$ ) in  $\text{Arc}(\bar{X}_{i-1})_{p_i}$ . This subset is stable under the action on  $\text{Arc}(\bar{X}_{i-1})_{p_i}$  of the stabilizer of  $p_i$  in  $\text{Aut}^\bullet(\bar{X}_{i-1}, F)$ . This gives the inclusion

$$\text{Stab}_{i-1}(\text{Arc}(\bar{X}_{i-1})_{p_i}) \subset \text{Stab}_{i-1}(\text{Pui}(p_i)).$$

The inverse inclusion is also clear, since the elements of  $\text{Stab}_{i-1}(\text{Pui}(p_i))$  fix the point  $p_i$ . Passing to the images of our subgroups under their natural embeddings in  $\text{Aut}^\bullet(\bar{X}_0, F)$ , which we denote by the same symbols, we obtain the equalities

$$\text{Aut}^\bullet(\bar{X}_i, F) = \text{Stab}_{i-1}(\text{Pui}(p_i)) = \text{Aut}^\bullet(\bar{X}_{i-1}, F) \cap \text{Stab}_0(\text{Pui}(p_i)).$$

This yields (32) for  $\text{Aut}^\bullet(\bar{X}_i, F)$ , since by the inductive conjecture, it holds for the group  $\text{Aut}^\bullet(\bar{X}_{i-1}, F)$ .  $\square$

### 8.3. Automorphism groups of $\mathbb{A}^1$ -fibrations.

**Notation 8.8.** We fix an  $(\text{Aut}_\mu B)$ -stable Zariski open subset  $\omega \subset B$  such that  $\mu$  admits a trivialization over  $\omega$ . We assume that  $\omega$  is maximal with these properties. We let  $X_\omega = \mu^{-1}(\omega) \cong \omega \times \mathbb{A}^1$ ; this is an  $(\text{Aut}(X, \mu))$ -stable dense open subset in  $X$ .

**Remark 8.9.** If the curve  $B$  is rational, then  $\omega = B \setminus \{\beta_1, \dots, \beta_{n_s}\}$ . Indeed,  $\mu|_{X_\omega} = \mu_0|_{(X_0)_\omega}$  is the projection of a line bundle (see Lemma 7.5), which is trivial if  $B$  is rational. As follows from Lemma 8.3, the open set  $B \setminus \{\beta_1, \dots, \beta_{n_s}\}$  is  $(\text{Aut}_\mu B)$ -stable.

In what follows we treat separately the cases  $\omega \cong \mathbb{A}_*^1, \mathbb{A}^1$  and  $\omega \not\cong \mathbb{A}_*^1, \mathbb{A}^1$ , see Theorems 8.12 and 8.24, respectively.

8.3.1. *Case*  $B = \mathbb{A}^1$ ,  $\omega \cong \mathbb{A}_*^1$ . If  $\omega = B \cong \mathbb{A}^1$  or  $\omega = B \cong \mathbb{A}_*^1$ , then  $X = X_\omega \cong \mathbb{A}^2$  and  $X = X_\omega \cong \mathbb{A}_*^1 \times \mathbb{A}^1$ , respectively, and the group  $\text{Aut}(X, \mu)$  is the usual de Jonquères group and its analog as in 3.3 and 3.5, respectively. Hence we assume in the sequel that  $B = \mathbb{A}^1$  and  $\omega = \mathbb{A}_*^1$ . This is the case, for instance, for any Gizatullin surface different from the plane  $\mathbb{A}^2$ . For the  $\text{ML}_1$ -surfaces, this case was studied in [8]; our Theorem 8.12 precises Corollary 2.3 in [8].

**8.10.** We let  $F = \bar{\mu}^{-1}(\infty)$  and  $\mathcal{T} = \mathcal{T}_1 = \bar{\mu}^{-1}(0)$ . Performing suitable elementary transformations with centers on  $F$  (see Definition 5.9) one can achieve the equality  $S^2 = 0$ . Then the linear pencil  $|S|$  on  $\bar{X}$  defines a second  $\mathbb{P}^1$ -fibration  $\bar{\nu}: \bar{X} \rightarrow \mathbb{P}^1$  such that  $S = \bar{\nu}^{-1}(\infty)$ . The birational morphism

$$\Phi: \bar{X} \rightarrow \bar{X}_0 = \mathbb{P}^1 \times \mathbb{P}^1, \quad w \mapsto (\bar{\mu}(w), \bar{\nu}(w)),$$

is biregular on  $\bar{X} \setminus \bar{\mu}^{-1}(0)$ . (In the notation of 5.12,  $\bar{\mu} = \Phi_0$  and  $\bar{\nu} = \Phi_1$ , while  $F = C_0$  and  $S = C_1$ .) The affine coordinate on  $\bar{B} \cong \mathbb{P}^1$  is chosen so that  $\bar{B} \setminus B = \{\infty\}$ ,  $B \setminus \omega = \{0\}$ , and so,  $F = \{\infty\} \times \mathbb{P}^1 \subset \mathbb{P}^1 \times \mathbb{P}^1$ .

**Remark 8.11.** On  $\bar{X}_0 = \mathbb{P}^1 \times \mathbb{P}^1$  we have (cf. 7.27)

$$\begin{aligned} \text{Aut}^\bullet(\bar{X}_0, F) &= \left\{ (x, y) \mapsto (ax, by + P(x)) \mid (a, b) \in \mathbb{G}_m^2, P = \sum_{i=0}^N c_i x^i \in \mathbb{K}[x] \right\} \\ &\cong \left( \bigoplus_{i=0}^{\infty} \mathbb{G}_a(i) \right) \rtimes \mathbb{T}, \end{aligned}$$

where

$$\mathbb{T} = \{(x, y) \mapsto (ax, by) \mid (a, b) \in (\mathbb{K}^\times)^2\} \cong \mathbb{G}_m^2$$

and

$$\mathbb{G}_a(i) = \{(x, y) \mapsto (x, y + c_i x^i) \mid c_i \in \mathbb{K}\} \cong \mathbb{G}_a.$$

The natural embedding  $\text{Aut}^\bullet(\bar{X}_0, F) \hookrightarrow \text{Aut} \text{Arc}(\bar{X}_0)_{T_0}^*$  is tautological in coordinates  $(a, b, c_0, \dots)$  and corresponds to the embedding of the polynomial ring into the ring of formal power series. So, all results of the previous section hold automatically for  $\text{Aut}^\bullet(\bar{X}, F)$ . In particular,  $\text{Stab}_0 W = \text{Stab}_{\text{fps}} W \cap \text{Aut}^\bullet(\bar{X}_0, F)$  for any subset  $W \subset \text{Arc}(\bar{X}_0)_{T_0}^*$ , and  $\text{Aut}^\bullet(\bar{X}, F) = \text{Stab}_{\text{fps}} \mathcal{T} \cap \text{Aut}^\bullet(\bar{X}_0, F)$ .

**Theorem 8.12.** *Let  $X$  be a normal affine surface, and let  $\mu: X \rightarrow \mathbb{A}^1$  be an  $\mathbb{A}^1$ -fibration with a unique special fiber  $\mu^{-1}(0)$ . Then the automorphism group  $\text{Aut}(X, \mu)$  is a finite extension of*

$$(33) \quad \text{Aut}^\bullet(\bar{X}, F) = \left( \bigoplus_{k=d}^{\infty} \mathbb{G}_a(k) \right) \rtimes \Lambda_\mu,$$

where  $d \in \mathbb{Z}_{>0}$  and  $\Lambda_\mu$  is conjugate to a closed subgroup of the standard torus  $\mathbb{T}$  by an element of  $\bigoplus_{k=0}^{d-1} \mathbb{G}_a(k)$ . Furthermore, if  $\dim \Lambda_\mu = 1$ , then the action of  $\Lambda_\mu$  on  $X$  is transversal, i.e., the intersection of a  $\Lambda_\mu$ -orbit and a  $\mu$ -fiber is always finite.

*Proof.* As mentioned in Remark 8.11, for any  $W \subset \text{Arc}(\bar{X}_0)_{T_0}$  we have

$$\text{Stab}_0 W = \text{Stab}_{\text{fps}} W \cap \text{Aut}^\bullet(\bar{X}_0, F).$$

Hence by Proposition 8.7,

$$\text{Aut}^\bullet(\bar{X}, F) = \text{Stab}_{\text{fps}} \mathcal{T} \cap \text{Aut}^\bullet(\bar{X}_0, F).$$

Now Proposition 7.30 implies (33). Finally,  $\text{Aut}(X, \mu)$  is a finite extension of  $\text{Aut}^\bullet(\bar{X}, F)$  by Lemma 8.5.  $\square$

8.3.2. Case  $\omega \not\cong \mathbb{A}_*^1, \mathbb{A}^1$ .

**8.13.** As before, the notation  $\text{Aut}_B$  with the base curve  $B$  as a subscript means passing to the subgroup of automorphisms of  $(X, \mu)$  which induces the identity on  $B$ . In particular,

$$\text{Aut}_B^\bullet(\bar{X}, F) = \text{Aut}^\bullet(\bar{X}, F) \cap \text{Aut}_B(\bar{X} \setminus F).$$

**Lemma 8.14.** *If  $\omega \not\cong \mathbb{A}_*^1, \mathbb{A}^1$  then  $\text{Aut}_B^\bullet(\bar{X}, F)$  is a normal subgroup of  $\text{Aut}(X, \mu)$  of finite index.*

*Proof.* By Lemma 8.5,  $\text{Aut}^\bullet(\bar{X}, F) \triangleleft \text{Aut}(X, \mu)$  is a normal subgroup of finite index. Due to (31),  $\text{Aut}_B(X, \mu) \triangleleft \text{Aut}(X, \mu)$ . Hence

$$\text{Aut}_B^\bullet(\bar{X}, F) = \text{Aut}^\bullet(\bar{X}, F) \cap \text{Aut}_B(X, \mu) \triangleleft \text{Aut}(X, \mu).$$

Likewise in (31) we have an exact sequence

$$(34) \quad 1 \rightarrow \text{Aut}_B^\bullet(\bar{X}, F) \rightarrow \text{Aut}^\bullet(\bar{X}, F) \rightarrow \text{Aut}_\mu B \subset \text{Aut} B.$$

Since  $\omega$  is  $\text{Aut}_\mu B$ -stable, there is an inclusion  $\text{Aut}_\mu B \subset \text{Aut} \omega$ , where  $\omega \not\cong \mathbb{A}_*^1, \mathbb{A}^1$  is an affine curve of non-exceptional type. Hence  $\text{Aut} \omega$  is a finite group. Now the assertion follows.  $\square$

**8.15.** Using the equivariant local trivialization  $X_\omega \cong \omega \times \mathbb{A}^1$  (see 8.8), for  $(x, y) \in \omega \times \mathbb{A}^1$  we can write

$$(35) \quad \text{Aut}_\omega X_\omega = \{(x, y) \mapsto (x, Qy + P) \mid Q \in \mathcal{O}_\omega^\times(\omega), P \in \mathcal{O}_\omega^+(\omega)\},$$

where  $A^\times$  stands as before for the multiplicative group of an algebra  $A$  and  $A^+$  for its additive group. Note that  $\mathcal{O}_\omega^+(\omega)$  is an infinite dimensional Abelian unipotent group. In fact,  $\mathcal{O}_\omega^+(\omega) = \mathbb{U}_\mu(X_\omega) \subset \text{Aut}_\omega X_\omega$  is the subgroup of automorphisms that act on the  $\mu$ -fibers by translations, and  $\mathcal{O}_\omega^\times(\omega) \subset \text{Aut}_\omega X_\omega$  is the subgroup of automorphisms that fix the zero section of the (trivial) line bundle  $\mu|_{X_\omega}: X_\omega \rightarrow \omega$ .

The functions  $Q \in \mathcal{O}_\omega^\times(\omega)$  have their zeros and poles in  $\bar{B} \setminus \omega$ . So, there is a homomorphism  $\mathcal{O}_\omega^\times(\omega) \rightarrow \mathbb{Z}^N$  with kernel  $\mathbb{K}^\times \subset \mathcal{O}_\omega^\times(\omega)$  consisting of the nonzero constants, where  $N$  is the number of punctures of  $\omega$ . Thus,  $\mathcal{O}_\omega^\times(\omega) \cong \mathbb{G}_m \times \mathbb{Z}^r$  for some  $r < N$ , and

$$\text{Aut}_\omega X_\omega \cong \mathcal{O}_\omega^+(\omega) \rtimes \mathcal{O}_\omega^\times(\omega) \cong \mathcal{O}_\omega^+(\omega) \rtimes (\mathbb{G}_m \times \mathbb{Z}^r).$$

**Notation 8.16.** We let  $X_0 = \bar{X}_0 \setminus (S \cup F)$ . By Lemma 7.5,  $\mu_0 = \bar{\mu}_0|_{X_0}: X_0 \rightarrow B$  has a structure of a line bundle, say,  $L_0$  with a zero section given in  $X_\omega \cong \omega \times \mathbb{A}^1$ ,  $X_\omega \subset X_0$ , by equation  $y = 0$ . Let  $\mathbb{U}_{\mu_0} \subset \text{Aut}_B(X_0, \mu_0)$  be the unipotent de Jonquères group of the automorphisms which restrict to translations on general  $\mu$ -fibers (see Definition 6.1).

**Proposition 8.17.** *For some  $s \geq 0$  there are decompositions*

$$(36) \quad \text{Aut}_B^\bullet(\bar{X}_0, F) = \text{Aut}_B(\bar{X}_0, F) \cong \mathbb{U}_{\mu_0} \rtimes \Lambda_B \cong \mathbb{U}_{\mu_0} \rtimes (\mathbb{G}_m \times \mathbb{Z}^s),$$

where  $\Lambda_B = \text{Aut}_B L_0 \cong \mathcal{O}_B^\times(B)$ .

*Proof.* The subgroup  $\mathbb{U}_{\mu_0} \subset \text{Aut}_\mu^\bullet(\bar{X}_0, \bar{\mu}_0)$  acts freely and transitively on the space  $H^0(B, \mathcal{O}_B(L_0))$  of global sections of  $L_0$ . Therefore,  $\text{Aut}_B^\bullet(\bar{X}_0, F)$  is generated by  $\mathbb{U}_{\mu_0} \cong H^0(B, \mathcal{O}_B(L_0))$  and the subgroup  $\Lambda_B \subset \text{Aut}_B^\bullet(\bar{X}_0, F)$  of automorphisms that fix the zero section. This leads to the first decomposition in (36). By our assumption, the trivialization  $X_\omega \cong \omega \times \mathbb{A}^1$  and the line bundle  $L_0$  share the same zero section. Hence there is an embedding  $\Lambda_B \hookrightarrow \mathcal{O}_\omega^\times(\omega) \cong \mathbb{G}_m \times \mathbb{Z}^r$  with  $\Lambda_B \supset \mathbb{G}_m$ , see 8.15. Now the second isomorphism in (36) follows.

To show the last assertion, it suffices to observe that the group  $\text{Aut}_B^\bullet(\bar{X}_0, F) = \mathbb{U}_{\mu_0} \rtimes \Lambda_B$  acts on  $H^0(B, \mathcal{O}_B(L_0))$  via

$$g = (P, Q): \Psi \mapsto Q\Psi + P \quad \forall \Psi \in H^0(B, \mathcal{O}_B(L_0)),$$

where  $P \in \mathbb{U}_{\mu_0} = H^0(B, \mathcal{O}_B(L_0))$  and  $Q \in \Lambda_B = \text{Aut}_B L_0 = \mathcal{O}_B^\times(B)$ .  $\square$

**Remark 8.18.** The neutral component  $\mathbb{G}_m$  of the group  $\mathcal{O}_B^\times(B)$  acts on  $L_0$  by homotheties; this defines a 1-torus  $\mathbb{T}_B \subset \text{Aut}_B L_0$  (cf. 7.27).

**8.19.** For every  $j = 1, \dots, n_s$  there is a natural embedding

$$\iota_j: \text{Aut}_B^\bullet(\bar{X}_0, F) \hookrightarrow \text{Aut}_B \text{Arc}(\bar{X}_0)_{T_0, \beta_j}^*$$

(see Remark 8.11), where  $\iota_j$  embeds the factors of the decomposition  $\text{Aut}_B(\bar{X}_0, F) \cong \mathbb{U}_{\mu_0} \rtimes \Lambda_B$  into the respective factors of the decomposition

$$\text{Aut}_B \text{Arc}(\bar{X}_0)_{T_0, \beta_j}^* = \left( \prod_{i=0}^{\infty} \mathbb{G}_a(i) \right) \rtimes \mathcal{H}_B,$$

see 7.27. Indeed, in both cases the first factor is the unipotent radical (acting by translations on fibers), and the second consists of the automorphisms preserving the zero section. Thus,  $\mathbb{U}_{\mu_0} \hookrightarrow \prod_{i=0}^{\infty} \mathbb{G}_a(i)$  and  $\Lambda_B \hookrightarrow \mathcal{H}_B$ .

For a special fiber  $\mathcal{T}' = \mathcal{T}_j \subset \mathcal{T}$  we have by (29),

$$\text{Stab}_{\text{fps}} \mathcal{T}_j = \left( \prod_{i=N_j}^{\infty} \mathbb{G}_a(i) \right) \rtimes \mathcal{H}_j^{h_j},$$

where in the notation of 7.31,  $h_j = h(\mathcal{T}_j) \in \bigoplus_{i=0}^{N_j-1} \mathbb{G}_a(i)$  with  $N_j = N(\mathcal{T}_j)$ , and  $\mathcal{H}_j = \mathcal{H}(\mathcal{T}_j) \subset \mathcal{H}$ .

In Proposition 8.23 below we gather these local data for different special fibers. In the proof we use the following lemma.

**Lemma 8.20.** *Given natural numbers  $N_1, \dots, N_{n_s} \in \mathbb{Z}_{>0}$  and a collection  $(\psi_j)_{j=1, \dots, n_s}$ , where  $\psi_j \in \mathcal{O}_{B, \beta_j} \forall j$ , there exists a section  $\Psi \in H^0(B, \mathcal{O}_B(L_0))$  such that*

$$\iota_j(\Psi) \equiv \psi_j \pmod{\mathfrak{m}_{\beta_j}^{N_j}} \quad \forall j = 1, \dots, n_s.$$

*Proof.* Indeed, consider the coherent ideal sheaf

$$\mathcal{I} = \left( \prod_{j=1}^{n_s} \mathfrak{m}_{\beta_j}^{N_j} \right) \cdot \mathcal{O}_B(L_0) \subset \mathcal{O}_B(L_0).$$

The local data  $(\psi_j)_j$  defines a section of the skyscraper sheaf  $\mathcal{O}_B(L_0)/\mathcal{I}$  on the affine curve  $B$ , which can be interpolated by a global section  $\Psi \in H^0(B, \mathcal{O}_B(L_0))$  due to the Serre Vanishing  $H^1(B, \mathcal{I}) = 0$ .  $\square$

The following corollary is immediate.

**Corollary 8.21.** *There exists  $h \in \mathbb{U}_{\mu_0} = H^0(B, \mathcal{O}_B(L_0))$  such that*

$$(37) \quad \iota_j(h) \equiv h_j \pmod{t^{N_j} \mathbb{K}[[t]]} \quad \forall j = 1, \dots, n_s,$$

where  $N_j$  and  $h_j$  are as in 8.19.

**Notation 8.22.** We let  $D_0 = \sum_{j=1}^{n_s} N_j[\beta_j]$  and

$$\Lambda_{\mu, B} = \bigcap_{j=1}^{n_s} \iota_j^{-1}(\iota_j(\Lambda_B) \cap \mathcal{H}_j) \subset \Lambda_B,$$

$$\mathbb{U}_{\mu} = \{P \in H^0(B, \mathcal{O}_B(L_0)) = \mathbb{U}_{\mu_0} \mid \text{div} P \geq D_0\} = H^0(B, \mathcal{O}_B(-D_0)).$$

**Proposition 8.23.** *We have*

$$(38) \quad \text{Aut}_B^\bullet(\bar{X}, F) = \mathbb{U}_\mu \rtimes \Lambda_{\mu, B}^h,$$

where  $h \in \mathbb{U}_{\mu_0}$  verifies (37).

*Proof.* Let  $h \in \mathbb{U}_{\mu_0}$  verifies (37), see Corollary 8.21. Letting  $h^{-1} \circ g \circ h = (P, Q) \in \mathbb{U}_{\mu_0} \rtimes \Lambda_B$  (that is, letting  $g = (P + h(1 - Q), Q)$ ), we let also  $P_j = \iota_j(P)$  and  $Q_j = \iota_j(Q)$ . We claim that the following are equivalent:

- (i)  $g \in \text{Aut}_B^\bullet(\bar{X}, F)$ ;
- (ii)  $\iota_j(g) \in \text{Stab}_{\text{fps}} \mathcal{T}_j$  for each  $j = 1, \dots, n_s$ ;
- (iii)  $\iota_j(h^{-1} \circ g \circ h) \in \left( \prod_{i=N_j}^\infty \mathbb{G}_a(i) \right) \rtimes \mathcal{H}_j \quad \forall j$ ;
- (iv)  $P_j \in \prod_{i=N_j}^\infty \mathbb{G}_a(i)$  and  $Q_j \in \mathcal{H}(\mathcal{T}_j) \quad \forall j$ ;
- (v)  $\text{div} P \geq D_0 = \sum_{j=1}^{n_s} N_j [\beta_j]$  and  $Q \in \bigcap_{j=1}^{n_s} \iota_j^{-1}(\iota_j(\Lambda_B) \cap \mathcal{H}_j)$ ;
- (vi)  $P \in \mathbb{U}_\mu$  and  $Q \in \Lambda_{\mu, B}$ .

Indeed, it follows from Propositions 7.30 and 8.7 that an element  $g \in \text{Aut}_B^\bullet(\bar{X}_0, F)$  belongs to  $\text{Aut}_B^\bullet(\bar{X}, F)$  if and only if  $\iota_j(g) \in \text{Stab}_{\text{fps}} \mathcal{T}_j$  for each  $j$ , where  $\text{Stab}_{\text{fps}} \mathcal{T}_j$  is defined as in (28). This proves the equivalence (i)  $\Leftrightarrow$  (ii).

After replacing  $h$  in (iii) by  $h_j$ , the equivalence (ii)  $\Leftrightarrow$  (iii) follows from Proposition 7.30. By Corollary 8.21 this holds even without this replacement.

The equivalence (iii)  $\Leftrightarrow$  (iv) is immediate, and (iv)  $\Leftrightarrow$  (v)  $\Leftrightarrow$  (vi) follow from our definitions, see 7.27 and 8.22. This proves the equivalence (i)  $\Leftrightarrow$  (vi). Now the proposition follows.  $\square$

**Theorem 8.24.** *Let  $\mu: X \rightarrow B$  be an  $\mathbb{A}^1$ -fibration on a normal affine surface  $X$  over a smooth affine curve  $B$ . If, in the notation of 8.2,  $\omega \not\cong \mathbb{A}_*^1, \mathbb{A}^1$ , then the automorphism group  $\text{Aut}(X, \mu)$  is a finite extension of*

$$(39) \quad \text{Aut}_B^\bullet(\bar{X}, F) \cong \mathbb{U}_\mu \rtimes (\Gamma_\mu \rtimes \mathbb{Z}^l) \quad \text{for some } l \geq 0,$$

where  $\mathbb{U}_\mu = H^0(B, \mathcal{O}_B(-D_0))$  with  $D_0$  as in 8.22, and  $\Gamma_\mu = \mathbb{T}_B \cap \Lambda_{\mu, B}$ , see 8.18. Furthermore, either  $\Gamma_\mu$  is a finite cyclic group, or  $\Gamma_\mu = \mathbb{T}_B \cong \mathbb{G}_m$  and each  $\mu$ -fiber  $\mu^{-1}(b)$ ,  $b \in B$ , is isomorphic to  $\mathbb{A}^1$ .

*Proof.* By Lemma 8.14,  $\text{Aut}_B^\bullet(\bar{X}, F) \cong \mathbb{U}_\mu \rtimes \Lambda_{\mu, B}$  is a normal subgroup of finite index in  $\text{Aut}(X, \mu)$ . To deduce (39) it suffices to apply Proposition 8.23, where

$$\Lambda_{\mu, B} \hookrightarrow \Lambda_B = \mathcal{O}_B^\times(B) \cong \mathbb{T}_B \times \mathbb{Z}^s \cong \mathbb{G}_m \times \mathbb{Z}^s,$$

see 8.17 and 8.18. For the last assertion, see [32, §3] and [33, Rem. 3.13(iii)].  $\square$

The following corollary is immediate from Corollary 6.4 and Theorems 8.12 and 8.24 (cf. Corollary 6.11).

**Corollary 8.25.** *For any  $\mathbb{A}^1$ -fibration  $\mu: X \rightarrow B$  on a normal affine surface  $X$  over a smooth affine curve  $B$ , the group  $\text{Aut}(X, \mu)$  is an extension of a metabelian nested ind-group of rank  $\leq 2$  by at most countable group. Any two maximal tori in  $\text{Aut}(X, \mu)$  are conjugated.*

**8.26.** If  $\mu: X \rightarrow B$  admits an effective  $\mathbb{G}_m$ -action along the fibers of  $\mu$ , that is,  $\Gamma_\mu \cong \mathbb{G}_m$ , then  $\mu$  is the projection of a parabolic  $\mathbb{G}_m$ -surface (see Definition 4.10).

**Proposition 8.27.** *Under the assumptions of Theorem 8.24,  $\Gamma_\mu \cong \mathbb{G}_m$  if and only if  $X$  is a parabolic  $\mathbb{G}_m$ -surface, if and only if the connected components of the dual graph  $\Gamma_{\mathcal{T}}$  are linear chains. In the case  $\omega \cong \mathbb{A}_*^1$  we have the equivalences*

$$\Gamma_\mu = \mathbb{G}_m \Leftrightarrow \Lambda_{\mu, B} \cong \mathbb{T} \Leftrightarrow \text{rk Aut } X = 2,$$

where the latter means that  $X$  is an affine toric surface.

*Proof.* By Proposition 8.27, the components of  $\Gamma_{\mathcal{T}}$  are linear if and only if  $\mathcal{H} = \mathcal{H}_j$  for every special fiber  $\mathcal{T}_j$ . The latter is true if and only if  $\mathcal{H}_j \supset \mathbb{T}_B$  for each  $j$ , which is equivalent to  $\Gamma_{\mathcal{T}} \supset \mathbb{T}_B$ . The assertions now follow easily.  $\square$

## REFERENCES

- [1] I. Arzhantsev, H. Flenner, S. Kaliman, F. Kutzschebauch, and M. Zaidenberg, *Flexible varieties and automorphism groups*, Duke Math. J. 162 (2013), 767–823.
- [2] I. Arzhantsev, M. Zaidenberg, *Acyclic curves and group actions on affine toric surfaces*, In: Affine Algebraic Geometry, Proceedings of the Conference dedicated to Prof. Miyanishi’s 70th birthday. Kayo Masuda, Hideo Kojima, Takashi Kishimoto (eds.), 1–41. World Scientific Publishing Co. 2013.
- [3] L. Bădescu, *Projective geometry and formal geometry*, Mathematics Institute of the Polish Academy of Sciences. Mathematical Monographs (New Series), 65. Birkhäuser Verlag, Basel, 2004.
- [4] T. Bandman, L. Makar-Limanov, *Affine surfaces with  $AK(S) = \mathbb{C}$* , Michigan Math. J. 49 (2001), 567–582.
- [5] T. Bandman, L. Makar-Limanov, *On  $\mathbb{C}$ -fibrations over projective curves*, Michigan Math. J. 56 (2008), 669–686.
- [6] W. P. Barth, K. Hulek, C. A. M. Peters, A. Van de Ven, *Compact complex surfaces*. Second edition, Springer-Verlag, Berlin, 2004.
- [7] H. Bass, *Algebraic group actions on affine spaces*, Group actions on rings (Brunswick, Maine, 1984), 1–23, Contemp. Math. 43, Amer. Math. Soc., Providence, RI, 1985.
- [8] J. Bertin, *Pinceaux de droites et automorphismes des surfaces affines*, J. Reine Angew. Math. 341 (1983), 32–53.
- [9] J. Blanc, A. Dubouloz, *Automorphisms of  $\mathbb{A}^1$ -fibered affine surfaces*, Trans. Amer. Math. Soc. 363 (2011), 5887–5924.
- [10] J. Blanc, A. Dubouloz, *Affine surfaces with a huge group of automorphisms*, arXiv:1302.3813 (2013), 18p.
- [11] J. Blanc, I. Stampfli, *Automorphisms of the plane preserving a curve*, Algebraic Geom. 2 (2) (2015), 193–213.
- [12] A. Borel, J. Tits, *Groupes réductifs*, Publ. IHES 27 (1965), 55–150.
- [13] S. Cantat, S. Lamy, *Groupes d’automorphismes polynomiaux du plan*, Geom. Dedicata 123 (2006), 201–221.
- [14] P. Cassou-Noguès, P. Russell, *Birational morphisms  $\mathbb{C}^2 \rightarrow \mathbb{C}^2$  and affine ruled surfaces*, In : Affine Algebraic Geometry, in honor of Prof. M. Miyanishi, 57–105. Osaka Univ. Press, Osaka, 2007.
- [15] R. Cicurel, *Sur les orbites affines des  $\mathbb{C}$ -variétés admettant une orbite ouverte*, CRAS, Paris, 274, ser. A (1972), 1316–1318.
- [16] D. Cox, *Lectures on toric varieties. Lecture 7: Toric surfaces*, Grenoble, Institut Fourier, École d’Été 2000: Géométrie des variétés toriques, <http://www-fourier.ujf-grenoble.fr/>
- [17] A. J. Crachiola, *On automorphisms of Danielewski surfaces*, J. Algebraic Geom. 15 (2006), 111–132.
- [18] D. Daigle, *Classification of weighted graphs up to blowing-up and blowing-down*. math.AG/0305029, 2003, 47p.
- [19] D. Daigle, *Classification of linear weighted graphs up to blowing-up and blowing-down*, Canad. J. Math. 60 (2008), 64–87.
- [20] D. Daigle, *Affine surfaces with trivial Makar-Limanov invariant*, J. Algebra 319 (2008), 3100–3111.
- [21] D. Daigle, R. Kolhatkar, *Complete intersection surfaces with trivial Makar-Limanov invariant*, J. Algebra 350 (2012), 1–35.
- [22] V. I. Danilov, M. H. Gizatullin, *Automorphisms of affine surfaces. I*, Math. USSR Izv. 9 (1975), 493–534.
- [23] V. I. Danilov, M. H. Gizatullin, *Automorphisms of affine surfaces. II*, Math. USSR Izv. 11 (1977), 51–98.
- [24] V. I. Danilov, M. H. Gizatullin, *Examples of non-homogeneous quasihomogeneous surfaces* (Russian), Math. USSR Izv. 38:1 (1974), 42–58.
- [25] I. Dimitrov, I. Penkov, J. A. Wolf, *A Bott-Borel-Weil theory for direct limits of algebraic groups*, Amer. J. Math. 124 (2002), 955–998.

- [26] A. Dubouloz, *Completions of normal affine surfaces with a trivial Makar-Limanov invariant*, Michigan Math. J. 52 (2004), 289–308.
- [27] A. Dubouloz, S. Lamy, *Automorphisms of open surfaces with irreducible boundary*, arXiv:1404.4838 (2014), 35p.
- [28] E. B. Dynkin, *Maximal subgroups of the classical groups* (Russian). Trudy Moskov. Mat. Obsc. 1 (1952), 39–166. English translation in: Amer. Math. Soc. Transl. 6 (1957), 245–378.
- [29] L. Ein, M. Mustață, *Jet Schemes and Singularities*, Algebraic geometry, Seattle 2005, 505–546, Proc. Sympos. Pure Math., 80, Part 2, Amer. Math. Soc., Providence, RI, 2009.
- [30] K.-H. Fieseler, *On complex affine surfaces with  $\mathbb{C}^+$ -action*, Comment. Math. Helv. 69 (1994), 5–27.
- [31] H. Flenner, M. Zaidenberg,  *$\mathbb{Q}$ -acyclic surfaces and their deformations*, Classification of algebraic varieties (L’Aquila, 1992), 143–208. Contemp. Math. 162, Amer. Math. Soc., Providence, RI, 1994.
- [32] H. Flenner, M. Zaidenberg, *Normal affine surfaces with  $\mathbb{C}^*$ -actions*, Osaka J. Math. 40 (2003), 981–1009.
- [33] H. Flenner, M. Zaidenberg, *Locally nilpotent derivations on affine surfaces with a  $\mathbb{C}^*$ -action*, Osaka J. Math. 42 (2005), 931–974.
- [34] H. Flenner, M. Zaidenberg, *On the uniqueness of  $\mathbb{C}^*$ -actions on affine surfaces*, Affine Algebraic Geometry, Contemp. Math. 369, Amer. Math. Soc., Providence, RI, 2005.
- [35] H. Flenner, M. Zaidenberg, *On a result of Miyanishi-Masuda*, Arch. Math. (Basel) 87 (2006), 15–18.
- [36] H. Flenner, S. Kaliman, M. Zaidenberg, *Birational transformations of weighted graphs*, in: Affine algebraic geometry, 107–147. Osaka Univ. Press, 2007. *Corrigendum*, in: Affine algebraic geometry, 35–38. CRM Proc. Lecture Notes, 54, Amer. Math. Soc., Providence, RI, 2011.
- [37] H. Flenner, S. Kaliman, M. Zaidenberg, *Completions of  $\mathbb{C}^*$ -surfaces*, in: Affine algebraic geometry, 149–200. Osaka Univ. Press, Osaka 2007.
- [38] H. Flenner, S. Kaliman, M. Zaidenberg, *Uniqueness of  $\mathbb{C}^*$ - and  $\mathbb{C}_+$ -actions on Gizatullin surfaces*, Transformation groups 13 (2008), 305–354.
- [39] H. Flenner, S. Kaliman, M. Zaidenberg, *Smooth Affine Surfaces with Non-Unique  $\mathbb{C}^*$ -Actions*, J. Algebraic Geom. 20 (2011), 329–398.
- [40] H. Flenner, S. Kaliman, M. Zaidenberg, *On the Danilov-Gizatullin isomorphism theorem*, Enseign. Math. (2) 55 (2009), 3–4, 275–283.
- [41] H. Flenner, S. Kaliman, M. Zaidenberg, *Corrigendum to Our Paper ‘Birational transformations of weighted graphs’*, in: Affine Algebraic Geometry, 35–38. CRM Proceedings & Lecture Notes 54. Providence, RI: Amer. Math. Soc. 2011.
- [42] H. Flenner, S. Kaliman, M. Zaidenberg, *Deformation equivalence of affine ruled surfaces*, arXiv:1305.5366 (2013), 35p.
- [43] M. Fraser, A. Mader, *The structure of the automorphism group of polynomial rings*, J. Algebra 25 (1973), 25–39.
- [44] G. Freudenburg, *Algebraic theory of locally nilpotent derivations*, Encyclopaedia of Mathematical Sciences, 136. Invariant Theory and Algebraic Transformation Groups, VII. Springer-Verlag, Berlin, 2006.
- [45] J.-Ph. Furter, S. Maubach, *A Characterization of Semisimple Plane Polynomial Automorphisms*, J. Pure Appl. Algebra 214 (2010), 574–583.
- [46] M. H. Gizatullin, *On affine surfaces that can be completed by a nonsingular rational curve*, Izvestia Akad. Nauk SSSR, Ser. Mat. 34 (1970), 778–802.
- [47] M. H. Gizatullin, *Invariants of incomplete algebraic surfaces that can be obtained by means of completions*, Izvestia Akad. Nauk SSSR, Ser. Mat. 35 (1971), 485–497.
- [48] M. H. Gizatullin, *Affine surfaces that are quasihomogeneous with respect to an algebraic group*, Math. USSR Izv. 5 (1971), 754–769.
- [49] M. H. Gizatullin, *Quasihomogeneous affine surfaces*, Math. USSR Izv. 5 (1971), 1057–1081.
- [50] R. V. Gurjar, K. Masuda, M. Miyanishi, P. Russell, *Affine lines on affine surfaces and the Makar-Limanov invariant*, Canad. J. Math. 60 (2008), 109–139.
- [51] R. V. Gurjar, M. Miyanishi, *Automorphisms of affine surfaces with  $\mathbb{A}^1$ -fibrations*, Michigan Math. J. 53 (2005), 33–55.
- [52] J. E. Humphreys, *Linear Algebraic Groups*, Springer, New York, 1975.
- [53] S. Itaka, *On logarithmic Kodaira dimension of algebraic varieties*, Compl. Anal. and Algebr. Geom., Tokyo, Iwanami Shoten, 1977, 175–189.
- [54] S. Ishii, *Jet schemes, arc spaces and the Nash problem*, C. R. Math. Acad. Sci. Soc. R. Can. 29 (2007), 1–21.

- [55] Z. Jelonek, *On the group of automorphisms of a quasi-affine variety*, arXiv:1403.6640 (2014), 7p.
- [56] Z. Jelonek, T. Lenarcik, *Automorphisms of affine smooth varieties*, Proc. Amer. Math. Soc. 142 (2014), 1157–1163.
- [57] T. Kambayashi. On the absence of nontrivial separable forms of the affine plane. J. Algebra 35 (1975), 449–456.
- [58] T. Kambayashi. Automorphism group of a polynomial ring and algebraic group action on an affine space. J. Algebra 60 (1979), 439–451.
- [59] T. Kambayashi, M. Miyanishi, *On flat fibrations of the affine line*, Illinois J. Math. 22 (1978), 662–671.
- [60] T. Kambayashi, D. Wright, *Flat families of affine lines are affine-line bundles*, Illinois J. Math. 29 (1985), 672–681.
- [61] T. Kishimoto, Yu. Prokhorov, M. Zaidenberg, *Group actions on affine cones*, Affine algebraic geometry, 123–163, CRM Proc. Lecture Notes, 54, Amer. Math. Soc., Providence, RI, 2011.
- [62] S. Kovalenko, *Transitivity of automorphism groups of Gizatullin surfaces*, Internat. Mathem. Res. Notices 2015, doi: 10.1093/imrn/rnv025, 34p.
- [63] S. Kovalenko, *Smooth Non-Homogeneous Gizatullin Surfaces*, arXiv:1311.3823 (2013), 30p.
- [64] H. Kraft, *Geometrische Methoden in der Invariantentheorie*. 2. Auflage. Aspekte der Mathematik D1. Vieweg Verlag (1985).
- [65] H. Kraft, *Automorphism groups of affine varieties and a characterization of affine  $n$ -space*, arXiv:1501.06362 (2015), 15p.
- [66] H. Kraft, G. Schwarz, *Finite automorphisms of affine  $n$ -space*, In: *Automorphisms of affine spaces*. Curaçao, 1994, 55–66, Kluwer Acad. Publ., Dordrecht, 1995.
- [67] S. Kumar, *Kac-Moody groups, their flag varieties and representation theory*, Progress in Mathematics 204, Birkhauser Boston Inc., Boston, MA, 2002.
- [68] S. Lamy, *L’alternative de Tits pour  $\text{Aut}[\mathbb{C}^2]$* , J. Algebra 239 (2001), 413–437.
- [69] S. Lamy, *Sur la structure du groupe d’automorphismes de certaines surfaces affines*, Publ. Mat. 49 (2005), 3–20.
- [70] A. Liendo, *Affine  $T$ -varieties of complexity one and locally nilpotent derivations*, Transform. Groups 15 (2010), 389–425.
- [71] V. Lin, M. Zaidenberg, *Configuration Spaces of the Affine Line and their Automorphism Groups*. In: *Automorphisms in birational and affine geometry. Levico Terme, Italy, October 2012*. Springer Proceedings in Mathematics and Statistics 79. Springer, 2014.
- [72] L. Makar-Limanov, *On groups of automorphisms of a class of surfaces*, Israel J. Mathem. 69 (1990), 250–256.
- [73] L. Makar-Limanov, *On the group of automorphisms of a surface  $xy = P(z)$* , Israel J. Mathem. 121 (2001), 113–123.
- [74] R. K. Markanda, J. Pascual, *Fixed rings of automorphisms of  $K[x, y]$* , Bull. Malaysian Math. Soc. (2) 4 (1981), 37–40.
- [75] H. Matsumura, F. Oort, *Representability of group functors, and automorphisms of algebraic schemes*, Invent. Math. 4 (1967), 1–25.
- [76] T. Matsusaka. *Polarized varieties, fields of moduli and generalized Kummer varieties of polarized Abelian varieties*, Amer. J. Math. 80 (1958), 45–82.
- [77] M. Miyanishi, *Open Algebraic Surfaces*, CRM Monogr. Ser., 12, Amer. Math. Soc., Providence, RI, 2001.
- [78] M. Miyanishi, *Recent developments in affine algebraic geometry: from the personal viewpoints of the author*, Affine algebraic geometry, 307–378, Osaka Univ. Press, Osaka, 2007.
- [79] M. Miyanishi, K. Masuda, *Affine pseudo-planes with torus actions*, Transform. Groups 11 (2006), 249–267.
- [80] M. Miyanishi, T. Sugie, *Affne surfaces containing cylinderlike open sets*, J. Math. Kyoto Univ. 20 (1980) 11–42.
- [81] G. D. Mostow, *Fully reducible subgroups of algebraic groups*, Amer. J. Math. 78 (1956), 200–221.
- [82] H. Neumann, *Generalized free products with amalgamated subgroups*. Amer. J. Math. 70 (1948), 590–625.
- [83] T. Petrie, *Algebraic automorphisms of smooth affine surfaces*. Invent. Math. 95 (1989), 355–378.
- [84] A. Perepechko, M. Zaidenberg, *Automorphisms of open surfaces and the Thompson group*, 2015 (a work in progress).
- [85] Yu. M. Polyakova, *A family of categories of log terminal pairs and automorphisms of surfaces*, Izv. RAN Ser. Mat. 74 (2010), 103–156.



- [86] V. L. Popov, *Quasihomogeneous affine algebraic varieties of the group  $SL(2)$* , Izv. Akad. Nauk SSSR Ser. Mat. 37 (1973), 792–832.
- [87] V. L. Popov, *Classification of affine algebraic surfaces that are quasihomogeneous with respect to an algebraic group*, Math. USSR Izv. 7 (1973), 1039–1055 (1975).
- [88] C. P. Ramanujam, *A note on automorphism groups of algebraic varieties*, Math. Ann. 156 (1964), 25–33.
- [89] R. Rentschler, *Opérations du groupe additif sur le plane affine*, C. R. Acad. Sci. 267 (1968), 384–387.
- [90] R. W. Richardson, Jr. *A rigidity theorem for subalgebras of Lie and associative algebras*, Illinois J. Math. 11 (1967), 92–110.
- [91] R. W. Richardson, Jr. *Conjugacy classes in Lie algebras and algebraic groups*, Annals of Mathem. 86 (1967), 1–15.
- [92] W. Rudin, *Preservation of level sets by automorphisms of  $\mathbb{C}^n$* , Indag. Math. (N.S.) 4 (1993), 489–497.
- [93] P. Samuel, *À propos du théorème des unités*, Bull. Sci. Math. (2) 90 (1966), 89–96.
- [94] Séminaire C. Chevalley, *Classification des Groupes de Lie algébriques* (two volumes), Paris, 1956–58.
- [95] J.-P. Serre, *Trees*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003.
- [96] I. R. Shafarevich, *On some infinite-dimensional groups*, Rend. Mat. e Appl. (5) 25 (1966), no. 1-2, 208–212.
- [97] I. R. Shafarevich, *On some infinite-dimensional groups. II*, Izv. Akad. Nauk SSSR Ser. Mat. 45 (1981), 214–226.
- [98] R. Steinberg, *Regular elements of semisimple algebraic groups*, Publ. IHES 25 (1965), 49–80.
- [99] M. Suzuki, *Sur les opérations holomorphes du groupe additif complexe sur l'espace de deux variables complexes*, Ann. Sci. École Norm. Sup. (4) 10 (1977), 517–546.
- [100] R. Tanimoto, *Algebraic torus actions on affine algebraic surfaces*. J. Algebra 285 (2005), 73–97.
- [101] D. Wright, *Abelian subgroups of  $\text{Aut}_k(k[X, Y])$  and applications to actions on the affine plane*, Illinois J. Math. 23 (1979), 579–634.
- [102] D. Wright, *Polynomial automorphism groups*, in: Polynomial automorphisms and related topics, 1–19, Publishing House for Science and Technology, Hanoi, 2007.
- [103] H. Yoshihara, *Projective plane curves and the automorphism groups of their complements*, J. Math. Soc. Japan 37 (1985), 87–113.

MATHEMATISCHES INSTITUT, ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, ALBERTSTRASSE  
19, 79104 FREIBURG, GERMANY

*E-mail address:* sergei.kovalenko@rub.de

DEPARTMENT OF MATHEMATICS AND MECHANICS, ST. PETERSBURG STATE UNIVERSITY,  
UNIVERSITETSKY PR. 28, STARY PETERHOF, 198504, RUSSIA

*E-mail address:* perepeal@gmail.com

UNIVERSITÉ GRENOBLE I, INSTITUT FOURIER, UMR 5582 CNRS-UJF, BP 74, 38402  
SAINT MARTIN D'HÈRES CÉDEX, FRANCE

*E-mail address:* Mikhail.Zaidenberg@ujf-grenoble.fr